

Crystal structure of (*Z*)-4-methylbenzyl 3-[1-(5-methylpyridin-2-yl)ethylidene]-dithiocarbazate

Thahira Begum S. A. Ravoof,^a Edward R. T. Tiekkink,^{b*}
Siti Aminah Omar,^a Sharifa Zaithun Begum^a and
Mohamed I. M. Tahir^a

^aDepartment of Chemistry, Universiti Putra Malaysia, 43400 Serdang, Malaysia, and

^bCentre for Crystalline Materials, Faculty of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia. *Correspondence e-mail: edwardt@sunway.edu.my

Received 2 December 2015; accepted 15 December 2015

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


In the title dithiocarbazate compound, $C_{17}H_{19}N_3S_2$, the central CN_2S_2 residue is essentially planar (r.m.s. deviation = 0.0288 Å) and forms dihedral angles of 9.77 (8) and 77.47 (7)° with the substituted-pyridyl and *p*-tolyl rings, respectively, indicating a highly twisted molecule; the dihedral angle between the rings is 85.56 (8)°. The configuration about the C≡N bond is *Z*, which allows for the formation of an intramolecular N—H···N(pyridyl) hydrogen bond. The packing features tolyl-methyl-C—H···N(imine), pyridyl-C—H···π(tolyl) and π—π interactions [between pyridyl rings with a distance = 3.7946 (13) Å], which generates jagged supramolecular layers that stack along the *b* axis with no directional interactions between them.

Keywords: crystal structure; hydrogen bonding; dithiocarbazate.

CCDC reference: 1442456

1. Related literature

For the structure of the 4-methylpyridin-2-yl derivative, with an *E* configuration for the C≡N bond, allowing for the formation of centrosymmetric {···HNCS}₂ synthons in the crystal, see: Omar *et al.* (2014). For the synthesis, see: Ravoof *et al.* (2010).



2. Experimental

2.1. Crystal data

$C_{17}H_{19}N_3S_2$	$V = 833.35 (3) \text{ \AA}^3$
$M_r = 329.47$	$Z = 2$
Monoclinic, P_c	$Cu K\alpha$ radiation
$a = 9.0073 (2) \text{ \AA}$	$\mu = 2.88 \text{ mm}^{-1}$
$b = 12.3856 (2) \text{ \AA}$	$T = 100 \text{ K}$
$c = 7.5553 (1) \text{ \AA}$	$0.31 \times 0.22 \times 0.18 \text{ mm}$
$\beta = 98.620 (2)^\circ$	

2.2. Data collection

Agilent Xcalibur, Eos, Gemini diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2011)
 $T_{\min} = 0.43$, $T_{\max} = 0.60$

16118 measured reflections
3195 independent reflections
3191 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.081$
 $S = 1.05$
3195 reflections
205 parameters
3 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$
Absolute structure: Flack x determined using 1558 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons *et al.*, 2013).
Absolute structure parameter:
−0.011 (13)

Table 1
Hydrogen-bond geometry (Å, °).

$Cg1$ is the centroid of the C3–C8 ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N1—H1N···N3	0.88 (2)	1.98 (3)	2.624 (3)	130 (3)
C12'—H12C···N2 ⁱ	0.98	2.58	3.483 (3)	154
C13—H13···Cg1 ⁱⁱ	0.95	2.76	3.582 (3)	145

Symmetry codes: (i) $x + 1, y, z$; (ii) $x + 1, -y + 2, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Acknowledgements

This research was funded by Universiti Putra Malaysia (UPM) under Research University Grant Schemes (RUGS No. 9419400), the Fundamental Research Grant Scheme (FRGS No. 5524425) and the Science Fund (Science Fund No: 06–01–04-SF810). SAO wishes to thank the UPM for the award of a Graduate Research Fellowship.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7554).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
Omar, S. A., Ravoof, T. B. S. A., Tahir, M. I. M. & Crouse, K. A. (2014).
Transition Met. Chem. **39**, 119–126.
Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst. B* **69**, 249–259.
Ravoof, T. B. S. A., Crouse, K. A., Tahir, M. I. M., How, F. N. F., Rosli, R. &
Watkins, D. J. (2010). *Transition Met. Chem.* **35**, 871–876.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, o1071–o1072 [doi:10.1107/S205698901502407X]

Crystal structure of (*Z*)-4-methylbenzyl 3-[1-(5-methylpyridin-2-yl)ethylidene]dithiocarbazate

Thahira Begum S. A. Ravoof, Edward R. T. Tiekkink, Siti Aminah Omar, Sharifa Zaithun Begum and Mohamed I. M. Tahir

S1. Refinement


S2. Experimental

The precursor molecule, *S*-4-methylbenzyl dithiocarbazate, was prepared by adapting the literature procedure of Ravoof *et al.* (2010). Thus, KOH (11.2 g, 0.2 mol) was dissolved in absolute ethanol (70 ml) and to this solution was added hydrazine hydrate (10 g, 0.2 mol) followed by cooling in an ice-salt bath. Drop wise addition of carbon disulfide (15.2 g, 0.2 mol) with constant stirring over 1 h followed. The two layers that subsequently formed were separated. The light-brown lower layer was dissolved in 40% ethanol (60 ml) below 268 K. The mixture was kept in an ice-bath and 4-methylbenzyl chloride (26.5 ml, 0.2 mol) was added drop wise with vigorous stirring. The major product, which was white and sticky was filtered and left overnight to dry over anhydrous silica gel in a desiccator. Recrystallization to yield analytically pure *S*-4-methylbenzyl dithiocarbazate was achieved from hot acetonitrile. Yield: 82%; *M.pt*: 160–161 °C.


S-4-Methylbenzyl dithiocarbazate (2.12 g, 0.01 mol) was dissolved in hot acetonitrile (100 ml) and added to an equimolar solution of 5-methyl-pyridine-2-aldehyde (1.21 g, 0.01 mol) in ethanol (25 ml). The mixture was then heated on a water bath until the volume has been reduced by half. A yellow precipitate formed after standing at room temperature for 1 h and this was washed with ethanol. Yellow prisms were deposited from its acetonitrile solution within a week. Yield: 78%. *M.pt*: 112–113 °C. Anal. Found (calc'd for C₁₇H₁₉N₃S₂): C, 62.32 (61.97); H, 5.59 (5.81); N 12.80 (12.75).

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H = 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$. The N—H atom was refined with N—H = 0.88±0.01 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.

**Figure 1**

The molecular structure of the title compound showing the atom-labelling scheme and displacement ellipsoids at the 70% probability level.

**Figure 2**

A view of the unit-cell contents in projection down the a axis. The tolyl-methyl-C—H···N(imine), pyridyl-C—H··· π (tolyl) and π — π interactions are shown as orange, pink and orange dashed lines, respectively.

(Z)-4-Methylbenzyl 3-[1-(5-methylpyridin-2-yl)ethylidene]dithiocarbazate

Crystal data

C₁₇H₁₉N₃S₂
 $M_r = 329.47$
Monoclinic, Pc
 $a = 9.0073 (2)$ Å
 $b = 12.3856 (2)$ Å
 $c = 7.5553 (1)$ Å
 $\beta = 98.620 (2)^\circ$
 $V = 833.35 (3)$ Å³
 $Z = 2$

$F(000) = 348$
 $D_x = 1.313 \text{ Mg m}^{-3}$
Cu $K\alpha$ radiation, $\lambda = 1.5418$ Å
Cell parameters from 12240 reflections
 $\theta = 4\text{--}71^\circ$
 $\mu = 2.88 \text{ mm}^{-1}$
 $T = 100$ K
Prism, yellow
 $0.31 \times 0.22 \times 0.18$ mm

Data collection

Agilent Xcalibur, Eos, Gemini
diffractometer
Radiation source: Enhance (Cu) X-ray Source
Graphite monochromator
Detector resolution: 16.1952 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(CrysAlis PRO; Agilent, 2011)
 $T_{\min} = 0.43$, $T_{\max} = 0.60$

16118 measured reflections
3195 independent reflections
3191 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 $\theta_{\max} = 71.5^\circ$, $\theta_{\min} = 3.6^\circ$
 $h = -11 \rightarrow 11$
 $k = -15 \rightarrow 15$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.081$
 $S = 1.05$
3195 reflections
205 parameters
3 restraints
Hydrogen site location: mixed
H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0663P)^2 + 0.1105P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$
Absolute structure: Flack x determined using
1558 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons *et al.*, 2013).
Absolute structure parameter: -0.011 (13)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.01004 (7)	0.74703 (4)	0.22309 (8)	0.01770 (17)
S2	0.20197 (7)	0.59121 (5)	0.07598 (9)	0.02369 (18)
N1	0.2382 (2)	0.80053 (17)	0.1071 (3)	0.0165 (4)
H1N	0.324 (2)	0.788 (3)	0.069 (4)	0.020*
N2	0.1881 (2)	0.90223 (17)	0.1402 (3)	0.0151 (4)
N3	0.4835 (2)	0.88995 (18)	0.0262 (3)	0.0171 (4)
C1	0.1524 (3)	0.7141 (2)	0.1305 (3)	0.0164 (5)
C2	-0.0843 (3)	0.6138 (2)	0.2616 (4)	0.0227 (6)

H2A	-0.1317	0.5817	0.1469	0.027*
H2B	-0.0025	0.5655	0.3160	0.027*
C3	-0.1993 (3)	0.6274 (2)	0.3865 (4)	0.0187 (5)
C4	-0.1541 (3)	0.6423 (2)	0.5698 (4)	0.0211 (5)
H4	-0.0502	0.6481	0.6153	0.025*
C5	-0.2590 (3)	0.6487 (2)	0.6864 (4)	0.0232 (5)
H5	-0.2259	0.6587	0.8107	0.028*
C6	-0.4129 (3)	0.6409 (2)	0.6239 (4)	0.0224 (6)
C6'	-0.5270 (4)	0.6430 (2)	0.7516 (4)	0.0309 (7)
H6'1	-0.5477	0.7180	0.7812	0.046*
H6'2	-0.4872	0.6038	0.8612	0.046*
H6'3	-0.6200	0.6085	0.6950	0.046*
C7	-0.4574 (3)	0.6281 (2)	0.4406 (4)	0.0239 (6)
H7	-0.5613	0.6238	0.3946	0.029*
C8	-0.3525 (3)	0.6216 (2)	0.3238 (4)	0.0221 (5)
H8	-0.3857	0.6130	0.1992	0.027*
C9'	0.1975 (3)	1.0925 (2)	0.1434 (4)	0.0174 (5)
H9'1	0.1047	1.0801	0.1937	0.026*
H9'2	0.1743	1.1322	0.0305	0.026*
H9'3	0.2675	1.1348	0.2282	0.026*
C9	0.2679 (3)	0.9857 (2)	0.1097 (3)	0.0149 (5)
C10	0.4164 (3)	0.9860 (2)	0.0464 (3)	0.0148 (5)
C11	0.6190 (3)	0.8899 (2)	-0.0256 (3)	0.0174 (5)
H11	0.6657	0.8221	-0.0378	0.021*
C12'	0.8495 (3)	0.9734 (2)	-0.1194 (3)	0.0198 (5)
H12A	0.8776	1.0426	-0.1679	0.030*
H12B	0.8470	0.9174	-0.2115	0.030*
H12C	0.9233	0.9538	-0.0156	0.030*
C12	0.6969 (3)	0.9832 (2)	-0.0632 (3)	0.0164 (5)
C13	0.6260 (3)	1.0811 (2)	-0.0445 (3)	0.0174 (5)
H13	0.6734	1.1468	-0.0692	0.021*
C14	0.4852 (3)	1.0830 (2)	0.0107 (3)	0.0166 (5)
H14	0.4363	1.1498	0.0239	0.020*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0175 (3)	0.0118 (3)	0.0255 (3)	-0.0019 (2)	0.0087 (2)	-0.0006 (2)
S2	0.0239 (3)	0.0140 (3)	0.0359 (4)	-0.0007 (3)	0.0131 (3)	-0.0039 (3)
N1	0.0152 (10)	0.0128 (10)	0.0220 (10)	-0.0006 (7)	0.0043 (8)	0.0006 (8)
N2	0.0138 (10)	0.0142 (11)	0.0165 (9)	-0.0007 (7)	0.0001 (8)	0.0002 (7)
N3	0.0147 (9)	0.0160 (10)	0.0198 (10)	-0.0023 (8)	0.0008 (8)	-0.0003 (8)
C1	0.0144 (11)	0.0196 (13)	0.0154 (11)	0.0002 (10)	0.0029 (9)	0.0003 (10)
C2	0.0276 (14)	0.0112 (12)	0.0326 (15)	-0.0066 (10)	0.0151 (11)	-0.0013 (10)
C3	0.0215 (12)	0.0098 (11)	0.0266 (13)	-0.0010 (10)	0.0096 (10)	0.0011 (9)
C4	0.0215 (12)	0.0131 (12)	0.0285 (13)	0.0016 (10)	0.0031 (10)	0.0021 (10)
C5	0.0324 (14)	0.0160 (13)	0.0219 (12)	0.0013 (10)	0.0061 (10)	0.0016 (10)
C6	0.0276 (14)	0.0090 (12)	0.0342 (14)	-0.0017 (10)	0.0162 (11)	0.0007 (10)

C6'	0.0382 (16)	0.0168 (13)	0.0434 (17)	-0.0032 (12)	0.0246 (13)	-0.0037 (12)
C7	0.0196 (13)	0.0149 (13)	0.0376 (15)	-0.0032 (10)	0.0057 (11)	-0.0035 (11)
C8	0.0252 (13)	0.0157 (13)	0.0256 (13)	-0.0042 (10)	0.0043 (10)	-0.0033 (10)
C9'	0.0166 (12)	0.0156 (13)	0.0196 (11)	0.0005 (8)	0.0012 (9)	-0.0007 (9)
C9	0.0147 (11)	0.0175 (12)	0.0115 (10)	-0.0010 (9)	-0.0019 (8)	0.0000 (8)
C10	0.0150 (11)	0.0167 (12)	0.0115 (10)	-0.0015 (9)	-0.0021 (8)	-0.0002 (9)
C11	0.0148 (11)	0.0171 (12)	0.0200 (12)	-0.0009 (9)	0.0018 (9)	-0.0001 (9)
C12'	0.0145 (11)	0.0229 (13)	0.0218 (12)	-0.0027 (9)	0.0023 (10)	0.0025 (10)
C12	0.0138 (12)	0.0217 (13)	0.0125 (10)	-0.0031 (9)	-0.0022 (9)	0.0002 (9)
C13	0.0171 (11)	0.0176 (12)	0.0166 (11)	-0.0040 (9)	-0.0005 (9)	0.0022 (9)
C14	0.0172 (11)	0.0155 (12)	0.0160 (12)	0.0002 (9)	-0.0008 (9)	-0.0001 (9)

Geometric parameters (Å, °)

S1—C1	1.762 (3)	C6'—H6'2	0.9800
S1—C2	1.820 (3)	C6'—H6'3	0.9800
S2—C1	1.655 (3)	C7—C8	1.389 (4)
N1—C1	1.348 (3)	C7—H7	0.9500
N1—N2	1.374 (3)	C8—H8	0.9500
N1—H1N	0.872 (14)	C9'—C9	1.505 (3)
N2—C9	1.299 (3)	C9'—H9'1	0.9800
N3—C11	1.336 (3)	C9'—H9'2	0.9800
N3—C10	1.353 (3)	C9'—H9'3	0.9800
C2—C3	1.512 (4)	C9—C10	1.486 (3)
C2—H2A	0.9900	C10—C14	1.397 (4)
C2—H2B	0.9900	C11—C12	1.403 (4)
C3—C8	1.391 (4)	C11—H11	0.9500
C3—C4	1.396 (4)	C12'—C12	1.503 (3)
C4—C5	1.387 (4)	C12'—H12A	0.9800
C4—H4	0.9500	C12'—H12B	0.9800
C5—C6	1.399 (4)	C12'—H12C	0.9800
C5—H5	0.9500	C12—C13	1.388 (4)
C6—C7	1.392 (4)	C13—C14	1.394 (4)
C6—C6'	1.511 (3)	C13—H13	0.9500
C6'—H6'1	0.9800	C14—H14	0.9500
C1—S1—C2	101.58 (12)	C6—C7—H7	119.4
C1—N1—N2	119.6 (2)	C7—C8—C3	121.0 (3)
C1—N1—H1N	117 (3)	C7—C8—H8	119.5
N2—N1—H1N	123 (3)	C3—C8—H8	119.5
C9—N2—N1	119.5 (2)	C9—C9'—H9'1	109.5
C11—N3—C10	118.5 (2)	C9—C9'—H9'2	109.5
N1—C1—S2	121.07 (18)	H9'1—C9'—H9'2	109.5
N1—C1—S1	113.26 (19)	C9—C9'—H9'3	109.5
S2—C1—S1	125.66 (16)	H9'1—C9'—H9'3	109.5
C3—C2—S1	107.59 (18)	H9'2—C9'—H9'3	109.5
C3—C2—H2A	110.2	N2—C9—C10	127.3 (2)
S1—C2—H2A	110.2	N2—C9—C9'	114.3 (2)

C3—C2—H2B	110.2	C10—C9—C9'	118.4 (2)
S1—C2—H2B	110.2	N3—C10—C14	121.0 (2)
H2A—C2—H2B	108.5	N3—C10—C9	118.3 (2)
C8—C3—C4	118.2 (2)	C14—C10—C9	120.7 (2)
C8—C3—C2	121.2 (2)	N3—C11—C12	124.5 (2)
C4—C3—C2	120.6 (2)	N3—C11—H11	117.8
C5—C4—C3	120.8 (2)	C12—C11—H11	117.8
C5—C4—H4	119.6	C12—C12'—H12A	109.5
C3—C4—H4	119.6	C12—C12'—H12B	109.5
C4—C5—C6	121.1 (2)	H12A—C12'—H12B	109.5
C4—C5—H5	119.4	C12—C12'—H12C	109.5
C6—C5—H5	119.4	H12A—C12'—H12C	109.5
C7—C6—C5	117.8 (2)	H12B—C12'—H12C	109.5
C7—C6—C6'	121.0 (3)	C13—C12—C11	116.6 (2)
C5—C6—C6'	121.2 (3)	C13—C12—C12'	123.6 (2)
C6—C6'—H6'1	109.5	C11—C12—C12'	119.8 (2)
C6—C6'—H6'2	109.5	C12—C13—C14	119.9 (2)
H6'1—C6'—H6'2	109.5	C12—C13—H13	120.1
C6—C6'—H6'3	109.5	C14—C13—H13	120.1
H6'1—C6'—H6'3	109.5	C13—C14—C10	119.6 (2)
H6'2—C6'—H6'3	109.5	C13—C14—H14	120.2
C8—C7—C6	121.1 (2)	C10—C14—H14	120.2
C8—C7—H7	119.4		
C1—N1—N2—C9	-176.9 (2)	C2—C3—C8—C7	176.4 (3)
N2—N1—C1—S2	174.81 (17)	N1—N2—C9—C10	-2.4 (4)
N2—N1—C1—S1	-5.6 (3)	N1—N2—C9—C9'	177.39 (19)
C2—S1—C1—N1	-173.23 (18)	C11—N3—C10—C14	1.3 (3)
C2—S1—C1—S2	6.4 (2)	C11—N3—C10—C9	-178.1 (2)
C1—S1—C2—C3	164.66 (18)	N2—C9—C10—N3	-4.0 (4)
S1—C2—C3—C8	104.6 (2)	C9'—C9—C10—N3	176.2 (2)
S1—C2—C3—C4	-77.7 (3)	N2—C9—C10—C14	176.6 (2)
C8—C3—C4—C5	1.4 (4)	C9'—C9—C10—C14	-3.1 (3)
C2—C3—C4—C5	-176.4 (2)	C10—N3—C11—C12	-0.9 (4)
C3—C4—C5—C6	-0.2 (4)	N3—C11—C12—C13	-0.1 (4)
C4—C5—C6—C7	-1.0 (4)	N3—C11—C12—C12'	179.6 (2)
C4—C5—C6—C6'	177.3 (3)	C11—C12—C13—C14	0.5 (3)
C5—C6—C7—C8	1.1 (4)	C12'—C12—C13—C14	-179.1 (2)
C6'—C6—C7—C8	-177.3 (3)	C12—C13—C14—C10	-0.1 (3)
C6—C7—C8—C3	0.1 (4)	N3—C10—C14—C13	-0.9 (3)
C4—C3—C8—C7	-1.3 (4)	C9—C10—C14—C13	178.5 (2)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C3—C8 ring.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···N3	0.88 (2)	1.98 (3)	2.624 (3)	130 (3)

C12'—H12C···N2 ⁱ	0.98	2.58	3.483 (3)	154
C13—H13···Cg1 ⁱⁱ	0.95	2.76	3.582 (3)	145

Symmetry codes: (i) $x+1, y, z$; (ii) $x+1, -y+2, z-1/2$.