



# The Development of Drugs against *Acanthamoeba* Infections

AQ: au Ruqaiyyah Siddiqui,<sup>a</sup> Yousuf Aqeel,<sup>b</sup> Naveed Ahmed Khan<sup>a</sup>

AQ: aff Department of Biological Sciences, Faculty of Science and Technology, Sunway University, Malaysia<sup>a</sup>; Department of Molecular and Cell Biology, School of Medicine, Boston University, Boston, Massachusetts, USA<sup>b</sup>

For the past several decades, there has been little improvement in the morbidity and mortality associated with *Acanthamoeba* keratitis and *Acanthamoeba* encephalitis, respectively. The discovery of a plethora of antiacanthamoebic compounds has not yielded effective marketed chemotherapeutics. The rate of development of novel antiacanthamoebic chemotherapies of translational value and the lack of interest of the pharmaceutical industry in developing such chemotherapies have been disappointing. On the other hand, the market for contact lenses/contact lens disinfectants is a multi-billion-dollar industry and has been successful and profitable. A better understanding of drugs, their targets, and mechanisms of action will facilitate the development of more-effective chemotherapies. Here, we review the progress toward phenotypic drug discovery, emphasizing the shortcomings of useable therapies.

AQ:A Antimicrobial chemotherapy is the most widely used method of treating infections due to *Acanthamoeba*. Despite advances in antimicrobial chemotherapy, the morbidity and mortality associated with *Acanthamoeba* keratitis and *Acanthamoeba* encephalitis, respectively, have remained high. For example, the mortality rate of granulomatous amoebic encephalitis due to pathogenic *Acanthamoeba* spp. is more than 90%, even with treatment with various combinations of drugs such as amphotericin B, rifampin, trimethoprim-sulfamethoxazole, ketoconazole, fluconazole, sulfadiazine, miltefosine, albendazole, etc. (reviewed in references 1, 2, 3, 4, 5, 6, 7, and 8). Current treatment of *Acanthamoeba* keratitis involves chlorhexidine, in combination with diamidines and neomycin, and can last up to a year, and even then infection recurrence occurs in approximately 10% of cases (reviewed in references 1, 2, 3, 4, 5, 6, 7, and 8). In part, this is due to our incomplete understanding of the biology of the parasite and of the pathogenesis and pathophysiology of the disease, as well as to the lack of effective chemotherapeutic agents and/or the lack of clinical testing of the potential targets that have been shown to play an important role in the virulence of pathogenic *Acanthamoeba*. This is despite the fact that a plethora of drugs, some of which show potent antiacanthamoebic effects, have been tested; however, their translational value in the treatment of *Acanthamoeba* infections remains unclear (reviewed in reference 9). Many of the drugs tested target functional aspects of *Acanthamoeba*, as it is “easier to erase function of an organism than its structure” (9). However, there are disadvantages to this approach. Being eukaryotes, *Acanthamoeba* species share functional homologies with mammalian cells. Consequently, many of the available drugs cannot be prescribed at effective concentrations due to their unwanted side effects. This is particularly relevant for treatment of amoebal brain infection, in which drugs are given intravenously and are expected to cross the blood-brain barrier to access the central nervous system to target the intracerebral parasite. In this process, drugs penetrate many tissues and can affect their physiology before reaching the target site at an effective concentration. Hence, there is a need to develop a targeted therapeutic approach, i.e., to identify drugs that can affect *Acanthamoeba* viability without affecting the host cells. The purpose of this review is to classify the tested antiacanthamoebic agents into functional groups to identify drugs and/or chemotherapeutic approaches of potential value for further work.

The knowledge of the mode of action of the majority of drugs tested against *Acanthamoeba* is largely derived from studies conducted in bacterial, fungal, or protozoan pathogens. These are indicated here for information; however, future studies are needed to determine and/or confirm their mechanism of action against *Acanthamoeba*.

## MEMBRANE-ACTING AGENTS

Being the outermost surface of the cell, the outer plasma membrane and its constituents provide a logical target(s) as it is easier to access. With the actively growing infective trophozoite that undergoes binary fission, the properties and charge of the cell membrane and its biosynthesis and modulation offer chemotherapeutic opportunities. Repurposing drugs with known modes of action for the aforementioned targets and/or agents with growth-inhibitory effects has been a useful avenue, but this approach lacks specificity, produces host cell toxicity, and may not be appropriate for the dormant cyst stage and thus poses a challenge in wider application of such drugs against *Acanthamoeba* infections. For example, chlorhexidine is positively charged and ionic with the negatively charged plasma membrane of the parasite, resulting in structural and permeability changes, ionic leakage, and cytoplasmic disruptions causing cellular damage and cell death, although those effects have not curtailed the use of the compound. It exhibits potent amoebicidal properties as well as cysticidal properties at 200 µg per ml (0.02%), and it is used clinically against *Acanthamoeba* keratitis but is not a drug of choice for ocular and brain infection (10–21) (Table 1). Similarly, biguanide compounds (polyhexamethylene biguanide [polyhexadine or polyaminopropyl biguanide]) are known to interact with membrane phospholipids, affecting membrane fluidity and conformation and leading to ionic leakage and cell death at 200 µg per ml (0.02%), and are used clinically against *Acanthamoeba* keratitis but may exhibit side

Accepted manuscript posted online ●●●

Citation Siddiqui R, Aqeel Y, Khan NA. 2016. The development of drugs against *Acanthamoeba* infections. *Antimicrob Agents Chemother* 60:000–000. doi:10.1128/AAC.00686-16.

Address correspondence to Naveed Ahmed Khan, naveed5438@gmail.com.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

TABLE 1 List of antiacanthamoebic agents

Classification and agent no.	Drug(s)	Mode of action	Description and effects on <i>Acanthamoeba</i>
1	Membrane-acting agents Chlorhexidine	<ul style="list-style-type: none"> <li>● Chlorhexidine is positively charged and ionic with the negatively charged plasma membrane of the parasite, resulting in structural and permeability changes, and ionic leakage, cytoplasmic disruptions causing cellular damage and cell death.</li> </ul>	<ul style="list-style-type: none"> <li>● Amoebicidal and cysticidal concn is 200 µg per ml (0.02%).</li> </ul>
2	Polyhexamethylenebiguanide (PHMB) (polyhexadine or polyaminopropylbiguanide; alexidine)	<ul style="list-style-type: none"> <li>● PHMB interacts with membrane phospholipids, affecting the membrane fluidity and conformation that leads to ionic leakage and cell death.</li> </ul>	<ul style="list-style-type: none"> <li>● PHMB exhibits amoebicidal and cysticidal properties at 200 µg per ml (0.02%), while alexidine shows antiacanthamoebic properties at 200 µg per ml but is less toxic <i>in vivo</i>.</li> <li>● Limited antiamoebic effects are observed.</li> </ul>
3	Antibacterials (polymyxin B; ceftazolin; meropenem)	<ul style="list-style-type: none"> <li>● Several antibacterials have been tested, including polymyxin B (binds to negatively charged membranes and disrupts the membrane integrity), ceftazolin (binds to penicillin binding proteins present in the cell wall, thus inhibiting cell wall synthesis), and meropenem (inhibits penicillin binding proteins).</li> </ul>	
4	Azole compounds (miconazole nitrate; ketoconazole; clotrimazole; itraconazole; fluconazole; voriconazole)	<ul style="list-style-type: none"> <li>● Azole compounds target 14-α demethylase, a cytochrome P-450 enzyme necessary to convert lanosterol to ergosterol, inhibiting phospholipid and triglyceride synthesis and affecting oxidative and peroxidative enzyme activities that cause deterioration subcellular components, leading to cell necrosis.</li> <li>● In addition to ergosterol biosynthesis, clotrimazole is known to inhibit endogenous respiration by impairing triglyceride and phospholipid biosynthesis as well as to inhibit cellular calcium homeostasis and calcium ATPases.</li> </ul> <p>Itraconazole inhibits ergosterol biosynthesis and has been shown to inhibit endogenous respiration, interact with membrane phospholipids, inhibit purine uptake, and impair triglyceride and/or phospholipid biosynthesis.</p> <p>Fluconazole inhibits ergosterol biosynthesis and is known to inhibit endogenous respiration, interact with membrane phospholipids, inhibit purine uptake, and impair triglyceride and/or phospholipid biosynthesis.</p>	<ul style="list-style-type: none"> <li>● The majority of azole compounds can show amoebicidal effects at as low as 10 µg per ml <i>in vitro</i>, but cysticidal properties and host cell toxicity are observed at a far higher concn of ~1,000 µg per ml.</li> <li>● For example, 200 mg ketoconazole twice daily is prescribed in <i>Acanthamoeba</i> keratitis cases, and the recommended concn in GAE<sup>2</sup> is 5 mg per kg, body wt.</li> <li>● In keratitis cases, clotrimazole is recommended at 10 mg per ml.</li> </ul>
5	Amphotericin B/natamycin	<ul style="list-style-type: none"> <li>● Amphotericin B, a polyene, binds irreversibly to ergosterol, resulting in disruption of membrane integrity and ultimately in cell death.</li> <li>● Natamycin binds to ergosterol in the plasma membrane, preventing ergosterol-dependent fusion of vacuoles, as well as membrane fusion and fission.</li> </ul>	<ul style="list-style-type: none"> <li>● Amphotericin B has been shown to possess antiacanthamoebic properties at 100 µg per ml, while natamycin exhibits effects at far higher concn (10–50 mg per ml).</li> </ul>

6	Caspofungin	<ul style="list-style-type: none"> <li>Caspofungin inhibits beta-(1,3)-glucan synthase, inhibiting the synthesis of beta-(1,3)-D-glucan.</li> <li>Mannose-linked or anti-MBP antibody-linked photoactivated agents can selectively bind to <i>Acanthamoeba</i>, and exposure to light of the appropriate wavelength produces reactive oxygen species, targeting the parasite.</li> </ul>	<ul style="list-style-type: none"> <li>Caspofungin shows amoebicidal effects at 250 µg per ml, and cysticidal properties are observed at 500 µg per ml.</li> <li>Amoebicidal properties were observed at &lt;50 µM, while excystation was abolished at this concn.</li> </ul>
7	Mannose-linked or anti-MBP antibody-linked photoactivated agents		
8	Intracellular targeting agents	<p>Quaternary ammonium compounds and their derivatives (cetyltrimethylammonium bromide; cetylpyridiniumbromide) (insertion of the ethyl phosphate group into the molecule of cetyltrimethylammonium bromide leads to the formation of alkylphosphocholines [hexadecylphosphocholine/miltefosine]).</p> <p>Quaternary ammonium compounds (benzethonium chloride)</p> <p>Calcium modulating agents (amlodipine; loperamide; amiodarone; trifluoperazine dihydrochloride; chlorpromazine dihydrochloride).</p>	<ul style="list-style-type: none"> <li>These quaternary ammonium compounds exhibit antiacanthamoebic activity (at &lt;20 µM) compared with alkylphosphocholines compounds, which show activity at &gt;60 µM.</li> <li>These quaternary ammonium compounds show antiacanthamoebic effects at &gt;500 µM.</li> <li>These agents showed antiacanthamoebic effects at &gt;500 µg per ml <i>in vitro</i>.</li> <li>Chlorpromazine and trifluoperazine showed amoebicidal and cysticidal effects at concn in the micromolar range.</li> </ul>
9			
10			
11	Nucleic acid-acting drugs	<p>Proflavine hemisulfate</p> <p>Hydroxystilbamide isethionate</p>	<ul style="list-style-type: none"> <li>Proflavine hemisulfate is reported to exhibit amoebicidal effects at 100 µg per ml and cysticidal effects at 1,000 µg per ml.</li> <li>Hydroxystilbamide isethionate exhibits amoebicidal properties at 100 µg per ml and cysticidal properties at 1,000 µg per ml.</li> </ul>
12			
13	Trimethoprim		<ul style="list-style-type: none"> <li>Trimethoprim has been shown to exhibit amoebicidal effects at 100 µg per ml.</li> </ul>
14	5-Fluorocytosine and moxifloxacin		<ul style="list-style-type: none"> <li>Limited anti-amoebic effects are observed.</li> </ul>

(Continued on following page)

TABLE 1 (Continued)

Classification and agent no.	Drug(s)	Mode of action	Description and effects on <i>Acanthamoeba</i>
15	Pyrimethamine-sulformethoxine combination and trimethoprim-sulfamethoxazole	<ul style="list-style-type: none"> <li>● Moxifloxacin is an inhibitor of DNA gyrase, a type II topoisomerase, and topoisomerase IV, required for DNA replication.</li> <li>● Pyrimethamine is dihydrofolate reductase inhibitor to block biosynthesis of purines and pyrimidines), sulformethoxine targets dihydropteroate synthase and dihydrofolate reductase and competes with para-aminobenzoic acid for incorporation into folic acid.</li> </ul>	<ul style="list-style-type: none"> <li>● Both combinations showed amoebicidal effects at 100 µg per ml.</li> </ul>
16	Pentamidine isethionate/propamidine isethionate	<ul style="list-style-type: none"> <li>● Pentamidine isethionate and propamidine isethionate inhibit synthesis of DNA, RNA, phospholipids, and proteins.</li> </ul>	<ul style="list-style-type: none"> <li>● Pentamidine isethionate and propamidine isethionate show amoebicidal and cysticidal properties at ~100–200 µg per ml, while propamidine isethionate is used clinically against keratitis at a concn of 1 mg per ml.</li> <li>● Diminazene aceturate has been shown to exhibit amoebicidal and cysticidal properties at ~100–200 µg per ml.</li> </ul>
17	Diminazene aceturate	<ul style="list-style-type: none"> <li>● Diminazene aceturate binds to the groove between the complementary strands of DNA at regular intervals and thus distorts the helical structure. It is also known to affect synthesis of phospholipids and also interferes with the glycolytic pathway.</li> </ul>	<ul style="list-style-type: none"> <li>● Diminazene aceturate has been shown to exhibit amoebicidal and cysticidal properties at ~100–200 µg per ml.</li> </ul>
18	Clinically useful protein synthesis inhibitors (including paromomycin sulfate, tobramycin, and neomycin sulfate)	<ul style="list-style-type: none"> <li>● Paromomycin sulfate inhibits initiation and elongation steps of protein synthesis.</li> <li>● Tobramycin inhibits protein synthesis by binding to ribosomes and preventing mRNA translation, leading to cell death, while neomycin sulfate binds to four nucleotides of 16S rRNA and a single amino acid of protein S12 and interferes with the initiation complex, leading to misreading of mRNA such that incorrect amino acids are inserted into the polypeptide, leading to nonfunctional or toxic peptides and the breakup of polysomes into nonfunctional monosomes.</li> </ul>	<ul style="list-style-type: none"> <li>● Paromomycin sulfate has been shown to exhibit amoebicidal and cysticidal properties at &gt;100 µg per ml, while the other agents exhibit anti-amoebic effects at &gt;250 µg per ml, but cysticidal properties are observed at &gt;500 µg per ml.</li> </ul>
19	Prednisolone, beta-methasone phosphate, linezolid, co-trimoxazole.	<ul style="list-style-type: none"> <li>● Prednisolone irreversibly binds with glucocorticoid receptors, inhibiting gene transcription for COX-2, cytokines, cell adhesion molecules, and inducible NO synthase, beta-methasone phosphate binds to plasma transcortin and becomes active when it is not bound to transcortin, linezolid inhibits the formation of subunits of ribosome, and co-trimoxazole is known to inhibit folic acid synthesis.</li> </ul>	<ul style="list-style-type: none"> <li>● These protein synthesis inhibitors show amoebicidal properties but have limited cysticidal effects and are of limited value as antiacanthamoebic agents.</li> </ul>

<sup>a</sup> GAE, granulomatous amebic encephalitis.

effects and are not ideal for the treatment of ocular or brain infections, albeit they can be used in combination with chlorhexidine (10–13, 22–36). More recently, alexidine, an amphipathic bis-guanide, has shown amoebicidal activity at 10 µg per ml and cysticidal activity at 100 µg per ml (37). The cytotoxic activities of alexidine are comparable to those of chlorhexidine; however, alexidine appeared less toxic *in vivo* (37). Several antibacterials have been tested in *Acanthamoeba* infection, including polymyxin B (binds to negatively charged membranes and disrupts membrane integrity) (18, 38–40), cefazolin (18), and meropenem (inhibits penicillin binding proteins) (41), but limited anti-amoebic effects were observed at physiologically tolerable concentrations.

Despite similarities to the host's cell plasma membrane, ergosterol, mannose-binding protein (MBP), and laminin-binding protein have been identified as useful components for the targeted killing of amoebae. As the presence of ergosterol is limited to fungi and protists, while human cells contain cholesterol, this is considered a useful target, but anti-ergosterol biosynthesis strategies have shown limited value against *Acanthamoeba* infections. For example, several azole compounds, including miconazole nitrate, ketoconazole, clotrimazole, fluconazole, and voriconazole, have been tested that target 14- $\alpha$  demethylase, a cytochrome P-450 enzyme necessary to convert lanosterol to ergosterol, inhibiting phospholipid and triglyceride synthesis and affecting oxidative and peroxidative enzyme activities, resulting in deterioration of subcellular components and leading to cell necrosis. Although several azole compounds showed amoebicidal effects at a concentration as low as 10 µg per ml *in vitro*, they showed cysticidal properties at a far higher concentration of 1,000 µg per ml (19, 20, 27, 31, 33, 40, 42–61). Among the ergosterol inhibitors, amphotericin B has been tested and has been shown to bind to ergosterol, forming a transmembrane channel that leads to monovalent ion leakage, which is the primary effect leading to cell death. The effective concentration against *Acanthamoeba* is reported to be 100 µg per ml (39, 40, 55, 59–63). Additionally, natamycin, which targets ergosterol in the plasma membrane, preventing ergosterol-dependent fusion of vacuoles as well as membrane fusion and fission, has shown limited amoebicidal properties at physiologically relevant concentrations (14, 27, 43, 64). In part, this is due to the fact that the present compounds targeting ergosterol or its synthetic pathway lack specificity and produce inconsistent results against various strains/species of *Acanthamoeba* and may also target the host cell P450 enzymes, resulting in side effects (60–65). Additionally, the effects of these compounds are often amoebistatic, rather than amoebicidal (65); hence, prolonged clinical application is needed, which could result in the emergence of drug-resistant strains, as seen in yeast, where azole resistance emerging through increased function of efflux mechanisms or through changes in the azole targets, e.g., C14 demethylase, or through changes in the biosynthetic steps of ergosterol synthesis has been observed. Similar mechanisms may explain variations in antimicrobial sensitivity among various isolates of *Acanthamoeba* (66). Overall, the ergosterol biosynthesis pathway is a potential target in the rational development of targeted therapeutic interventions against *Acanthamoeba*, as long as specificity is achieved to optimize the antiparasitic effects.

Within the plasma membrane, mannose-binding protein (MBP) has been identified as a key adhesin in *Acanthamoeba*-mediated host cell binding and cytotoxicity. The MBP protein consists of a signal peptide (amino acids 1 to 21), an extracellular

cysteine (C)-rich region covering amino acid positions 274 to 615, and a single predicted transmembrane region (amino acids 733 to 755). Expression of MBP is linked with the pathogenicity of *Acanthamoeba* and is associated with binding to and cytotoxicity of host cells. Notably, immunization with recombinant MBP (rMBP) protects animals against subsequent challenge with pathogenic *Acanthamoeba* species (5). Although rMBP is not applicable as a vaccine tool given the rarity of the disease, these findings have highlighted MBP as an important chemotherapeutic target. For example, recent studies showed that MBP-targeted chemotherapy can effectively eradicate amoebae *in vitro* and protect host cells against amoeba-mediated damage (66). Although studies are needed to prove the value of this approach *in vivo*, it has been suggested that mannose-conjugated porphyrin may have an application for targeted photodynamic chemotherapy against *Acanthamoeba* infections and should be explored for potential clinical applications in future investigations (66). Other drugs targeting the membrane include caspofungin, which is known to inhibit the synthesis of beta-(1,3)-D-glucan (59, 67). Caspofungin shows amoebicidal properties *in vitro* at 250 µg per ml and is cysticidal at 500 µg per ml (59, 67) and is thus of limited value in clinical applications.

#### INTRACELLULAR TARGETING AGENTS

Calcium channels play a critical role in the viability of *Acanthamoeba*. For example, the viability of trophozoites depends on their amoeboid movement in search and uptake of food particles, encystation or excystation, and asexual reproduction. These processes involve myosin contractility, activation of actin filament, inhibition of actin cross-linking by alpha-actinin, or binding to calmodulin. Other low-molecular-weight calcium-binding proteins and calpain, actophorin, actobindin, calcium-sensitive actin gelation protein, actin bundling protein (AhABP), and calcium-dependent extracellular proteases play important roles in its physiology. Thus, drugs affecting these functions would have deleterious effects on the viability of *Acanthamoeba*. Notably, calcium antagonists such as amlodipine (inhibits the transmembrane influx of calcium ions), loperamide (inhibits calcium channel activity and calmodulin binding), amiodarone (calcium blocker-like activity), and trifluoperazine dihydrochloride and chlorpromazine dihydrochloride (inhibit calmodulin) exhibit amoebicidal effects *in vitro* (68). Although the majority of drugs are used clinically, they exhibit anti-amoebic effects at a relatively high concentration of 500 µg per ml. However, two neuroleptic agents, chlorpromazine and trifluoperazine, show amoebicidal and cysticidal effects in the micromolar range of concentrations *in vitro* (69). Furthermore, the combination of chlorpromazine with rokitamycin or amphotericin B enhances protection of host cells against the parasite (69), suggesting the need for future studies to test the clinical relevance of these drugs against *Acanthamoeba* infections in experimental models as well as in patients.

Quaternary ammonium compounds, including cetyltrimethylammonium bromide and cetylpyridinium bromide, have been tested for anti-*Acanthamoeba* properties (54). Their effects are known to result in inactivation of energy-producing enzymes, denaturation of essential cell proteins, and disruption of the cell membrane. Insertion of the ethyl phosphate group into the molecule of cetyltrimethylammonium bromide leads to the formation of alkylphosphocholines (hexadecylphosphocholine/miltefosine). *In vitro* studies showed that quaternary ammonium compounds ex-

AQ: C

hibit higher antiacanthamoebic properties (at concentrations of  $<20 \mu\text{M}$ ) than hexadecylphosphocholine/miltefosine ( $>60 \mu\text{M}$ ) (54) and are promising agents. Another quaternary ammonium compound, benzethonium chloride (mode of action, induction of apoptosis) shows amoebicidal effects at  $>500 \mu\text{g per ml}$  (54) and is of limited utility. More recently, prochlorperazine (a known antagonist of dopamine [D<sub>2</sub>] receptor, muscarinic receptor, and histamine antagonist) and corticosteroids were shown to exhibit amoebicidal effects at  $250 \mu\text{g per ml}$  *in vitro*, but determinations of their usefulness require further studies (47, 64, 68, 70).

### NUCLEIC ACID-ACTING DRUGS

Nucleic acid inhibitors inhibit DNA/RNA synthesis, prevent DNA from functioning as a template, affect the function of polymerases involved in the replication and transcription of DNA, or intercalate into the DNA. The antibacterial properties of nucleic acid inhibitors are well known, making this pathway a useful target, but the lack of specificity against eukaryotic *Acanthamoeba*, together with the high concentrations required to target cysts and the observation that the majority of nucleic acid inhibitors are toxic or carcinogenic, suggests that, with the exception of few compounds, nucleic acid inhibitors are clinically inappropriate or not ideal candidates as antiacanthamoebic compounds. For example, proflavine hemisulfate exhibits mutagenic effects on DNA by intercalating between nucleic acid base pairs and causes base pair deletions and insertions. It has been reported to exhibit amoebicidal effects at  $100 \mu\text{g per ml}$  and cysticidal effects at  $1,000 \mu\text{g per ml}$  (40). The mode of action of hydroxystilbamidine isethionate involves binding extensively and selectively to kinetoplastic DNA, inhibiting cell division and reproduction. It has also been shown to bind to RNA and is a significant inhibitor of ribonucleases, and it is taken up in lysosomes, leading to a significant increase in the number of lysosome-like bodies and secretion granules. It exhibits amoebicidal properties at  $100 \mu\text{g per ml}$  and cysticidal properties at  $1,000 \mu\text{g per ml}$  (40). Other compounds tested include trimethoprim, which binds to dihydrofolate reductase and inhibits the reduction of dihydrofolic acid to tetrahydrofolic acid, which is important in the thymidine synthesis pathway for DNA synthesis. It has been shown to exhibit amoebicidal effects at  $100 \mu\text{g per ml}$  (39), while 5-fluorocytosine (mode of action, competitive inhibition of purine and pyrimidine uptake) and moxifloxacin (inhibitor of DNA gyrase, a type II topoisomerase, and topoisomerase IV, required for DNA replication), have shown limited value in the treatment of granulomatous amoebic encephalitis (41, 63). Given the nonselective nature of these compounds and their associated toxicity, several studies have tested combinations of nucleic acid synthesis inhibitors against *Acanthamoeba*. When pyrimethamine (a dihydrofolate reductase inhibitor blocking biosynthesis of purines and pyrimidines) and sulfamethoxazole (targeting dihydropteroate synthase and dihydrofolate reductase and competing with para-aminobenzoic acid for incorporation into folic acid) were tested in combination, amoebicidal properties were observed at  $100 \mu\text{g per ml}$  (39). Similarly, trimethoprim plus pyrimethamine and trimethoprim plus sulfamethoxazole (inhibitor of folic acid synthesis) showed amoebicidal effects at  $100 \mu\text{g per ml}$  (39, 50, 51, 71, 72).

Among the effective compounds tested, pentamidine isethionate inhibited synthesis of DNA, RNA, phospholipids, and proteins, with amoebicidal and cysticidal properties seen at  $\sim 100$  to

$200 \mu\text{g per ml}$  with variable results (11, 19, 73), while propamidine isethionate (DNA synthesis inhibitor) is used clinically against keratitis at a concentration of up to  $1 \text{ mg per ml}$  (13, 16, 17, 23, 27, 32, 33, 35, 36, 38, 42, 46, 53, 70, 74–79). Other drugs tested included diminazene aceturate, which binds to the groove between the complementary strands of DNA at regular intervals and thus distorts the helical structure. It is also known to affect phospholipids synthesis and also interferes with the glycolytic pathway of the parasite. It has been shown to exhibit amoebicidal and cysticidal properties at  $\sim 100$  to  $200 \mu\text{g per ml}$  (11). For treatment of brain infections, rifampin is promising as an additive drug, as it is lipophilic, a property that makes it a good candidate for treatment of infections of the central nervous system, which requires distribution to the central nervous system by penetration through the blood-brain barrier. The mode of action is inhibition of DNA-dependent RNA polymerase by binding to its beta-subunit, thus preventing transcription of RNA and subsequent translation to proteins. It has been shown to exhibit amoebicidal properties but is of limited value in treatments (50, 51, 64, 71).

### PROTEIN SYNTHESIS-INHIBITING DRUGS

Inhibition of protein synthesis has been one of the key targets for many of the available antibiotics, mostly taking advantage of differences in prokaryotic and eukaryotic ribosome structures and functions. The majority of protein synthesis inhibitors block mRNA translation into proteins, e.g., initiation, elongation (including aminoacyl tRNA entry, proofreading, peptidyl transfer, and ribosomal translocation), and termination. As for other pathways and structures of eukaryotes, in homology with host mammalian cells, selective targeting of protein synthesis remains a challenge and the use of protein synthesis inhibitors is often associated with host cell toxicity. Given that amoebae are actively growing in their infective states, such compounds can be used to block reproduction with tolerable toxicities. For example, paromomycin sulfate (inhibitor of the initiation and elongation steps of protein synthesis) has been shown to exhibit amoebistatic, amoebicidal, and cysticidal properties at more than  $100 \mu\text{g per ml}$  (33, 40, 54). Tobramycin (inhibitor of protein synthesis by binding to ribosomes and preventing mRNA translation, leading to cell death) has shown amoebicidal properties at more than  $250 \mu\text{g per ml}$  (25, 80).

Similarly, neomycin sulfate, which binds to four nucleotides of 16S rRNA and a single amino acid of protein S12 and interferes with the initiation complex, leading to misreading of mRNA such that incorrect amino acids are inserted into the polypeptide, resulting in nonfunctional or toxic peptides and the breakup of polysomes into nonfunctional monosomes, has been shown to exhibit antiamoebic effects at  $250 \mu\text{g per ml}$ , but cysticidal properties are observed at  $>500 \mu\text{g per ml}$  (30, 32, 35, 38, 40, 70, 74). Among the drug combinations tested, neomycin plus polymyxin B (33, 42, 45, 46, 52) and neomycin sulfate plus polymyxin B sulfate plus gramicidin (cation detergent) exhibited amoebicidal properties (42, 45, 53). In contrast, the combination of neomycin plus polymyxin B plus bacitracin exhibited amoebistatic and amoebicidal as well as cysticidal properties (18, 40, 51, 65, 76).

Several other protein synthesis inhibitors tested show amoebicidal properties but have limited cysticidal effects. These include prednisolone (irreversibly binds with glucocorticoid receptors, inhibiting gene transcription for cytochrome oxidase 2 [COX-2], cytokines, cell adhesion molecules, and inducible NO synthase)

(52), beta-methasone phosphate (binds to plasma transcortin and becomes active when it is not bound to transcortin) (81), and linezolid (inhibits the formation of subunits of ribosome) (39, 41, 50, 51, 71, 72).

### ENZYME-ACTING AGENTS

As described above, quaternary ammonium compounds (such as cetyltrimethylammonium bromide and cetylpyridinium bromide) and alkylphosphocholines (such as miltefosine) are promising candidates against *Acanthamoeba* infections. The mode of action of miltefosine is induction of apoptosis-like cell death by acting as an inhibitor of proteinase kinase B. It has been shown to exhibit amoebicidal properties (61, 82–84). Notably, miltefosine (at 65.12 µg per ml) was tested in combination with polyhexamethylene biguanide, chlorhexidine, and propamide isethionate in a rat model for the topical treatment of *Acanthamoeba* keratitis (83). The results revealed that the miltefosine-polyhexamethylene biguanide combination gave the best treatment results, and approximately 86% of the eyes were cleared of amoebae. It is also recommended as part of the treatment regimen against human brain infection due to *Acanthamoeba* (85). Future studies of the combination and effective delivery of quaternary ammonium compounds and their derivatives to the target site will determine the clinical usefulness. Other drugs tested showed amoebicidal effects but limited cysticidal effects. These include sulfadiazine (inhibitor of dihydropteroate synthetase) (86, 87), flurbiprofen (nonselective cytochrome oxidase [COX] inhibitor of pathway responsible for the conversion of arachidonic acid into prostaglandin G2 into prostaglandin H2) (77), riboflavin (targeting riboflavin hydrogenase, riboflavin kinase, and riboflavin synthase) (88), diclofenac (inhibitor of prostaglandin synthesis by inhibiting COX) (13), albendazole (targeting the colchicine-sensitive site of tubulin, inhibiting its polymerization into microtubules and leading to impaired uptake of glucose and depletion of glycogen stores) (89), and digoxin, which binds to the sodium/potassium-transporting ATPase alpha-1 chain (68).

Given the rarity of the disease and availability of a number of compounds with various effects against *Acanthamoeba* trophozoites and cysts, there is a need to test various combinations to prove their clinical usefulness with tolerable toxicities and acceptable pharmacokinetics profiles, safety margins, etc. In the absence of targeted therapy, this would provide the logical avenues for further research in clinical practice that may provide strategies for chemotherapy against this difficult-to-treat infection.

### THE WAY FORWARD

The search for safe and effective antiacanthamoebic drugs remains a challenge. Research over the past few decades has identified a large number of compounds that have therapeutic potential, but their translational value has not been explored. As discussed above, independent laboratories have done the groundwork in identifying several molecular targets and have identified several drugs of potential therapeutic value and used lead compounds for synthesis of derivatives; however, that work did not gain the attention of the major pharmaceutical companies, whose participation is needed to carry out the expensive *in vivo* studies as well as the clinical trials. Although the lack of interest of the pharmaceutical industry in finding cures for parasitic infections is well known, it is worth noting that eye infection due to the *Acanthamoeba* parasite occurs in contact lens wearers. The number of con-

tact lens wearers was estimated at 125 million in 2004, worldwide, with approximately 35 million contact lens wearers in the United States alone (90). The contact lens market was estimated at \$6.1 billion in 2010, and it was estimated that the global market would reach \$11.7 billion by 2015 (91). For a multi-billion-dollar industry, it is puzzling that pharmaceutical companies are not investing in this research, especially as novel molecules/inhibitors/drugs and their clinical applications can be patented, which offers tremendous commercial value. Notably, several companies have agreed to pay billions of dollars to settle lawsuits and have also withdrawn contact lenses/disinfectants from the market for being ineffective against *Acanthamoeba* or *Fusarium*. This makes no financial sense. It is far more economical to develop effective contact lens disinfectants against *Acanthamoeba*. Although the ability of amoebae to switch phenotypes into a dormant cyst form is a major hindrance in the development of effective contact lens disinfectants and/or chemotherapeutic approaches, recent studies have shown that the addition of cellulase enzyme to disrupt cyst wall structure renders amoeba cysts susceptible to the effects of antiacanthamoebic drugs (92). The combination of antiacanthamoebic agent and cellulase enzyme was shown to abolish the viability of both cysts and trophozoites. Notably, none of the agents, when tested alone, completely destroyed cysts and trophozoites, suggesting that the use of cellulose-degrading molecules is a useful avenue for targeted killing of amoebae. As cellulose synthesis is absent in mammalian cells, the use of cellulose-degrading molecules in contact lens disinfectants as well as in drug formulations in the treatment of *Acanthamoeba* infection needs to be explored.

It is hoped that the recent completion of the *Acanthamoeba* genome (93) will expedite identification of novel drug targets further, through genomics, proteomics, and bioinformatics. Among the existing drugs/disinfectants, given that they are limited in efficacy, there is a need to find ways to enhance their efficacy. The constituents of contact lens disinfectants must be carefully selected to target the cyst stage of amoebae. For example, the recall of Complete MoisturePlus contact lens disinfectant (AMO, Santa Ana, CA) following an outbreak of *Acanthamoeba* keratitis revealed that one of the constituents of the solution, propylene glycol, induced encystation in *Acanthamoeba*, resulting in the formation of cysts, which are resistant to the majority of contact lens disinfectants. Similarly, treatment is problematic due to specificity and parasite dormancy. The use of a carrier for antiacanthamoebic drug delivery is an important avenue that could yield promising results without affecting host cell viability. For eye infections, mannose- or antibody- or Fab-conjugated drugs should allow specific targeting of drugs to *Acanthamoeba*. Notably, recent studies showed that conjugation of mannose with photodynamic compounds allows specific targeting of *Acanthamoeba* and enhances their antiacanthamoebic effects (66). These findings suggested that specific antibodies or antiacanthamoebic agents, coupled with selective cytotoxic agents, could be useful in the treatment of *Acanthamoeba* infections, as they can be minimally invasive and minimally toxic to the host cells. Alternatively, parasite-specific pathways, such as ergosterol biosynthesis or cellulose biosynthesis, and the underlying enzymes that are required for the makeup of these molecules offer important targets for the rational development of therapeutic interventions.

Additionally, liposome-complexed antiacanthamoebic drugs have shown promising *in vitro* results in enhanced killing of pathogenic *Acanthamoeba* compared with the use of the drugs alone (94).

Moreover, liposomal ergosterol-pentamidine proved effective in preventing parasite-mediated host cell cytotoxicity *in vitro* (94), suggesting that ergosterol-formulated liposomes hold promise in the targeted delivery of drugs. The pace of research in identifying and characterizing novel targets has yielded promising results; however, the translational value for therapeutic interventions requires further investigation. The recent research shift to phenotypic screening against the whole parasite, as well as to repurposing of drugs, i.e., screening of FDA-approved drugs to identify those with antiacanthamoebic activity (68), is auspicious and has the potential to open several avenues for further research. Once active compounds are identified, the approval process can be expedited, as the drugs are already being used for clinical applications against other diseases. Moreover, several animal-based, plant-based, and microbe-based molecules have been identified that show antiacanthamoebic effects. Some of the aforementioned components represent appealing therapeutic targets that need to be exploited in future studies. With the availability of relevant disease models and of assays for target validation, there is an urgent need to develop translational research by encouraging academia-industry partnerships, which offer tremendous opportunities of commercial and scientific value.

## ACKNOWLEDGMENTS

This work was supported by Sunway University, Malaysia.

We declare that there is no conflict of interests regarding the publication of this paper.

## FUNDING INFORMATION

AQ: fund This work, including the efforts of Naveed Khan, was funded by Sunway University.

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## REFERENCES

- Khan NA. 2006. *Acanthamoeba*: biology and increasing importance in human health. *FEMS Microbiol Rev* 30:564–595. <http://dx.doi.org/10.1111/j.1574-6976.2006.00023.x>.
- Visvesvara GS, Moura H, Schuster FL. 2007. Pathogenic and opportunistic free-living amoebae: *Acanthamoeba* spp., *Balamuthia mandrillaris*, *Naegleria fowleri*, and *Sappinia diploidea*. *FEMS Immunol Med Microbiol* 50:1–26. <http://dx.doi.org/10.1111/j.1574-695X.2007.00232.x>.
- Marciano-Cabral F, Cabral G. 2003. *Acanthamoeba* spp. as agents of disease in humans. *Clin Microbiol Rev* 16:273–307. <http://dx.doi.org/10.1128/CMR.16.2.273-307.2003>.
- Niederhorn JY, Alizadeh H, Leher H, McCulley JP. 1999. The pathogenesis of *Acanthamoeba* keratitis. *Microb Infect* 1:437–443. [http://dx.doi.org/10.1016/S1286-4579\(99\)80047-1](http://dx.doi.org/10.1016/S1286-4579(99)80047-1).
- Panjwani N. 2010. Pathogenesis of *Acanthamoeba* keratitis. *Ocul Surf* 8:70–79. [http://dx.doi.org/10.1016/S1542-0124\(12\)70071-X](http://dx.doi.org/10.1016/S1542-0124(12)70071-X).
- Lloyd D. 2014. Encystment in *Acanthamoeba castellanii*: a review. *Exp Parasitol* 145(Suppl):S20–S27.
- Lorenzo-Morales J, Khan NA, Walochnik J. 2015. An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis and treatment. *Parasite* 22:10. <http://dx.doi.org/10.1051/parasite/2015010>.
- Carnt N, Stapleton F. 2016. Strategies for the prevention of contact lens-related *Acanthamoeba* keratitis: a review. *Ophthalmic Physiol Opt* 36:77–92. <http://dx.doi.org/10.1111/opo.12271>.
- Khan NA. 2015. *Acanthamoeba*: biology and pathogenesis, 2nd ed. Caister Academic Press, Linton, Cambridge, United Kingdom. ISBN: 978-1-908230-50-8.
- Balashubramanya R, Garg P, Sharma S, Vemuganti GK. 2006. *Acanthamoeba* keratitis after LASIK. *J Refract Surg* 22:616–617.
- Hay J, Kirkness CM, Seal DV, Wright P. 1994. Drug resistance and *Acanthamoeba* keratitis: the quest for alternative antiprotozoal chemotherapy. *Eye (Lond)* 8(Pt 5):555–563. <http://dx.doi.org/10.1038/eye.1994.137>.
- Lee JE, Hahn TW, Oum BS, Choi HY, Yu HS, Lee JS. 2007. *Acanthamoeba* keratitis related to orthokeratology. *Int Ophthalmol* 27:45–49. <http://dx.doi.org/10.1007/s10792-007-9055-8>.
- Stapleton F, Ozkan J, Jalbert I, Holden BA, Petsoglou C, McClellan K. 2009. Contact lens-related *Acanthamoeba* keratitis. *Optom Vis Sci* 86: E1196–E1201. <http://dx.doi.org/10.1097/OPX.0b013e3181baae11>.
- Kitagawa K, Nakamura T, Takahashi N, Oikawa Y, Ikeda T. 2003. A novel combination treatment of chlorhexidine gluconate, natamycin (pimaricin) and debridement for a *Acanthamoeba* keratitis. *Jpn J Ophthalmol* 47:616–617. <http://dx.doi.org/10.1016/j.jjo.2003.08.005>.
- Kosirukvongs P, Wanachiwanwin D, Visvesvara GS. 1999. Treatment of *Acanthamoeba* keratitis with chlorhexidine. *Ophthalmology* 106:798–802. [http://dx.doi.org/10.1016/S0161-6420\(99\)0169-0](http://dx.doi.org/10.1016/S0161-6420(99)0169-0).
- Lorenzo-Morales J, Morcillo-Laiz R, Martín-Navarro CM, López-Vélez R, López-Arencibia A, Arnalich-Montiel F, Maciver SK, Valladares B, Martínez-Carretero E. 2011. *Acanthamoeba* keratitis due to genotype T11 in a rigid gas permeable contact lens wearer in Spain. *Cont Lens Anterior Eye* 34:83–86. <http://dx.doi.org/10.1016/j.clae.2010.10.007>.
- Seal D, Hay J, Kirkness C, Morrell A, Booth A, Tullo A, Ridgway A, Armstrong M. 1996. Successful medical therapy of *Acanthamoeba* keratitis with topical chlorhexidine and propamidine. *Eye (Lond)* 10(Pt 4):413–421. <http://dx.doi.org/10.1038/eye.1996.92>.
- Sharma R, Jhanji V, Satpathy G, Sharma N, Khokhar S, Agarwal T. 2013. Coinfection with *Acanthamoeba* and *Pseudomonas* in contact lens-associated keratitis. *Optom Vis Sci* 90:e53–e55. <http://dx.doi.org/10.1097/OPX.0b013e31827f15b4>.
- Slater CA, Sichel JZ, Visvesvara GS, Pabico RC, Gaspari AA. 1994. Brief report: successful treatment of disseminated *Acanthamoeba* infection in an immunocompromised patient. *N Engl J Med* 331:85–87. <http://dx.doi.org/10.1056/NEJM199407143310204>.
- Wynter-Allison Z, Lorenzo Morales J, Calder D, Radlein K, Ortega-Rivas A, Lindo JF. 2005. *Acanthamoeba* infection as a cause of severe keratitis in a soft contact lens wearer in Jamaica. *Am J Trop Med Hyg* 73:92–94.
- Xuguang S, Lin C, Yan Z, Zhiqun W, Ran L, Shiyun L, Xiuying J. 2003. *Acanthamoeba* keratitis as a complication of orthokeratology. *Am J Ophthalmol* 136:1159–1161. [http://dx.doi.org/10.1016/S0002-9394\(03\)00635-4](http://dx.doi.org/10.1016/S0002-9394(03)00635-4).
- Gray TB, Gross KA, Cursons RT, Shewan JF. 1994. *Acanthamoeba* keratitis: a sobering case and a promising new treatment. *Aust N Z J Ophthalmol* 22:73–76. <http://dx.doi.org/10.1111/j.1442-9071.1994.tb01700.x>.
- Guerrero S, La Tegola MG, Monno R, Apruzzese M, Cantatore A. 2009. A case of descemet's membrane rupture in a patient affected by *Acanthamoeba* Keratitis. *Eye Contact Lens* 35:338–340. <http://dx.doi.org/10.1097/ICL.0b013e3181b912d6>.
- Kaur H, Maguire LJ, Salomao DR, Cameron JD. 2007. Rapid progression of amebic keratitis 1 week after corneal trauma and 1 year after LASIK. *Cornea* 26:212–214. <http://dx.doi.org/10.1097/ICO.0b013e31820eb136>.
- Kim EC, Kim MS. 2010. Bilateral *Acanthamoeba* keratitis after orthokeratology. *Cornea* 29:680–682. <http://dx.doi.org/10.1097/ICO.0b013e3181861bf9>.
- Larkin DF, Kilvington S, Dart JK. 1992. Treatment of *Acanthamoeba* keratitis with polyhexamethylene biguanide. *Ophthalmol* 99:185–191. [http://dx.doi.org/10.1016/S0161-6420\(92\)31994-3](http://dx.doi.org/10.1016/S0161-6420(92)31994-3).
- Lin HC, Hsiao CH, Ma DH, Yeh LK, Tan HY, Lin MY, Huang SC. 2009. Medical treatment for combined *Fusarium* and *Acanthamoeba* keratitis. *Acta Ophthalmol* 87:199–203. <http://dx.doi.org/10.1111/j.1755-3768.2008.01192.x>.
- Mills RA, Wilhelmus KR, Osato MS, Pyron M. 1993. Polyhexamethylene biguanide in the treatment of *Acanthamoeba* keratitis. *Aust N Z J Ophthalmol* 21:277–278. <http://dx.doi.org/10.1111/j.1442-9071.1993.tb00971.x>.
- Murdoch D, Gray TB, Cursons R, Parr D. 1998. *Acanthamoeba* keratitis in New Zealand, including two cases with *in vivo* resistance to polyhexamethylene biguanide. *Aust N Z J Ophthalmol* 26:231–236. <http://dx.doi.org/10.1111/j.1442-9071.1998.tb01317.x>.
- Rama P, Matuska S, Viganò M, Spinelli A, Paganoni G, Brancato R. 2003. Bilateral *Acanthamoeba* keratitis with late recurrence of the infection in a corneal graft: a case report. *Eur J Ophthalmol* 13:311–314.
- Rivasi F, Longanesi L, Casolari C, Croppo GP, Pierini G, Zunarelli E, Visvesvara GS. 1995. Cytologic diagnosis of *Acanthamoeba* keratitis. Report of a case with correlative study with indirect immunofluorescence and scanning electron microscopy. *Acta Cytol* 39:821–826.



32. Schnaidt AG, Gatzoufas Z, Schirra F, Hasenus AK, Seitz B. 2013. Delayed course of *Acanthamoeba* keratitis. *Ophthalmologie* 110:164–168. <http://dx.doi.org/10.1007/s00347-012-2707-8>.
33. Skarin A, Florén I, Kiss K, Mjörner H, Stenevi U. 1996. *Acanthamoeba* keratitis in the south of Sweden. *Acta Ophthalmol Scand* 74:593–597.
34. Tien SH, Sheu MM. 1999. Treatment of *Acanthamoeba* keratitis combined with fungal infection with polyhexamethylene biguanide. *Kaohsiung J Med Sci* 15:665–673.
35. Varga JH, Wolf TC, Jensen HG, Parmley VC, Rowsey JJ. 1993. Combined treatment of *Acanthamoeba* keratitis with propamidine, neomycin, and polyhexamethylene biguanide. *Am J Ophthalmol* 115:466–470. [http://dx.doi.org/10.1016/S0002-9394\(14\)74448-4](http://dx.doi.org/10.1016/S0002-9394(14)74448-4).
36. Wong VW, Chi SC, Lam DS. 2007. Good visual outcome after prompt treatment of *Acanthamoeba* keratitis associated with overnight orthokeratology lens wear. *Eye Contact Lens* 33(Pt 1):329–331. <http://dx.doi.org/10.1097/ICL.0b013e318030d5cf>.
37. Alizadeh H, Neelam S, Cavanagh HD. 2009. Amoebicidal activities of alexidine against 3 pathogenic strains of *Acanthamoeba*. *Eye Contact Lens* 35:1–5. <http://dx.doi.org/10.1097/ICL.0b013e3181909a66>.
38. Beattie AM, Slomovic AR, Rootman DS, Hunter WS. 1990. *Acanthamoeba* keratitis with two species of *Acanthamoeba*. *Can J Ophthalmol* 25:260–262.
39. Casemore DP. 1970. Sensitivity of *Hartmannella* (*Acanthamoeba*) to 5-fluorocytosine, hydroxystilbamidine, and other substances. *J Clin Pathol* 23:649–652. <http://dx.doi.org/10.1136/jcp.23.8.649>.
40. Nagington J, Richards JE. 1976. Chemotherapeutic compounds and *Acanthamoebae* from eye infections. *J Clin Pathol* 29:648–651. <http://dx.doi.org/10.1136/jcp.29.7.648>.
41. Lackner P, Beer R, Broessner G, Helbok R, Pfausler B, Brenneis C, Auer H, Walochnik J, Schmutzhard E. 2010. Acute granulomatous *Acanthamoeba* encephalitis in an immunocompetent patient. *Neurocrit Care* 12:91–94. <http://dx.doi.org/10.1007/s12028-009-9291-z>.
42. Berger ST, Mondino BJ, Hoft RH, Donzis PB, Holland GN, Farley MK, Levenson JE. 1990. Successful medical management of *Acanthamoeba* keratitis. *Am J Ophthalmol* 110:395–403. [http://dx.doi.org/10.1016/S0002-9394\(14\)77020-5](http://dx.doi.org/10.1016/S0002-9394(14)77020-5).
43. Inoue T, Asari S, Tahara K, Hayashi K, Kiritoshi A, Shimomura Y. 1998. *Acanthamoeba* keratitis with symbiosis of *Hartmannella* amoeba. *Am J Ophthalmol* 125:721–723. [http://dx.doi.org/10.1016/S0002-9394\(98\)00026-9](http://dx.doi.org/10.1016/S0002-9394(98)00026-9).
44. Ishibashi Y, Matsumoto Y, Kabata T, Watanabe R, Hommura S, Yasuraoka K, Ishii K. 1990. Oral itraconazole and topical miconazole with debridement for *Acanthamoeba* keratitis. *Am J Ophthalmol* 109:121–126. [http://dx.doi.org/10.1016/S0002-9394\(14\)75974-4](http://dx.doi.org/10.1016/S0002-9394(14)75974-4).
45. John T, Lin J, Sahm DF. 1990. *Acanthamoeba* keratitis successfully treated with prolonged propamidine isethionate and neomycin-polymyxin-gramicidin. *Ann Ophthalmol* 22:20–23.
46. Moore MB, McCulley JP, Luckenbach M, Gelender H, Newton C, McDonald MB, Visvesvara GS. 1985. *Acanthamoeba* keratitis associated with soft contact lenses. *Am J Ophthalmol* 100:396–403. [http://dx.doi.org/10.1016/0002-9394\(85\)90500-8](http://dx.doi.org/10.1016/0002-9394(85)90500-8).
47. Nakagawa H, Kazami N, Izai K, Iwasaki M, Uchida Y, Yamaura H, Shirasaka R, Horikami H, Ishii K. 1993. Two cases of early *Acanthamoeba* keratitis. *Nippon Ganka Gakkai Zasshi* 97:544–550.
48. Takatsu M, Nada T, Yamamoto H, Ichiyama S. 1995. A case of *Acanthamoeba* keratitis after operation for cataract. *Kansenshogaku Zasshi* 69:1159–1161. <http://dx.doi.org/10.11150/kansenshogakuzasshi1970.69.1159>.
49. Ofori-Kwakye SK, Sidebottom DG, Herbert J, Fischer EG, Visvesvara GS. 1986. Granulomatous brain tumor caused by *Acanthamoeba*. *Case report. J Neurosurg* 64:505–509.
50. Saxena A, Mittal S, Burman P, Garg P. 2009. *Acanthamoeba* meningitis with successful outcome. *Indian J Pediatr* 76:1063–1064. <http://dx.doi.org/10.1007/s12098-009-0205-z>.
51. Singhal T, Bajpai A, Kalra V, Kabra SK, Samantaray JC, Satpathy G, Gupta AK. 2001. Successful treatment of *Acanthamoeba* meningitis with combination oral antimicrobials. *Pediatr Infect Dis J* 20:623–627. <http://dx.doi.org/10.1097/00006454-200106000-00016>.
52. Yeung EY, Huang SC, Tsai RJ. 2002. *Acanthamoeba* keratitis presenting as dendritic keratitis in a soft contact lens wearer. *Chang Gung Med J* 25:201–206.
53. Driebe WT, Jr, Stern GA, Epstein RJ, Visvesvara GS, Adi M, Komadina T. 1988. *Acanthamoeba* keratitis. Potential role for topical clotrimazole in combination chemotherapy. *Arch Ophthalmol* 106:1196–1201.
54. Yip KW, Mao X, Au PY, Hedley DW, Chow S, Dalili S, Mocanu JD, Bastianutto C, Schimmer A, Liu FF. 2006. Benzethonium chloride: a novel anticancer agent identified by using a cell-based small-molecule screen. *Clin Cancer Res* 12:5557–5569. <http://dx.doi.org/10.1158/1078-0432.CCR-06-0536>.
55. Ben Salah S, Makni F, Cheikrouhou F, Ben Zina Z, Mlik M, Feki J, Colin J, Ayadi A. 2007. *Acanthamoeba* keratitis: about the first two Tunisian cases. *Bull Soc Pathol Exot* 100:41–42.
56. Gupta S, Shrivastava RM, Tandon R, Gogia V, Agarwal P, Satpathy G. 2011. Role of voriconazole in combined *Acanthamoeba* and fungal corneal ulcer. *Cont Lens Anterior Eye* 34:287–289. <http://dx.doi.org/10.1016/j.clae.2011.06.004>.
57. Kaul DR, Lowe L, Visvesvara GS, Farmen S, Khaled YA, Yanik GA. 2008. *Acanthamoeba* infection in a patient with chronic graft-versus-host disease occurring during treatment with voriconazole. *Transpl Infect Dis* 10:437–441. <http://dx.doi.org/10.1111/j.1399-3062.2008.00335.x>.
58. Tu EY, Joslin CE. 2010. Recent outbreaks of atypical contact lens-related keratitis: what have we learned? *Am J Ophthalmol* 150:602–608. <http://dx.doi.org/10.1016/j.ajo.2010.06.045>.
59. Vernon SE, Acar BC, Pham SM, Fertel D. 2005. *Acanthamoeba* infection in lung transplantation: report of a case and review of the literature. *Transpl Infect Dis* 7:154–157. <http://dx.doi.org/10.1111/j.1399-3062.2005.00113.x>.
60. Walia R, Montoya JG, Visvesvera GS, Booton GC, Doyle RL. 2007. A case of successful treatment of cutaneous *Acanthamoeba* infection in a lung transplant recipient. *Transpl Infect Dis* 9:51–54. <http://dx.doi.org/10.1111/j.1399-3062.2006.00159.x>.
61. Webster D, Umar I, Kolyvas G, Bilbao J, Guiot MC, Duplisea K, Qvarnstrom Y, Visvesvara GS. 2012. Treatment of granulomatous amoebic encephalitis with voriconazole and miltefosine in an immunocompetent soldier. *Am J Trop Med Hyg* 87:715–718. <http://dx.doi.org/10.4269/ajtmh.2012.12-0100>.
62. Nachega JB, Rombaux P, Weynand B, Thomas G, Zech F. 2005. Successful treatment of *Acanthamoeba* rhinosinusitis in a patient with AIDS. *AIDS Patient Care STDS* 19:621–625. <http://dx.doi.org/10.1089/apc.2005.19.621>.
63. Sheng WH, Hung CC, Huang HH, Liang SY, Cheng YJ, Ji DD, Chang SC. 2009. First case of granulomatous amoebic encephalitis caused by *Acanthamoeba castellanii* in Taiwan. *Am J Trop Med Hyg* 81:277–279.
64. Ma P, Willaert E, Juechter KB, Stevens AR. 1981. A case of keratitis due to *Acanthamoeba* in New York, New York, and features of 10 cases. *J Infect Dis* 143:662–667. <http://dx.doi.org/10.1093/infdis/143.5.662>.
65. Schuster FL, Visvesvara GS. 2004. Opportunistic amoebae: challenges in prophylaxis and treatment. *Drug Resist Updat* 7:41–51. <http://dx.doi.org/10.1016/j.drug.2004.01.002>.
66. Aqeel Y, Siddiqui R, Anwar A, Shah MR, Khoja S, Khan NA. 2015. Phototherapeutic strategy against *Acanthamoeba* infections. *Antimicrob Agents Chemother* 59:3031–3041. <http://dx.doi.org/10.1128/AAC.05126-14>.
67. Bouyer S, Imbert C, Daniault G, Cateau E, Rodier MH. 2007. Effect of caspofungin on trophozoites and cysts of three species of *Acanthamoeba*. *J Antimicrob Chemother* 59:122–124.
68. Baig AM, Iqbal J, Khan NA. 2013. *In vitro* efficacies of clinically available drugs against growth and viability of an *Acanthamoeba castellanii* keratitis isolate belonging to the T4 genotype. *Antimicrob Agents Chemother* 57:3561–3567. <http://dx.doi.org/10.1128/AAC.00299-13>.
69. Ondarza RN, Iturbe A, Hernández E. 2006. *In vitro* antiproliferative effects of neuroleptics, antimycotics and antibiotics on the human pathogens *Acanthamoeba polyphaga* and *Naegleria fowleri*. *Arch Med Res* 37:723–729. <http://dx.doi.org/10.1016/j.arcmed.2006.02.007>.
70. Draulans E, Maudgal PC. 1992. *Acanthamoeba* keratitis. *Bull Soc Belge Ophthalmol* 243:115–121.
71. Fung KT, Dhillion AP, McLaughlin JE, Lucas SB, Davidson B, Rolles K, Patch D, Burroughs AK. 2008. Cure of *Acanthamoeba* cerebral abscess in a liver transplant patient. *Liver Transpl* 14:308–312. <http://dx.doi.org/10.1002/lt.21409>.
72. Khurana S, Mewara A, Verma S, Totadri SK. 2012. Central nervous system infection with *Acanthamoeba* in a malnourished child. *BMJ Case Rep* <http://dx.doi.org/10.1136/bcr-2012-007449>.
73. Ardjomand N, Faschinger C, Haller-Schober EM, Scarpatetti M, Faulborn J. 2002. A clinico-pathological case report of necrotizing ulcerating keratopathy due to topical anaesthetic abuse. *Ophthalmologie* 99:872–875. <http://dx.doi.org/10.1007/s00347-002-0623-z>.

74. Brasseur G, Favennec L, Perrine D, Chenu JP, Brasseur P. 1994. Successful treatment of *Acanthamoeba* keratitis by hexamidine. *Cornea* 13:459–462. <http://dx.doi.org/10.1097/00003226-199409000-00015>.
75. McClellan K, Coster DJ. 1987. Acanthamoebic keratitis diagnosed by paracentesis and biopsy and treated with propamidine. *Br J Ophthalmol* 71:734–736. <http://dx.doi.org/10.1136/bjo.71.10.734>.
76. Moore MB, McCulley JP. 1989. *Acanthamoeba* keratitis associated with contact lenses: six consecutive cases of successful management. *Br J Ophthalmol* 73:271–275. <http://dx.doi.org/10.1136/bjo.73.4.271>.
77. Voyatzis G, McElvanney A. 2007. Bilateral *Acanthamoeba* keratitis in an experienced two-weekly disposable contact lens wearer. *Eye Contact Lens* 33:201–202. <http://dx.doi.org/10.1097/01.icl.0000252567.06446.7b>.
78. Wright P, Warhurst D, Jones BR. 1985. *Acanthamoeba* keratitis successfully treated medically. *Br J Ophthalmol* 69:778–782. <http://dx.doi.org/10.1136/bjo.69.10.778>.
79. Ziak P, Ondriska F, Mrva M. 2003. *Acanthamoeba* keratitis after use of soft contact lenses—case report. *Cesk Slov Oftalmol* 59:352–358.
80. Kim EC, Kim MS. 2009. Bilateral *Acanthamoeba* keratitis after orthokeratology. *Cornea* 28:348–350. <http://dx.doi.org/10.1097/ICO.0b013e31816b6a0b>.
81. Toshida H, Murakami A. 2009. An atypical case of microcysts associated with silicone hydrogel contact lens: findings on in vivo confocal laser microscopy. *Eye Contact Lens* 35:156–158. <http://dx.doi.org/10.1097/ICL.0b013e3181998dec>.
82. Aichelburg AC, Walochnik J, Assadian O, Prosch H, Steuer A, Perneczky G, Visvesvara GS, Aspöck H, Vetter N. 2008. Successful treatment of disseminated *Acanthamoeba* sp. infection with miltefosine. *Emerg Infect Dis* 14:1743–1746. <http://dx.doi.org/10.3201/eid1411.070854>.
83. Polat ZA, Obwaller A, Vural A, Walochnik J. 2012. Efficacy of miltefosine for topical treatment of *Acanthamoeba* keratitis in Syrian hamsters. *Parasitol Res* 110:515–520.
84. Walochnik J, Duchêne M, Seifert K, Obwaller A, Hottkowitz T, Wiedermann G, Eibl H, Aspöck H. 2002. Cytotoxic activities of alkylphosphocholines against clinical isolates of *Acanthamoeba* spp. *Antimicrob Agents Chemother* 46:695–701. <http://dx.doi.org/10.1128/AAC.46.3.695-701.2002>.
85. Cope JR. 2013. Investigational drug available directly from CDC for the treatment of infections with free-living amoebae. *MMWR Morb Mortal Wkly Rep* 62:666.
86. Paltiel M, Powell E, Lynch J, Baranowski B, Martins C. 2004. Disseminated cutaneous acanthamebiasis: a case report and review of the literature. *Cutis* 73:241–248.
87. Seijo Martinez M, Gonzalez-Mediero G, Santiago P, Rodriguez De Lope A, Diz J, Conde C, Visvesvara GS. 2000. Granulomatous amebic encephalitis in a patient with AIDS: isolation of *Acanthamoeba* sp. group II from brain tissue and successful treatment with sulfadiazine and fluconazole. *J Clin Microbiol* 38:3892–3895.
88. Morén H, Malmjö M, Mortensen J, Ohrström A. 2010. Riboflavin and ultraviolet A collagen crosslinking of the cornea for the treatment of keratitis. *Cornea* 29:102–104. <http://dx.doi.org/10.1097/ICO.0b013e31819c4e43>.
89. Hamide A, Sarkar E, Kumar N, Das AK, Narayan SK, Parija SC. 2002. *Acanthamoeba* meningoencephalitis: a case report. *Neurol India* 50:484–486.
90. Barr J. 2005. Contact Lens Spectrum's annual report of major corporate and product developments and events in the contact lens industry in 2004, as well as predictions for 2005. *Contact Lens Spectrum* Article ID 12733. Contact Lens Spectrum, Ambler, PA.
91. Nichols JJ. 2011. Market and survey data show that the industry remained largely unaffected in 2010 by the state of the economy. *Contact Lens Spectrum*, Article ID 105083. Contact Lens Spectrum, Ambler, PA.
92. Abjani F, Khan NA, Yousuf FA, Siddiqui R. 7 December 2015. Targeting cyst wall is an effective strategy in improving the efficacy of marketed contact lens disinfecting solutions against *Acanthamoeba castellanii* cysts. *Cont Lens Anterior Eye* <http://dx.doi.org/10.1016/j.clae.2015.11.004>.
93. Clarke M, Lohan AJ, Liu B, Lagkouvardos I, Roy S, Zafar N, Bertelli C, Schilde C, Kianianmomeni A, Bürglin TR, Frech C, Turcotte B, Kopec KO, Synnott JM, Choo C, Paponov I, Finkler A, Heng Tan CS, Hutchins AP, Weinmeier T, Rattei T, Chu JS, Gimenez G, Irimia M, Rigden DJ, Fitzpatrick DA, Lorenzo-Morales J, Bateman A, Chiu CH, Tang P, Hegemann P, Fromm H, Raoult D, Greub G, Miranda-Saavedra D, Chen N, Nash P, Ginger ML, Horn M, Schaap P, Caler L, Loftus BJ. 2013. Genome of *Acanthamoeba castellanii* highlights extensive lateral gene transfer and early evolution of tyrosine kinase signaling. *Genome Biol* 14:R11.
94. Siddiqui R, Syed A, Tomas S, Prieto-Garcia J, Khan NA. 2009. Effect of free versus liposomal-complexed pentamidine isethionate on biological characteristics of *Acanthamoeba castellanii* *in vitro*. *J Med Microbiol* 58:327–330. <http://dx.doi.org/10.1099/jmm.0.006494-0>.

**Naveed Ahmed Khan** received a B.Sc. degree (Pakistan), an M.Sc. degree (University of London, United Kingdom), and a Ph.D. degree (University of Hull, United Kingdom), followed by several years of research at Tufts University, Boston, MA, and Johns Hopkins University School of Medicine, Baltimore, MD. Dr. Khan joined the University of London as a faculty member, followed by the University of Nottingham, Nottingham, United Kingdom, as an Associate Professor of Molecular Microbiology and then the Aga Khan University as Professor and Chair of the Department of Biological and Biomedical Sciences and is now at Sunway University as Professor and Head of Biological Sciences. He maintains a broad interest in all aspects of infectious diseases, in particular, finding novel targets/sources for rational development of therapeutic interventions against parasitic infections.



## AUTHOR QUERIES

### AUTHOR PLEASE ANSWER ALL QUERIES

1

AQau—Please confirm the given-names and surnames are identified properly by the colors.

■ = Given-Name, ■ = Surname

AQaff—Please confirm the following full affiliations or correct here as necessary. This is what will appear in the online HTML version:

<sup>a</sup>Department of Biological Sciences, Faculty of Science and Technology, Sunway University, Malaysia

<sup>b</sup>Department of Molecular and Cell Biology, School of Medicine, Boston University, Boston, Massachusetts, USA

AQaff—This affiliation line will appear in the PDF version of the article and matches that on page 1 of the proof; corrections to this affiliation line may be made here **or** on page 1 of the proof:

Department of Biological Sciences, Faculty of Science and Technology, Sunway University, Malaysia<sup>a</sup>; Department of Molecular and Cell Biology, School of Medicine, Boston University, Boston, Massachusetts, USA<sup>b</sup>

AQfund—The Funding Information section includes information that you provided on the submission form when you submitted the manuscript. Any specific funding source(s) that you identified, along with its associated author(s) and, if applicable, grant number(s), appears in a paragraph (each funder-grant-author combination identified has a dedicated sentence). Any funding statement that you provided on the submission form appears in a separate paragraph. Please check the Funding Information section for accuracy, since the information may be restyled by ASM staff during preparation of the proofs. The final style of presentation is at the discretion of ASM staff.

Funder	Grant(s)	Author(s)
Sunway University		Naveed Khan

AQA—To ensure sequential order, references have been renumbered in the text and References.

Please check and correct the renumbering if necessary. If any reference should be deleted from the References list, please mark “Reference deleted” in the margin next to that entry; do not renumber subsequent references.

AQB—If “although those effects have not curtailed the use of the compound” is not as meant for “although used,” please clarify.

AQC—Please specify the yeast species meant here.

AQD—(i) The text of your original biography was trimmed per AAC policy. (ii) Please clarify what is meant by “Pakistan” here (by, e.g., specifying the name of the institution where the

## AUTHOR QUERIES

**AUTHOR PLEASE ANSWER ALL QUERIES**

**2**

degree was earned).

---