



IJUP

ENCONTRO 10ª EDIÇÃO
DE INVESTIGAÇÃO
JOVEM

UNIVERSIDADE
DO PORTO

BOOK OF ABSTRACTS

10TH MEETING OF YOUNG RESEARCHERS
OF UNIVERSITY OF PORTO

U. PORTO

IJUP

ENCONTRO
DE INVESTIGAÇÃO
JOVEM

104ª EDIÇÃO

UNIVERSIDADE
DO PORTO

U.PORTO

ORAL SESSIONS



- [12524 | Dispersion of plasmid-mediated colistin resistance by MCR-1 in pork-associated clinically relevant clones of Salmonella from Portugal](#)

Campos, Joana, UCIBIO/REQUIMTE. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Portugal

Cristino, Luís, Faculdade de Ciências da Nutrição e Alimentação, Portugal

Ribeiro, Sofia, Faculdade de Ciências, Portugal

Mourão, Joana, UCIBIO/REQUIMTE. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Portugal

Peixe, Luísa, UCIBIO/REQUIMTE. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Portugal

Antunes, Patrícia, UCIBIO/REQUIMTE. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Portugal; Faculdade de Ciências da Nutrição e Alimentação, Portugal

The first transferable gene conferring colistin (Col) resistance, *mcr-1* gene, was recently described in Enterobacteriaceae from several sources in China. Further worldwide reports of *mcr-1*, and of its variant *mcr-2* until now sporadic, have been associated mostly with animal sources, suggesting that its emergence is due to extensive Col use in food-producing animals. In this study, the presence of *mcr* genes in several Salmonella serotypes/clones obtained from different sources and regions of Portugal (2002-2016) was studied.

We analysed 1351 Salmonella isolates (66 serotypes), from several sources/regions (2002-2016) for the presence of *mcr-1/mcr-2* genes (PCR/sequencing). The susceptibility to Col and other antibiotics (AB) (microdilution/disk diffusion methods-EUCAST/CLSI), detection of other AB/metal (MT) resistance genes (PCR/sequencing), conjugation assays, plasmid characterization (ISAp1/PCRPBRT/pMLST/sequencing), genomic location (I-Ceul/S1-PFGE-hybridization) and clonality (XbaI-PFGE/MLST) were also performed.

MCR-2 was not detected, but MCR-1 was present in 1.3% of the isolates, (clinical/CL-n=5, pork/P-n=11 and animal-production/FP-n=1) from several regions (2011-2016). All presented resistance MICCol=4-8 mg/L and co-resistance to other AB and mostly carried several MT genes. They belonged to clinically relevant MDR serotypes/clones of S.1,4,[5],12:i:- [n=13/European clone-CL+P], S.Typhimurium [n=2-European clone-P/n=1-DT104 clone-CL] or S.Rissen (n=2-P), frequently associated with pig production and human infections. The *mcr-1* was located in 2 plasmid types, IncX4 or IncHI2 (ST4/non-typeable), most transferable.

Our study evidenced the acquisition of *mcr-1* by clinically relevant MDR Salmonella clones, with probable origin in pigs, in Portugal since at least 2011. The dissemination of those successful MDR clones and/or plasmids with *mcr-1* is of concern, since Col is frequently the only therapeutic alternative in infections with MDR Gram-negative bacteria.