

SUBAQUEOUS VOLCANICLASTIC SUCCESSIONS IN THE MIDDLE TRIASSIC OF WESTERN HUNGARY

BUDAI, T. 1, NEMETH, K. 1,2, MARTIN, U. 3, PIROS, O. 1

1. Geological Institute of Hungary, 14 Stefánia út, Budapest, Hungary
 2. Eötvös University, Department of Regional Geology, 14 Stefánia út, Budapest, Hungary
 3. Heidelberg, Germany

Budai, T., Németh, K., Martin, U. & Piros, O. (2004) Subaqueous volcanoclastic successions in the Middle Triassic of Western Hungary. In: Németh, K., Martin, U., Goth, K., & Lexa, J. (Eds) Abstract Volume of the Second International Maar Conference 21 - 26. September 2004., Lajosmizse/Kecskemét, Hungary, Occasional Papers of the Geological Institute of Hungary 203: p. 46.


Abstract

In western Hungary Middle Triassic sedimentation was steady from the Permian/Triassic till the end of the Early Anisian that was followed by carbonate platform disruption. Alkaline acidic volcanism started due to Late Anisian tectonism. The Middle Anisian Lofer cyclic platform carbonates are sharply overlain by reddish, grey or greenish crinoidal volcanoclastic limestone with ammonites. This sequence is overlain by a few m thick altered calcareous tuff ("pietra verde"). These beds are montmorillonitised, bentonitic, with green, yellowish, red matrix hosting vitro-, lithoclasts, and micro-holocrySTALLINE crystals. The K-rich trachyte became rhyolitic upwards with increasing calc-alkalinity. These beds are thicker in the Anisian basins (18 m) than above the platforms (5-8 m).


The Upper Ladinian sequence consists of silicified thickly bedded, red, grey, limestone with tuff layers, and with alternations of tuff, marl and thinly bedded limestone ("posidonia beds"). This sequence (as Buchenstein Formation) deposited in a pelagic basin, where carbonate deposition was ended by volcanism. The deposition of this sequence (30 m) occurred during the Longobardian substage in condensed sedimentation. In contrast in the Southern Alps the much thicker Upper Ladinian is represented by a volcanoclastic sandstone-silty-marl.

In western Hungary the Upper Anisian to Lower Ladinian volcanics are thick while they are of subordinate in the Upper Ladinian. Similarity does not exist in the thickness of the sequences between volcanoclastic rocks of the Lower Ladinian of western Hungary (tens of m) and the Livinalongo Formation (Dolomites, Italy) (180-200 m). The wide distribution of Lower Ladinian pyroclastics related to the higher explosivity of the magma and/or subaqueous reworking/redeposition.

The volcanism became basic and effusive during the Late Ladinian in the Southern Alps. In Hungary this sequence consists of volcanoclastic sandstones („wengen group”, Southern Alps). With decrease of silica content of the magma, its viscosity and explosivity decreased resulting limited dispersal of the deposits.



Middle Triassic Stratigraphy in Western Hungary



Middle Triassic Volcanoclastic Successions in Western Hungary

Middle Triassic volcanoclastic rocks from western Hungary are generally thin (dm-scale) units of inverse-to-normal graded tuff and lapilli tuff. The beds are altered by strong epidotisation and their colour reaches dark green with alternating pinkish fine tuff units. The volcanoclastic beds are intercalated with carbonate mudstone units and sandwiched between turbidite and debris beds of carbonate rich clastic units. The volcanoclasts are inferred to have been transported by turbulent volcanoclastic gravity currents. The microtexture of the volcanoclastic rocks are generally well-packed with angular to subrounded glassy volcanic fragments.

Strong epidote alteration of juvenile volcanic material replace the original texture of the rock completely in those locations where the volcanoclastic rocks reaches a total of few dm thickness.

The microtexture of the volcanoclastic rocks show a gradual transition from a primary pyroclastic to a reworked volcanoclastic character. Rocks that have well-packed and glassy fragment rich texture inferred to have more primary affinity.

A few cm to a dm thick volcanoclastic units embedded in pelagic limy turbidite beds are characteristically reworked in texture indicated by the abundance of altered coloured minerals and the strong abrasion of the glassy mafic clasts. Typical traction features indicate that the volcanoclastic interbeds are not suspension deposited fall units but horizontally transported volcanoclastic density currents transported to the pelagic regions by various subaqueous currents and/or the mechanical energy dispersed by the eruption nearby.


Facies Relationship to Southern Alps / Latemar

Typical phreatomagmatic succession with alternation of tuff and lapilli tuff rich in angular carbonatic accidental lithic fragments, impact sags, antidunes, scour fillings and traction depositional features indicate that shallow subaqueous (Surtseyan) to sunaerial (maar/diatreme) volcanism may have been responsible for the generation of these units in the Southern Alps.

A classic volcanoclastic bed called Tc interbedded in a pelagic carbonate mud rich siliciclastic turbidite unit (XXXXX). The volcanoclastic units bears textural characteristics indicating that it has been deposited in a similar way from physical point of view to those beds that are under- and overlain it. This bed documents a phase when more volcanic detritus got introduced to the pelagic depositional system due to distal volcanic eruption.

In the centre of the Latemar (Southern Alps, Italy) 4 massive volcanoclastic breccia zones, with fluidisation channels, collapsed wall rock blocks, entrapped fine tuff units and semicircular geometry have been identified recently and interpreted to be diatremes of phreatomagmatic volcanoes. The link between the distal "pietra verde" and such diatreme pipes is not established yet, however a casual link is seemingly obvious.

Eruptive Mechanism



Middle Triassic volcanoclastic rocks in Western Hungary are interpreted to be altered mafic volcanoclastic rocks deposited from small volume pyroclastic eruption generated turbidity currents in subaqueous environment. Magma/water interaction is inferred to have been an important fragmentation mechanism in the formation of pyroclasts however the exact transportation mechanism of the volcanoclasts is difficult to establish due to the limited drill core data. A casual relationship between the location of diatreme-like settings, near vent phreatomagmatic pyroclastic successions, and distal reworked mafic volcanoclastic units identified from the Southern Alps maybe a good analogy in the interpretation of the volcanoclastic rocks of similar age in the Western Hungarian realm.

Subaqueous volcanoclastic successions in the Middle Triassic of Western Hungary

Budai, Tamas

2004

<http://hdl.handle.net/10179/7635>

20/01/2020 - Downloaded from MASSEY RESEARCH ONLINE