

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.



"And when the big moment comes,
here's the nursery Robert and I have fixed up."

FRONT PIECE.

**HABITAT AND SEASONAL EFFECTS ON BLOWFLY ECOLOGY IN POSSUM
CARCASSES IN THE MANAWATU.**

A thesis presented in partial fulfilment for the
requirements for the degree of Master of Science
in Ecology at Massey University.

Clive Appleton

1993.

ABSTRACT.

Flies were trapped on Keeble farm and in Keeble forest reserve between May 1992 to February 1993. Fewer Calliphoridae were trapped in pasture during summer than in autumn or winter and different fly species showed differing preferences for bush or pasture at different times of the year. More flies were trapped around the bush margin than in either bush or pasture during spring and summer. In pasture the most numerous calliphorid trapped was *Calliphora stygia* (Fabricius, 1781), whereas in bush during summer *Calliphora hilli* Patton, 1925 was the predominant species. *Lucilia sericata* (Meigen, 1826) was never trapped in bush.

Decaying possum carcasses were examined in bush and pasture between February 1992 and February 1993. The number of maggots that left these carcasses was affected by the time of the year and whether the carcasses were in bush or pasture. More maggots left bush located possum carcasses during spring and summer while more maggots left possum carcasses in pasture during autumn and winter. Decay rates were most rapid during spring and summer. Temperatures were warm enough for adult fly activity during winter but the possum carcasses decayed much slower, especially in bush.

C. stygia and *Calliphora vicina* Robineau-Desvoidy, 1830 were generally the first species of maggot to leave possum carcasses and they were the only species present throughout the year in both habitats. *L. sericata* and *Chrysomya ruficacies* (Macquart, 1843) maggots were restricted to possum carcasses in pasture during spring and summer while *Calliphoroides antennatis* (Hutton, 1881) was restricted to possum carcasses in bush. *Hydrotaea rostrata* Robineau-Desvoidy, 1830 was the only species restricted to summer.

In spring 1992, 3400 flies emerged from a 2.5 kg possum carcass placed in pasture and 4200 flies emerged from a 3.1 kg possum carcass placed in bush. Estimated emergence success of adult flies was 22.0% in pasture and 20.2% in bush. This was 15% lower on average than the emergence level of comparable maggots raised in the laboratory. Flies began emerging in the bush 10 days after those in pasture.

L. sericata was restricted to pasture and *C. antennatis* was restricted to bush. Overall the major flystrike species to emerge were *C. stygia* and *L. sericata*. *C. stygia* made

stygia made up 46% of the total number of flies that emerged in pasture and 78% of those in bush, while *L. sericata* comprised 15.5% of the total number of flies that emerged in pasture.

The emergence success of field-collected maggots in the laboratory was significantly correlated with the estimated average total number of maggots that left the carcasses in the field. Maggots showed the highest emergence success in winter when larval competition in the carcass was lowest.

The temperature of a possum carcass containing fly maggots in bush reached 19.7°C during spring 1992 whereas the temperature exceeded 35°C on occasions in another carcass in pasture at the same time. Proportionally fewer maggots emerged from the possum carcass in pasture possibly because the high temperatures killed many maggots, especially those of *C. stygia*.

Maggots and bacterial decay both elevated carcass temperatures above ambient. Bacteria raised the temperature by about 2°C whereas maggots raised the temperature by 18.3°C to 26°C above ambient. The carcass temperature significantly affected the number of maggots that left the possum carcasses, and the number of maggots that left was directly related to the total maggot biomass that left the carcass. A minimum of 24% to 26% of the fresh weight of the possum carcasses was converted to maggot weight.

Acknowledgements.

I would like to firstly thank my supervisors Dr Ian Stringer and Dr Murray Potter (Ecology Dept Massey University) for their valuable contribution during research and writing of this thesis. Dr Allen Heath, David Cole and Dallas Bishop (AgResearch, Upper Hutt) for research advice and identification of specimens. Greg Arnold (Statistics Dept, Massey University) and Dr Ed Minot (Ecology Dept, Massey University) for statistical advice. Jens Jorgensen for helping with design suggestions and construction of carcass cages and emergence traps. Ellis Udy who supplied me with most of the possums for this study. Frank Arbon, for allowing me to shoot possums on his land. Keeble farm manager, Gerard Poff and Keeble Trust chairman, Mike Greenwood for permission to conduct this study on Keeble farm and in Keeble reserve. To my family for their support during this study. My wife Colleen, for her help in the field and encouragement during the writing of this thesis.

TABLE OF CONTENTS.

	Page.
Title page	i
Abstract	ii
Acknowledgements	iv
List of figures	ix
List of tables	xi
List of plates	xii
Chapter one. INTRODUCTION.	1
Chapter two. STUDY SITE.	3
2.1 Location.	3
2.2 Topography and vegetative cover.	3
2.3 Climate.	3
2.4 Collection of meterological data.	6
2.5 Climatic conditions during the research period.	7
Chapter three. HABITAT AND SEASONAL INFLUENCES ON FLY NUMBERS IN THE MANAWATU REGION, NEW ZEALAND.	8
3.1 INTRODUCTION.	8
3.2 METHODS.	9
3.3 Statistical analysis.	9
3.4 RESULTS.	12
3.5 DISCUSSION.	19
3.6 Habitat influence.	19
3.7 Seasonal influence.	20
3.8 Further research.	21

	Page.	
Chapter four.		
MAGGOT DEVELOPMENT, MAGGOT PRODUCTION AND FLY SPECIES SUCCESSION IN POSSUM CARCASSES.	22	
4.1	INTRODUCTION.	22
4.2	METHODS.	23
4.3	Possum carcasses.	23
4.4	Field sites.	23
4.5	Possum carcass cages.	23
4.6	Additional protection from attack by rats in the bush.	25
4.7	Sampling of maggots.	25
4.8	Estimating fly species succession in possum carcasses.	26
4.9	Estimating the number of maggots that escaped from the collection trays.	26
4.10	Statistical analyses.	27
4.11	RESULTS.	27
4.12	Diptera.	27
4.13	Predators and parasitoids of maggots.	32
4.14	Species presence and succession.	32
4.15	DISCUSSION.	36
4.16	Arrival of adult flies at possum carcasses.	36
4.17	Succession.	36
4.18	Maggot dispersal.	37
4.19	Habitat restriction.	38
4.20	Species competition.	38
4.21	Conclusion.	39
Chapter five.		
SPECIES EMERGENCE AND EMERGENCE TRENDS FROM POSSUM CARCASSES LOCATED IN BUSH & PASTURE.	40	
5.1	INTRODUCTION.	40

	Page.	
5.2	METHODS.	41
5.3	Statistical analysis.	44
5.4	RESULTS.	44
5.5	Emergence in the field.	44
5.6	Emergence in the laboratory.	45
5.7	DISCUSSION.	50
5.8	Number of flies produced by possum carcasses in the field.	50
5.9	Habitat influence on fly emergence.	50
5.10	Emergence success.	51
5.11	Further research.	52
Chapter six.	INFLUENCE OF TEMPERATURE ON THE NUMBER OF MAGGOTS LEAVING DECAYING POSSUM CARCASSES.	53
6.1	INTRODUCTION.	53
6.2	METHODS.	54
6.3	Statistical analyses.	56
6.4	RESULTS.	56
6.5	The temperature of decaying possum carcasses containing maggots in pasture and bush.	56
6.6	Temperatures of possum carcasses in bush with and without maggots present.	56
6.7	The relationship between the number of maggots that left the possum carcass per day and the carcass temperature and ambient temperature.	59
6.8	The amount of material remaining after decay of possum carcasses.	59
6.9	DISCUSSION.	64
6.10	Solar heating effects.	65
6.11	Influence of temperature on the number of maggots leaving a possum carcass.	65

	Page.
6.12 Conversion of carcase weight to maggot weight.	66
Chapter seven. CONCLUSIONS.	67
7.1 Habitat and seasonal effects on blowfly ecology.	67
7.2 Temperatures of possum carcases containing maggots and those without maggots.	68
7.3 Fly emergence in the field and laboratory.	68
7.4 FURTHER RESEARCH.	69
References.	70
Appendix.	
1. Rainfall recorded during the study period.	75
2. Mean ambient temperature during the study.	76
3. Relative humidity for winter, spring and summer 2 runs.	77
4. Light intensity (microeinsteins $\mu\text{Em}^{-2}\text{s}^{-1}$) at carcase sites.	78
5. Light intensity (microeinsteins $\mu\text{Em}^{-2}\text{s}^{-1}$) at flytraps.	78
6. Wind readings (Km/hr) taken in bush and pasture.	79
7. Number of flies trapped for each sample during each trapping run.	80
8. The estimated total number of maggots produced by each possum carcase during each seasonal run.	85
9. The estimated average total number of maggots that left possum carcases in bush and pasture for each seasonal run.	86

List of figures.

		Page.
2.1	Map of Keeble farm and Keeble native bush reserve.	4
3.1	Map of Keeble farm and bush showing location of fly traps.	11
3.2	Numbers of flies caught for each species on each sample occasion in traps in bush and pasture.	16-17
3.3	Relative abundance of flies at each trapping site during spring and summer.	18
4.1	Map of Keeble farm and Keeble bush showing location of sites where possum carcasses were placed. The sites are labelled 1 to 10. Sites 1-5 were located in pasture. Sites 6-10 were located in bush.	24
4.2	The estimated mean number of maggots leaving possum carcasses per day in bush and pasture, for each seasonal run.	30
4.3	The estimated average total number of maggots that left possum carcasses in bush and pasture for each seasonal run.	31
4.4	Diptera species succession in possum carcasses located in bush and pasture during each seasonal run. Each dot signifies the presence of a species on a particular sampling occasion. Five possum carcasses were sampled in each habitat at each sampling time.	34
5.1	Total number of flies emerging per day.	46
5.2	The total number of flies emerging per day for each major species.	47
5.3	Species emergence succession from possum carcasses.	48
5.4	Mean ambient temperature during fly emergence.	48
5.5	Correlation between percentage of flies emerging in the laboratory and the estimated average total number of maggots that left possum carcasses during the seasonal runs.	49
6.1	Daily mean temperatures during decay of a possum carcass in pasture.	57
6.2	Daily mean temperatures during decay of a possum carcass in bush.	57

6.3	Daily mean temperatures during decay of two possum carcasses containing maggots in bush.	58
6.4	Daily mean temperatures of two possum carcasses without maggots in bush.	58
6.5	Relationship between the internal temperatures of possum carcasses in pasture and bush and ambient temperature in these situations.	60
6.6	Relationship between in internal temperature of possum carcasses and the estimated number of maggots that left these carcasses per day.	60
6.7	Relationship between the ambient temperature and the estimated number of maggots that left the possum carcasses per day.	61
6.8	The relationship between the estimated number of maggots that left the possum carcase and the total weight of these maggots.	62

List of tables.

	Page.	
3.1	Number of flies trapped in pasture for each trapping occasion.	14
3.2	Number of flies trapped in bush for each trapping occasion.	15
4.1	Time taken for events to occur and species to leave possum carcasses after they were placed in the field (in days).	28
4.2	Species ranking in order of appearance as maggots leave possum carcasses placed in pasture (P) and bush (B). For example. <i>C. quadrimaculata</i> was the second, third or fourth species to leave possum carcasses in autumn in bush.	35
5.1	Species percentage composition and the number of flies that emerged from possum carcasses between the 4.12.92 and 22.2.93.	45
5.2	The percentage of flies successfully emerging in the laboratory taken from each seasonal run.	46
6.1	The estimated number of maggots that left the possum carcasses and weight differences of the carcasses in bush between 24.3.93 and 16.4.93.	61

List of plates.

	Page.
2.1 Keeble reserve.	5
2.2 Meterological station on the farm.	5
3.1 Western Australian fly trap, fully assembled. Note opening on the side for allowing flies to enter the trap.	10
3.2 Fly trap dismantled, showing bait chamber top left, excluder bottom left, collection chamber top right (with funnel) and trap top, bottom right.	10
4.1 Photograph showing the amount of material that had been removed from possum carcass at site 2 after 30 days during spring.	33
4.2 The amount of material removed from possum carcass at site 2 after 32 days during autumn. Compare this photo to the one above and note the difference in state of decay.	33
5.1 Cage set-up with metal maggot inclosure at the pasture site.	42
5.2 Pasture located carcass after 13 days, just before emergence trap was erected.	42
5.3 Emergence trap in place over the possum carcass located in pasture.	43
6.1 Two carcass cages in the bush site. One is insect-proof and the other on the left allows free access to flies.	55
6.2 Possum carcass 1 (table 6.1) showing the temperature probe in place.	55
6.3 Possum carcass 3 (table 6.1) that had contained maggots at the end of the study.	63
6.4 Possum carcass 1 (table 6.1) that was not infested with maggots at the end of the study.	63