

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**THE IMPACT OF HONEY BEES ON
MONTANE ECOSYSTEMS WITHIN
TONGARIRŌ NATIONAL PARK**

A thesis presented in partial fulfilment
of the requirements for the degree of
Masters of Science in Ecology
at Massey University

Claire Lisa Murphy
1996

ABSTRACT

A study of the effects of honey bees on montane ecosystems was conducted during the summers of 1993/1994 and 1994/1995 at Tongariro National Park. Three possible effects of the introduced honey bee were examined. The primary aims of the study were to identify areas with and without honey bees and to identify differences in the pollination success of a weed species, heather (*Calluna vulgaris*) and a native species, the New Zealand flax (*Phormium tenax*) under different pollinator regimes, and to examine differences in the composition of native pollinator communities in these different bee areas.

The impact of honey bees on the reproductive success of heather, an important weed species in Tongariro National Park was examined over two flowering seasons. Insect visitation rates on heather flowers were low at each of the four study sites. Bagging plants to exclude insect flower visitors had little effect on female fitness. The potential of other pollen vectors, wind and thrips, as pollinators of heather was also examined. Both were determined to have a negative effect on several measures of female success, including pollen deposition, pollen tube formation, and pollination levels. However it appears that none of the pollen vectors (honey bees, wind or thrips) significantly effect the overall fitness of heather in terms of the viable seed produced.

The second part of the study examined the impact of honey bees on the pollination systems of a native plant species. Flax is thought to be predominantly bird pollinated, however, the floral resources are also utilised by a variety of native and introduced insect species. At some sites birds were either not present or rarely used the flowers. Seed set in flax was highest in heavily bird pollinated sites. The results also suggest, however, that flax has a flexible pollinating system that enables it to maintain a range of fruit and seed set levels under the different pollinator regimes.

The abundance and diversity of insect flower visitors on manuka and *Hebe stricta*, two common subalpine shrubs, was highly variable between sites, and between observation periods. Some of this variation may be described to differences in the weather or to altitude. However, I have shown that the abundance and diversity of diptera appears to be strongly influenced by levels of honey bee activity. This indicates that honey bees do play a role in determining the structure of pollinator communities and may be displacing a significant component of the native pollinating fauna.

ACKNOWLEDGEMENTS

I would like to thank my supervisor, Dr Alastair Robertson, for his support and encouragement throughout the course of this study. Without him I would never have got to where I am today. Alastair was always there if I needed him, particularly over the last few months, and he never gave up on me, for that I am extremely grateful.

To the Department of Conservation, in particular, the Tongariro /Taupo Conservancy and the Science and Research Division, your contribution and support on this project was greatly appreciated. Funding was also provided by the Massey University Post-Graduate Research Fund.

I would also like to thank the staff and fellow post-graduates of the Ecology Department, Massey University for all their help, practical and emotional, in completing this thesis. To the staff of the Whakapapa Field Station, thank you for making field work not only tolerable, but also interesting.

I would like to thank my parents, Shirley and Craig , for their support over the past years, - I have finally finished!. And to my husband, Phil thank you for putting up with me and for keeping things going at home when I was too busy to be there.

TABLE OF CONTENTS

TITLE PAGE	i
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
LIST OF FIGURES	vi
LIST OF TABLES	ix
GENERAL INTRODUCTION	
The Introduction of Species and Competitive Interactions	1
Tongariro National Park	6
CHAPTER 1.	
Honey Bees and Their Distribution Within Tongariro National Park	8
CHAPTER 2.	
Heather Pollen Vectors and Female Reproductive Success	15
CHAPTER 3.	
Variable Pollinator Regimes and Their Effect on Female Reproductive Success in Flax	36
CHAPTER 4.	
The Influence of Honey Bee Abundance on Native Pollinator Communities	64
REFERENCES	83

LIST OF FIGURES

GENERAL INTRODUCTION.

**CHAPTER 1. HONEY BEES AND THEIR DISTRIBUTION WITHIN
TONGARIRO NATIONAL PARK.**

Figure 1. The distribution of feeding stations and their utilisation by honey bees within Tongariro National Park. 12

Figure 2. Feeding station design. 13

**CHAPTER 3. HEATHER POLLEN VECTORS AND THEIR EFFECT ON
FEMALE REPRODUCTIVE SUCCESS**

Figure 1. Range of heather expansion from 1912 to 1993 within the Tongariro/Taupo Conservancy, DoC. 17

Figure 2. Map of Tongariro National Park showing the locations of heather study sites. 19

Figure 3. Proportion of set, aborted and unfertilised ovules, 1994 flowering season. 26

Figure 4. Proportion of set, aborted and unfertilised ovules, 1995 flowering season 31

**CHAPTER 4. VARIABLE POLLINATOR REGIMES AND THEIR EFFECT ON
FEMALE REPRODUCTIVE SUCCESS IN FLAX**

Figure 1. Diagrammatic representation of flax (*P tenax*) 40.

- Figure 2.** Flowering stages of *P. tenax* , showing characteristics used to determine floral age. 41
- Figure 3.** Map of Tongariro National Park and surrounding areas. 43
- Figure 4.** Nectar standing crops (mean \pm s.e) of Stage 2 flowers expressed as milligrams sugar equivalents. 49
- Figure 5.** Nectar production (mean \pm s.e) of bagged (●) and unbagged (○) flax flowers at Erua and Lake Rotoaira. All flowers were bagged at State Highway. 52
- Figure 6.** Fruit set (mean \pm s.e) at each site, calculated as a proportion of total flower production. 55
- Figure 7.** Seed set (mean \pm s.e) at each site, calculated as being the proportion of large seeds produced. 55
- Figure 8.** Insect damaged pods (mean \pm s.e) at each site. 57

CHAPTER 4. THE INFLUENCE OF HONEY BEE ABUNDANCE ON NATIVE POLLINATOR COMMUNITIES.

- Figure 1.** Tongariro National Park and surrounding areas, showing location of study sites. 70

Figure 2. Census results for manuka, expressed as encounter rate per minute of flower visits by guild. 75

Figure 3. Analysis of manuka pollinator communities 76

Figure 4. Census results for *Hebe stricta*, expressed as encounter rate per minute of flower visits by guild. 78

Figure 5. Census results for *Hebe stricta* at Mangawhero, expressed as encounter rate per minute of flower visits by guild, showing changes in pollinator communities over time. 79

Figure 6. Analysis of *Hebe stricta* pollinator communities 80

LIST OF TABLES

GENERAL INTRODUCTION	page
Table 1. Numbers of endemic and established alien species in major groups in New Zealand	2
CHAPTER 1 THE INFLUENCE OF HONEY BEE ABUNDANCE ON NATIVE POLLINATOR COMMUNITIES	
CHAPTER 2. HEATHER POLLEN VECTORS AND FEMALE REPRODUCTIVE SUCCESS	
Table 1. Insect Visitation Rates on Heather at four sites within Tongariro National Park.	23
Table 2 Pollen grain deposition under two pollinator regimes, 1994 flowering season.	24
Table 3 Frequency table showing number of pollen tubes formed at each site in bagged and unbagged heather flowers during the 1994 flowering season.	25
Table 4 ANOVA table showing the effects of site, bagging and individual plant variation on the proportion of heather ovules apparently fertilised, 1994 flowering season.	27
Table 5 ANOVA table showing the effects of site, bagging and individual plant variation on seed set in heather, 1994 flowering season.	28

Table 6 Pollen grain deposition under four pollination regimes, 1995 flowering season. 29

Table 7. Effects of trial, wind, thrips and individual plant variation on pollen tube formation, 1995 flowering season. 30

Table 8 ANOVA table showing the effects of wind, thrips and individual plant variation on the pollination of heather ovules, 1995 flowering season. 32

Table 9 ANOVA table showing the effects of wind, thrips and individual plant variation on seed set in heather, 1995 flowering season. 32

CHAPTER 3. VARIABLE POLLINATOR REGIMES AND THEIR EFFECT ON FEMALE REPRODUCTIVE SUCCESS IN FLAX.

Table 1. Flax flower visitation rates, expressed as # visits/flower/hour, 47 for (a) insects and (b) birds at six sites around Tongariro National Park.

Table 2. Census results for insects, expressed as encounter rates/minute 48 of flax floral visitors.

Table 3. Nectar Standing Crop (n=204) averaged over all study sites. 49

Table 4. Comparisons of mean nectar production in some bird 50 pollinated species.

Table 5. ANOVA showing the effects of day, bagging and individual 53 plant variation on nectar sugar(mg) production at three study sites.

Table 6. Mean and range of pollen tubes formed per stigma for each site.	54
Table 7. ANOVA showing the effect of site and individual plant variation on fruit set at six study sites.	56
Table 8. ANOVA showing the effects of site, and individual plant and inflorescence variations on seed set at six study sites.	56
Table 9. ANOVA showing the effects of site variation on insect damage at six study sites.	57

CHAPTER 4. THE INFLUENCE OF HONEY BEE ABUNDANCE ON NATIVE POLLINATOR COMMUNITIES.

Table 1. Flower visitor diversity for two plant species (a) manuka and (b) <i>Hebe stricta</i> in relation to honey bee densities at each study site.	72
Table 2. Temporal changes in flower visitor diversity for <i>Hebe stricta</i> at Mangawhero	73
Table 3. Insect visitation rates by guild on manuka flowers, expressed as the number of visits/flower/hour, at four sites around Tongariro National Park.	73
Table 4. Census results for manuka.	74
Table 5. Census results for <i>Hebe stricta</i>	77