

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**EFFECT OF PLANT NUTRITION, TIME AND METHOD OF
HARVESTING ON SEED YIELD AND QUALITY OF WRINKLED AND
SMOOTH-SEEDED PEA(*Pisum sativum* L.) VARIETIES.**

A thesis presented in partial fulfilment of the requirements
for the degree of Master of Agricultural Science
in Plant Science(Seed Technology)
at
Massey University, Palmerston North
New Zealand

JUMPA PADRIT

1996

ABSTRACT

Effects of nitrogen(N), phosphorus(P) and time and method of harvesting on seed yield and quality were examined for smooth(Maple) and wrinkled(Pania)-seeded pea cultivars. These cultivars were grown under field conditions, and at different rates of N(0, 100 and 200 kg N/ha) and P(0 and 250 kg superphosphate/ha). Seed was harvested at 3 different times(35%, 25% and 15%SMC), and shelled either by hand or by a combine-harvester(at 1,350rpm). For hand-shelling, 120 plants were sampled from each plot of which 40 plants were used to determine the number of pods/plant, seeds/pod and 1000-seed weight, and subsequently used to determine seed quality i.e. Standard Germination, Accelerated Aging, Conductivity and incidence of Hollow Heart. The remaining 80 plants from each plot were used for separation into bottom, middle and top pods and subsequently used to determine 1000-seed weight, seed germination, conductivity and the incidence of hollow heart.

For machine-threshing, 120 plants were sampled from each plot, threshed by combine-harvester and seed subsequently used to determine seed quality by 1000-seed weight, Standard Germination Test, Accelerated Aging, Conductivity Test and incidence of Hollow Heart. Samples from each plot, following machine-threshing, were also used to determine seed damage by visual observation and by the Ferric Chloride Test.

Pea cv. Pania produced higher seed yield than cv. Maple in both hand-shelling and machine-threshing due to a much higher numbers of seeds/pod(5.76 and 3.57 seeds/pod, respectively) and much lower mechanical damage(10.32% and 27.98%, respectively), but had a much lower capacity to produce seed of high vigour than cv. Maple.

Application of nitrogen increased seed yield of both hand-shelled and machine-threshed seeds due to increased numbers of pods/plant, whereas yield was not directly affected by phosphorus addition. However, interaction between 100kg N/ha and 250 kg superphosphate/ha decreased seed weight. Application of nitrogen also increased seed vigour as expressed by increased seed germination percentage after accelerated aging, decreased hollow heart incidence and decreased conductivity value, particularly in cv.Pania. Application of phosphorus had only a small effect on

seed vigour compared with that of nitrogen. Neither seed germination percentage nor mechanical damage was affected by application of nitrogen or phosphorus.

Hand-shelling at different seed moisture contents did not affect seed germination percentage or conductivity value of either cultivar, but delaying the harvest (at the lower seed moisture content) decreased seed vigour in cv. Pania, as expressed by decreased seed germination percentage after accelerated ageing and increased hollow heart incidence. Machine-threshing at different seed moisture contents resulted in different degrees of seed damage, and decreased seed vigour in both cultivars. The most severe damage in cv. Maple occurred when machine-threshed at 15%SMC, whereas in cv. Pania it occurred at 35%SMC. Least damage occurred at 35% and 25%SMC in cv. Maple and cv. Pania, respectively. Unlike hand-shelling, machine-threshing at lower seed moisture content resulted in higher seed vigour in both cultivars, suggesting that bruising which occurs mainly at the high seed moisture content is more harmful than splitting which mainly occurs at the low seed moisture content threshing, in terms of decreasing seed vigour.

The top pods on pea plants produced seeds with lower seed weight in both cultivars, with higher hollow heart incidence in cv. Pania and with higher conductivity value in cv. Maple, than middle and bottom pods. Application of 200 kg N/ha and 250 kg superphosphate/ha improved vigour of seed from different pod positions to a similar and high level.

I would like to express my gratitude to my chief supervisor, Prof. Murray J. Hill, for his assistance, guidance, supervision and encouragement throughout this study and his understanding in correcting my English. I am also greatly indebted to him for his direct concern about my personal welfare during the whole period of study and stay in New Zealand.

I would like to express my indebtedness to Prof. B.R. Watkin, my co-supervisor, for his close and continuous assistance in writing this thesis, endless patience in correcting my English, direct guidance and concern for my personal welfare and study.

I would also like to express my sincere thanks to Assoc. Prof. John G. Hampton, my co-supervisor, for his valuable suggestions and constructive criticism in this work.

My sincere gratitude and appreciation are also extended to:

Dr Chulhathep Pongsroyech, a special expert of Department of Agricultural Extension, Bangkok, Thailand, for selecting and encouraging me to study in New Zealand.

Assoc. Prof.S.J.Haslett of the Department of Statistics for his statistical assistance.

Mrs Karen A. Hill, for her technical assistance in the laboratory work.

Mr Craig R. McGill, for his help in computer analysis of the experimental data.

Mr Robert C. Southward for his technical assistance in field work.

Finally, I am grateful to the New Zealand Ministry of Foreign Affairs and Trade for their financial support, and the Royal Thai Government for allowing me to study in New Zealand.

TABLE OF CONTENTS

		Page
ABSTRACT		i
ACKNOWLEDGMENTS		iii
LIST OF PLATES		vii
LIST OF APPENDICES		viii
LIST OF TABLES		ix
CHAPTER 1.	INTRODUCTION & OBJECTIVES	1
CHAPTER 2.	LITERATURE REVIEW	
	2.1. SMOOTH-AND WRINKLED-SEEDED PEAS	5
	2.2. SEED VIGOUR	7
	2.2.1. Genetic Constitution	9
	2.2.2. Nutrition of Parent Plant	10
	2.2.3. Position on The Parent Plant	11
	2.2.4. Environment During Seed Development and Maturation	12
	2.2.5. Mechanical Damage	14
	2.2.6. Storage Condition	16
	2.3. SEED VIGOUR MEASUREMENT IN PEAS	17
	2.3.1. Hollow Heart	17
	2.3.2. Conductivity Test	19
CHAPTER 3.	EXPERIMENTAL DESIGN & MATERIALS AND METHODS	
	3.1. TREATMENTS	23
	3.2. EXPERIMENTAL DESIGN	23
	3.3. THE FIELD LAY-OUT	24
	3.4. MATERIALS AND METHODS	

3.4.1. Experimental Field	25
3.4.2. Land Preparation	25
3.4.3. Seed Lots	25
3.4.4. Time and Method of Sowing	25
3.4.5. Crop Management	26
3.4.6. Seed Moisture Content Determination	26
3.4.7. Time of Harvest	26
3.4.8. Method of Harvest	27
3.4.9. Pod Position Determination	27
3.5. DETERMINATION OF SEED QUALITY	
3.5.1. Seed Moisture Content Test	28
3.5.2. Germination Test	28
3.5.3. Conductivity Test	28
3.5.4. Hollow Heart Test	29
3.5.5. Accelerated Ageing Test	29
3.5.6. Thousand Seed Weight	29
3.5.7. Ferric Chloride Test	30

CHAPTER 4. RESULTS

4.1. SEED YIELD AND YIELD COMPONENTS	
4.1.1. Seed Yield	31
4.1.2. Yield Component	37
4.1.2.1. Pods/Plant and Seeds/Pod	37
4.1.2.2. Seed Weight	37
4.2. HARVESTING AND MECHANICAL DAMAGE	
4.2.1. Seed Damage	42
4.2.3. Ferric Chloride Test	46
4.3. SEED QUALITY	
4.3.1. Seed Germination	51
4.3.2. Accelerated Ageing Test	58
4.3.3. Electroconductivity	66

4.3.4. Hollow Heart	71
4.4. EFFECT OF POD POSITION ON SEED QUALITY	
4.4.1. Seed Weight	74
4.4.2. Seed Germination	76
4.4.3. Hollow Heart	78
4.4.4. Electroconductivity	81
CHAPTER 5. DISCUSSION	
5.1. SEED YIELD AND YIELD COMPONENTS	84
5.2. SEED DAMAGE	85
5.3. FERRIC CHLORIDE TEST	88
5.4. SEED GERMINATION	89
5.5. HOLLOW HEART	89
5.6. SEED VIGOUR	94
CONCLUSION	97
REFERENCES	99
APPENDICES	109

LIST OF PLATES

Plate		page
1.	Type of seed damage.	87
2.	Imbibition characteristics of Maple and Pania seeds.	92
3.	Hollow heart in garden pea.	93
4.	Normal seed	93

LIST OF APPENDICES

Appendix	Page
1. Proportion of split and bruised seed, following machine-threshing.	109
2. Harvesting dates and rainfall data.	110
3. Soil fertility data.	111

LIST OF TABLES

Table		Page
1.	Interaction effects of cultivar and seed moisture content on seed yield.	31.
2.	Interaction effects of cultivar and threshing method on seed yield.	32.
3.	Interaction effects of seed moisture content at threshing and threshing method on seed yield.	33.
4.	Interaction effect of nitrogen application and threshing method on seed yield.	34.
5.	Interaction effects of cultivar, addition of nitrogen and seed moisture content on seed yield.	35.
6.	Interaction effects of cultivar, seed moisture content at threshing and threshing method on seed yield.	36.
7.	Effect of cultivar and nitrogen-level on number of pods/plant and seeds/pod.	37.
8.	Interaction effects of cultivar and seed moisture content at threshing on seed weight.	38.
9.	Interaction effects of cultivar and threshing method on seed weight.	39.

10. Interaction effects of cultivar, seed moisture content and threshing method on seed weight. 40.
11. Interaction effects of nitrogen and phosphorus applications on seed weight. 41.
12. Interaction effects of cultivar and seed moisture content at threshing on seed damage. 42.
13. Interaction effects of cultivar and threshing method on seed damage. 43.
14. Interaction effects of seed moisture content at threshing and threshing method on seed damage. 44.
15. Interaction effects of cultivar, seed moisture content at threshing and addition of nitrogen on seed damage. 45.
16. Interaction effects of cultivar, seed moisture content at threshing and threshing method on seed damage. 46.
17. Interaction effects of cultivar and seed moisture content at threshing on seed damage. 47.
18. Interaction effects of cultivar and threshing method on seed damage. 48.
19. Interaction effects of threshing method and seed moisture content at threshing on seed damage. 49.

20. Interaction effects of cultivar, threshing method and seed moisture content at threshing on seed damage. 50.
21. Interaction effects of cultivar and addition of nitrogen on seed germination. 51.
22. Interaction effect of cultivar and seed moisture content at threshing on seed germination. 52.
23. Interaction effects of cultivar and threshing method on seed germination. 53.
24. Interaction effects of threshing method and seed moisture content at threshing on seed germination. 54.
25. Interaction effects of cultivar, threshing method and seed moisture content at threshing on seed germination. 55.
26. Interaction effects of cultivar, threshing method and seed moisture content at threshing on seed germination. 56.
27. Interaction effects between cultivar, threshing method, nitrogen application rate and seed moisture content at threshing on seed germination. 58.
28. Interaction effects of cultivar and addition of nitrogen on seed germination after accelerated ageing. 59.
29. Interaction effects of cultivar and seed moisture content at threshing on seed germination after accelerated ageing. 60.

30. Interaction effects of cultivar and threshing method on seed germination after accelerated ageing. 61.
31. Interaction effect of threshing method and seed moisture content at threshing on seed germination after accelerated ageing. 62.
32. Interaction effects of cultivar, addition of phosphorus and seed moisture content at threshing on seed germination after accelerated ageing. 63.
33. Interaction effects of cultivar, threshing method and seed moisture content at threshing on seed germination after accelerated ageing. 64.
34. Interaction of addition of nitrogen and phosphorus, and seed moisture content at threshing on seed germination after accelerated ageing. 65.
35. Interaction effect of cultivar and seed moisture content at threshing on the conductivity of pea seeds. 66.
36. Interaction effects of cultivar and threshing method on the conductivity of pea seeds. 68.
37. Interaction effects of threshing method and seed moisture content at threshing on the conductivity of pea seeds. 68.
38. Interaction effects of cultivar, threshing method and seed moisture content at threshing on the conductivity of pea seeds. 69.
39. Interaction effects of cultivar, addition of nitrogen and threshing method on the conductivity of pea seeds. 70.

- | | | |
|-----|--|-----|
| 40. | Effect of threshing method on the incidence of hollow heart. | 71. |
| 41. | Interaction effects of cultivar and nitrogen application rate on the incidence of hollow heart. | 72. |
| 42. | Interaction effect of the addition of phosphorus and threshing method on the incidence of hollow heart. | 73. |
| 43. | Interaction effects of cultivar and seed moisture content at threshing on the incidence of hollow heart. | 74. |
| 44. | Interaction effects of nitrogen application and pod position on seed weight. | 75. |
| 45. | Interaction effects of phosphorus application rate and pod position on seed weight. | 76. |
| 46. | Interaction effects of seed moisture content at harvest and pod position on seed germination. | 77. |
| 47. | Interaction effects of nitrogen application rate and pod position on seed germination. | 78. |
| 48. | Interaction effects of nitrogen application rate and pod position on the incidence of hollow heart in cv. Pania. | 79. |
| 49. | Interaction effects of phosphorus application rate and pod position on hollow heart incidence in cv. Pania. | 80. |
| 50. | Interaction effects of cultivar and pod position on electroconductivity. | 81. |

51. Interaction effects of pod position and seed moisture content at harvest on electroconductivity. 82.
52. Interaction effects of cultivar, nitrogen application rate and pod position on electroconductivity. 83.