Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

CAUSES OF MORTALITY AND CHARACTERISATION OF MYCOBACTERIOSIS IN ADULT NEW ZEALAND SEA LIONS (*PHOCARCTOS HOOKERI*) AT ENDERBY ISLAND

A thesis presented in partial fulfilment of the requirements for the degree of

Master of Veterinary Science

in

Wildlife Health

at Massey University, Manawatū, Palmerston North

New Zealand.

Baukje Mirjam Lenting

2017

ABSTRACT

The New Zealand sea lion (*Phocarctos hookeri*) is classified as endangered and "Nationally Critical" due to a declining population and restricted population range. There have been recent bacterial epizootic events at the breeding colonies of this species, however the role of disease in the population decline is not known. As part of the investigation into the population decline, the species management plan recommends investigation of disease agents affecting this species, their epidemiology and their long-term effects on population dynamics.

Since the 1998/1999 breeding season, post mortem examinations have been performed on deceased New Zealand sea lions at the Enderby Island breeding colonies during each breeding season, including the collection of samples for histology and bacteriology.

This study describes the causes of mortality in New Zealand sea lions one year of age and older at Enderby Island between the 1998/99 to 2010/11 breeding seasons inclusive, using the archived post mortem reports, histology samples and bacteriology samples. Conspecific trauma was found to be a significant cause of mortality (34.3%), as were various infectious causes (35.7%). The organism *Klebsiella pneumoniae* was isolated from non-pup New Zealand sea lions both from individuals that died from other causes and individuals that showed apparent morbidity as a result of this bacteria. These findings suggest that older animals may be reservoirs of infection for *K. pneumoniae*, which causes significant mortality in neonatal New Zealand sea lions.

Another important infectious agent that was described in non-pup New Zealand sea lions was *Mycobacterium pinnipedii*, which caused both subclinical and clinical disease. Mycobacteriosis of the lymph nodes, lungs, pleura, liver, peritoneum and reproductive tract was described in this study. Strain determination of the *M. pinnipedii* isolates grown show only minor strain variation among isolates, which may reflect the isolated geographic distribution of these animals. There was no apparent association between the individual strains of *M. pinnipedii* and their pathogenicity as indicated by the pathology present in infected animals.

ACKNOWLEDGEMENTS

Many individuals gave time and expertise to help make the research presented in this thesis possible. First and foremost, I cannot thank enough my supervisors Brett Gartrell and Wendi Roe, for all their time, support and teaching as this thesis progressed. Thank you to Brett for teaching me statistics, to Wendi for many hours of teaching me histology, and to both for teaching me to be a more ruthless editor.

The gross post mortem examinations and the collection of samples for histology and bacteriology used by this study were performed by researchers in Department of Conservation (DOC) field expeditions to Enderby Island, as part of ongoing DOC monitoring of the endangered New Zealand sea lion. Many thanks to all those that performed post mortems on New Zealand sea lions over the breeding seasons included in this thesis, in the sometimes wild subantarctic weather and field conditions. Thank you to Kelly Buckle, Wendi Roe, Sarah Michael, Katja Geschke, Padraig Duignan, Nick Gales, Simon Childerhouse, Jacinda Amey and Amelie Auge, and to all other field researchers involved in the post mortem examinations performed during this study period.

Louise Chilvers from the Department of Conservation led and organised the expedition for the 2009/10 breeding season when I went to Enderby Island for 6 weeks in the position of field veterinarian, as well as many other expeditions. A big thank you to Louise for an extremely well planned and enjoyable expedition, and for much helpful advice on the fieldwork that was performed and day-to-day subantarctic island life. Thanks to my fellow expedition members Nathan, Elaine and Eric for moral and nutritional support, and a lot of rounds of "Uno".

The IVABS histology and pathology departments allowed me to use their work spaces for the trimming of tissues and storage of samples, and performed the processing of tissues for histology. Thank you to Mike Hogan and Evelyn Lupton especially.

Thank you to Stuart Hunter for helping with the interpretation of the histology for some of the cases.

Aerobic bacteriology cultures were performed by the ever helpful and skilled team at NZVP, Palmerston North, and by Hamish at IVABS. Mycobacterial culture, PCR and *Mycobacterium pinnipedii* strain determination were performed by the team at AgResearch Wallaceville, with thanks especially to Geoff de Lisle and Marian Price-Carter. For your many hours of lab work and all your expertise I am very grateful.

And lastly thanks to the wonderful Jon and to my family and friends, for all your support throughout. And to Nicky Denning, for letting me "thesis" by the fire under a handmade quilt on many occasions.

TABLE OF CONTENTS

Abstract	ii
Acknowledgements	iii
Table of Contents	v
List of Figures	viii
List of Tables	xi
CHAPTER 1: Literature Review	1
1.1 New Zealand Sea Lion Distribution and Biology	1
1.2 Causes of Mortality in Pinnipeds	7
1.2.1 Non-infectious causes of mortality	7
1.2.1.1 Predation	7
1.2.1.2 Conspecific trauma	8
1.2.1.3 Biotoxins	10
1.2.2 Infectious disease	10
1.2.2.1 Mass mortality events in New Zealand sea lions	10
1.2.2.2 Other disease agents of potential significance to the NZSL	12
1.3 Disease Ecology in Pinnipeds	15
1.3.1 Disease transmission in pinnipeds	15
1.3.2 Effects of disease on pinniped populations	16
1.4 Tuberculosis	17
1.4.1 The Mycobacterium tuberculosis complex	17
1.4.2 Pathophysiology of tuberculosis	18
1.4.3 Mycobacterium pinnipedii	20
1.4.3.1 Taxonomy	20
1.4.3.2 Distribution of Mycobacterium pinnipedii	21
1.4.3.3 Mycobacterial infection in New Zealand pinnipeds	22
1.4.3.4 Pathology of Mycobacterium pinnipedii infection in pinnipeds	23
1.4.3.5 Diagnosis of Mycobacterium pinnipedii infection in pinnipeds	25
1.4.3.6 Zoonotic implications	27

1.4.3.7 Implications for other species	27
1.5 Research Aims	29
CHAPTER 2: Post Mortem Findings in Non-Pup New Zealand Sea	
(Phocarctos hookeri) at Enderby Island	
2.1 Introduction	30
2.2 Materials and Methods	32
2.3 Results	34
2.3.1 General comments	34
2.3.2 Causes of mortality	39
2.3.2.1 Bacterial and parasitic infections (n = 25)	39
2.3.2.2 Conspecific trauma (n = 24)	42
2.3.2.3 Predation (n = 2)	44
2.3.2.4 Unknown causes of death (n = 19)	44
2.3.3 Concurrent and incidental diseases	45
2.3.3.1 Bacterial infections	45
2.3.3.2 Other bacterial isolates	45
2.3.3.3 Parasitism	48
2.3.3.4 Predation injuries	49
2.3.3.5 Neoplasia	49
2.4 Discussion	49
2.5 Conclusions	57
CHAPTER 3: Characterisation of Mycobacterium pinnipedii infecti	on in New
Zealand Sea Lions (<i>Phocarctos hookeri</i>) at Enderby Island	
3.1 Introduction	59
3.2 Materials and Methods	61
3.2.1 Archived samples and data	
3.2.2 Sample preparation, lesion review and laboratory analysis	
3.3 Results	
3.3.1 Gross and histological findings	
J.J.1 GIU33 AIU III3WIUZICAI IIIUIIIZS	

3.3.2 Identification and characterisation of mycobacterial isolates	64
3.3.3 Characterisation of pathology associated with confirmed mycoba NZSLs	
3.4 Discussion	87
3.4.1 Pathology	87
3.4.2 Strain variation in Mycobacterium pinnipedii from NZSLs	92
3.4.3 Importance of mycobacteriosis to the NZSL	93
3.4.4 Study limitations	99
3.4.5 Implications for livestock health and zoonotic disease	102
3.5 Conclusions	103
CHAPTER 4: General Discussion	104
APPENDIX 1: Post Mortem, Histology and Bacteriology Findings	110
LITERATURE CITED	150

LIST OF FIGURES

Figure 1.1	A map of the Auckland Islands, showing their location in relation to New Zealand and other sites of interest with regards to the current distribution of the New Zealand sea lion. Figure included with permission from the Department of Conservation
Figure 2.1	Causes of mortality of non-pup NZSLs at Enderby Island for the breeding seasons 1998/99 to 2010/11 inclusive
Figure 2.2	Number of tag resights and number of dead NZSLs that had post morter examinations performed at Enderby Island for the breeding seasons 1998/99 to 2010/11 inclusive
Figure 3.1	A case of mycobacterial infection in a NZSL examined at IVABS Massey University Palmerston North showing severe consolidation and multifocal coalescing granulomas of various sizes in the lung on cursurface
Figure 3.2	Haematoxylin and eosin stain. Scale bar represents 2mm. Lung tissue from case E08/09-53Ph showing severe chronic multifocal to coalescing granulomatous bronchopneumonia. There are large areas of necrosis surrounded by granulomatous inflammation. In parts of the lungs that were less severely affected, smaller discrete granulomas were present with or without necrotic centres
Figure 3.3	Haematoxylin and eosin stain. Scale bar represents 200µm. Lung tissue from case E08/09-53Ph showing lung granulomas consisting central necrosis surrounded by an inflammatory infiltrate of predominantly macrophages and a small number of lymphocytes and plasma cells71
Figure 3.4	Ziehl-Neelsen stain. Scale bar represents 50µm. Lung granulomas from case E00/01-42Ph showing a small number of visible AFOs intracellularly in macrophages and in necrotic debris
Figure 3.5	An enlarged middle tracheobronchial lymph node from case E09/10-13Ph, showing a cream-coloured nodular mass 20mm in diameter on cu surface
Figure 3.6	Haematoxylin and eosin stain. Scale bar represents 2mm. A lymph node from case E09/10-13Ph showing large well-circumscribed and ofter coalescing areas of necrosis, surrounded by a rim of granulomatous inflammation with a small number of lymphocytes

Figure 3.7	Ziehl-Neelsen stain. Scale bar represents 50µm. A lymph node from case E09/10-13Ph showing occasional visible AFOs in granulomas. In this case, AFOs were seen in the cytoplasm of macrophages and in necrotic debris
Figure 3.8	Gingival mucosa of case E09/10-17Ph showing severe jaundice76
Figure 3.9	Haematoxylin and eosin stain. Scale bar represents $500\mu m$. Liver tissue from case E09/10-17Ph showing a severe chronic granulomatous hepatitis. There are multifocal discrete and occasionally coalescing aggregations of macrophages randomly distributed throughout all examined liver sections. Normal hepatic architecture is disrupted77
Figure 3.10	Haematoxylin and eosin stain. Scale bar represents $200\mu m$. Liver tissue from case E09/10-17Ph showing a discrete aggregation of macrophages with a small number of associated lymphocytes and neutrophils. There is mild degeneration and necrosis of hepatic tissue surrounding the granulomatous inflammation
Figure 3.11	Ziehl-Neelsen stain. Scale bar represents $50\mu m$. Liver tissue from case E09/10-17Ph showing large numbers of intracellular AFOs within the macrophages of hepatic granulomas
Figure 3.12	Haematoxylin and eosin stain. Scale bar represents 500µm. Peritoneum from case E00/01-42Ph, showing moderate-severe, fibrinous and lymphocytic peritonitis, resulting in marked thickening of the peritoneum
Figure 3.13	Haematoxylin and eosin stain. Scale bar represents 200µm. Peritoneum from case E00/01-42Ph showing a moderate-severe, fibrinous and lymphocytic peritonitis with small multi-focal accumulations of macrophages throughout areas of thickened peritoneum, forming small granulomas
Figure 3.14	Haematoxylin and eosin stain. Scale bar represents 500µm. Uterus from case E00/01-42Ph showing a moderate granulomatous endometritis affecting approximately 30% of the endometrium of the examined section of uterus
Figure 3.15	Haematoxylin and eosin stain. Scale bar represents 200µm. Uterus from case E00/01-42Ph showing a large discrete accumulation of macrophages within the endometrium, with some lymphocytes and plasma cells in the periphery. The larger of these granulomas in other sections of uterus showed central necrosis and mineralisation

Opened pleural cavity of case E09/10-10Ph showing severe thickening of
the pleura and pericardium. The pleura is 2-3mm thick and white, with a
cobblestoned surface of soft villous nodules measuring 3-10mm in
diameter. There is a large volume of clear yellow-green viscous fluid in
the pleural space. There are multiple soft white round bodies on average
5mm in diameter floating free within the pleural effusion84
Haematoxylin and eosin stain. Scale bar represents 2mm. Lung and
pleura from case E09/10-10Ph showing severe chronic granulomatous
pleuritis, resulting in a markedly thickened pleura with villous
projections. There was a concurrent severe chronic granulomatous
pericarditis
Haematoxylin and eosin stain. Scale bar represents 200μm. Pleura from
case E09/10-10Ph showing a severe chronic granulomatous pleuritis
consisting of a thick layer of macrophages, fibrous tissue and occasional
plasma cells and lymphocytes with no distinct organisation, and with
small focal areas of necrosis86

LIST OF TABLES

Table 2.1	Causes of mortality in NZSLs at Enderby Island, showing the frequency of causes of mortality by cohort
Table 2.2	Bacterial isolates grown from post mortem tissues of non-pup NZSLs sampled at Enderby Island for the breeding seasons 1998/99 to 2010/11 inclusive
Table 3.1	Mycobacterial culture results from NZSL post mortem cases with confirmed mycobacterial infection
Table 3.2	Table of nine variable number of tandem repeat (VNTR) and two direct repeat (DR) loci for the isolates grown from six cases of <i>Mycobacterium pinnipedii</i> infection in New Zealand sea lions
Table 3.3	Histological findings, culture results and PCR results in nine New Zealand sea lions with confirmed mycobacterial infection67