

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

A COMPARISON OF FOLIAR AND SOIL UPTAKE

OF NUTRIENTS IN FRENCH BEAN

(PHASEOLUS VULGARIS L.).

A THESIS PRESENTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF HORTICULTURAL SCIENCE IN SOIL SCIENCE MASSEY UNIVERSITY

Susan Elizabeth Jolly

1986

ABSTRACT

An aspect of each of three factors relating to efficiency of fertilizer use were studied in glasshouse experiments using beans <u>Phaseolus vulgaris</u> var. Gallatin 50. These three factors were: the quantities that can be applied; physiological aspects of nutrient utilization following foliar uptake; and interactions with other sources of nutrient supply.

Distribution patterns of S^{35} , P^{32} and Zn^{65} were examined following application to soil and foliage of beans. It was found that a greater proportion of P^{32} and Zn^{65} was present in the fruit following foliar uptake than was the case following root uptake. This difference was not evident for S^{35} .

Retention of a commercial nutrient spray on the foliage of bean plants was measured and found to correlate well with both leaf area and leaf fresh weight.

The effect of sprays on leaf chlorophyll was also examined. Environmental effects were found to have more influence on leaf chlorophyll than nutrient sprays.

Root uptake of P³² was increased by spraying the foliage with either nutrient solution or water. It was concluded that the effect was water related and not connected with nutrient application.

The implications of the above findings were discussed in the

context of efficiency of fertilizer use.

I gratefully acknowledge the assistance of the following people:

Professor J.K. Syers, for supervision and encouragement in this study.

Mr R.W. Tillman, for his supervision, inspiration and understanding throughout this study.

Mr A.G. Robertson, for many helpful discussions and comments on aspects relating to plant physiology.

Jacqueline Rowarth and Howard Nicholson for their invaluable aid . with proof reading.

Martin Lewis, for his help with computing and graphics.

Members of the Soil Science department, especially Margaret Wallace whose help with analytical methods was much appreciated.

Hoescht (NZ) Ltd, for funding this project; and Massey University for the Helen E. Akers Scholarship, the Johannes August Anderson Scholarship, the Farmers Union Scholarship, the Sydney Campbell Memorial Scholarship and the Yates Corporation Bursary.

Finally, and most importantly, my parents, for their support and encouragement.

Table of Contents

p	a	g	e

ABSTRACT	 ii
ACKNOWLEDGEMENTS	 iv
TABLE OF CONTENTS	 v
LIST OF FIGURES	 x
LIST OF TABLES	 xii

CHAPTER 1

INTRODUCTION		1
--------------	--	---

REVIEW OF LITERATURE
2.1 Introduction
2.2 Pathway of nutrient movement during uptake6
2.2.1 Roots
2.2.1.1 Supply of nutrients to the root
2.2.1.2 Movement into and across the root
2.2.2 Leaves
2.2.2.1 Supply of nutrients to the leaf
2.2.2.2 Movement into and across the leaf

2.3 Translocation11
2.3.1 Root absorbed nutrients
2.3.2 Leaf absorbed nutrients14
2.4 Factors affecting uptake and translocation
2.4.1 Environmental factors15
2.4.1.1 Light16
2.4.1.2 Temperature16
2.4.1.3 Moisture17
2.4.1.4 Oxygen18
2.4.2 Solution Factors19
2.4.2.1 Composition19
2.4.2.2 Concentration20
2.4.2.3 pH21
2.4.2.4 Other solution factors
2.4.3 Plant factors22
2.4.3.1 Age and position of absorbing tissue22
2.4.3.2 Nutrient status of the plant24
2.4.3.3 Plant species
2.5 Interactions
2.6 Crop responses to fertilizers applied to soil and foliage27
2.6.1 Fertilizers applied to the soil27
2.6.2 Fertilizers applied to the Foliage
2.6.2.1 Nutrient levels and visual deficiency30
2.6.2.2 Vegetative Growth

		(i) 1	lacronu	trients					••		•••	• •	•••			• •		3	2
		(ii)	Micron	utrient	s		•••		•••		•••	•••	•••	• • •	• •	• •		3	3
		2.6.2.3	Yield:	Quanti	ty	• • •					•••		• •	•••	• •	•		3	3
		(i) !	Macronu	trients						•••					• •	• 1	•••	3	4
		(ii)	Micron	utrient	s	• • • •			•••	•••	•••	•••	••			•	•••	3	7
		2.6.2.4	Yield:	Qualit	у							••	••	•••	• •		•••	3	9
		2.6.2.5	Other	respons	es.				••	•••					• •	•		4	1
2.7	Beans	(Phaseo	lus vul	garis)	and	Fol	iar	Fe	ert	il:	zer	s.	••	•••	•••			4	3
	2.7.1	Botany.						•••	••	•••						•		4	3
	2.7.2	Nutriti	onal re	quireme	nts			•••	•••	•••			••			•	•••	4	4
	2.7.3	Foliar	fertili	zers an	id be	eans		•••	•••	••	•••	••	••	•••	•	•		4	5
2.8	Conclu	usions		· · · <i>· · · · ·</i>				•••								•		4	6

MATERIALS AND METHODS48
3.1 Preparation of plants
3.2 Nutrient applications50
(i)Spraying
(ii)Spot applications50
3.3 Plant tissue preparation50
3.4 Nutrient analyses
3.4.1 Total Sulphur
3.4.2 Total Phosphorus52

	3.4.3 Total Zinc
3.5	Isotope analyses
	3.5.1 Sulphur-3553
	3.5.2 Phosphorus-32
	3.5.3 Zinc-65

CHAPTER 4

DIST	TRIBUTION	PATTERNS	OF	S ³⁵	IN	BEANS	PHASEOLUS	VULGARIS59
4.1	Introduct	tion						60
4.2	Method				• • • •			61
4.3	Results.				• • • •			64
4.4	Discussio	on			• • • •			68

CHAPTER 5

DIST	RIBUTION	PATTERNS	OF	P ³²	AND	ZN ⁶⁵	IN	PHASE	EOLUS	VULGAR	IS	.74
5.1	Introduct	ion					•••					.75
5.2	Methods.		• • • •	• • • •			•••		• • • • •			.76
5.3	Results.			••••	• • • •	••••						.77
5.4	Discussio	on		••••								.79

THE	CONTRIBUTION	OF	SPRAY	RUNOFF	Τ0	PLANT	RESPONSES	TO	NUTRIENT	
SPRA	AYS								8	7

6.1	Introduction
6.2	Methods
6.3	Results
6.4	Discussion

EFFI	ECT	OF	FO	LIAR	S	PI	RA	YS		ON		TH	ΙE	ι	IP'	ГA	KI	Ξ	OI	F	P	3 2	E	BY	Τ	HI	Ξ	RC)0	TS		• •	٠	• •	•	104
7.1	In	troc	luc	tion		•			•		•		•		•		•		•	• •	•	• •		a•.		• •		• •	•		•		•		•	105
7.2	Me	thoo	ł		••	•		••	•	•••	•		•				•	•••	•	•••	•	• •		•				• •	•		•		•		•	106
7.3	Res	sult	ts.			•	• •		•		•		•		•	• •	•		•	• •	•			•		•	•••				•		•			107
7.4	Dis	scus	ssi	on	•••	•	• •		•	• •		•••	••	• •	•		•		•	• •	•			•		• •	• •				•		•		•	109
CON	CLU	SION	۱		• •	•	• •	• •	•		•	•	•	• •	•		•		•	• •		• •	•••	•		•		•••	•	•••	•	• •	•		•	115
SUM	MAR	Y	•••		• •	•	• •		•		•	• •	••	• •	•	• •	•		•	•••	•	• •	• •	•	• •	•		• •	•		•		•	• •	•	118
BIB	LIO	GRAI	PHY			•			•																						•					122

LIST OF FIGURES

2-1	Contact angles of droplets on a surface9
2-2	Generalised crop response curve for fertilizer application showing ranges of positive response (I), no response (II) and negative response (III)28
3-1	Flow sequence for the determination of
	phosphorus in plant material54
3-2	Curve relating H-number and counting efficiency for S ³⁵ 57
3-3	Curve relating H-number and counting efficiency for Zn ⁶⁵ 58
4-1	Position of application of K_2SO_4 (0.05M) to soil and the first and third trifoliate leaves62
4-2	Autoradiographs of beans (<u>Phaseolus vulgaris</u>) following uptake of S ³⁵ applied to a leaf (a) and the soil (b)72
6-1	Chlorophyllometer calibration curve for <u>Phaseolus vulgaris</u> leaves grown under glasshouse conditions (high range setting). Scale factor = 1.391

- 6-3 Minimum day humidities over the growing period.....97
- 6-4 Maximum day temperatures over the growing period.....99
- 6-5 Retention of Complesal solution on <u>Phaseolus</u> vulgaris as a function of fresh weight (r=0.76)....102
- 6-6 Retention of Complesal solution on <u>Phaseolus</u> <u>vulgaris</u> as a function of leaf area (r=0.78).....103
- 7-1 Diagram of pot showing position of labelled soil at time of transplanting.....107

LIST OF TABLES

4-1	Total Sulphur in plant parts (µg $g^{-1})\ldots\ldots\ldots65$
4-2	Recovery of applied S ³⁵ in plant parts and
	distribution of S ³⁵ within the plants67
4-3	Ratios (α) of specific activity of plant
	parts to the average specific activity
	of the whole plant69
5-1	Total P and Zn ($\mu g g^{-1}$) in bean
	plants (average over treatments)78
F 0	
5-2	Ratios (α) of specific activity of plant
	parts to the average specific activity of
	the whole plant for P^{32} and Zn^{65} 80
5-3	Recoveries of applied P ³² in plant
	parts and distribution within the plants
5-4	Recoveries of applied Zn ⁶⁵ in plant
	parts and distribution within the plants
5-5	Quantities of S and P supplied by one
	application of Complesal 12-2-5 and Zn
	supplied by one application of multimicro
	at recommended rates and assuming retention
	of 5 ml of solution per plant and plant dry

	weight of 5 g86
6-1	Dry weights of tops (g)93
6-2	Estimate of the nutrient that can be
	supplied to the surface of the leaf
	in a single application of Complesal
	12-2-5 solution, as a proportion
	of the total plant requirements100
6-3	Estimate of the nutrient that can be
	supplied to the surface of the leaf
	in a single application of Complesal
	multimicro solution, as a proportion
	of the total plant requirements101
7-1	Analysis of plant tops109
7-2	ANOVA for response to spraying111
7-3	Dry weights and root length estimates
	for sprayed and unsprayed plants
	(average of three replicates)111