

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

# **Characterisation of Serine Proteinase Inhibitors in Dry Seeds of Cultivated Pasture Grass Species**

A thesis presented in  
partial fulfilment of the requirements for  
the degree of Master of Philosophy in  
Plant Biology and Biotechnology at  
Massey University  
Palmerston North  
New Zealand

**Mohammad Tasneem**

**1994**

## Abstract

A chymotrypsin inhibitor has been partially purified from *Lolium perenne* cv. Grasslands Ruanui that is stable at pH 3 and after heating at 80°C for 10 minutes (acid/heat stable). The inhibitor had a native molecular weight of ca. 20-22 kDa determined using a 2.5 cm x 100 cm Sephadex G-75 column and gel electrophoresis indicated that it may be comprised of two subunits that had molecular weights of ca. 11 kDa and 12 kDa. To determine the occurrence of such inhibitors in other grass species, a survey of seeds from several cultivars of pasture grass species was conducted. Seeds of two cultivars of *Festuca arundinaceae*, cv. Grasslands Garland and cv. Grasslands Roa have been found to contain the most potent chymotrypsin inhibitory activity of the species surveyed. Seeds of two cultivars of an economically-important genus, *Lolium*, *L. perenne* cv Grasslands Ruanui and *L. x boucheanum* cv Grasslands Greenstone also exhibited significant chymotrypsin inhibitory activity, and so these four species were studied further. Using a 2.5 cm x 100 cm gel filtration column, chymotrypsin inhibitory activity eluted as two peaks in all four cultivars examined which had native molecular weights of ca. 20-22 kDa for peak I and ca. 10-12 kDa for peak II. A ca. 12 kDa chymotrypsin inhibitor was observed in peak 1 of *F. arundinaceae* cv. Grasslands Garland during gel electrophoresis, and was further purified by ion exchange chromatography. Twelve amino acid residues were sequenced from the N-terminal and the protein was found to be homologous with members of the cereal  $\alpha$ -amylase inhibitor family. To purify this activity further, a 10 cm x 100 cm gel filtration column was used. This time trypsin inhibitory activity was also assayed and for all four cultivars, both chymotrypsin and trypsin inhibitory activities co-eluted within two peaks, both with similar molecular weights (peak I, 20-22 kDa and peak II, 10-12 kDa) to that observed previously using the smaller gel filtration column. Gel electrophoresis of peak I from all the four cultivars revealed at least six comparatively lower molecular weight chymotrypsin inhibitors ranging from ca. 12 kDa to 3 kDa for the *Festuca* cultivars, and at least two ranging from ca. 8 kDa and 12 kDa for the *Lolium* cultivars. However, higher molecular weight trypsin inhibitors were present in Peak I with 5 inhibitors

ranging from ca. 12-18 kDa for *Festuca*, and three ranging from ca. 12-16 kDa for *Lolium*. Peak II from all the four cultivars contained only one 12 kDa inhibitor band. The 12 kDa inhibitor is active against both trypsin and chymotrypsin. The 12 kDa inhibitor from peak II of *F. arundinaceae* cv. Grasslands Garland was purified further using an anhydro-trypsin affinity chromatography and then to homogeneity by reverse-phase HPLC. Two inhibitory peaks were separated that had dual trypsin/chymotrypsin inhibitory activity and had an identical 20 amino acid residues at the N-terminal. The two inhibitor polypeptides were found to be homologous to members of the barley trypsin inhibitor family, but did not share homology with the ca. 12 kDa inhibitor protein purified by ion exchange from gel filtration peak I.

## ACKNOWLEDGEMENTS

First of all I would like to thank Almighty ALLAH who enabled me to complete this thesis. With out His help I am nothing. Next is my family who supported and encouraged me morally and financially through out my life. Many thanks to my wife Luna for her patience and support during research and writing up. I would like to thank all my friends in New Zealand and in Pakistan for their help, encouragement and suggestions particularly Tariq Javed, Abdul Khaliq, Manzoor Awan, S. Wajid Hussain, M. Anwar, S.G. Abbas, Babar Mahmood, Shahab, M. Idris. and Dr. M.A. Choudhury.

I would like to thank New Zealand Ministry of External Relation and Trade for tuition fees scholarship, the Miss E.L. Hellaby Trust and Pakistan Banking Council, for financial support.

I wish to thank all the staff of Plant Molecular Genetics Lab, AgResearch Grasslands for their help and cooperation throughout this research project especially Dr. D. W. White for permission to work in his Lab. I also thank all the staff of Department of Plant Biology and Biotechnology, Massey University for their help and cooperation, particularly Mrs Jacky McDonald and Ms E. Nickless.

Thanks are also due to Richard Scott for his cheerful technical help, Dr. Richard Biggs, AgResearch, Palmerston North for the use of anhydro-trypsin affinity column and Dr. W. Laing and Dr. John Christeller, HortResearch, Palmerston North for their assistance with reverse phase HPLC column and technical advice and HortResearch, Palmerston North Library staff for their help with literature search.

I gratefully acknowledge the help of Dr. Nick Elison, AgResearch, Palmerston North, for sequence alignment, Dr. C. Moore, Department of Biochemistry, Massey University, Palmerston North and Dr. David Christie, Department of Biochemistry, University of Auckland for the peptide sequencing, and Photographic Unit, Massey University, Palmerston North for the photography.

Finally I thank to my supervisors Dr. Michael McManus, Dr. Clive Cornford and Dr. David Fountain for their tireless and patient supervision, advice and encouragement.



## Table of Contents

Chapter 1 .....	1
Introduction .....	1
1.1 Proteinases and their Inhibitor Groups .....	2
1.1.1 Serine proteinases .....	2
1.1.2 Cysteine proteinases .....	2
1.1.3 Metallo-proteinases .....	3
1.1.4 Aspartic proteinases .....	3
1.2 Serine Proteinase Inhibitors .....	4
1.2.1 Mechanism of action .....	4
1.2.2 Families of serine proteinase inhibitors .....	5
1.2.2.1 Soybean (Kunitz) trypsin inhibitor family (STI) .....	5
1.2.2.2 Soybean proteinase inhibitor (Bowman-Birk inhibitor) family (BBI) .....	7
1.2.2.3 Potato inhibitor I and II .....	8
1.2.2.3.1 Potato inhibitor I .....	9
1.2.2.3.2 Potato inhibitor II .....	10
1.2.2.4 Squash trypsin inhibitor family .....	11
1.2.2.5 Barley trypsin inhibitor family (BTI) .....	12
1.2.2.6 White mustard/oil-rape trypsin/chymotrypsin inhibitor family .....	13
1.2.2.7 Serine proteinase inhibitors from <i>Brassica</i> ..	13
1.3 Inhibitors of Other Proteinases .....	14
1.3.1 Cysteine proteinase inhibitors .....	14
1.3.2 Metallo-carboxypeptidase inhibitors .....	14
1.3.3 Aspartic proteinase inhibitors .....	15
1.4 Localization of Proteinase Inhibitors in Plant Tissues .....	15
1.5 Implication for Animal Health .....	16
1.5.1 Nutritional significance .....	16
1.5.2 Allergens .....	16

	vi
1.6 Functions of Proteinase Inhibitors .....	17
1.6.1 Storage proteins .....	17
1.6.2 Regulation of endogenous proteinases .....	18
1.6.3 Plant Protection .....	19
1.7 Serine Proteinase Inhibitors from the Graminae Family .....	22
1.7.1 Serine proteinase inhibitors from cereals .....	22
1.7.2 Serine proteinase inhibitors from the grasses .....	26
1.8 Objectives .....	27
 Chapter 2 .....	 29
 Materials and Methods .....	 29
2.1 Plant Material .....	29
2.2 Chemicals .....	29
2.3 Extraction Protocols .....	29
2.3.1 Small scale extraction .....	29
2.3.2 Large scale extraction .....	30
2.3.3 Acid and heat treatment .....	30
2.4 Gel Filtration .....	30
2.4.1 Sephadex G-25 gel filtration .....	30
2.4.1.1 Spin column method .....	30
2.4.1.2 Large scale purification method .....	31
2.4.2 Sephadex G-75 gel filtration .....	31
2.4.2.1 Purification I .....	31
2.4.2.2 Purification II .....	32
2.4.3 Calibration of Sephadex G-75 column .....	32
2.5 Ion Exchange Chromatography .....	34
2.5.1 Anion exchange chromatography .....	34
2.5.2 Cation exchange chromatography .....	34
2.5.3 Anhydro-trypsin affinity column chromatography .....	35
2.5.4 Reverse-phase column chromatography .....	35



2.6 Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) and Western Blotting .....	36
2.6.1 SDS-PAGE .....	36
2.6.2 Minigels .....	36
2.6.3 Protein staining .....	37
2.6.4 Proteinase inhibition activity staining .....	37
2.6.5 Electroblotting on polyvinylidene difluoride membrane (PVDF) for sequencing .....	37
2.7 Sequencing and Identity with other Proteins .....	38
2.7.1 N-Terminal amino acid sequencing .....	38
2.7.2 Sequence homology search .....	38
2.8 Assays .....	39
2.8.1 Protein assay .....	39
2.8.1.1 Standard assay .....	39
2.8.1.2 Column fractions .....	39
2.8.2 Serine proteinase inhibitor (Pi) assays .....	40
 Chapter 3 .....	 41
 Results .....	 41
3.1 Identification of Serine Proteinase Inhibitory activity in dry seeds of <i>Lolium perenne</i> Cultivar Grasslands Ruanui (perennial ryegrass) .....	41
3.2 Survey of Serine Proteinase Inhibitory Activity in Seeds of Pasture Grass Species .....	46
3.3 Preliminary Purification of Proteinase Inhibitory Activity .....	48
3.3.1 Stability at low pH and high temperature .....	50
3.3.2 Initial Purification .....	50
3.3.3 Partial purification .....	52
3.3.4 Ion exchange chromatography of the chymotrypsin inhibitor from <i>F. arundinaceae</i> cultivar Grasslands Garland .....	56

3.3.5 N-terminal amino acid sequence of the chymotrypsin inhibitor purified by ion exchange chromatography . . .	58
3.4 Purification of a Proteinase Inhibitor to Homogeneity . . . . .	62
3.4.1 Partial purification of proteinase inhibitors . . . . .	63
3.4.2 Anhydro-trypsin affinity chromatography of a ca. 12 kDa dual trypsin/chymotrypsin inhibitor from <i>F. arundinaceae</i> cultivar Grasslands Garland . . . . .	69
3.4.3 Reverse-phase column chromatography of the ca. 12 kDa dual trypsin/chymotrypsin inhibitory peptide. . . . .	73
3.4.4 N-terminal amino acid sequence of reverse-phase HPLC purified inhibitor from <i>F. arundinaceae</i> cv. Grasslands Garland . . . . .	73
Chapter 4 . . . . .	79
Discussion . . . . .	79
References . . . . .	92

## List of Figures

Figure 2.1 Calibration curves obtained with a mixture of molecular weight standards (MW-GF-70; Sigma) . . . . .	33
Figure 3.1 Titration of bovine chymotrypsin with crude inhibitor protein from dry seeds of <i>Lolium perenne</i> cv. Grasslands Ruanui. . . . .	42
Figure 3.2 A typical Sephadex G-25 chromatograph of an acid/heat treated extract from dry seeds of <i>L. perenne</i> cv. Grasslands Ruanui. . . . .	42
Figure 3.3 Sephadex G-75 chromatography of an acid/heat treated extract from dry seeds of <i>L. perenne</i> cv. Grasslands Ruanui . . . . .	44
Figure 3.4 Titration of bovine chymotrypsin with a Sephadex G-75 purified inhibitor protein from <i>L. perenne</i> cv. Grasslands Ruanui . . . . .	44
Figure 3.5 SDS-PAGE of the chymotrypsin inhibitor preparation purified from <i>L. perenne</i> cv. Grasslands Ruanui after gel filtration . . . . .	45
Figure 3.6 chromatography through a 2.5 cm x 100 cm Sephadex G-75 column of acid/heat treated extracts from dry seeds of four pasture grass species . . . . .	53
Figure 3.7 Titration of bovine chymotrypsin with chymotrypsin inhibitor preparations from Sephadex G-75 purified inhibitor proteins from four pasture grass species . . . . .	55
Figure 3.8 SDS-PAGE of the chymotrypsin inhibitors preparation purified from <i>F. arundinaceae</i> cv. Grasslands Roa after gel filtration . . . . .	57

Figure 3.9 Elution profile of chymotrypsin inhibitory activity from <i>F. arundinaceae</i> cv. Grasslands Garland after ion exchange column .....	59
Figure 3.10 SDS-PAGE of chymotrypsin inhibitory activity purified from <i>F. arundinaceae</i> cv. Grasslands Garland by ion exchange chromatography .....	60
Figure 3.11 The N-terminal sequence of the 12 kDa chymotrypsin inhibitory peptide purified from <i>F. arundinaceae</i> cv. Grasslands Garland by ion exchange .....	61
Figure 3.12 Chromatography through a 10 cm x 100 cm Sephadex G-75 column of acid/heat treated extracts from dry seeds of four pasture grass species .....	64
Figure 3.13 SDS-PAGE of serine proteinase inhibitors purified by Sephadex G-75 chromatography from dry seeds of four pasture grass species .....	66
Figure 3.14 Titration of bovine chymotrypsin and bovine trypsin with Sephadex G-75 chromatography purified inhibitor preparations from four pasture grass species .....	68
Figure 3.15 Elution of dual trypsin/chymotrypsin inhibitory activity from <i>F. arundinaceae</i> cv. Grasslands Garland from an anhydro-trypsin affinity column. ....	71
Figure 3.16 SDS-PAGE of a dual trypsin/chymotrypsin inhibitor from <i>F. arundinaceae</i> cv. Grasslands Garland, purified by anhydro-trypsin affinity chromatography .....	72

- Figure 3.17 Titration of bovine chymotrypsin and bovine trypsin with the affinity-purified dual trypsin/chymotrypsin inhibitor from *Festuca arundinaceae* cv. Grasslands Garland ..... 72
- Figure 3.18 Reverse-phase column chromatography of affinity-purified dual trypsin/chymotrypsin inhibitor from *F. arundinaceae* cv. Grasslands Garland ..... 74
- Figure 3.19 The N-terminal sequence of the two dual trypsin/chymotrypsin inhibitory peptides purified from *F. arundinaceae* cv. Grasslands Garland by reverse-phase HPLC .. 77



## List of Tables

Table 1.1 Families of protein inhibitors of serine proteinases .....	6
Table 1.2 Survey of serine proteinase inhibitors from cereals .....	23
Table 3.1 Survey of trypsin and chymotrypsin inhibitory activity in extracts of seeds from various pasture grass .....	47
Table 3.2 Chymotrypsin inhibitory activity in crude extracts of different cultivars of <i>F. arundinaceae</i> .....	47
Table 3.3 Chymotrypsin inhibitory activity in crude extracts of <i>F.</i> <i>arundinacea</i> (cv. Grasslands Roa) seed representing different years of harvest .....	49
Table 3.4 Chymotrypsin inhibitory activity in crude extracts of <i>L. perenne</i> (cv. Grasslands Ruanui) seed representing different years of harvest .....	49
Table 3.5 Purification table of four cultivars for stability at low pH and high temprature .....	51

## LIST OF ABBREVIATIONS

BApNA	N-benzoyl-DL-arginine-p-nitroanalide
BBI	Soybean (Bowman-Birk) trypsin inhibitor
BTI	Barley trypsin inhibitor
BTpNA	N-benzoyl-L-tyrosine-p-nitroanalide
Ca.	Circa
CI-1	Barley chymotrypsin inhibitor 1
CI-2	Barley chymotrypsin inhibitor 2
CMe	Chloroform methanol soluble
CPTI	Cow pea trypsin inhibitor
ELISA	Enzyme-linked immunosorbent assay
FMTI	Foxtail millet trypsin inhibitor
h	Hour
HPLC	High pressure liquid chromatography
kDa	Kilo dalton
Ki	Enzyme inhibition constant
LPC	Leaf protein concentrate
min	Minute
OD	Optical density
P1	Proteinase inhibitor reactive site
PIIF	Proteinase inhibitor inducing factor
PVDF	Polyvinylidene difluoride membrane
RBTI	Rice bran trypsin inhibitor
SDS-PAGE	Sodium dodecyl sulphate-polyacrylamide gel electrophoresis
STI	Soybean (Kunitz) trypsin inhibitor
TFA	Trifluoroacetic acid
Tris	Tris (hydroxymethyl) amino-methane
cv.	Cultivar