

# FOREST CONNECTIVITY RESTORATION THROUGH REFORESTATION

## AN INTEGRATED METHODOLOGY FOR PRIORITIZING AGRICULTURAL LANDS AND SELECTING REFORESTATION SPECIES



Aitor Gastón<sup>1</sup>, Celia García-Feced<sup>1</sup>, Santiago Saura<sup>2</sup>

<sup>1</sup> Technical University of Madrid, ECOGESFOR research group, Madrid, Spain

<sup>2</sup> Technical University of Madrid, Department of Forest Management and Economics, ETSI Montes, Madrid, Spain



### OBJECTIVES


Provide land planners and managers effective tools and methodologies...

1. To identify in advance those landscapes where connectivity should be really treated as a critical concern for the conservation goals.
2. To optimize the reforestation of agricultural patches in order to favor the enhancement of forest connectivity.
3. To make a more reliable selection of reforestation species.

### STEP 1

Quantify forest connectivity within landscape units


- Discriminate and map the landscape types according to abiotic and biotic variables (García-Feced et al., 2008).
- Use the software Conefor Sensinode 2.2 (Saura and Torné, 2009; available at <http://www.conefor.org>), a powerful tool for analyzing potential landscape connectivity, to calculate within each landscape unit the Integral Index of Connectivity (IIC, Pascual-Hortal and Saura, 2006) at a specified dispersal distance (as an example, the dispersal distance of the figures is 1000 m).



### STEP 2

Identify priority agricultural patches for reforestation in order to enhance forest connectivity within the landscape units

- Calculate the increase of IIC (dIIC) that would result from the conversion of each agricultural patch into a forest (García-Feced et al., 2011).
- Classify dIIC values into five categories using natural breaks of the whole district values in order to prioritize patches for reforestation.



### STEP 3

Identify suitable tree species and order them by probability of occurrence

- Estimate occurrence probability of each tree species within the priority patches for reforestation using:
  - Ecological niche models fitted with penalized logistic regression (Gastón & García-Viñas, 2011).
  - Native tree species distribution data from the Spanish Forest Map as response variable and climatic and lithological variables as predictors.

3

| Species                     | Occurrence probability |
|-----------------------------|------------------------|
| <i>Quercus ilex</i>         | 0.60                   |
| <i>Q. faginea</i>           | 0.43                   |
| <i>Juniperus oxycedrus</i>  | 0.32                   |
| <i>Pinus nigra</i>          | 0.26                   |
| <i>P. halepensis</i>        | 0.22                   |
| <i>J. phoenicea</i>         | 0.19                   |
| <i>J. thurifera</i>         | 0.15                   |
| <i>Q. coccifera</i>         | 0.14                   |
| <i>Crataegus monogyna</i>   | 0.12                   |
| <i>J. communis</i>          | 0.09                   |
| <i>Acer monspessulanum</i>  | 0.02                   |
| <i>Amelanchier ovalis</i>   | 0.01                   |
| <i>Pistacia terebinthus</i> | 0.01                   |

Suitable tree species in a patch important for connectivity

### CONCLUSIONS

The major outputs of this combined methodology are: 1) A map of the agricultural patches that would contribute most to uphold forest connectivity if they were reforested. 2) A list of suitable tree species for those patches ordered by occurrence probability. Therefore this methodology may be useful for suitable and efficient forest planning and landscape designing.

### REFERENCES

- GARCÍA-FECED C., SAURA S., ELENA-ROSSELLÓ R., 2011. Improving landscape connectivity in forest districts: A two-stage process for prioritizing agricultural patches for reforestation. *For.Ecol.Manage.*, 261(1), 154-161.
- GARCÍA-FECED C., GONZÁLEZ-ÁVILA S., ELENA-ROSSELLÓ R., 2008. Metodología para la tipificación y caracterización estructural de paisajes en comarcas forestales españolas. *Forest Systems (Formerly Invest Agrar: Sist Recur For)* 17, 130-142. [In Spanish].
- GASTÓN A., GARCÍA-VIÑAS J.I., 2011. Modelling species distributions with penalised logistic regressions: A comparison with maximum entropy models. *Ecol.Model.* 222(13), 2037-2041.
- PASCUAL-HORTAL L., SAURA S., 2006. Comparison and development of new graph-based landscape connectivity indices: towards the prioritization of habitat patches and corridors for conservation. *Landscape Ecology* 21 (7): 959-967.
- SAURA S., TORNÉ J., 2009. Conefor Sensinode 2.2: a software package for quantifying the importance of habitat patches for landscape connectivity. *Environmental Modelling & Software* 24: 135-139.



RESTORING FORESTS:

Advances in Techniques and Theory

27.28.29 September 2011 | MADRID, SPAIN

IUFRO CONFERENCE