

106-4 Using a Large Ensemble of Crop Models to Simulate the Climate Sensitivity of Wheat Yields Across a European Transect.

Reimund P. Roetter¹, Nina K Pirttioja², Stefan K Fronzek², Timothy R Carter², Marco Bindi³, Roberto Ferrise³, Taru I Palosuo¹, Fulu Tao¹, Holger Hoffmann⁴, Frank Ewert⁵, Margarita Ruiz-Ramos⁶, Alfredo Rodriguez⁶, Maria Ines Minguez Tudela⁷, Miroslav Trnka⁸, Petr Hlavinka⁹, Marco Acutis¹⁰, Senthold Asseng¹¹, Piotr Baranow¹², Bruno Basso¹³, Per Bodin¹⁴, Samuel Buis¹⁵, Davide Cammarano¹⁶, Paola Deligios¹⁷, Marie-France Destain¹⁸, Benjamin Dumont¹⁹, Louis Francois²⁰, Ingrid Jacquemin²⁰, Chris Kollas²¹, Kurt C. Kersebaum²¹, Jaromir Kryszczak¹², Ignacio J Lorite²², Julien Minet²³, Manuel Montesino²⁴, Marco Moriondo²⁵, Christoph Müller²⁶, Claas Nendel²⁷, Isik Ozturk²⁸, Alessia Peregio²⁹, Sanna Mattia³⁰, Mikhail Semenov³¹, Czesary Slawinski³², Pierre Stratonovitch³¹, Iwan Supit³³, Alex Ruane³⁴, Francoise Ruget³⁵, Katharina Waha²⁶, Enli Wang³⁶, Lianhai Wu³⁷ and Zhigan Zhao³⁸, (1)Climate Impacts Group, Natural Resources Institute Finland (Luke), Helsinki, Finland

(2)Climate Change Programme, Finnish Environment Institute (SYKE), Helsinki, Finland

(3)University of Florence, Florence, Italy

(4)Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Bonn, Germany

(5)Institute of Crop Science and Resource Conservation (INRES), University of Bonn, Bonn, GERMANY

(6)Universidad Politecnica de Madrid, Madrid, Spain

(7)Polytechnic University of Madrid, Madrid, Spain

(8)Mendel University Brno, Brno, Czech Republic

(9)Institute of Agrosystems and Bioclimatology, Mendel University in Brno, Brno, Czech Republic

(10)University of Milan, Milano, Italy

(11)Agricultural and Biological Engineering, University of Florida, Gainesville, FL

(12)Institute of Agrophysics, Polish Academy of Sciences, Lublin, Poland

(13)Michigan State University, Michigan State University, East Lansing, MI

(14)Environmental Sciences, Lund University, Lund, Sweden

(15)INRA, Avignon, France

(16)Invergowrie, James Hutton Institute, Dundee, Scotland

(17)Agriculture, University of Sassari, Sassari, Italy

(18)Dpt AgroBioChem and BioSE, ULg - Gembloux Agro-Bio Tech, Gembloux, Belgium

(19)Geological Sciences, Michigan State University, East Lansing, MI

(20)Agro, Université de Liège, Liège, Belgium

(21)Leibniz Centre for Agricultural Landscape Research, Muencheberg, Germany

(22)IFAPA, IFAPA Junta de Andalucia, Córdoba, Spain

(23)Université de Liège, Liège, Belgium

(24)Agriculture, University of Copenhagen, Taastrup, Denmark

(25)CNR-IBIMET, University of Florence, Florence, Poland

(26)Potsdam Institute for Climate Impact Research, Potsdam, Germany

(27)ZALF - Leibniz Centre for Agricultural Landscape Research, Muencheberg, Germany

(28)Aarhus University, Tjele, (Non U.S.), DENMARK

(29)Agricultura, University of Milan, Milan, Italy

(30)University of Milan, Milan, Italy

(31)Rothamsted Research, Harpenden, England

(32)Polish Academy of Sciences, Lublin, Poland

(33)Wageningen University, Wageningen, Netherlands

(34)NASA Goddard Institute for Space Studies, New York, NY

(35)UAPV, UMR1114 EMMAH, Avignon, France

(36)CSIRO, Canberra, ACT 2601, Australia

(37)Rothamsted Research, Okehampton, United Kingdom

(38)Agriculture, China Agricultural University, Beijing, China

We studied the sensitivity of a 26-member ensemble of process-based wheat models to perturbations in baseline temperature and precipitation along a transect in Europe to construct impact response surfaces (IRs) of simulated wheat yields.

Resultant IRSs show the simulated yield response to systematic changes to the 1981-2010 baseline in temperature (ranging from -2 to +9°C) and precipitation (-50 to +50%). IRSs were calculated for spring and winter wheat cultivated at four contrasting sites: in southwestern Finland, Germany (winter wheat in the west; spring wheat in the east) and north-eastern Spain. Simplified assumptions were made on CO₂ level, management and soils with the aim to distinguish differences in model response attributable to climate.

Results show that simulated absolute yield levels vary considerably between models under baseline and perturbed conditions. Across the ensemble, there is general agreement among models that the dominant sensitivity changes along the transect. Wheat yields are more sensitive to temperature changes at the Finnish site, sensitive to a combination of temperature and precipitation at the German sites, and more sensitive to precipitation at the Spanish site. Yields benefit from cooling at the Spanish and German sites, while temperatures are close to optimal for the baseline in Finland. Reasons for these site-specific patterns of response can partly be attributed to differences in baseline climate, local cultivars and management practices. Standardized anomalies of simulated yield series match observed regional yield anomalies more closely for spring wheat in Germany than at other sites, mainly due to better resolved observations.

The IRS approach appears suitable for illustrating model behaviour under changing climate, as well as for comparing results from multi-model ensemble simulations.

This study was conducted as part of the impact uncertainty research of the crop modelling (CropM) component of MASUR, a project funded by national funding agencies and the EC.