

EXTENDED REPORT

Implication of *IL-2/IL-21* region in systemic sclerosis genetic susceptibility

Lina-Marcela Diaz-Gallo,¹ Carmen P Simeon,² Jasper C Broen,³ Norberto Ortego-Centeno,⁴ Lorenzo Beretta,⁵ Madelon C Vonk,³ Patricia E Carreira,⁶ Sofia Vargas,¹ José Andrés Román-Ivorra,⁷ Miguel A González-Gay,⁸ Carlos Tolosa,⁹ Francisco Javier López-Longo,¹⁰ Gerard Espinosa,¹¹ Esther F Vicente,¹² Roger Hesselstrand,¹³ Gabriela Riemekasten,¹⁴ Torsten Witte,¹⁵ Jörg H W Distler,¹⁶ Alexandre E Voskuyl,¹⁷ Annemie J Schuerwegh,¹⁸ Paul G Shiels,¹⁹ Annika Nordin,²⁰ Leonid Padyukov,²⁰ Anna-Maria Hoffmann-Vold,²¹ Raffaella Scorza,⁵ Claudio Lunardi,²² Paolo Airo,²³ Jacob M van Laar,²⁴ Nicolas Hunzelmann,²⁵ Birgit S Gathof,²⁶ Alexander Kreuter,²⁷ Ariane Herrick,²⁸ Jane Worthington,²⁸ Christopher P Denton,²⁹ Xiaodong Zhou,³⁰ Frank C Arnett,³⁰ Carmen Fonseca,²⁹ Bobby PC Koeleman,³¹ Shervin Assasi,³⁰ Timothy R D J Radstake,³² Maureen D Mayes,³⁰ Javier Martín,¹ The Spanish Scleroderma Group

► Additional supplementary data are published online only. To view these files please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2012-202357>).

For numbered affiliations see end of article.

Correspondence to

Lina-Marcela Diaz-Gallo, Cellular Biology and Immunology Department, Instituto de Parasitología y Biomedicina López-Neyra (IPBLN-CSIC), Parque Tecnológico Ciencias de la Salud, Avenida del Conocimiento s/n 18100-Armilla, Granada 18100, Spain; lina.diaz@ipb.csic.es

Accepted 21 October 2012
Published Online First
21 November 2012

ABSTRACT

Objective The interleukin 2 (*IL-2*) and interleukin 21 (*IL-21*) locus at chromosome 4q27 has been associated with several autoimmune diseases, and both genes are related to immune system functions. The aim of this study was to evaluate the role of the *IL-2/IL-21* locus in systemic sclerosis (SSc).

Patients and methods The case control study included 4493 SSc Caucasian patients and 5856 healthy controls from eight Caucasian populations (Spain, Germany, The Netherlands, USA, Italy, Sweden, UK and Norway). Four single nucleotide polymorphisms (rs2069762, rs6822844, rs6835457 and rs907715) were genotyped using TaqMan allelic discrimination assays.

Results We observed evidence of association of the rs6822844 and rs907715 variants with global SSc ($p_c=6.6E-4$ and $p_c=7.2E-3$, respectively). Similar statistically significant associations were observed for the limited cutaneous form of the disease. The conditional regression analysis suggested that the most likely genetic variation responsible for the association was the rs6822844 polymorphism. Consistently, the rs2069762A-rs6822844T-rs6835457G-rs907715T allelic combination showed evidence of association with SSc and limited cutaneous SSc subtype ($p_c=1.7E-03$ and $p_c=8E-4$, respectively).

Conclusions These results suggested that the *IL-2/IL-21* locus influences the genetic susceptibility to SSc. Moreover, this study provided further support for the *IL-2/IL-21* locus as a common genetic factor in autoimmune diseases.

INTRODUCTION

Interleukin 2 (*IL-2*) and interleukin 21 (*IL-21*) are equally attractive biological candidates that may influence the pathogenesis of autoimmune diseases. Both are cytokines involved in the proliferation of

T and B lymphocytes and different immunological activation pathways.¹ Moreover, the *IL-2* and *IL-21* genes cover a region of approximately 200 kb that maps in the 4q27 locus. *IL-2* has an important role in the maintenance of immune system homeostasis and self-tolerance. This cytokine has two paradoxical roles: promoting T cell proliferation and terminating T cell responses. Moreover, *IL-2* facilitates the production of immunoglobulins through B cells and induces the differentiation and proliferation of natural killer cells.^{1–2} *IL-21* is a potent immunomodulatory cytokine with pleiotropic effects on both innate and adaptive immune responses. These actions include the following positive effects: enhanced proliferation of lymphoid cells, increased cytotoxicity of CD8 T cells and natural killer cells, and differentiation of B cells into plasma cells. *IL-21* is also produced by T helper 17 (Th17) cells and is a critical regulator of Th17 development.^{1–3} Genetic association studies have demonstrated that several *IL-2/IL-21* polymorphisms influence the risk for autoimmune diseases (AIDs). The first evidence of this association was found in type 1 diabetes, Graves' disease, coeliac diseases and rheumatoid arthritis.^{4–7} These results have been confirmed through replication studies in different populations and extended to other autoimmune diseases, such as inflammatory bowel diseases, giant cell arthritis, psoriasis and systemic lupus erythematosus (SLE).^{8–17}

Systemic sclerosis (SSc) is a chronic fibrotic autoimmune disease in which patients are commonly classified into the following two major subgroups that are related to the specific autoantibodies against several nuclear and/or nucleolar antigens: (i) limited cutaneous SSc (lcSSc), which is related to the positive status of anticentromere autoantibodies (ACA) and (ii) diffuse cutaneous (dcSSc), which is related to the positive status of antitopoisomerase

autoantibodies (ATA).^{18–22} More than 40 susceptibility loci to SSc have been identified during the last 10 years. Half of these variants need to be replicated in different populations and many of these variants are shared among different AIDs, especially SLE.^{22–25} In this regard, one single nucleotide polymorphism (SNP) of the *IL-2* gene was proposed as risk factor to lcSSc subtype,²⁶ but this association has not been confirmed by other studies. Moreover the *IL-21* gene has been implicated as a potential driver of AIDs and recently a fine-mapping in SLE demonstrated that variants of the *IL-2/IL-21* region are implicated in the genetic susceptibility to SLE.^{12 16} Thus, the aim of this study was to evaluate the influence of the *IL-2/IL-21* region in SSc genetic susceptibility.

PATIENTS AND METHODS

Subjects

This case-control association study was comprised of 4493 SSc patients and 5896 controls of Caucasian ancestry. The discover cohort included the Spanish group, which consisted of 1176 SSc patients and 1721 healthy controls. The follow-up phase consisted of the following subjects: 609 SSc cases and 426 controls from Germany, 365 SSc cases and 734 controls from the Netherlands, 916 SSc cases and 884 controls from USA, 595 SSc cases and 1107 controls from Italy, 225 SSc cases and 273 controls from Sweden, 374 SSc cases and 436 controls from the UK and 102 SSc cases and 278 controls from Norway. There was an overlapping of 1726 SSc and 2578 controls with the previous GWAS in SSc.²⁵ The patients fulfilled the 1980 American College of Rheumatology classification criteria for SSc²⁷ or the criteria proposed for early SSc.²¹ In addition, the patients were classified as having lcSSc or dcSSc as described by LeRoy *et al.*²¹ The following clinical data were collected for the ascertainment of the clinical phenotype of the SSc patients: age, gender and presence of SSc-specific autoantibodies (Ab; ACA and ATA). The control population consisted of unrelated healthy individuals recruited in the same geographical regions as the SSc patients, and they were matched by age, sex and ethnicity with the SSc patient groups. The study was approved by local ethical committees from all the participating centres. Both patients and controls were included in the study after written informed consent was obtained.

SNP Selection and genotyping

Four SNPs of the *IL-2/IL-21* region were selected for this study. The rs2069762 SNP was selected because it has been suggested to be a genetic factor of lcSSc subtype susceptibility by a study in a small Italian cohort.²⁶ SSc and SLE share some immunogenetic pathways; thus, the rs6822844, rs6835457 and rs907715 *IL-2/IL-21* polymorphisms were studied because they are the most associated variants in a recent fine-mapping of the region in SLE.¹²

DNA from the patients and the controls were extracted from peripheral white blood cells following standard procedures. The samples were genotyped for the rs2069762, rs6822844, rs6835457 and rs907715 *IL-2/IL-21* region polymorphisms using predesigned SNP genotyping assays from Applied Biosystems (Assay IDs: C_15859930_10, C_28983601_10, C_1597475_10 and C_8949748_10, respectively). TaqMan SNP genotyping was performed using a 7900HT Real-Time PCR system from Applied Biosystems following the manufacturer's suggestions (Foster City, California, USA). In all the cohorts, the genotyping success rate was greater than 95%, and randomly selected samples were genotyped twice to verify the genotyping accuracy. Ninety-nine per cent of the genotypes were identical.

Statistical analysis

The Hardy-Weinberg equilibrium was tested for all the SNPs in all the studied populations. Significance was calculated using 2×2 contingency tables and Fisher's exact test or the χ^2 test when necessary to obtain p values, ORs and 95% CIs using PLINK (V1.07) software (<http://pngu.mgh.harvard.edu/purcell/plink/>).²⁸ The p values less than 0.05 were considered to be statistically significant. The Bonferroni correction was applied to the significant p values and referred in the text as p_c ($p_{corrected}$). Cochran-Mantel-Haenszel meta-analysis was performed to control the differences among populations as implemented by the PLINK software. In addition, the Breslow-Day test (BD test) and the Higgins' test (I^2) were performed using the PLINK software in each meta-analysis. The random-effects model was checked in the significant BD P_{values} analysis. The dependency of the association between each SNP and every studied genetic variant was determined by a conditional logistic regression analysis (considering the different cohorts as covariates) using the PLINK software. Linkage disequilibrium (LD) patterns between the four studied SNPs were estimated by the expectation-maximisation algorithm using HAPLOVIEW (V4.2; Broad Institute of MIT and Harvard) and PLINK software. To evaluate the allelic combination difference between cases and controls, the conditional haplotype-based associations test was applied using the PLINK software.²⁹ The statistical power of the combined analysis was between 91% and 99% for all the SNPs, allowing for the detection of associations with an OR equal to 1.2 at a 5% significance level and the lowest minor allelic frequency, according to the Power Calculator for Genetic Studies 2006 software, which uses the methods described by Skol *et al.*³⁰

RESULTS

The cases and controls of the eight Caucasian populations were in Hardy-Weinberg equilibrium at a 5% significance level. Additionally, the minor allelic frequencies of the four studied SNPs were similar to those reported by the HapMap project for the Utah residents with ancestry from northern and western Europe (CEU) population (<http://hapmap.ncbi.nlm.nih.gov/>). The LD structure of the eight cohorts is shown in the supplemental material (see online supplementary figure S1).

First, an association study was conducted in a Spanish case-control set, and a significant association was observed between the rs907715 SNPs minor allele and the global SSc ($p_c=0.03$, OR=0.85 95% CI 0.8 to 0.9) and the lcSSc subtype ($p_c=0.04$, OR=0.83 95% CI 0.7 to 0.9). A trend of association was observed between the minor allele of the rs6822844 SNP and the global SSc ($p_{value}=0.04$, OR=0.84 95% CI 0.7 to 1) and lcSSc subtype ($p_{value}=0.04$, OR=0.79 95% CI 0.7 to 0.9). Also a trend of association was detected between the minor allele of rs6835457 and lcSSc subtype in this population ($p_{value}=0.03$, OR=0.87 95% CI 0.8 to 1). In contrast, no association was observed with the rs2069762 SNP ($p_{value}=0.8$ for both SSc and lcSSc) (see online supplementary tables S1–S3). Based on these observations, we decided to evaluate other Caucasian cohorts and to perform a meta-analysis.

Table 1 shows the meta-analysis results for the *IL-2/IL-21* SNPs, the global SSc, the main SSc subtypes, the ACA and the ATA antibodies positive status. The combined analysis showed that the minor allele frequencies of the rs6822844 and rs907715 SNPs were significantly higher in controls than in SSc ($p_c=6.6E-04$ OR=0.86 95% CI 0.79 to 0.93 and $p_c=7.2E-3$ OR=0.91 95% CI 0.85 to 0.96, respectively)

Table 1 Genotype and minor allele frequencies of meta-analysis of four *IL-2/IL-21* SNPs located in SSc patients and healthy controls from European and US populations

SNP	1/2	Subgroup (N)	Genotype, N (%)			MAF (%)	Allele test		
			1/1	1/2	2/2		p Value*	p _c †	OR (CI 95%)‡
rs2069762	C/A	Controls (n=5482)	510 (9.30)	2266 (41.34)	2706 (49.36)	29.97			
		SSc (n=4281)	429 (10.02)	1778 (41.53)	2074 (48.45)	30.79	0.08	NA	1.06 (0.99 to 1.13)
		lcSSc (n=2897)	295 (10.18)	1203 (41.53)	1399 (48.29)	30.95	0.09	NA	1.06 (0.99 to 1.14)
		dcSSc (n=1384)	134 (9.68)	575 (41.55)	675 (48.77)	30.46	0.31	NA	1.05 (0.96 to 1.15)
		ACA+ (n=1736)	170 (9.79)	730 (42.05)	836 (48.16)	30.82	0.25	NA	1.05 (0.97 to 1.14)
rs6822844	T/G	ATA+ (n=1031)	94 (9.12)	428 (41.51)	509 (49.37)	29.87	0.98	NA	1.00 (0.90 to 1.11)
		Controls (n=5792)	149 (2.57)	1475 (25.47)	4168 (71.96)	15.31			
		SSc (n=4407)**	98 (2.22)	996 (22.60)	3313 (75.18)	13.52	1.7E-04	6.6E-04	0.86 (0.79 to 0.93)
		lcSSc (n=2977)***	67 (2.25)	659 (22.14)	2251 (75.61)	13.32	1.5E-04	6.0E-04	0.84 (0.76 to 0.92)
		dcSSc (n=1430)	31 (2.17)	337 (23.57)	1062 (74.27)	13.95	0.06	NA	0.89 (0.79 to 1)
rs6835457	G/A	ACA+ (n=1763)	38 (2.16)	395 (22.40)	1330 (75.44)	13.36	0.01	0.06	0.87 (0.78 to 0.97)
		ATA+ (n=1074)	29 (2.70)	257 (23.93)	788 (73.37)	14.66	0.67	NA	0.97 (0.85 to 1.11)
		Controls (n=5720)	668 (11.68)	2507 (43.83)	2545 (44.49)	33.59			
		SSc (n=4392)****	445 (10.13)	1908 (43.44)	2039 (46.43)	31.85	0.013	0.05	0.93 (0.87 to 0.98)
		lcSSc (n=2965)*****	312 (10.52)	1255 (42.33)	1398 (47.15)	31.69	0.014	0.06	0.92 (0.86 to 0.98)
rs907715	T/C	dcSSc (n=1427)	133 (9.32)	653 (45.76)	641 (44.92)	32.20	0.28	NA	0.95 (0.87 to 1.04)
		ACA+ (n=1765)	186 (10.54)	756 (42.83)	823 (46.63)	31.95	0.12	NA	0.94 (0.86 to 1.02)
		ATA+ (n=1064)	113 (10.62)	481 (45.21)	470 (44.17)	33.22	0.99	NA	1.00 (0.90 to 1.10)
		Controls (n=5644)	670 (11.87)	2491 (44.14)	2483 (43.99)	33.94			
		SSc (n=4341)*****	437 (10.07)	1883 (43.38)	2021 (46.56)	31.76	1.8E-03	7.2E-03	0.91 (0.85 to 0.96)
rs907715	T/C	lcSSc (n=2929)*****	307 (10.48)	1236 (42.20)	1386 (47.32)	31.58	2.7E-03	0.01	0.90 (0.84 to 0.96)
		dcSSc (n=1412)	130 (9.21)	647 (45.82)	635 (44.97)	32.12	0.14	NA	0.93 (0.85 to 1.02)
		ACA+ (n=1744)	180 (10.32)	754 (43.23)	810 (46.44)	31.94	0.05	NA	0.92 (0.85 to 1)
		ATA+ (n=1056)	109 (10.32)	475 (44.98)	472 (44.70)	32.81	0.48	NA	0.96 (0.87 to 1.07)

*All p values have been calculated for the allelic model.

**Breslow-Day p_{value}=0.29. Higgins' test (I²)=17.3%. Random-effects model p_{value}=8.8E-04 p_c=3.5E-3 Random-effects OR=0.86.

***Breslow-Day p_{value}=0.16. I²=33.9%. Random-effects model p_{value}=4.1E-03. Random-effects OR=0.84.

****Breslow-Day p_{value}=0.06. I²=48.6%. Random-effects model p_{value}=0.1. Random-effects OR estimate=0.93.

*****Breslow-Day p_{value}=0.09. I²=43.4%. Random-effects model p_{value}=0.11. Random-effects OR estimate=0.92.

*****Breslow-Day p_{value}=0.02. I²=58%. Random-effects model p_{value}=0.08. Random-effects OR estimate=0.91.

*****Breslow-Day p_{value}=0.09. I²=43.7%. Random-effects model p_{value}=0.05. Random-effects OR estimate=0.91.

†If it is applicable, Bonferroni correction is shown.

‡OR for the minor allele.

ACA, anticentromere autoantibodies; ATA, antitopoisomerase autoantibodies; dcSSc, diffuse cutaneous SSc; NA, not applicable; SNP, single nucleotide polymorphisms; SSc, systemic sclerosis.

and lcSSc patients (p_c=6E-4 OR=0.84 95% CI 0.76 to 0.92 and p_c=0.01 OR=0.9 95% CI 0.84 to 0.96, respectively). A trend of association was observed in the meta-analysis for the rs6822844 and rs6835457 variants and ACA positive

status (p_{value}=0.01 OR=0.87 95% CI 0.78 to 0.97 and p_{value}=0.05 OR=0.92 95% CI 0.85 to 1, respectively). The rs6835457 SNP also had a trend of association with global SSc and lcSSc (p_{value}=0.01 OR=0.93 95% CI 0.87 to 0.98 and

Table 2 Conditional logistic regression analysis for the *IL-2/IL-21* SNPs located in SSc considering the eight European and US populations as covariate

Group of analysis	SNP	MAF Cases	MAF Controls	p value of each SNP conditioned by rs6822844	p value of rs6822844 conditioned by each SNP	r ² with rs6822844							
						Spain	Germany	The Netherlands	USA	Italy	Sweden	UK	Norway
SSc	rs2069762	0.31	0.30	0.69	1.30E-03	0.06	0.06	0.09	0.07	0.07	0.11	0.06	0.09
	rs6835457	0.32	0.34	0.43	0.024	0.25	0.37	0.38	0.36	0.28	0.48	0.37	0.49
	rs907715	0.32	0.34	0.19	0.026	0.26	0.37	0.39	0.36	0.29	0.39	0.37	0.49
lcSSc	rs2069762	0.31	0.30	0.69	9.07E-04	–	–	–	–	–	–	–	–
	rs6835457	0.32	0.34	0.53	0.014	–	–	–	–	–	–	–	–
	rs907715	0.32	0.34	0.3	0.015	–	–	–	–	–	–	–	–
ACA+	rs2069762	0.31	0.30	0.64	0.015	–	–	–	–	–	–	–	–
	rs6835457	0.32	0.34	0.81	0.061	–	–	–	–	–	–	–	–
	rs907715	0.32	0.34	0.56	0.063	–	–	–	–	–	–	–	–

ACA, anticentromere autoantibodies; lcSSc, limited cutaneous SSc; MAF, minor allelic frequencies; SNP, single nucleotide polymorphisms; SSc, systemic sclerosis.

Basic and translational research

$p_{\text{value}}=0.01$ OR=0.92 95% CI 0.86 to 0.98, respectively). We did not detect any significant association between the rs6835457 or rs2069762 SNPs and the global SSc diseases or its different phenotypes (for detailed information see supplementary tables S1 through S3). It is worth noting that the minor I^2 percentage was observed in the meta-analysis for the rs6822844 SNP with SSc (17.3%) and lcSSc (33.9%), suggesting that the variation between the populations is moderate. Moreover, these analyses were the only ones that remained significant in the random-effect model (rs6822844 and SSc $p_c=3.5E-3$, rs6822844 and lcSSc $p_c=0.016$).

A conditional logistic regression analysis was used to identify which SNP could be the causal SNP for the observed associations between the studied polymorphisms. The association of each SNP was evaluated using the populations as covariates, and the association was conditioned to the rs6822844 SNP because the lowest p_{value} and strongest effect (OR) were observed in this locus. Pairwise conditional analysis showed that the association of the rs907715 SNP was explained by the rs6822844 effect, because only the latter SNP remained significant after conditioned to each other (rs907715 conditioned $p_{\text{value}}=0.19$; rs6822844 conditioned $p_{\text{value}}=0.026$). Moreover, the rs2069762 and rs6835457 SNPs exhibited significance only when conditioned to the rs6822844 SNP. These results suggested that the rs6822844 signal could explain the association observed in the *IL-2/IL-21* locus (table 2).

Finally, the results of the conditional haplotype-based association testing are shown in table 3. The allelic combination formed by the rs2069762 major allele and the rs6822844, rs6835457 and rs907715 minor alleles was significantly increased in the controls compared with the global SSc ($p_c=1.7E-3$, OR=0.89 95% CI 0.81 to 0.98), the lcSSc subtype ($p_c=8E-4$, OR=0.86 95% CI 0.77 to 0.96) and the ACA positive status ($p_c=2.7E-2$, OR=0.86 95% CI 0.75 to 0.98). Interestingly, the OR observed for this analysis was not different from the one observed in the allelic test. Moreover, the significant effect of the omnibus analyses for SSc, lcSSc and ACA positive status disappeared when they were controlled by the rs6822844 SNP (p_{values} of the likelihood ratio test were: $p_{\text{value}}=0.66$ for global SSc, $p_{\text{value}}=0.74$ for lcSSc and $p_{\text{value}}=0.93$ for ACA+).

DISCUSSION

Our study suggests for the first time the influence of the rs6822844 polymorphism of the *IL-2/IL-21* region in susceptibility to SSc. This variant also influences the lcSSc subtype of

the diseases and probably the ACA positive status due to the trend of association observed between the rs6822844 polymorphism and this phenotype. Although, our study had sufficient statistical power for both dcSSc and ATA analysis (95% and 91%, respectively), we observed that there were not significant associations between the four *IL-2/IL-21* SNPs and dcSSc or ATA positive status. The ORs exhibited the same direction as the significant associations with SSc and lcSSc, suggesting that an increment in the sample size with future studies could show a significant relation between the *IL-2/IL-21* SNPs and dcSSc or ATA. Interestingly, the rs6822844 variant was associated in the same OR direction as that observed in SLE. The minor allele of this variant is more frequent in healthy donors than in SSc patients, lcSSc subtype subjects and SLE patients.^{12 16} The logistic regression and the allelic combination analyses support that the rs6822844 SNP association was responsible for the observed associations. The rs2069762A-rs6822844T-rs6835457G-rs907715T allelic combination was associated as a protective factor to SSc, lcSSc subtype and ACA positive status, which is the same effect observed for the T allele of the rs6822844. Importantly, the ORs observed for this allelic combination were not different from the ORs observed for the rs6822844 SNP analysis. These observations were slightly different from the results of the SLE study performed by Hughes *et al*¹² where the observed association between *IL-2/IL-21* region and SLE could be explained by the rs6835457 and rs907715 SNPs. Together, these results support the idea that the common genetic factors in autoimmune diseases may be associated at a regional level but differ in the specific SNPs associated with each disease, including the magnitude and direction of the association.^{31 32} Although, the logistic regression test and the allelic combination analyses conditioned by the rs6822844 SNP suggest that this variant is responsible for the association observed in the region; we cannot totally discard a slight role of the rs6835457 and rs907715 polymorphisms in SSc due to the moderate LD between them and the rs6822844 SNP.

The rs6822844 and rs6835457 SNPs are located in the flanking 3'-untranslated region of *IL-21*, and the rs907715 polymorphism is located in intron 3 of the *IL-21* gene. In contrast, the rs2069762 SNP is located in the flanking 5'-untranslated region of *IL-2*, which did not exhibit significant association with SSc, the subtypes of the disease or the antibodies' status. The rs2069762 minor allele has been previously associated with the lcSSc subtype.²⁶ Our study has a considerably larger sample size than the previous study; therefore the previously reported

Table 3 Conditional haplotype-based association analysis of four *IL-2/IL-21* SNPs located according to diseases, lcSSc diseases subtype and ACA status and considering the eight European and US populations as covariate

Allelic combination†	Frequency				Frequency				Frequency				
	Controls	SSc	OR (CI 95%)	P*	Pc‡	lcSSc	OR (CI 95%)	P*	Pc‡	ACA+	OR (CI 95%)	P*	Pc‡
AGGT	0.182	0.181	-ref-	---	-	0.181	-ref-	---	-	0.183	-ref-	----	-
ATGT	0.152	0.133	0.89 (0.81 to 0.98)	4.12E-04	1.65E-03	0.131	0.86 (0.77 to 0.96)	2.01E-04	8.04E-04	0.132	0.86 (0.75 to 0.98)	6.81E-03	2.72E-02
CGAC	0.298	0.306	1.04 (0.96 to 1.12)	0.18	NA	0.308	1.03 (0.94 to 1.13)	0.2	NA	0.308	1.02 (0.92 to 1.14)	0.27	NA
AGAC	0.369	0.379	1.04 (0.96 to 1.13)	0.1	NA	0.38	1.03 (0.94 to 1.13)	0.12	NA	0.377	1.02 (0.91 to 1.13)	0.4	NA

†The order of the SNPs is rs2069762, rs6822844, rs6835457, rs907715.

* p_{value} of the likelihood ratio test. Based on WHAP method.²⁹

‡If it is applicable, Bonferroni correction is shown. Not applicable (NA).

***Omnibus test $X^2=14.5$ (df=3); $p_{\text{value}}=2.35E-03$; $P_c=9.4E-03$.

****Omnibus test $X^2=14.5$ (df=3); $p_{\text{value}}=2.32E-03$; $P_c=9.28E-03$.

*****Omnibus test $X^2=8.92$ (df=3); $p_{\text{value}}=0.03$; $P_c=0.12$.

ACA, anticentromere autoantibodies; lcSSc, limited cutaneous SSc; SNP, single nucleotide polymorphisms; SSc, systemic sclerosis.

significant association for rs2069762 might stem from type 1 statistical error. This fact together with the location of the associated SNP suggests a highlighted role of the *IL-21* cytokine. By examining the expression and regulation of *IL-21* and the *IL-21* receptor (*IL-21R*) in patients with SSc, a previous study demonstrated an upregulation of *IL-21R* in epidermis samples.³³ However, a recent study has demonstrated that the scleroderma burden in allogeneic haemopoietic stem cell transplantations is driven by Th17 induction via *IL-21* and *IL-23* signalling.³⁴ Together, these results suggest that *IL-21/IL-21R* signalling has a pathogenic function in SSc.

The role of *IL-2* and *IL-21* in the immune system makes these genes plausible candidates for the genetic component of autoimmune diseases.^{1 35 36} Our results increase the evidence that have showed that the rs6822844 is significantly associated with multiple autoimmune diseases.¹⁰⁻¹³ According to the HapMap project for the CEU population (<http://hapmap.ncbi.nlm.nih.gov/>), the rs682284 polymorphism tags seven other variants located along the *IL-2/IL-21* region (rs13132245, rs13122573, rs4459999, rs13151961, rs13140464, rs6814280 and rs2069778), but clear evidence that connects any of these variants with the *IL-2* and/or *IL-21* regulation is lacking. Together, all point out these genetic variants as good candidates for functional studies in SSc pathogenesis and in other autoimmune diseases.

Although, the combined analyses of the rs6822844 polymorphism did not show heterogeneity (BD $p_{\text{value}}=0.29$, $I^2=17.3\%$) between the eight European populations, a weak point of our study is that we did not have enough data available to control the association by principal component. Furthermore, as we mentioned before, an increment in the sample size for the stratified analysis could define in an accurate way the role of the studied variants in different clinical manifestations of SSc as their influence in the presence of coautoimmunity. Consequently, it is necessary to replicate the actual observation.

To conclude, consistent with previous studies on autoimmune diseases, the *IL-2/IL-21* region is a susceptibility genetic factor for SSc and its lcSSc subtype. The rs6822844 polymorphism confers the best association signal for SSc. It is also worth mentioning that this study shows the importance of the study of different populations and broad collaboration to find the missing heritability for relatively rare diseases like SSc.

Author affiliations

¹Cellular Biology and Immunology Department, Instituto de Parasitología y Biomedicina López-Neyra, IPBLN-CSIC, Granada, Spain

²Department of Internal Medicine, Hospital Vall d'Hebron, Barcelona, Spain

³Department of Rheumatology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

⁴Department of Internal Medicine, Hospital Clínico Universitario San Cecilio, Granada, Spain

⁵Department of Allergy, Clinical Immunology and Rheumatology, IRCCS Fondazione Policlinico-Mangiagalli-Regina Elena & University of Milan, Milan, Italy

⁶Department of Rheumatology, Hospital Universitario 12 de Octubre, Madrid, Spain

⁷Department of Rheumatology, Hospital Universitario y Politécnico La Fe, Valencia, Spain

⁸Department of Rheumatology, Hospital Marqués de Valdecilla, IFIMAV, Santander, Spain

⁹Department of Internal Medicine, Hospital Parc Taulí, Sabadell, Spain

¹⁰Department of Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid, Spain

¹¹Department of Systemic Autoimmune Diseases, Hospital Clinic de Barcelona, Barcelona, Spain

¹²Department of Rheumatology, Hospital Universitario La Princesa, Madrid, Spain

¹³Department of Rheumatology, Lund University Hospital, Lund, Sweden

¹⁴Department of Rheumatology and Clinical Immunology, Charité Universitätsmedizin and German Rheumatism Research Centre, a Leibniz institute, Berlin, Germany

¹⁵Department of Clinical Immunology and Rheumatology, Hannover Medical School, Hannover, Germany

¹⁶Department of Internal Medicine 3, Institute for Clinical Immunology, University of Erlangen-Nuremberg, Erlangen, Germany

¹⁷Department of Rheumatology, VU University Medical Center, Amsterdam, The Netherlands

¹⁸Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands

¹⁹Institute of Cancer Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, Scotland

²⁰Rheumatology Unit, Department of Medicine, Karolinska Institutet, Stockholm, Sweden

²¹Department of Rheumatology, Rikshospitalet, Oslo University Hospital, Oslo, Norway

²²Department of Medicine, Università degli Studi di Verona, Verona, Italy

²³UO Reumatologia ed Immunologia Clinica, Spedali Civili, Brescia, Italy

²⁴Musculoskeletal Research Group, Institute of Cellular Medicine, Newcastle University, Newcastle, UK

²⁵Department of Dermatology, University of Cologne, Cologne, Germany

²⁶Division of Transfusion Medicine, University Hospital of Cologne, Cologne, Germany

²⁷Department of Dermatology, Allergology, and Venereology, Ruhr University of Bochum, Bochum, Germany

²⁸Arthritis Research UK Epidemiology Unit, The University of Manchester, Manchester Academic Health Science Centre, Manchester, UK

²⁹Centre for Rheumatology, Royal Free and University College Medical School, London, UK

³⁰Department of Rheumatology and Clinical Immunogenetics, The University of Texas Health Science Center at Houston, Houston, Texas, USA

³¹Section Complex Genetics, Department of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands

³²Department of Rheumatology & Clinical Immunology, University Medical Center Utrecht, Utrecht, The Netherlands

Acknowledgements We thank Sonia García and Gema Robledo for their excellent technical assistance and all the patients and control donors for their essential collaboration. We thank Banco Nacional de ADN (University of Salamanca, Spain). We are also thankful to EUSTAR (The EULAR Scleroderma Trials and Research group) and the German Network of Systemic Sclerosis for the facilitation of this project.

Spanish Scleroderma Group Jose Luis Callejas and Raquel Ríos, Unidad de Enfermedades Sistémicas Autoinmunes, Servicio de Medicina Interna, Hospital Clínico Universitario San Cecilio, Granada; Nuria Navarrete, Servicio de Medicina Interna, Hospital Virgen de las Nieves, Granada; Rosa García Portales, Servicio de Reumatología, Hospital Virgen de la Victoria, Málaga; María Teresa Camps, Servicio de Medicina Interna, Hospital Carlos Haya, Málaga; Antonio Fernández-Nebro, Servicio de Reumatología, Hospital Carlos Haya, Málaga; María F. González-Escribano, Servicio de Inmunología, Hospital Virgen del Rocío, Sevilla; Julio Sánchez-Román, Francisco J. García-Hernández and M^a Jesús Castillo, Servicio de Medicina Interna, Hospital Virgen del Rocío, Sevilla; M^a Ángeles Aguirre and Inmaculada Gómez-Gracia, Servicio de Reumatología, Hospital Reina Sofía, Córdoba; Benjamín Fernández-Gutiérrez and Luis Rodríguez-Rodríguez, Servicio de Reumatología, Hospital Clínico San Carlos, Madrid; José Luis Andreu, Servicio de Reumatología, Hospital Puerta del Hierro, Madrid; Paloma García de la Peña, Servicio de Reumatología, Hospital Madrid Norte Sancharinaro, Madrid; Lina Martínez, Servicio de Reumatología, Hospital General Universitario Gregorio Marañón, Madrid; María Ángeles Robles, Natividad Oreiro, Servicio de Reumatología, INIBIC-Hospital Universitario A Coruña, La Coruña; Vicente Fonollosa, Servicio de Medicina Interna, Hospital Valle de Hebrón, Barcelona; Anna Pros, Servicio de Reumatología, Hospital Del Mar, Barcelona; Mónica Rodríguez Carballeira, Servicio de Medicina Interna, Hospital Universitari Mútua Terrasa, Barcelona; Francisco Javier Narváez, Servicio de Reumatología, Hospital Universitari de Bellvitge, Barcelona; Bernardino Díaz, Luis Tripiella and María Gallego, Servicio de Medicina Interna, Hospital Central de Asturias, Oviedo; María del Carmen Freire and Inés Vaquero, Unidad de Trombosis y Vasculitis, Servicio de Medicina Interna, Hospital Xeral-Complejo Hospitalario Universitario de Vigo, Vigo; María Victoria Egurbide, Servicio de Medicina Interna, Hospital de Cruces, Barakaldo; Luis Sáez-Comet, Unidad de Enfermedades Autoinmunes Sistémicas, Servicio de Medicina Interna, Hospital Universitario Miguel Servet, Zaragoza; Federico Díaz and Vanesa Hernández, Servicio de Reumatología, Hospital Universitario de Canarias, Tenerife; Emma Beltrán, Hospital General Universitario, Valencia Spain.

Contributors All the authors listed participated in all or at list one of these activities: Conception and design, acquisition of data or analysis and interpretation of data. Drafting the article or revising it critically for important intellectual content. Final approval of the version published.

Funding This work was supported by the following grants: LMDG was funded by the 'Ayudas Predoctorales de Formación en Investigación en Salud (PFIS-FI09/00544)' from the Instituto de Salud Carlos III. JM was funded by GEN-FER from the Spanish

Basic and translational research

Society of Rheumatology, SAF2009-11110 from the Spanish Ministry of Science, CTS-4977 from Junta de Andalucía, Spain, in part by Redes Temáticas de Investigación Cooperativa Sanitaria Program, RD08/0075 (RIER) from Instituto de Salud Carlos III (ISCIII), Spain and by Fondo Europeo de Desarrollo Regional (FEDER). TRDJR was funded by the VIDÍ laureate from the Dutch Association of Research (NWO) and Dutch Arthritis Foundation (National Reumafonds). JM and TRDJR were sponsored by the Orphan Disease Program grant from the European League Against Rheumatism (EULAR). BPCK is supported by the Dutch Diabetes Research Foundation (grant 2008.40.001) and the Dutch Arthritis Foundation (Reumafonds, grant NR 09-1-408). TW was granted by DFG WI 1031/6.1 and DFG KFO 250 TP03. NOC was funded by PI-0590-2010, Consejería de Salud, Junta de Andalucía, Spain. US National Institutes of Health and National Institute of Arthritis and Musculoskeletal Diseases (NIH-NIAMS) R01-AR-055258, Two-Stage Genome Wide Association Study in Systemic Sclerosis and by the NIH-NIAMS Center of Research Translation (CORT) in SSc (P50AR054144), K23AR061436, and the Department of Defense Congressionally Directed Medical Research Programs (W81XWH-07-01-0111), (NIH/NIAD) 1U01AI09090.

Competing interests None.

Ethics approval The local ethical committees in accordance with the tenets of the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- McGuire HM, Vogelzang A, Hill N, *et al*. Loss of parity between IL-2 and IL-21 in the NOD Idd3 locus. *Proc Natl Acad Sci USA* 2009;**106**:19438–43.
- Yamanouchi J, Rainbow D, Serra P, *et al*. Interleukin-2 gene variation impairs regulatory T cell function and causes autoimmunity. *Nat Genet* 2007;**39**:329–37.
- Bubier JA, Sproule TJ, Foreman O, *et al*. A critical role for IL-21 receptor signaling in the pathogenesis of systemic lupus erythematosus in BXSB-Yaa mice. *Proc Natl Acad Sci USA* 2009;**106**:1518–23.
- Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;**447**:661–78.
- Todd JA, Walker NM, Cooper JD, *et al*. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat Genet* 2007;**39**:857–64.
- van Heel DA, Franke L, Hunt KA, *et al*. A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. *Nat Genet* 2007;**39**:827–9.
- Zhernakova A, Alizadeh BZ, Bevova M, *et al*. Novel association in chromosome 4q27 region with rheumatoid arthritis and confirmation of type 1 diabetes point to a general risk locus for autoimmune diseases. *Am J Hum Genet* 2007;**81**:1284–8.
- Barton A, Eyre S, Ke X, *et al*. Identification of AF4/FMR2 family, member 3 (AFF3) as a novel rheumatoid arthritis susceptibility locus and confirmation of two further pan-autoimmune susceptibility genes. *Hum Mol Genet* 2009;**18**:2518–22.
- Festen EA, Goyette P, Scott R, *et al*. Genetic variants in the region harbouring IL2/IL21 associated with ulcerative colitis. *Gut* 2009;**58**:799–804.
- Hollis-Moffatt JE, Chen-Xu M, Topless R, *et al*. Only one independent genetic association with rheumatoid arthritis within the KIAA1109-TENR-IL2-IL21 locus in Caucasian sample sets: confirmation of association of rs6822844 with rheumatoid arthritis at a genome-wide level of significance. *Arthritis Res Ther* 2010;**12**:R116.
- Hollis-Moffatt JE, Geary RB, Barclay ML, *et al*. Consolidation of evidence for association of the KIAA1109-TENR-IL2-IL21 rs6822844 variant with Crohn's disease. *Am J Gastroenterol* 2010;**105**:1204–5.
- Hughes T, Kim-Howard X, Kelly JA, *et al*. Fine-mapping and transethnic genotyping establish IL2/IL21 genetic association with lupus and localize this genetic effect to IL21. *Arthritis Rheum* 2011;**63**:1689–97.
- Maiti AK, Kim-Howard X, Viswanathan P, *et al*. Confirmation of an association between rs6822844 at the IL2-IL21 region and multiple autoimmune diseases: evidence of a general susceptibility locus. *Arthritis Rheum* 2010;**62**:323–9.
- Marquez A, Orozco G, Martinez A, *et al*. Novel association of the interleukin 2-interleukin 21 region with inflammatory bowel disease. *Am J Gastroenterol* 2009;**104**:1968–75.
- Rodriguez-Rodriguez L, Castaneda S, Vazquez-Rodriguez TR, *et al*. Role of the rs6822844 gene polymorphism at the IL2-IL21 region in biopsy-proven giant cell arteritis. *Clin Exp Rheumatol* 2011;**29**(1 Suppl 64):S12–16.
- Sawalha AH, Kaufman KM, Kelly JA, *et al*. Genetic association of interleukin-21 polymorphisms with systemic lupus erythematosus. *Ann Rheum Dis* 2008;**67**:458–61.
- Warren RB, Smith RL, Flynn E, *et al*. A systematic investigation of confirmed autoimmune loci in early-onset psoriasis reveals an association with IL2/IL21. *Br J Dermatol* 2011;**164**:660–4.
- Gabrielli A, Avvedimento EV, Krieg T. Scleroderma. *N Engl J Med* 2009;**360**:1989–2003.
- Katsumoto TR, Whitfield ML, Connolly MK. The pathogenesis of systemic sclerosis. *Annu Rev Pathol* 2011;**6**:509–37.
- Koenig M, Dieude M, Senecal JL. Predictive value of antinuclear autoantibodies: the lessons of the systemic sclerosis autoantibodies. *Autoimmun Rev* 2008;**7**:588–93.
- LeRoy EC, Black C, Fleischmajer R, *et al*. Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol* 1988;**15**:202–5.
- Martin JE, Bossini-Castillo L, Martin J. Unraveling the genetic component of systemic sclerosis. *Hum Genet* 2012;**131**:1023–37.
- Assassi S, Mayes MD, Arnett FC, *et al*. Systemic sclerosis and lupus: points in an interferon-mediated continuum. *Arthritis Rheum* 2010;**62**:589–98.
- Carmona FD, Gutala R, Simeon CP, *et al*. Novel identification of the IRF7 region as an anticentromere autoantibody propensity locus in systemic sclerosis. *Ann Rheum Dis* 2011;**71**:114–9.
- Radstake TR, Gorlova O, Rueda B, *et al*. Genome-wide association study of systemic sclerosis identifies CD247 as a new susceptibility locus. *Nat Genet* 2010;**42**:426–9.
- Mattuzzi S, Barbi S, Carletto A, *et al*. Association of polymorphisms in the IL1B and IL2 genes with susceptibility and severity of systemic sclerosis. *J Rheumatol* 2007;**34**:997–1004.
- Preliminary criteria for the classification of systemic sclerosis (scleroderma). Subcommittee for scleroderma criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. *Arthritis Rheum* 1980;**23**:581–90.
- Purcell S, Neale B, Todd-Brown K, *et al*. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;**81**:559–75.
- Purcell S, Daly MJ, Sham PC. WHAP: haplotype-based association analysis. *Bioinformatics* 2007;**23**:255–6.
- Skol AD, Scott LJ, Abecasis GR, *et al*. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet* 2006;**38**:209–13.
- Cho JH, Gregersen PK. Genomics and the multifactorial nature of human autoimmune disease. *N Engl J Med* 2011;**365**:1612–23.
- Sirota M, Schaub MA, Batzoglu S, *et al*. Autoimmune disease classification by inverse association with SNP alleles. *PLoS Genet* 2009;**5**:e1000792.
- Distler JH, Jungel A, Kowal-Bielecka O, *et al*. Expression of interleukin-21 receptor in epidermis from patients with systemic sclerosis. *Arthritis Rheum* 2005;**52**:856–64.
- Hill GR, Olver SD, Kuns RD, *et al*. Stem cell mobilization with G-CSF induces type 17 differentiation and promotes scleroderma. *Blood* 2010;**116**:819–28.
- Crispin JC, Tsokos GC. Transcriptional regulation of IL-2 in health and autoimmunity. *Autoimmun Rev* 2009;**8**:190–5.
- Ettinger R, Kuchen S, Lipsky PE. Interleukin 21 as a target of intervention in autoimmune disease. *Ann Rheum Dis* 2008;**67**(Suppl 3):iii83–6.