
Selection of Islands for Conservation in the Urban Archipelago of Helsinki, Finland

PERTTI RANTA,*§ ANTTI TANSKANEN,† JARI NIEMELÄ,* AND ARTO KURTTO‡

*Department of Ecology and Systematics, P.O. Box 17, University of Helsinki, FIN-00014, Helsinki, Finland

†Metsätähti Ltd. Hitsajankatu 1, FIN-00810, Helsinki, Finland

‡Botanical Museum, Finnish Museum of Natural History, P.O. Box 7, University of Helsinki, FIN-00014, Helsinki, Finland

Abstract: *The occurrence of vascular plants was surveyed on 207 islands (size range 0.01–390.2 ha, number of plant species 1–449) offshore from the city of Helsinki in the Baltic Sea to examine the conservation value of these islands. We calculated a rarity score for each species (1/number of islands occupied by the species) and a biodiversity score for each island (sum of the rarity scores of each species present on the island). Positive correlations between species number and biodiversity score ($r_s = 0.97$, $p < 0.001$) and between biodiversity score and island area ($r_s = 0.87$, $p < 0.001$) indicated that these parameters are heavily dependent on island size. With the goal of including at least one occurrence (island) of all plant species, an iterative selection algorithm chose a set of 41 islands whose average size (29.3 ha) was four times the average size of all existing islands (7.0 ha). Strong nestedness ($N < 54$) explains the concentration of plant species diversity on large islands. An operational strategy for selection of sites for protection is to complement the set produced by a selection algorithm with target species not yet included (e.g., endangered species with several occurrences). Comprehensive mapping and analysis of a taxonomic group will help integrate conservation biology into land-use planning and increase the quality of the networks of protected areas.*

Selección de Islas para Conservación en el Archipiélago Urbano de Helsinki, Finlandia

Resumen: *Muestreamos la presencia de plantas vasculares en 207 islas (rango de tamaño 0.01–390.2 ha, número de plantas 1–449) en las costas de la ciudad de Helsinki, en el mar Báltico para examinar el valor de conservación de estas islas. Calculamos un índice de rareza para cada especie (1/número de islas ocupadas por la especie) y un índice de biodiversidad para cada isla (suma de los índices de rareza de cada especie presente en la isla). Correlaciones positivas entre número de especies y el índice de biodiversidad ($r_s = 0.97$, $p < 0.001$) y el índice de biodiversidad y el área de la isla ($r_s = 0.87$, $p < 0.001$) indicaron que estos parámetros son altamente dependientes del tamaño de la isla. Con la meta de incluir al menos una ocurrencia (isla) para todas las especies de plantas, un algoritmo de selección escogió 41 islas, cuyo tamaño promedio (29.3 ha) fue cuatro veces el tamaño promedio de todas las islas (7.0 ha). La fuerte anidación ($N = 54$) explica la concentración de la diversidad de especies de plantas en islas grandes. Una estrategia operacional para la selección de sitios para su protección complementaría el juego de sitios producido por la selección de algoritmo con especies blanco aún no incluidas (e.g., especies amenazadas con varias ocurrencias). El mapeo comprensivo y el análisis de un grupo taxonómico podría ayudar a integrar la biología de la conservación en la planeación de uso del suelo e incrementar la calidad de las redes de áreas protegidas.*

§email pertti.ranta@metsatahti.inet.fi, jari.niemela@helsinki.fi

Paper submitted June 8, 1998; revised manuscript accepted March 10, 1999.

Introduction

A crucial issue in planning representative networks of protected areas is how to select the sites. Several selection algorithms have been published (e.g., Nicholls & Margules 1993; Pressey et al. 1993, 1997; Underhill 1994; Williams et al. 1996; Freitag et al. 1997), but usually the selection units have been grid cells of equal size and the algorithms have been mostly applied at national or regional level. Only a few local-level applications have been offered (Game & Peterken 1984; Rapoport et al. 1986). Furthermore, selection methods have been developed and tested in floristically rich areas with many endemic species, such as southern Africa and Australia (Margules et al. 1988; Rebelo & Siegfried 1992; Lombard et al. 1997; Pressey et al. 1997).

In the species-poor northern boreal region, on the other hand, there are no globally important hotspots of endemism. For instance, the entire flora of Finland does not include a single endemic vascular plant species. Our focus is on protecting nationally or locally significant biodiversity. To achieve this there is a need to develop methods and tools for local conservation-oriented planning because municipalities, the basic land-use planning units, frequently include biodiversity conservation in their planning principles.

The purpose of our study was to establish which islands of a group of 207 in an archipelago would need to be protected to maintain the diversity of vascular plants. This represents a typical planning situation in which real land units instead of grid cells are used. The selection was based on a comprehensive database of vascular plants.

Methods

Study Area and Databases

The archipelago offshore from the city of Helsinki in the eastern Baltic consists of 315 islands (City of Helsinki Urban Facts 1998). The 207 islands included in our study (size range 0.01–390.2 ha) comprised practically all the "real" islands inside Helsinki city limits; small rocks protruding from the sea were excluded because they lack vascular plants. Most of the islands studied are uninhabited, but several large ones close to the mainland have permanent human settlement. The occurrence (presence-absence) of vascular plants on the islands was mapped by the author A.K. and his collaborators between 1990 and 1995.

Islands in the Baltic have risen from the sea since the last glaciation. Land uplift of approximately 2.5 mm/year makes new islands appear continuously and existing ones expand. The islands studied consist of Precambrian bedrock with a thin layer of topsoil. The natural vegetation on the larger islands typically consists of low Scots pine (*Pinus sylvestris*) forest with heather (*Calluna vul-*

garis), bilberry (*Vaccinium myrtillus*), and cowberry (*Vaccinium vitis-idaea*) as dominant species in the field layer. In protected bays, common alder (*Alnus glutinosa*) groves are found with more luxuriant vegetation. Small islands lack forests and are characterized by stony shore-meadows and patches of meadow vegetation in depressions and rock fissures. Bare rock surfaces are common on exposed shores on both large and small islands. The ecological features of the study area are comparable to those of other archipelagos in the Baltic (Ås et al. 1992).

Calculations

We calculated a rarity score for each plant species: $1/c_i$, $\{c_i > 0, 1 \leq i \leq n\}$, where c_i is the number of islands occupied by species i . The biodiversity score of an island was the sum of the rarity scores of all species occurring on it.

Several algorithms have been developed for reserve selection (Saetersdal et al. 1993; Lombard et al. 1995; Csuti et al. 1997; Pressey et al. 1997). Common to all these algorithms is that they start from an empty selection, adding new areas in the most effective way until the predetermined goal is reached. For instance, the goal may be to include all species at least once.

We used an inverted method to select a near-minimum set of islands that includes all vascular plant species present in the archipelago. The algorithm "most common" (MC) starts with all the islands included and eliminates one island in every cycle until the selected stop level has been reached. Initially, MC calculates the number of occurrences—the number of islands on which the species occurs—for every species (step 1) and identifies the rarest species on each island (step 2; Fig. 1). The algorithm then ranks the islands according to the number of occurrence of the rarest species. Other species on each island are ignored (step 3). At this stage, each island is represented by only one species, the rarest. Thereafter, MC finds the island with the rarest species with the highest number of occurrences (step 4). The rarest species on each island is the species with fewest occurrences. The MC then starts eliminating islands where the number of occurrences of the rarest species is highest (condition 1). This island may be interpreted as least valuable. The eliminating order follows the number of occurrences of species as the number of islands included gradually decreases (step 5). The MC recalculates the frequencies of all species on the remaining islands and again identifies the rarest species on each island.

Cyclical elimination continues until the rarest species on each of the remaining islands occurs on only one island. This means every island in the set has at least one unique species (condition 1). Setting the stop level to zero produces a numerical order of all islands. This is an




Figure 1. Flow diagram of the algorithm "most common" for reserve selection.

advantage of MC that makes it possible to compare islands according to the relative rarity of their species.

The stop level may be set to any nonnegative value. For instance, stop level one produces a set of islands that contains all species at least once. If species occur on fewer than or on a number of islands equal to the selected stop level, those islands will be automatically included in the near-minimum set. The MC also gives the numerical rank of those islands not included in the near-minimum set (Fig. 1). If the rarest species have the same number of occurrences on several islands, the island with the lowest biodiversity score will be chosen (step 4). Other elimination rules could also be applied, such as the largest island (lower cost of establishing protection) or most remote island (maximization of continuous area).

To investigate nestedness among species assemblages we used the index N (Patterson & Atmar 1986), which measures how much the observed presence-absence matrix differs from perfect nestedness ($N = 0$). The maximum value of N (i.e., minimum nestedness) depends on the size of the matrix and the number of species present.

For the analysis, we divided the islands into four size classes: 0.01–0.09 ha (28 islands), 0.1–0.99 ha (83 islands), 1.0–9.9 ha (76 islands), and >10.0 ha (20 islands). The analyses were also made without size classes.

Results

Occurrence and Nestedness of Species

We made 22,364 observations of 686 vascular plant species (hybrids were excluded) on the 207 islands (1–449 species/island). Most of the species were rare in our data; about 400 species occurred on <20 islands (<10% of islands), and the number of species occurring only on one island was 108. Rare species were clearly concentrated on a small group of islands: 11 islands hosted more than 100 rare species each. Common species were few in number, with only 20 species occurring on more than 155 islands (Fig. 2).

The island-specific biodiversity score varied from 0.1 to 40.3. Fifty islands had a biodiversity score of <1.0 (Table 1). The value of the biodiversity score depended both on the rarity of species on an island and on the island's species richness. Because one unique species has the same weight as 100 relatively common species (occurring on >100 islands), score values increased rapidly with the presence of rare species.

Occurrence of the plant species showed strong nestedness ($N = 54$). The value of the index N would be 640 if the plant species were randomly distributed among the islands, and maximum $N = 1056$. The species assemblages of the small island size classes were included within the large island size classes, and rare species occurred almost exclusively on islands falling into the largest size class. Without size classes, nestedness was weaker ($N = 46,975$), and maximum $N = 119,640$. This was due to a great number of small islands of almost the same size and only slight variation of species composition.

Selection of Minimum Sets of Islands

"MOST COMMON" ALGORITHM

Forty-one islands were needed in a minimum set of islands that included all the plant species at least once (Table 2). The set represents 20% of the 207 islands studied but as much as 83% (1200.4 ha) of the total island area. The average size of an island included in the minimum set (29.3 ha) was over four times the average island size among all 207 (7.0 ha).

Because the preservation of a single occurrence of a species does not guarantee its long-term survival, minimum sets of islands were established for 1–178 occurrences (number of islands occupied) for each species. The total island area included in the minimum set in-




Figure 2. Occurrence of vascular plant species on the islands studied. Most species are rare: over 400 species grow on only 10% or less of all islands.

Table 1. Distribution of land area, number of species, and sums of biodiversity scores among size classes of islands offshore from Helsinki.

Size class (ha)	Number of islands	Total area (ha)	Number of species	Sum of biodiversity scores
0.01-0.09	28	1.8	162	8.7
0.1-0.99	83	38.6	323	67.5
1.0-9.9	76	239.4	555	259.6
>10.0	20	1172.3	648	350.2
Total	207	1452.2	686	686.0

creases rapidly to almost 100% as the number of predetermined species occurrences increases (Fig. 3). For instance, a minimum set of islands consisting of more than 20 occupied islands for each species included practically the entire island area but only approximately 150

Table 2. The 41 islands off Helsinki included in the minimum set selected by the "most common" algorithm for reserve selection and six other islands^a selected by target species (threatened species).

Island	Island area (ha)	Number of plant species	Biodiversity score	Land-use type ^b
Pieni Leikosaari*	1.16	97	1.2	pri
Kuminapaasi	0.52	59	1.5	pub
Matalahara	2.52	84	1.8	res
Koirasaari*	3.24	136	2.4	pub
Päntäri	3.60	93	2.4	pub
Kalkkipaasi	0.29	78	2.4	pub
Harmaja*	1.84	115	1.8	mil
Korkeakupu	0.53	54	2.8	pub
Pikku Lehdeasaari	1.60	141	3.3	pri
Malkasaari	3.44	141	3.5	pub
Jänissaari*	3.90	160	3.6	mil
Tiirasaari	3.14	167	3.9	pri
Luoto (Klippan)	1.80	139	4.0	pub
Pikku Kuivasaari*	2.08	185	4.0	mil
Vallisaaren Pukkisaari	4.76	160	4.5	mil
Pihlajaluoto	3.28	185	4.7	pub
Uunisaaret	2.20	157	4.7	pub
Iso Leikosaari	5.26	192	4.8	pri
Pikku-Musta	4.00	178	5.7	pub (sf)
Kotiluoto	3.36	185	6.4	pub
Rysäkari	10.04	225	7.0	mil
Hylkysaari	4.00	223	7.3	pub
Kalkkisaari	3.33	181	7.3	pri
Länsi-Musta	8.50	240	8.7	pub
Kuivasaari	11.33	244	8.8	mil
Reposaari	2.10	206	9.0	pri
Itäinen Pihlajasaari	6.84	254	9.1	pub
Itä-Villinki	33.30	276	9.3	mil
Läntinen Pihlajasaari	19.00	267	10.3	pub
Melkki	40.50	298	10.8	mil
Lammassaari	8.70	248	11.0	pub
Pikku Niinisaari	31.88	257	11.2	pri + pub
Mustikkamaa*	37.50	304	12.8	pub
Harakka	8.80	307	14.2	res
Vasikkasaari	17.94	295	14.7	pri
Korkeasaari	25.30	265	14.8	pub
Kuninkaansaari	37.20	322	14.9	mil
Kustaanmeikka	33.50	275	15.0	pub (sf)
Palosaari	2.00	195	15.3	pri
Seurasaari	40.60	310	18.3	pub
Susisaari	33.50	310	19.4	pub (sf)
Villinki	134.50	375	21.8	pri + pub
Isosaari	73.35	355	22.0	mil
Iso Mustasaari	24.40	270	23.5	pub (sf)
Vartiosaari	89.00	352	31.7	pri + pub
Santahamina	390.20	449	38.9	mil
Vallisaari	76.70	415	40.2	mil

^aThese six other islands are marked with an asterisk.

^bLand-use type: pub, public area; res, nature reserve; pri, private area; sf, Suomenlinna sea fortress; mil, military or coast guard area.

islands. This implies that the islands not included in this minimum set were small and had few plant species.

On the other hand, the number of occurrences of many plant species remains below the selected number




Figure 3. Cumulative increase in area and number of islands at selection stop limits from 1 to 178 (minimum sets with 1–178 occurrences).

of occupied islands because they do not occur on so many islands. Consequently, the increase in the number of islands selected does not have a positive effect on the future persistence of very rare species in that a species occurring on a certain number of islands will not occur on more islands regardless of the number of islands selected for protection.

SELECTION BY TARGET SPECIES

To examine the usefulness of the most common algorithm (MC) we compared sets of islands selected by it and sets selected by other methods. A simple method of selecting sites for protection is to use target species. We made the selection using two kinds of pre-defined target species: (1) unique species (108 species with only one occurrence) and (2) nationally or provincially threatened species (14 species; Rassi et al. 1992).

Unique species were found on 37 islands. The five most unique species-rich islands had nearly half (53) of the 108 species, and they were also among the most species-rich islands. Because islands with a unique species were automatically included in the minimum set of 41, only four of the minimum set did not harbor unique species. Thus, selection using MC with stop level 1 and selecting the islands harboring unique species gave nearly the same set of islands.

Selecting islands based on the occurrence of threatened species produced a somewhat different set. The 14 threatened species occurred on 24 islands, 18 of which were the same as in the minimum set produced by MC. Threatened species were found mostly on large, species-rich islands. The combination of the two target species groups as a selection criterion produced a set of 43 islands with 6 different islands than were in the minimum set produced by MC (Table 2).

BIODIVERSITY SCORE, ISLAND SIZE, OR SPECIES NUMBER

The sum of the biodiversity scores of the minimum set of 41 islands selected using MC was 466.2, 68% of the value of the whole archipelago. Comparing the MC-produced set of 41 with a set of the same number of islands with the highest biodiversity score revealed that 32 islands are the same in each group. Thus, selection using the biodiversity score produces a set fairly similar to that using the algorithm.

Other criteria for selection might include island size and species richness. A comparison between the set of 41 islands produced by MC and the 41 largest islands revealed that 25 islands are in both groups. A similar comparison between the MC set and the 41 most species-rich islands revealed that 31 are the same in each set. In both comparisons, practically all of the 20 highest-ranked islands are also included in the minimum set produced by MC.

The observation that island size, species richness, and biodiversity score produced a set of islands similar to that produced by MC indicates the high correlation between these variables. Species number and island size correlated positively ($r_s = 0.88$, $n = 207$, $p < 0.001$), as did island size and biodiversity score ($r_s = 0.87$, $n = 207$, $p < 0.001$). Furthermore, there was a positive correlation between species number and biodiversity score ($r_s = 0.97$, $n = 207$, $p < 0.001$). These correlations indicate that the conservation value of an island increases with its size.

Discussion

Ecological Features Important for Conservation Planning

Our study demonstrates that biodiversity score, island size, and species richness reflect the conservation value

of the islands studied fairly well. But these parameters do not provide information about species composition and are therefore not useful criteria for conservation evaluation by themselves. The use of species identities in conservation planning is thus preferable.

Furthermore, the strong nestedness among vascular plants in the Helsinki archipelago implies that species-rich islands also harbor the rarer species. Nestedness among species assemblages is a common feature on islands (Patterson 1987, 1990; Blake 1991; Cutler 1991, 1994; Simberloff & Martin 1991; Wright & Reeves 1992; Atmar & Patterson 1993; Cook 1995; Kadmon 1995), but its causes are often unclear (Yiming et al. 1998). In the Helsinki archipelago, nestedness of species is to a large extent explained by the geological history of the islands. As island size increases constantly due to land uplift, pioneer communities invade the islands as soon as suitable habitat appears, and the same communities are present along the shores of the larger islands. Large islands also contain species that occur in habitats that do not exist on smaller islands. Colonization from the mainland or

other, larger islands could therefore be the main factor that has generated this nestedness (Patterson 1990; Cook & Quinn 1995).

Strategy for Conservation Selection

The "most common" algorithm used here was tested with the Finnish Bird Atlas data (250,000 observations, 3,800 grid cells of 10×10 km). The algorithm found about the same number of grid cells as the simple greedy algorithm 1 (type 1) presented by Csuti et al. (1997), but the set was different (Tanskanen 1996a, 1996b). Using plant data from the Helsinki archipelago, the progressive rarity algorithm 7 (Csuti et al. 1997) produced a set of exactly the same number of islands as did MC (41 islands), only one island being different. The sea-kale (*Crambe maritima*) occurred only on these two islands; consequently, one of them had to be included in the set.

The minimum set of 41 islands that included all the species at least once serves as a rough guide to where




Figure 4. The group of islands selected for conservation (in black) includes the 41 islands of the minimum set and 6 other islands with threatened species. The numbered islands (1-5) with dense urban settlement were not included in the study area.

the essential plant diversity exists in the Helsinki archipelago. The selection method needs several refinements, however, to produce results that can be applied in conservation planning. The first consideration is the need for filtering of species. For example, the occurrence of peanut (*Arachis hypogaea*) is clearly irrelevant for conservation purposes because it is an accidental introduction and has no viable populations on Baltic islands. Peanut and similar species should be filtered out before the analysis.

Another essential refinement is the inclusion of desirable target species. Because the minimum set may include only one occurrence per species, it does not necessarily include all the islands on which target species (e.g., threatened species) occur. A solution may be a post-selection of the islands harboring target species but not included in the minimum set. A procedure for the selection of islands or other sites for conservation could thus proceed as follows. First, irrelevant species are removed from the database. Second, the selection algorithm selects a set of islands with the required number of occurrences per species. Third, occurrences of target species are ensured by post-selection.

Following this procedure, we selected a network of 41 islands produced by MC on which each species occurred at least once. Thereafter, an additional six islands harboring endangered species not selected by the algorithm were included in the proposed network of protected islands (Fig. 4). Several of the islands found by the algorithm are protected for reasons other than their ecological value (e.g., the World Heritage site of Suomenlinna Sea Fortress). Species found on these already protected islands could be used as a starting point in seeking islands that add new species to the combination. This procedure follows a pragmatic selection strategy, combining reserve selection algorithms with other reserve-design criteria (Bedward et al. 1992).

In an archipelago off a large city with heavy recreational use such as Helsinki, a network of protection that covers one-fifth of the islands, and as much as four-fifths of the total area, would hardly be realistic. For maintaining biodiversity, however, there is no need to protect entire islands or to exclude other types of land use from the islands selected, if essential habitat patches are preserved. These patches may cover only a small fraction of the island area. For example, a habitat-based plant mapping was done on the island of Isosaari (73 ha). The island was included in the minimum network (355 species, 7 of them unique and 3 threatened), but the habitat patches supporting these species represented only about 5% of the total area of the island. If a comprehensive habitat map is not available, the list of habitat specialists with a low size range could be used for the identification of the most important habitats for protection.

Our study emphasises the need for comprehensive, high quality ecological databases for conservation evalu-

ation. Unfortunately, these are relatively rare, especially for larger areas. Furthermore, as mentioned above, there are strong arguments that each habitat patch should also be analyzed separately (Deshaye and Morisset 1988; Worthen 1996). Experienced professionals are needed to carry out the fieldwork, which may take several field seasons. This requires a substantial investment in research work, but the total cost of comprehensive mapping of a taxonomic group (e.g., vascular plants) in a city like Helsinki (land area 187 km²) is modest compared with investments needed in urban infrastructure planning in general. Consequently, the fundamental issue is recognition of the importance of the incorporation of conservation biology into the planning process and the acceptance of the associated costs.

Acknowledgments

This study was financially supported by the City of Helsinki Environment Centre and the Finnish Biodiversity Research Programme (grant 39715). We are thankful to L. Helynranta for help in the field and to J. Kotze for constructive comments on the manuscript.

Literature Cited

- Ås, S., J. Bengtsson, and T. Ebenhard. 1992. Archipelagoes and theories of insularity. Pages 201-251 in L. Hansson, editor. *Ecological principles of nature conservation*. Elsevier Applied Science, London.
- Atmar, W., and B. D. Patterson. 1993. The measure of order and disorder in the distribution of species in fragmented habitat. *Oecologia* 96:373-382.
- Bedward, M., R. L. Pressey, and D. A. Keith. 1992. A new approach for selecting fully representative reserve networks: addressing efficiency, reserve design and land suitability with an iterative analysis. *Biological Conservation* 62:115-125.
- Blake, J. G. 1991. Nested subsets and the distribution of birds on isolated woodlots. *Conservation Biology* 5:58-66.
- City of Helsinki Urban Facts. 1998. *Environmental statistics of Helsinki*. City of Helsinki, Helsinki.
- Cook, R. R. 1995. The relationship between nested subsets, habitat subdivision, and species diversity. *Oecologia* 101:204-210.
- Cook, R. R., and J. F. Quinn. 1995. The influence of colonisation in nested species subsets. *Oecologia* 102:413-424.
- Csuti, B., S. Polasky, P. H. Williams, R. L. Pressey, J. D. Camm, M. Kershaw, A. R. Kiestler, B. Downs, R. Hamilton, M. Huso, and K. Sahr. 1997. A comparison of reserve selection algorithms using data on terrestrial vertebrates in Oregon. *Biological Conservation* 80:83-97.
- Cutler, A. 1991. Nested faunas and extinction in fragmented habitats. *Conservation Biology* 5:496-505.
- Cutler, A. 1994. Nested biotas and biological conservation: metrics, mechanisms, and meaning of nestedness. *Landscape and Urban Planning* 28:73-82.
- Deshaye, J., and P. Morisset. 1988. Floristic richness, area, and habitat diversity in a hemiarctic archipelago. *Journal of Biogeography* 15:747-757.
- Freitag, S., A. S. van Jaarsveld, and H. C. Biggs. 1997. Ranking priority biodiversity areas: an iterative conservation value-based approach. *Biological Conservation* 82:263-272.
- Game, M., and G. F. Peterken. 1984. Nature reserve selection strategies

- in the woodlands of central Lincolnshire, England. *Biological Conservation* 29:157-181.
- Kadmon, R. 1995. Nested species subsets and geographic isolation: a case study. *Ecology* 76:458-465.
- Lombard, A. T., A. O. Nicholls, and P. V. August. 1995. Where should nature reserves be located in South Africa? A snake's perspective. *Conservation Biology* 9:363-372.
- Lombard, A. T., R. M. Cowling, R. L. Pressey, and P. J. Mustart. 1997. Reserve selection in a species-rich and fragmented landscape on the Agulhas Plain, South Africa. *Conservation Biology* 11:1101-1116.
- Margules, C. R., A. O. Nicholls, and R. L. Pressey. 1988. Selecting networks of reserves to maximise biological diversity. *Biological Conservation* 43:663-676.
- Nicholls, A. O., and C. R. Margules. 1993. An upgraded reserve selection algorithm. *Biological Conservation* 64:165-169.
- Patterson, B. D. 1987. The principle of nested subsets and its implications for biological conservation. *Conservation Biology* 1:323-334.
- Patterson, B. D. 1990. On the temporal development of nested subset patterns of species composition. *Oikos* 59:330-342.
- Patterson, B. D., and W. Atmar. 1986. Nested subsets and the structure of insular mammalian faunas and archipelagos. *Biological Journal of the Linnean Society* 28:65-82.
- Pressey, R. L., C. J. Humphries, C. R. Margules, R. I. Vane-Wright, and P. H. Williams. 1993. Beyond opportunism: key principles for systematic reserve selection. *Trends in Ecology and Evolution* 8:124-128.
- Pressey, R. L., H. P. Possingham, and J. R. Day. 1997. Effectiveness of alternative heuristic algorithms for identifying indicative minimum requirements for conservation reserves. *Biological Conservation* 80:207-219.
- Rapoport, E. H., G. Borioli, J. A. Monjeau, J. E. Puntieri, and R. D. Oviedo. 1986. The design of nature reserves: a simulation trial for assessing specific conservation value. *Biological Conservation* 37:269-290.
- Rassi, P., H. Kaipainen, I. Mannerkoski, and G. Ståhls, editors. 1992. Uhanalaisten eläinten ja kasvien seurantatoimikunnan mietintö. [Report on the monitoring of threatened animals and plants in Finland.] Komiteamietintö 1991:30. Ympäristöministeriö, Helsinki.
- Rebelo, A. G., and W. R. Siegfried. 1992. Where should nature reserves be located in the Cape Floristic Region, South Africa? Models for the spatial configuration of a reserve network aimed at maximizing the protection of floral diversity. *Conservation Biology* 6:243-252.
- Saetersdal, M., J. M. Line, and H. J. B. Birks. 1993. How to maximize biological diversity in nature reserve selection: vascular plants and breeding birds in deciduous woodlands, Western Norway. *Biological Conservation* 66:131-138.
- Simberloff, D., and J.-L. Martin. 1991. Nestedness of insular avifaunas: simple summary statistics masking complex species patterns. *Ornis Fennica* 68:178-192.
- Tanskanen, A. 1996a. Näkökulmia lintuatlakseen 1. Mistä bongaan Suomen pesimälinnut? *Linnut* 5:12-15.
- Tanskanen, A. 1996b. Näkökulmia lintuatlakseen 2. Säästäkää edes nämä! *Linnut* 6:31-33.
- Underhill, L. G. 1994. Optimal and suboptimal reserve selection algorithms. *Biological Conservation* 70:85-87.
- Williams, P., D. Gibbons, C. Margules, A. Rebelo, C. Humphries, and R. Pressey. 1996. A comparison of richness hotspots, rarity hotspots, and complementary areas for conserving diversity of British birds. *Conservation Biology* 10:155-174.
- Worthen, W. B. 1996. Community composition and nested-subset analyses: basic descriptors for community ecology. *Oikos* 76:417-426.
- Wright, D. H., and J. H. Reeves. 1992. On the meaning and measurement of nestedness of species assemblages. *Oecologia* 92:416-428.
- Yiming, L., J. Niemelä, and L. Dianmo. 1998. Nested distribution of amphibians in Zhoushan archipelago, China: can selective extinction cause nested subsets of species? *Oecologia* 113:557-564.

