

<https://helda.helsinki.fi>

---

## The effect of pH and potassium phosphate buffer on the toxicity of cadmium for bacteria

Korkeala, H.

BioMed Central  
1978

---

Acta Veterinaria Scandinavica. 1978. 19: 93 101.

---

<http://hdl.handle.net/1975/735>

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

From the Department of Food Hygiene, College of Veterinary  
Medicine, Helsinki, Finland.

## THE EFFECT OF pH AND POTASSIUM PHOSPHATE BUFFER ON THE TOXICITY OF CADMIUM FOR BACTERIA

By

*H. Korkeala and T. J. Pekkanen*

KORKEALA, H. and T. J. PEKKANEN: *The effect of pH and potassium phosphate buffer on the toxicity of cadmium for bacteria.* Acta vet. scand. 1978 19, 93—101. — An increase in the pH of Plate Count Agar led to increased toxicity of cadmium (Cd) for *Micrococcus luteus* ATCC 9341, *Staphylococcus aureus* ATCC 12600, *Clostridium perfringens* ATCC 8798, *Escherichia coli* ATCC 11775 and *Pseudomonas aeruginosa* ATCC 10145. The effect of the pH on toxicity was not clearly observed with *Streptococcus bovis* ATCC 9809 and was absent with *Bacillus subtilis* ATCC 8473.

When potassium phosphate buffer was added to the growth medium, the Cd toxicity for *M. luteus* and *B. subtilis* was enhanced. The toxicity of Cd for *S. bovis* decreased when potassium phosphate buffer was added to the medium. When the pH increased at differing phosphate concentrations, the sensitivity of *M. luteus* to Cd decreased.

In the authors' opinion, the effect of pH on Cd toxicity for bacteria is due to a continuously increasing negative charge towards an alkaline value by most bacteria which increases the affinity of cations towards the cell wall.

cadmium; pH; phosphate; bacteria.

Cadmium (Cd) is a potentially hazardous pollutant in the environment. Much attention has been given to Cd toxicity studies on animals (e. g. *Friberg et al.* 1974), but less attention to the effects of Cd on plants and microorganisms.

Some studies have shown that the pH and the phosphorus content of the soil influence the Cd uptake by plants. *Miller et al.* (1976) and *Williams & David* (1976) observed that extensive Cd accumulation and growth inhibition of plants were especially apparent at a low pH. Increasing available soil phosphorus was related to increased Cd accumulation (*Miller et al.*).

In the studies of *Abelson & Aldous* (1950) the toxic effects of Cd on the metabolism of bacteria were apparent. Later *Bernheim* (1973) described the effect of pH on the actions of Cd salts on *Pseudomonas aeruginosa*. *Babich & Stotzky* (1977) found that for bacteria, in contrast to plants, Cd toxicity was generally potentiated at a high pH.

The present study was undertaken to investigate the sensitivity of certain bacteria to Cd by the minimum inhibitory concentration technique and the effects of the pH and a phosphate buffer on that sensitivity.

## MATERIAL AND METHODS

### *Organisms*

The microbial strains used in the study were *Micrococcus luteus* ATCC 9341, *Staphylococcus aureus* ATCC 12600, *Bacillus subtilis* ATCC 8473, *Streptococcus bovis* ATCC 9809, *Clostridium perfringens* ATCC 8798, *Escherichia coli* ATCC 11775 and *Pseudomonas aeruginosa* ATCC 10145.

### *Chemicals and water*

$\text{KH}_2\text{PO}_4$ ,  $\text{K}_2\text{HPO}_4$  and  $\text{CdCl}_2 \cdot 2\frac{1}{2} \times \text{H}_2\text{O}$  were of pro analysis grade and obtained from J. T. Baker (Chemical Co. Deventer, Holland and Phillipsburg, N. J., USA). The water used was double distilled and deionized. Before the experiments the Cd content of the water was determined by flameless atomic absorption spectrophotometry and found to be below the lowest limit of detection (0.2 ng). The apparatus used was a Perkin-Elmer 303 atomic absorption spectrophotometer equipped with a graphite furnace and graphite cell power supply HGA 72.

### *Media and culture conditions*

The basic medium used throughout the experiments was Plate Count Agar (Difco Laboratories, Detroit, Michigan, USA). The incubation time was 48 hrs. and the incubation temperature for *M. luteus* and *B. subtilis* 30°C and for the other bacteria 35°C. All the experiments were conducted on cells from overnight cultures in Nutrient Broth (Difco). The Cd content of the Plate Count Agar and the Nutrient Broth was determined before the experiments, as above. The greatest concentrations found were 0.0006 mg of Cd/l substrate and 0.0057 mg of Cd/l substrate, re-

spectively. The studies on *C. perfringens* were carried out in anaerobic conditions (BBL GasPak jars equipped with GasPak H + CO<sub>2</sub> generator envelopes, BBL, Cockeysville, Maryland, USA) and on the other bacteria in aerobic conditions.

### *Sensitivity testing*

Minimum inhibitory concentration (M. I. C.) for Cd (used as CdCl<sub>2</sub>) was determined by a plate dilution test (Barber & Waterworth 1964) as the lowest concentration of Cd which inhibits growth completely. A barely visible haze or a single colony was disregarded. Sterile 10 ml volumes of CdCl<sub>2</sub> solutions of certain Cd concentrations were added to 90 ml of the autoclaved Plate Count Agar to give the desired Cd concentrations and the recommended amounts of water in the media. The final concentrations of Cd were 0, 0.001, 0.002, 0.004, 0.006, 0.008, 0.01, 0.02, 0.04, 0.06, 0.08, 0.1, 0.2, 0.4, 0.6, 0.8, 1, 2, 4, 6, 8, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350 and 400 mg of Cd/l medium. Three replicate plates were employed for each concentration of Cd. The inoculum from the diluted overnight cultures was spread on the agar surface with a standard 1-mm loop. The volume of one loop was used for each test. The cultures in the Nutrient Broth were first diluted with sterile isotonic saline so as to contain  $170 \times 10^4$  colony forming units (CFU)/ml of *M. luteus*,  $60 \times 10^4$  CFU/ml of *S. aureus*,  $75 \times 10^4$  CFU/ml of *B. subtilis*,  $366 \times 10^4$  CFU/ml of *S. bovis*,  $395 \times 10^4$  CFU/ml of *C. perfringens*,  $63 \times 10^4$  CFU/ml of *E. coli* or  $70 \times 10^4$  CFU/ml of *P. aeruginosa*.

### *The effect of pH of the growth medium on Cd toxicity*

The experiments were carried out by determining the M.I.C. values of Cd as described above. The pH of the Plate Count Agar was adjusted to 6.0 and 6.5 with 4 N-HCl and 7.5 and 8.0 with 4 N-NaOH. Experiments were also made at the normal pH of Plate Count Agar, which is pH 7.0.

### *The effect of phosphate buffer at different pH values on Cd toxicity*

The experiments were carried out by determining the M.I.C. values of Cd as above. Plate Count Agar was buffered with potassium phosphate buffer to pH 6.0, 7.0 and 8.0. Potassium phos-

phate was selected instead of sodium phosphate because it is known that some heavy metal compounds (mercurials) increase the swelling rate of *Pseudomonas aeruginosa* to a lesser extent in potassium phosphate than in sodium phosphate (Bernheim 1970). The phosphate concentrations used at each pH value were 0.05 M, 0.1 M and 0.2 M. The phosphate concentration of the Plate Count Agar was found to be 0.001 M, as determined by the method of Neraal & Hamm (1972). This is not taken into account when the final phosphate concentrations of the media are given. Sterile  $\text{CdCl}_2$  solution was added to the media after sterilization in order to avoid the possible precipitation of Cd with phosphate on autoclaving (Donald *et al.* 1952).

Because a precipitate was observed when  $\text{CdCl}_2$  was mixed with phosphate buffer to give Cd concentrations high enough to inhibit the growth of the strains of *S. aureus*, *C. perfringens*, *E. coli* and *P. aeruginosa* used in the sensitivity testing without added phosphate, the effect of phosphate buffer at different pH values on Cd toxicity was studied only on *M. luteus*, *B. subtilis* and *S. bovis*.

## RESULTS

The effect of the pH on the toxicity of Cd with *M. luteus*, *B. subtilis* and *S. bovis* is presented in Table 1 and with *S. aureus*, *C. perfringens*, *E. coli* and *P. aeruginosa* in Fig. 1. The results of the studies on the effects of potassium phosphate buffer and pH on the toxicity of Cd for *M. luteus*, *B. subtilis* and *S. bovis* are given in Table 1. The three replicate plates for each Cd concentration always gave the same M.I.C. value with all used media.

## DISCUSSION

When the results presented in Table 1 and Fig. 1 are examined, it is obvious that the *C. perfringens* ATCC 8798, *S. aureus* ATCC 12600, *E. coli* ATCC 11775 and *P. aeruginosa* ATCC 10145 strains used in the experiments are more resistant to the toxic effects of Cd at most pH values and phosphate concentrations than *M. luteus* ATCC 9341, *B. subtilis* ATCC 8473 or *S. bovis* ATCC 9809, but because it is well known that within a bacterial species there exist more or less Cd resistant strains, no general conclusions concerning interspecies sensitivity can be made from the results (Novick & Roth 1968, Kondo *et al.* 1974, Tokuyama &

Table 1. The effect of pH and potassium phosphate buffer at different pH values on the toxicity of Cd for *Micrococcus luteus* ATCC 9341, *Bacillus subtilis* ATCC 8473 and *Streptococcus bovis* ATCC 9809.

Medium	<i>M. luteus</i>	<i>B. subtilis</i>	<i>S. bovis</i>
Plate Count Agar. No added phosphate			
pH 6.0	2 <sup>1</sup>	4	4
pH 6.5	4	4	4
pH 7.0	2	4	4
pH 7.5	0.6	4	4
pH 8.0	0.6	4	2
Plate Count Agar prepared in 0.05 M phosphate buffer solution			
pH 6.0	0.08	2	8
pH 7.0	0.4	2	8
pH 8.0	2	1	4
Plate Count Agar prepared in 0.1 M phosphate buffer solution			
pH 6.0	0.008	0.8	8
pH 7.0	0.2	2	8
pH 8.0	0.8	1	6
Plate Count Agar prepared in 0.2 M phosphate buffer solution			
pH 6.0	— <sup>2</sup>	0.6	8
pH 7.0	0.1	2	10
pH 8.0	0.2	1	—

<sup>1</sup> The minimum concentrations of Cd (mg/l) in the test medium which inhibited the growth of the test microbe (M.I.C. value).

<sup>2</sup> No visible growth without added Cd in the medium.

Asano 1974, Tynecka *et al.* 1975, Nakahara *et al.* 1977). It can further be seen from the results that most of the bacteria used in the experiments are generally more sensitive to Cd at a high than at a low pH when phosphate is not added to the substrate. This is in accordance with the results obtained by Babich & Stotzky (1977), who showed the toxicity of Cd for eubacteria, actinomycetes and fungi to be similarly pH-dependent. On the other hand this is in contradiction to the observations made on plants, where Cd is most toxic at low pH values of the soil (Miller *et al.* 1976). According to Miller *et al.* this is due to the increased solubility of Cd compounds, e.g. phosphates, at low pH values. Babich & Stotzky explained the pH effect of Cd referring to




Figure 1. The effect of the pH of the growth medium on the toxicity of Cd for *Staphylococcus aureus* ATCC 12600, *Clostridium perfringens* ATCC 8798, *Escherichia coli* ATCC 11775 and *Pseudomonas aeruginosa* ATCC 10145. The pH of the Plate Count Agar which contained 1, 2, 4, 6, 8, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350 and 400 mg of Cd/l medium was adjusted to pH 6.0, 6.5, 7.5 and 8.0 with 4 N-HCl or 4 N-NaOH. The normal pH of the Plate Count Agar is pH 7.0. Each point represents the lowest concentration of Cd which inhibits the growth of the test microbe in the respective medium (M.I.C. value).

*Hahne & Kroontje* (1973), who observed that in an aqueous system most Cd at pH values under 8 is in an ionic  $\text{Cd}^{2+}$  state. According to *Babich & Stoltzky*,  $\text{Cd}^{2+}$  should readily react with the constituents of the growth medium and thus be less available to the microbe than the  $\text{CdOH}^+$  ion which begins to form at pH values higher than 7.5. On the other hand, *Ramamoorthy & Kushner* (1975) showed that the constituents of bacterial growth mediums do not bind large amounts of the  $\text{Cd}^{2+}$ . According to the present authors the greater toxicity of Cd for bacteria in an alkaline medium than in an acid medium might be more due to a continuously increasing negative charge towards an alkaline value by most bacteria (*Lamanna et al.* 1973), which increases the affinity of cations towards the cell wall, than to other factors.

The effect of pH on the toxicity of Cd was not clearly observed with *S. bovis* and was absent with *B. subtilis*. The reason for the lack of the effect remains obscure. A reversal of or difference in the negative charge on the bacteria might explain the difference.

When potassium phosphate buffer was added to the growth medium the toxic effects of Cd on *M. luteus* were enhanced at pH 6.0 and 7.0 of the medium. A similar increase in toxicity was observed for *B. subtilis* at the pH values studied, but not for *S. bovis*, for which the toxicity of Cd decreased when potassium phosphate buffer was added to the medium. It was also observed that when the pH increased at differing phosphate concentrations the sensitivity of *M. luteus* to Cd decreased. Obviously the stress of the low water activity at an acid pH increases the sensitivity of the microbe to Cd. Finally, when the phosphate concentration is high enough, i.e. 0.2 M at pH 6.0, the growth of *M. luteus* is inhibited even without added Cd in the substrate (Table 1). *M. luteus* is relatively resistant to high solute concentrations in the growth medium (Baird-Parker 1974) as is also *B. subtilis* (Gibson & Gordon 1974). Differences in the structure and composition of the cell wall, e.g. the absence of teichoic acids in *M. luteus* (Baird-Parker) but their presence in *B. subtilis* and also in *S. bovis* (Heptinstall *et al.* 1970, Lamanna *et al.*) might explain the differences observed regarding the sensitivity of the microbes to high phosphate concentrations at varying pH values. It is also well known that there are fundamental differences in the intake of inorganic phosphates between bacteria (Mitchell 1953).

#### REFERENCES

- Abelson, P. H. & E. Aldous: Ion antagonisms in microorganisms. Interference of normal magnesium metabolism by nickel, cobalt, cadmium, zinc, and manganese. *J. Bact.* 1950, 60, 401—413.
- Babich, H. & G. Stotzky: Sensitivity of various bacteria, including actinomycetes, and fungi to cadmium and the influence of pH on sensitivity. *Appl. Environ. Microbiol.* 1977, 33, 681—695.
- Baird-Parker, A. C.: Genus I. *Micrococcus*. In Buchanan, R. E. & N. E. Gibbons, eds.: *Bergey's Manual of Determinative Bacteriology*, 8th Ed. The Williams & Wilkins Company, Baltimore, USA 1974, 478—483.
- Barber, M. & P. M. Waterworth: Penicillinase-resistant penicillins and cephalosporins. *Brit. med. J.* 1964, 2, 344—349.



- Bernheim, F.*: The effect of sulfhydryl compounds and mercurials on the swelling rate of cells of *Pseudomonas aeruginosa* in sodium and potassium buffers. *Proc. Soc. exp. Biol. (N.Y.)* 1970, 133, 841—844.
- Bernheim, F.*: Effect of Na salts of organic acids on acid induced swelling and K loss in *Pseudomonas aeruginosa* and effect of pH on actions of mercuric, cadmium and zinc salts. *Cytobios* 1973, 8, 7—13.
- Donald, C., B. I. Passey & R. I. Swaby*: A comparison of methods for removing trace metals from microbiological media. *J. gen. Microbiol.* 1952, 7, 211—220.
- Friberg, L., M. Piscator, G. F. Nordberg & T. Kjellström*: Cadmium in the Environment, CRC Press, Cleveland, USA 1974.
- Gibson, T. & R. E. Gordon*: Genus I. *Bacillus*. In Buchanan, R. E. & N. E. Gibbons, eds.: *Bergey's Manual of Determinative Bacteriology*. 8th Ed. The Williams & Wilkins Company, Baltimore, USA 1974, 529—550.
- Hahne, H. C. H. & W. Kroontje*: Significance of pH and chloride concentration on behavior of heavy metal pollutants: mercury(II), cadmium(II), and lead(II). *J. Environ. Qual.* 1973, 2, 444—450.
- Heptinstall, S., A. R. Archibald & J. Baddiley*: Teichoic acids and membrane function in bacteria. *Nature (Lond.)* 1970, 225, 519—520.
- Kondo, I., T. Ishikawa & H. Nakahara*: Mercury and cadmium resistances mediated by the penicillinase plasmid in *Staphylococcus aureus*. *J. Bact.* 1974, 117, 1—7.
- Lamanna, C., M. F. Mallette & L. N. Zimmerman*: *Basic Bacteriology — its Biological and Chemical Background*. 4th Ed. The Williams & Wilkins Company, Baltimore, USA 1973.
- Miller, J. E., J. J. Hassett & D. E. Koeppe*: Uptake of cadmium by soybeans as influenced by soil cation exchange capacity, pH, and available phosphorus. *J. Environ. Qual.* 1976, 5, 157—160.
- Mitchell, P.*: Transport of phosphate across the surface of *Micrococcus pyogenes*: Nature of the cell 'inorganic phosphate'. *J. gen. Microbiol.* 1953, 9, 273—287.
- Nakahara, H., T. Ishikawa, Y. Sarai, I. Kondo, H. Kozukue & S. Silver*: Linkage of mercury, cadmium and arsenate and drug resistance in clinical isolates of *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 1977, 33, 975—976.
- Neraal, R. & R. Hamm*. Methode zur quantitativen Bestimmung von zugesetztem Diphosphat und Tripolyphosphat in zerkleinertem Fleisch. (Method for quantitative determination of added diphosphate and tripolyphosphate in comminuted meat). *Die Fleischwirtschaft* 1972, 52, 1171—1174.
- Novick, R. P. & C. Roth*: Plasmid-linked resistance to inorganic salts in *Staphylococcus aureus*. *J. Bact.* 1968, 95, 1335—1342.
- Ramamoorthy, S. & D. J. Kushner*: Binding of mercuric and other heavy metal ions by microbial growth media. *Microb. Ecol.* 1975, 2, 162—176.

- Tokuyama, T. & K. Asano*: Studies on cadmium-ion-resistant bacteria. I. Pure isolation and bacteriological characteristics. Bull. Coll. agric. vet. Med. Nihon Univ. 1974, 31, 182—190.
- Tynecka, Z., J. Zajac & Z. Goś*: Plasmid dependent impermeability barrier to cadmium ions in *Staphylococcus aureus*. Acta Microbiol. polon. Ser. A 1975, 7, 11—20.
- Williams, C. H. & D. J. David*: The accumulation in soil of cadmium residues from phosphate fertilizers and their effect on the cadmium content of plants. Soil Sci. 1976, 121, 86—93.

### SAMMANFATTNING

#### *Inverkan av pH och kaliumfosfatbuffer på toxiciteten av kadmium för bakterier.*

Toxiciteten av kadmium (Cd) för *Micrococcus luteus* ATCC 9341, *Staphylococcus aureus* ATCC 12600, *Clostridium perfringens* ATCC 8798, *Escherichia coli* ATCC 11775 och *Pseudomonas aeruginosa* ATCC 10145 ökade vid stigande pH i bakterieräkningsmediet. Inverkan av pH på toxiciteten var mindre utpräglad för *Streptococcus bovis* ATCC 9809 och utblev för *Bacillus subtilis* ATCC 8473. När kaliumfosfatbuffer tillsattes i odlingsmediet tilltog toxiciteten av Cd för *M. luteus* och *B. subtilis* men minskade för *S. bovis*. När pH-värdet i mediet steg vid olika fosfatkoncentrationer minskade känsligheten av *M. luteus* för Cd. Författarna är av den åsikten att effekten av pH på toxiciteten av Cd är för de flesta mikrober förorsakad av den ökade negativa laddningen vid tilltagande alkalitet, vilket ökar cellväggens affinitet för kationer.

(Received September 8, 1977).

Reprints may be requested from: the Department of Food Hygiene, College of Veterinary Medicine, Hämeentie, 57, 00550 Helsinki 55, Finland.