

Thamnobryum capense comb. nov. (Neckeraceae, Musci) Author(s): Johannes Enroth Source: *Novon*, Vol. 1, No. 3 (Autumn, 1991), p. 110 Published by: <u>Missouri Botanical Garden Press</u> Stable URL: <u>http://www.jstor.org/stable/3391365</u> Accessed: 03/05/2011 04:53

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=mobot.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Missouri Botanical Garden Press is collaborating with JSTOR to digitize, preserve and extend access to Novon.

Thamnobryum capense comb. nov. (Neckeraceae, Musci)

Johannes Enroth

Department of Botany, University of Helsinki, Unioninkatu 44, SF-00170 Helsinki, Finland

Thamnobryum capense (Broth. & Dix.) Enroth, comb. nov. Basionym: Thamnium capense Broth. & Dix. in Dix., Bull. Torrey Bot. Club 43: 71. 1 f. 2. 1916. Porothamnium capense (Broth. & Dix.) Sim, Trans. Roy. Soc. South Africa 15: 404. 1926. TYPE: South Africa. "Habitat: in packing from Cape Town, 710, communicated by G. Webster" (Dixon, 1916) (holotype, BM not seen; isotype, H-Broth, "ex herb. H. N. Dixon"). Illustrations: Dixon (1916: pl. 1, fig. 2); Sim (1926: 404).

De Sloover (1983) noted that Porothamnium capense (Broth. & Dix.) Sim may belong to the genus Thamnobryum Nieuwl.: "D'après les descriptions et illustrations, cette espèce semble appartenir au genre Thamnobryum." However, he did not make a new combination, possibly because he had not seen the type specimen. In connection with preliminary studies on the Neckeraceae for the Bryologia Africana Project, I examined the isotype deposited in H-Broth. The habit, leaf areolation, and the abaxially strongly toothed costa clearly indicate the generic position of *Thamnobryum capense* (Broth. & Dix.) Enroth.

Sim (1926) cited a specimen (Wood 285, NYS) from Inanda, Natal in eastern South Africa. I have not seen that specimen, but Sim's illustrations clearly represent this species; he had also examined the type material deposited in Dixon's herbarium at BM.

Literature Cited

- De Sloover, J. L. 1983. Note de bryologie africaine. XII. Porotrichum et Porothamnium. Bull. Jard. Bot. Natl. Belgique 53: 97-152.
- Dixon, H. N. 1916. New and rare African mosses, from Mitten's herbarium and other sources. Bull. Torrey Bot. Club 43: 63-81 + pl. 1.
- Sim, T. R. 1926. The Bryophyta of South Africa, comprising Sphaerocarpales, Marchantiales, Jungermanniales, Anthocerotales, Sphagnales, Andreaeales, Polytrichales, Bryales. Trans. Roy. Soc. South Africa 15: I-III,