

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Efficacy, safety and economics of bracing after spine surgery: a systematic review of the literature
Author(s)	Zhu, Mary P.; Tetreault, Lindsay A.; Sorefan-Mangou, Fatimah; Garwood, Philip; Wilson, Jefferson R.
Publication date	2018-01-17
Original citation	Zhu, M. P., Tetreault, L. A., Sorefan-Mangou, F., Garwood, P. and Wilson, J. R. (2018) 'Efficacy, safety and economics of bracing after spine surgery: a systematic review of the literature', Spine Journal. doi: 10.1016/j.spinee.2018.01.011.
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.thespinejournalonline.com/article/S1529-9430(18)30014-7/abstract http://dx.doi.org/10.1016/j.spinee.2018.01.011 Access to the full text of the published version may require a subscription.
Rights	© 2018, Elsevier Inc. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. https://creativecommons.org/licenses/by-nc-nd/4.0/
Embargo information	Access to this article is restricted until 12 months after publication by request of the publisher.
Embargo lift date	2019-01-17
Item downloaded from	http://hdl.handle.net/10468/5356

Downloaded on 2021-11-27T05:15:28Z

Accepted Manuscript

Title: Efficacy, safety and economics of bracing after spine surgery: a systematic review of the literature

Author: Mary P. Zhu, Lindsay A. Tetreault, Fatimah Sorefan-Mangou, Philip Garwood, Jefferson R. Wilson

PII: S1529-9430(18)30014-7
DOI: <https://doi.org/10.1016/j.spinee.2018.01.011>
Reference: SPINEE 57576

To appear in: *The Spine Journal*

Received date: 14-9-2017
Revised date: 28-11-2017
Accepted date: 10-1-2018



Please cite this article as: Mary P. Zhu, Lindsay A. Tetreault, Fatimah Sorefan-Mangou, Philip Garwood, Jefferson R. Wilson, Efficacy, safety and economics of bracing after spine surgery: a systematic review of the literature, *The Spine Journal* (2018), <https://doi.org/10.1016/j.spinee.2018.01.011>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Efficacy, Safety and Economics of Bracing after Spine Surgery: A Systematic** 2 **Review of the Literature**

3 Mary P. Zhu^{1*}, Lindsay A. Tetreault^{2,3*}, Fatimah Sorefan-Mangou³, Philip Garwood³, Jefferson
4 R. Wilson^{1,4}

5 ¹Division of Neurosurgery, St. Michael's Hospital, Toronto, Canada

6 ²Division of Neurosurgery, Toronto Western Hospital, Toronto, Canada

7 ³Graduate Entry Medicine, University College Cork, Cork, Ireland

8 ⁴Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Canada

9 * Contributed equally to this paper

10 **Corresponding author:** Jefferson R. Wilson, jeffersonwilson7@gmail.com, 30 Bond Street,
11 Toronto, Ontario, Canada M5W 1W8

12 Abstract

13 **Background Context:** Bracing is often used after spinal surgery to immobilize the
14 spine, improve fusion, and relieve pain. However, controversy exists regarding the
15 efficacy, necessity and safety of various bracing techniques in the post-surgical setting.

16 **Purpose:** In this systematic review, we aimed to compare the effectiveness, safety and
17 cost-effectiveness of postoperative bracing versus no postoperative bracing following
18 spinal surgery in patients with several common operative spinal pathologies.

19 **Study Design/Setting:** Systematic Review

20 **Patient Sample:** N/A

21 **Outcome Measures:** N/A

22 **Methods:** A systematic search was conducted of MEDLINE, Embase and the Cochrane
23 Collaboration Library from 1970 to May 2017, supplemented by manual searching of the
24 reference list of relevant studies and previously published reviews. Studies were
25 included if they compared disability, quality of life, functional impairment, radiographic
26 outcomes, cost-effectiveness and/or complications between patients treated with
27 postoperative bracing versus those not receiving any postoperative bracing. Each article
28 was critically appraised independently by 2 reviewers, and the overall body of evidence
29 was rated using guidelines outlined by the Grading of Recommendation Assessment,
30 Development and Evaluation (GRADE) Working Group.

1 **Results:** Of the 858 retrieved citations, 5 studies met inclusion criteria and were
2 included in this review, consisting of 4 randomized controlled trials and 1 prospective
3 cohort study. Low to moderate evidence suggests that there are no significant
4 differences in most measures of disability, pain, quality of life, functional impairment,
5 radiographic outcomes, and safety between groups. Isolated studies reported
6 statistically significant and inconsistent differences between groups with respect to Neck
7 Disability Index at 6 weeks postoperatively and/or Short Form-36 Physical Component
8 Score at 1.5, 3, 6, and 12 months postoperatively.

9 **Conclusions:** Based on limited evidence, postoperative bracing does not result in
10 improved outcomes following spinal surgery. Future high quality randomized trials will
11 be required to confirm these findings.

12 Keywords: outcomes; postoperative bracing; surgery; spinal pathology; complications.

13 **Introduction**

14 Bracing is routinely used after surgery for a number of spinal pathologies, including degenerative
15 disease of the lumbar and cervical spine, thoracolumbar fractures and scoliosis.[1] The intended
16 goal of this practice is to immobilize the spine, relieve pain, improve fusion rates, and remind
17 patients to avoid certain activities that may compromise their recovery.[1, 2] However, a number
18 of important complications can arise from bracing, including dysphagia, nerve palsies, pressure
19 ulcers, and skin rashes.[1] Furthermore, braces can be uncomfortable for some patients as well as
20 costly.[2]

21 Given the paucity of high-quality comparative studies, it is unclear whether postoperative
22 bracing can effectively limit and restrict spinal movements, reduce rates of pseudoarthrosis and
23 optimize patient recovery. Certain advances in spinal surgery have allowed for rigid internal
24 stabilization of the spine and, arguably, have decreased the requirement for external
25 immobilization. Although these techniques may be sufficient to achieve successful fusion, there
26 may still be a role for postoperative bracing in higher risk patients, including those who smoke,
27 suffer from osteoporosis or require an extensive multilevel surgery.

28 As a result of the limited body of evidence available, spine surgeons often base their decision to
29 use postoperative bracing on their own clinical experience and training.[1] This finding is
30 supported by a survey that highlighted substantial disagreement among spinal surgeons with
31 respect to the appropriate type, duration and indication for use of postoperative bracing after
32 anterior cervical spine surgery.[3] Given the heterogeneity in management strategies, there is a
33 need to synthesize results from high quality studies and establish recommendations surrounding
34 care following spinal surgery.

35 This systematic review addresses 4 key questions (KQs). KQ1: What is the efficacy and
36 effectiveness of postoperative bracing compared with no bracing based on disability, pain,

1 quality of life and functional outcomes? KQ2: What is the impact of postoperative bracing
2 compared with no bracing on radiographic outcomes? KQ3: What is the safety profile of
3 postoperative bracing compared with no bracing? KQ4: What is the cost-effectiveness of
4 postoperative bracing? **Importantly, this systematic review will assess the overall strength of**
5 **the evidence using methodology developed by the Grading of Recommendation**
6 **Assessment, Development and Evaluation (GRADE) working group.**

7 **Methods**

8 **Eligibility Criteria**

9 *Population*

10 Our review targeted studies including patients undergoing surgery for any spinal pathology,
11 including cervical and lumbar degenerative disease, trauma, oncology and adolescent idiopathic
12 scoliosis. Studies were excluded if patients under study were treated non-operatively (Table 1).

13 *Intervention and Comparison*

14 This review focused on studies that had an intervention group who received postoperative
15 bracing, and a control group who received standard of care and no postoperative bracing (Table
16 1).

17 *Outcomes*

18 For KQ1, we sought studies that considered the clinical efficacy of postoperative bracing by
19 measuring patient disability, pain, quality of life, and/or functional outcomes. For KQ2, we
20 focused on studies that assessed radiographic outcomes, including fusion rate, sagittal alignment,
21 and range of motion. For KQ3, we sought studies that compared complication rates and adverse
22 events between the intervention and control groups. For KQ4, we focused on studies that
23 examined various measures of cost-effectiveness including incremental cost-effectiveness ratio
24 and cost per unit of outcome (Table 1).

25 **Study Characteristics**

26 For KQ1, 2 and 3, we sought comparative studies (i.e. randomized controlled trials, cohort
27 studies) designed to evaluate differences between a postoperative bracing group and a control
28 group. To be included, studies needed to have at least 10 patients per group. Case reports,
29 nonclinical studies, and animal studies were excluded. For KQ4, we focused on full economic
30 studies. For all KQs, abstracts, editorials, letters, narrative and systematic reviews were
31 excluded. Duplicate publications of the same study that did not report on different outcomes
32 were also excluded.

33 **Information Sources**

1 A systematic search of MEDLINE, Embase and Cochrane Collaboration Library was conducted
2 to identify relevant studies. Manual searching of the reference lists of included studies and
3 previously published reviews was also conducted to ensure all relevant studies were located.

4 **Search Strategy**

5 The search strategy was first developed in MEDLINE and then appropriately modified for the
6 other databases. We used the following search terms to search all databases: Orthotic Devices
7 AND Spinal Diseases AND Postoperative Complications/Care AND Treatment Outcome or
8 Outcome Assessment. Only studies involving humans, written in English and published in peer-
9 review journals between 1970 and May 2017 were considered for inclusion, with no other limits
10 applied. **A detailed search strategy is provided in the supplemental digital material.**

11 **Study Selection**

12 All abstracts and titles were reviewed and sorted by our predefined inclusion criteria. Studies
13 were classified as relevant, possibly relevant, or irrelevant. Full text investigation of all relevant
14 and possibly relevant studies was done for further clarification.

15 **Data Extraction and Synthesis**

16 The following data were extracted from each included article: study design; patient sample and
17 characteristics, including diagnosis, surgical summary and type of bracing; outcome assessment
18 tools; follow-up schedule; drop-out rate; and results of association, including standard deviation,
19 odds ratio, confidence intervals, and p-values.

20 **Risk of Bias in Individual Studies**

21 The class of evidence for each article was rated (Class I, II, III, IV) independently by 2 reviewers
22 using criteria outlined by the Journal of Bone and Joint Surgery for therapeutic studies and
23 modified to encompass both methodological quality and risk of bias. Randomized controlled
24 trials were rated based on patient allocation, intention to treat analysis, independent or blinded
25 assessment, whether co-interventions were applied equally, rates of follow-up, statistical power,
26 and control for possible confounding. Prospective cohort studies were rated based on
27 independent or blinded assessment, whether co-interventions were applied equally, rates of
28 follow-up, statistical power, and control for possible confounding. Due to the nature of the
29 intervention, studies were rated as having independent or blinded assessment if surgeons were
30 blinded to the randomization group until after surgery, patients were blinded to the
31 randomization group until day of admission of surgery or after surgery, and/or radiologists
32 reviewing radiographs were blinded to the randomization group.

33 **Risk of Bias Across Studies**

1 The overall body of evidence was assessed using a scoring system developed by the **GRADE**
2 working group with recommendations from the Agency for Healthcare Research and Quality
3 (AHRQ). **This methodology allows for an assessment of the overall strength of the evidence**
4 **and is particularly valuable for highlighting critical knowledge gaps.**

5 The initial strength of the overall body of evidence was graded as “high” if half or more of the
6 studies were randomized controlled trials and “low” if the majority of studies were observational
7 studies. The body of evidence was downgraded 1, 2, or 3 levels if there was risk of bias, results
8 were inconsistent or consistency was unknown, the evidence was indirect, the effect estimates
9 were imprecise (e.g. wide confidence intervals), or if there was publication bias. If no
10 downgrades were made, the body of evidence was upgraded 1, 2 or 3 levels based on large
11 magnitude of effect, dose-response gradient or if all plausible biases would decrease the
12 magnitude of an apparent effect.

13 The final rating of the body of evidence expresses our confidence in the estimate of effect and
14 the impact of further research on this topic. An overall strength of “high” means we have high
15 confidence that the evidence reflects the true effect. Further research is very unlikely to change
16 our confidence in the estimate of effect. The overall strength of “moderate” means we have
17 moderate confidence that the evidence reflects the true effect. Further research may change our
18 confidence in the estimate of effect and may change the estimate. A grade of “low” means we
19 have low confidence that the evidence reflects the true effect. Further research is likely to change
20 the confidence in the estimate of effect and likely to change the estimate. A grade of
21 “insufficient” means that evidence is either unavailable or does not permit a conclusion.

22 **Results**

23 **Study Selection**

24 The initial electronic search yielded a total of 853 citations. Five additional citations were
25 identified through reference scanning. After initial review of abstracts and titles, 841 studies did
26 not meet our inclusion criteria. Following full text investigation, an additional 12 studies were
27 excluded because 1) they were not comparative studies; 2) patients were not treated surgically; 3)
28 there was no postoperative comparison of intervention and control groups; 4) they had a different
29 outcome of interest; 5) they had no control group; 6) they were a duplicate publication with no
30 new results; and/or 7) they were not in English. A total of 5 studies were deemed relevant
31 following this review process.

32 **Study Characteristics**

33 For KQ1, we identified 4 studies (3 randomized controlled trials, 1 prospective cohort)
34 discussing the effect of postoperative bracing on disability, pain, quality of life and functional
35 outcomes.[4-7] Sample sizes ranged from 33 to 257 surgical patients with mean ages between
36 43.9 and 72.7 years. All patients were diagnosed with degenerative cervical myelopathy or

1 radiculopathy, or degenerative disease of the lumbar spine. Bracing included Philadelphia
 2 collars, cervical collars, and lumbar corsets for differing lengths of time. Various outcome
 3 measures were used across the studies, with the Short Form-36 (SF-36) Physical Component
 4 Score (PCS) reported the most frequently (n = 4),[4-7] followed by the SF-36 Mental
 5 Component Score (MCS) (n = 3),[4, 6, 7] SF-36 subscales (n = 3),[4, 6, 7] Neck Disability Index
 6 (NDI) (n = 2),[4, 5] neck pain (n = 2),[4, 5] and arm pain (n = 2).[4, 5]

7 For KQ2, a total of 5 studies (4 randomized controlled trials, 1 prospective cohort) met our
 8 inclusion criteria.[4-8] These studies were designed to assess the impact of postoperative bracing
 9 compared with no bracing on radiographic outcomes. Sample sizes ranged from 33 to 257
 10 surgical patients with mean ages between 14.3 and 72.7 years. Patients were diagnosed with
 11 degenerative cervical myelopathy or radiculopathy, degenerative disease of the lumbar spine, or
 12 adolescent idiopathic scoliosis. Bracing included Philadelphia collars, cervical collars, body
 13 casts, and lumbar corsets for various lengths of time. Fusion rate was the most frequently
 14 reported outcome measure (n = 3).[4, 5, 7]

15 For KQ3, we identified 3 studies (2 randomized controlled trials, 1 prospective cohort)
 16 examining the safety profile of postoperative bracing compared with no bracing.[5-7] Sample
 17 sizes ranged from 50 to 257 surgical patients with mean ages between 43.9 and 72.7 years.
 18 Patients were diagnosed with degenerative cervical myelopathy or radiculopathy, or degenerative
 19 disease of the lumbar spine. Bracing included Philadelphia collars, cervical collars, and lumbar
 20 corsets for various lengths of time. Outcome measures included complications (n = 3)[5-7] and
 21 revision surgery or second procedure (n = 2).[5, 7]

22 No studies met our inclusion criteria for KQ4 on the cost-effectiveness of postoperative bracing.

23 Risk of Bias

24 We critically appraised the 5 studies included in our review. **The inter-rater reliability was**
 25 **80%; disagreement on the fifth study, by Yee et al., surrounding the “intention to treat”**
 26 **analysis was resolved through discussion. It was noted that although 90 patients were**
 27 **randomized, only 72 were included in their analysis, and therefore the study did not use**
 28 **intention to treat analysis.**[7] **Of the studies included,** 4 were considered Class II and 1 was
 29 rated Class III. The 4 Class II studies were randomized controlled trials, and were downgraded
 30 from Class I because they did not include intent-to-treat analysis, independent or blind
 31 assessment, adequate sample size, random sequence generation and/or statement of concealed
 32 allocation, had unreported follow-up rates or follow-up rates < 80%, and/or did not control for
 33 possible confounders. The Class III study was a prospective cohort study that was downgraded
 34 from Class II because co-interventions were not applied equally and follow-up was not reported.

35 Results of Individual Studies

36 *KQ1: What is the efficacy and effectiveness of postoperative bracing compared with no bracing*
 37 *based on disability, pain, quality of life and functional outcomes?*

1 Degenerative cervical myelopathy or radiculopathy

2 Three studies compared disability, pain, quality of life and/or functional outcomes between the
3 postoperative bracing and non-bracing groups in patients with degenerative cervical myelopathy
4 or radiculopathy.[4-6]

5 According to Abbott et al., patients who received postoperative bracing had better NDI scores at
6 6 weeks after surgery compared to patients who did not receive postoperative bracing (mean
7 difference between groups -4.4, 95% CI -8.6 to -0.2, $p = 0.042$).[4] However, Campbell et al.,
8 reported that patients in the non-braced group had better NDI scores at 6 weeks after surgery
9 compared to patients in the braced group ($p = 0.008$).[5]

10 Abbott et al. also found that patients in the control group had better SF-36 PCS scores at 6 weeks
11 (mean difference between groups 5.8, 95% CI 0.8-10.7, $p = 0.025$), 3 (mean difference between
12 groups 6.8, 95% CI 0.4-13.1, $p = 0.038$), 6 (mean difference between groups 7.4, 95% CI 1.4-
13 13.4, $p = 0.017$), and 12 months (mean difference between groups 7.5, 95% CI 0.3-14.6, $p =$
14 0.041) after surgery compared to patients in the postoperative bracing group.[4] In contrast,
15 Campbell et al. and Hida et al. found no significant differences in SF-36 PCS scores between the
16 two groups at all time points assessed (Campbell et al. 6 months $p = 0.481$, 12 months $p = 0.260$,
17 24 months $p = 0.279$; Hida et al. $p = 0.537$).[5, 6]

18 Two studies evaluated differences in various subscales of the SF-36 between a postoperative
19 bracing and a control group.[4, 6] A single study by Abbott et al. reported significantly better
20 SF-36 Bodily Pain (BP) scores in the postoperative bracing group at 6 (mean difference between
21 groups 21.4, 95% CI 4.4-38.5, $p=0.016$) and 12 (mean difference between groups 17.5, 95% CI
22 1.7-33.2, $p=0.031$) months, as well as better SF-36 Social Functioning (SF) scores at 12-months
23 (mean difference between groups 16.5, 95% CI 0.1-32.9, $p=0.049$) than in the non-bracing
24 group.[4] The other six subscales (Physical Functioning (PF), General Health (GH), Role
25 Limitations Physical (RP), Vitality (VT), Role Limitations Emotional (RE) and Mental Health
26 (MH)) were not significantly different between treatment groups at all time points assessed (1.5,
27 3, 6, 12 and 24 months).[4] A second study by Hida et al., identified no differences between the
28 collar-fixation group and the control group with respect to SF-36 BP subscale ($p=0.848$).[6]

29 Other measures of disability, pain, quality of life, and functional impairment, including SF-36
30 MCS, Visual Analog Scale (VAS), Japanese Orthopedic Association (JOA) recovery rate, Falls
31 Efficacy Scale (FES), unipedal balance standing test, and neck and arm pain were not
32 significantly different between the postoperative bracing and non-bracing groups (Table 3).[4-6]

33 Degenerative disease of the lumbar spine

34 A single study by Yee et al. examined pain and quality of life outcomes for patients with a
35 degenerative disease of the lumbar spine.[7] There were no significant differences in the Dallas
36 Pain Questionnaire (DPQ), daily activity category $p = 0.34$, work/leisure category $p = 0.67$,

1 anxiety-depression category $p = 0.17$, social category $p = 0.40$), SF-36 PCS ($p = 0.30$), SF-36
2 MCS ($p = 0.57$) and SF-36 subscales (PF $p = 0.38$, BP $p = 0.28$, GH $p = 0.23$, RP $p = 0.41$, VT p
3 $= 0.25$, SF $p = 0.79$, RE $p = 0.86$, MH $p = 0.30$) between the braced and non-braced groups at all
4 time points assessed.[7]

5 *KQ2: What is the impact of postoperative bracing compared with no bracing on radiographic*
6 *outcomes?*

7 Degenerative cervical myelopathy or radiculopathy

8 Three studies assessed the radiographic outcomes between the postoperative bracing and non-
9 bracing groups in patients with degenerative cervical myelopathy or radiculopathy.[4-6]

10 Abbott et al. reported no significant differences in cervical range of motion (ROM) ($p > 0.05$),
11 fusion rate ($p = \text{NR}$), and sagittal alignment ($p = \text{NR}$) between the two groups.[4] Campbell et al.
12 also found no significant differences in fusion success between the bracing and non-bracing
13 groups ($p = \text{NR}$).[5] Similarly, Hida et al. concluded that there were no significant differences in
14 ROM ($p = 0.61$) or decrease in lordotic angle C2-7 ($p = 0.82$) between groups.[6]

15 Degenerative disease of the lumbar spine

16 Yee et al. found no significant differences in fusion rate at 12 months ($p = 0.8$) or 24 months ($p =$
17 0.9) postoperatively between patients who wore a lumbar corset and those who did not.[7]

18 Adolescent idiopathic scoliosis

19 Based on a single study, there was no significant difference in mean loss of spinal curve
20 correction between the braced and non-braced groups at all time points assessed in patients with
21 adolescent idiopathic scoliosis ($p = \text{NR}$).[8]

22 *KQ3: What is the safety profile of postoperative bracing compared with no bracing?*

23 Degenerative cervical myelopathy or radiculopathy

24 Campbell et al. reported rates of instrumentation failure, graft extrusion, and second procedures
25 including revisions, removals, reoperations, supplemental fixations, and external bone growth
26 stimulators,[5] while Hida et al. considered all perioperative complications including surgical
27 site infection, epidural hematoma, and C5 palsy.[6]

28 Both Campbell et al. and Hida et al. identified no significant differences in the incidence of
29 complications between the bracing and non-bracing groups (Campbell et al. no events of
30 instrumentation failure or graft extrusion in either group, $p = \text{NC}$; Hida et al. $p = 0.53$).[5, 6] In
31 addition, Campbell et al. found no significant differences between groups in the rate of revision
32 surgery ($p=0.653$), removals ($p=0.724$), reoperations ($p=1.000$), supplemental fixations
33 ($p=0.286$) or any second operation ($p = 0.184$).[5]

1 Degenerative disease of the lumbar spine

2 Based on a single study, there were no significant differences in rates of revision surgery, a
3 second procedure or complications between the braced and non-braced groups at all time points
4 assessed ($p = 0.8$).^[7] Revision surgery included later-stage revision surgery due to symptomatic
5 nonunion or later-stage hardware removal due to prominence/bursitis, and complications
6 included intraoperative incidental durotomy, intraoperative pedicle-screw-placement issues, new
7 postoperative radiculopathy, early postoperative pulmonary embolism, wound
8 seroma/hematoma, deep wound infection or persistent lumbar radiculopathy postoperatively.

9 *KQ4: What is the cost-effectiveness of postoperative bracing?*

10 No studies were identified that evaluated the cost-effectiveness of postoperative bracing.

11 **Summary of Evidence**

12 The overall quality of evidence ranged from “insufficient” to “moderate”. Evidence was
13 downgraded due to risk of bias, unknown or inconsistent consistency of results, and/or imprecise
14 effect estimates (e.g. wide confidence intervals).

15 Based on low evidence, postoperative bracing in patients with degenerative cervical myelopathy
16 or radiculopathy does not result in improved SF-36 MCS, VAS, JOA recovery rate, sagittal
17 alignment, ROM, decrease in lordotic angle C2-7, and rate of revision surgery or second
18 procedure outcomes. Based on moderate evidence, postoperative bracing does not result in
19 improved neck or arm pain, fusion rate, and incidence of complications in this patient
20 population.

21 Low evidence suggests no improvement in DPQ, SF-36 PCS, SF-36 MCS, SF-36 subscales,
22 fusion rate, incidence of complications, and rate of revision surgery or second procedure
23 following postoperative bracing in patients with a degenerative disease of the lumbar spine.

24 Based on low evidence, postoperative bracing is not associated with improved loss of spinal
25 curve correction in adolescents with idiopathic scoliosis.

26 **Discussion**

27 The use of bracing after surgery for a variety of spinal pathologies remains controversial with
28 limited evidence available to the surgeon to make an informed decision. Historically cited
29 reasons to use bracing include to limit mobility and stabilize the spine, improve fusion rates,
30 prevent graft dislodgement or subsidence, reduce postoperative pain and optimize outcomes.
31 Bracing, however, can be uncomfortable, lead to social isolation and be associated with
32 complications such as dysphagia, nerve palsies and pressure ulcers. This review summarizes the
33 current literature on the efficacy, safety and cost-effectiveness of bracing after spinal surgery
34 **using rigorous methodology and is the first to synthesize results using methodology**

1 **proposed by the GRADE working group.** This knowledge is valuable in a clinical setting, and
2 can be used by clinicians to determine the most appropriate postoperative management
3 strategies. Clinical judgement, however, is still required to determine whether a patient may
4 benefit from additional external immobilization.

5 *Degenerative cervical myelopathy or radiculopathy*

6 Based on this review, postoperative bracing of the cervical spine has no impact on 1) most
7 measures of pain, disability, functional impairment, and quality of life; 2) radiographic outcomes
8 such as fusion rate and range of motion; and 3) rates of complications and reoperations. In the
9 study by Abbott et al., patients who received postoperative bracing exhibited superior
10 improvements in neck disability (NDI at 6 weeks) and various metrics of quality of life (SF-36
11 PCS at 6 weeks to 12 months, SF-36 SF at 12-months and SF-36 BP at 6 to 12 months) than
12 those who did not.[4] These findings can be partly explained by psychological factors, including
13 a sense of security provided by the brace, increased coping mechanisms, improved functional
14 self-efficacy, and less fear avoidance.[4] In contrast, Campbell et al identified that patients with
15 postoperative bracing had worse NDI scores, likely due to the discomfort and disability
16 associated with wearing a brace.[5]

17 Previous studies in healthy subjects have demonstrated that cervical bracing reduces velocity of
18 eye movements and causes deterioration in the anterior to posterior body sway induced by
19 vibration of the calf muscles.[9, 10] Given these findings, it is hypothesized that restricting
20 cervical motion through external immobilization may significantly impair static postural control
21 and disturb balance during dynamic movement. In the study by Abbott et al., however, there
22 were no differences between a bracing group and a control group with respect to the unipedal
23 balance standing test.[4]

24 Biomechanical studies have indicated that cervical collars help to restrict motion during routine
25 activities and stabilize the spine.[11-13] However, early mobilization exercises can prevent spine
26 contracture and improve range of motion after surgery.[6] In studies by Hida et al. and Abbott et
27 al., there were no significant differences in cervical range of motion between the postoperative
28 bracing group and the control group.[4, 6] Although range of motion often decreases following
29 surgery, this is more likely due to fusion and fixation techniques, damage to the cervical flexors
30 and extensors and injury to the facet joints. Postoperative bracing may also help to decrease the
31 risk of graft or cage migration, maintain spinal alignment and improve fusion rates.

32 Advancements in surgical procedures, however, have allowed for internal stabilization of the
33 spine and may have decreased the requirement for external immobilization. For example, the use
34 of anterior plates has shown to increase fusion and decrease subsidence rates by limiting motion
35 between the graft and vertebral bodies.[14, 15] This finding was confirmed by Campbell et al.
36 who reported no significant difference in rates of fusion between a bracing and non-bracing
37 group;[5] these results question the need for postoperative bracing, especially in patients
38 undergoing internal stabilization. There may still be a role for postoperative bracing in patients at

1 a higher risk of pseudoarthrosis and disease progression, including those who smoke, have had a
2 previous spine operation, and/or are treated without rigid internal fixation.[16-21] Furthermore,
3 surgeons are more likely to use postoperative bracing following a multilevel anterior cervical
4 discectomy and fusion (ACDF) (76%) compared to a single level ACDF (55%).[3]

5 *Degenerative Disease of the Lumbar Spine*

6 Based on this review, the use of a lumbar corset following surgery for degenerative lumbar
7 disease has no impact on pain, disability, functional impairment, quality of life, radiographic
8 outcomes, incidence of complications, and rate of reoperations. Postoperative bracing is often
9 used in this population to relieve pain, limit mobility, improve fusion rates and optimize
10 outcomes; however, the study by Yee et al. indicated no advantage or disadvantage to the use of
11 a lumbar corset.[7] There may still be a role for bracing in patients at a higher risk of nonunion
12 or pseudoarthrosis, such as those who smoke or require a multilevel fusion.

13 *Adolescent Idiopathic Scoliosis*

14 Historically, molded plaster braces were used to correct the curve following posterior fusions
15 without instrumentation and maintain this correction until solid bony fusion. Techniques
16 proposed in the study by Christodoulou et al., including the use of Harrington distraction rods,
17 however, has decreased the need for postoperative external bracing due to more rigid internal
18 fixation.[8] In this review, a single study examined postoperative bracing in an adolescent
19 idiopathic scoliosis population and found no differences in radiographic outcomes between
20 patients who received bracing versus those who did not. Further investigation, however, is
21 needed to determine the effectiveness of bracing in this population based on other outcome
22 measures.

23 Our finding that bracing may not confer additional benefits following spine surgery will have
24 relevant applications in a clinical setting. First, complications such as skin reactions, dysphagia,
25 pressure ulcers and nerve palsies, as well as costs associated with bracing can be eliminated.
26 Second, if postoperative bracing is not required, there may be less of an impact on activities of
27 daily living, decreased social isolation, body anxiety, self-perception and body image issues and
28 an improved ability to return to work or school following surgery.

29 **Strengths and Limitations**

30 To our knowledge, no other reviews have evaluated the merits of bracing in the post-surgical
31 setting for patients with various spinal pathologies **using the GRADE approach. This**
32 **methodology allows for rigorous evaluation of the overall strength of the evidence and**
33 **helps to identify critical knowledge gaps in the literature (e.g a lack of high-quality**
34 **comparative studies and limited data on the cost-effectiveness of postoperative bracing).**
35 Furthermore, the majority of current studies published on this topic have moderately high to high

1 risk of bias and imprecise estimates of effect (or estimates with unknown precision). Consistency
2 of results is also largely unknown as results are based on single studies.

3 Our review also has its limitations. First, our search was restricted to studies published in English
4 and, as a result, some articles with relevant titles or abstracts were excluded. Second, although
5 results were separated based on patient population, the type and length of bracing, as well as
6 surgical technique varied substantially among studies, preventing pooling of data and meta-
7 analysis.

8 **Conclusions**

9 Based on the results of this review, postoperative bracing does not result in improved outcomes
10 after spine surgery in patients with various spinal pathologies. Although some outcomes were
11 significantly different between bracing and non-bracing groups, firm conclusions cannot be made
12 due to small sample sizes, risk of bias and low quality of evidence. Finally, given the paucity of
13 studies available, no conclusions can be made regarding the cost-effectiveness of bracing after
14 surgery.

15 **References**

- 16 1. Connolly, P.J. and D. Grob, *Bracing of patients after fusion for degenerative problems of*
17 *the lumbar spine--yes or no?* Spine (Phila Pa 1976), 1998. **23**(12): p. 1426-8.
- 18 2. Soliman, H.A., et al., *The Early Impact of Postoperative Bracing on Pain and Quality of*
19 *Life Following Posterior Instrumented Fusion for Lumbar Degenerative Conditions: A*
20 *Randomized Trial.* Spine (Phila Pa 1976), 2017.
- 21 3. Bible, J.E., et al., *Postoperative bracing after spine surgery for degenerative conditions:*
22 *a questionnaire study.* Spine J, 2009. **9**(4): p. 309-16.
- 23 4. Abbott, A., M. Halvorsen, and A. Dederig, *Is there a need for cervical collar usage post*
24 *anterior cervical decompression and fusion using interbody cages? A randomized*
25 *controlled pilot trial.* Physiother Theory Pract, 2013. **29**(4): p. 290-300.
- 26 5. Campbell, M.J., et al., *Use of cervical collar after single-level anterior cervical fusion*
27 *with plate: is it necessary?* Spine (Phila Pa 1976), 2009. **34**(1): p. 43-8.
- 28 6. Hida, T., et al., *Collar Fixation Is Not Mandatory After Cervical Laminoplasty: A*
29 *Randomized Controlled Trial.* Spine (Phila Pa 1976), 2017. **42**(5): p. E253-E259.
- 30 7. Yee, A.J., et al., *Use of a postoperative lumbar corset after lumbar spinal arthrodesis for*
31 *degenerative conditions of the spine. A prospective randomized trial.* J Bone Joint Surg
32 Am, 2008. **90**(10): p. 2062-8.
- 33 8. Christodoulou, A.G., et al., *Adolescent idiopathic thoracic scoliosis. A prospective trial*
34 *with and without bracing during postoperative care.* J Bone Joint Surg Br, 1987. **69**(1): p.
35 13-6.
- 36 9. Burl, M.M., J.G. Williams, and U.S. Nayak, *Effects of cervical collars on standing*
37 *balance.* Arch Phys Med Rehabil, 1992. **73**(12): p. 1181-5.
- 38 10. Karlberg, M., M. Magnusson, and R. Johansson, *Effects of restrained cervical mobility*
39 *on voluntary eye movements and postural control.* Acta Otolaryngol, 1991. **111**(4): p.
40 664-70.

- 1 11. DiPaola, C.P., et al., *Comparing cervical spine motion with different halo devices in a*
2 *cadaveric cervical instability model*. Spine (Phila Pa 1976), 2009. **34**(2): p. 149-55.
- 3 12. Koller, H., et al., *In vivo analysis of atlantoaxial motion in individuals immobilized with*
4 *the halo thoracic vest or Philadelphia collar*. Spine (Phila Pa 1976), 2009. **34**(7): p. 670-
5 9.
- 6 13. Schneider, A.M., et al., *Reduction in head and intervertebral motion provided by 7*
7 *contemporary cervical orthoses in 45 individuals*. Spine (Phila Pa 1976), 2007. **32**(1): p.
8 E1-6.
- 9 14. Yue, W.M., W. Brodner, and T.R. Highland, *Long-term results after anterior cervical*
10 *discectomy and fusion with allograft and plating: a 5- to 11-year radiologic and clinical*
11 *follow-up study*. Spine (Phila Pa 1976), 2005. **30**(19): p. 2138-44.
- 12 15. Wang, J.C., et al., *The effect of cervical plating on single-level anterior cervical*
13 *discectomy and fusion*. J Spinal Disord, 1999. **12**(6): p. 467-71.
- 14 16. Hilibrand, A.S., et al., *Impact of smoking on the outcome of anterior cervical arthrodesis*
15 *with interbody or strut-grafting*. J Bone Joint Surg Am, 2001. **83-A**(5): p. 668-73.
- 16 17. Yan, C., N.G. Avadhani, and J. Iqbal, *The effects of smoke carcinogens on bone*. Curr
17 Osteoporos Rep, 2011. **9**(4): p. 202-9.
- 18 18. Kusin, D.J., U.M. Ahn, and N.U. Ahn, *The Effect of Smoking on Spinal Cord Healing*
19 *Following Surgical Treatment of Cervical Myelopathy*. Spine (Phila Pa 1976), 2015.
20 **40**(18): p. 1391-6.
- 21 19. Tetreault, L.A., et al., *A clinical prediction model to determine outcomes in patients with*
22 *cervical spondylotic myelopathy undergoing surgical treatment: data from the*
23 *prospective, multi-center AOSpine North America study*. J Bone Joint Surg Am, 2013.
24 **95**(18): p. 1659-66.
- 25 20. Lau, D., et al., *The effects of smoking on perioperative outcomes and pseudarthrosis*
26 *following anterior cervical corpectomy: Clinical article*. J Neurosurg Spine, 2014. **21**(4):
27 p. 547-58.
- 28 21. Lee, J.C., et al., *Adjacent segment pathology requiring reoperation after anterior cervical*
29 *arthrodesis: the influence of smoking, sex, and number of operated levels*. Spine (Phila Pa
30 1976), 2015. **40**(10): p. E571-7.

31

32

33 Figure 1. Flow diagram of study selection

34

1 **Table 1. Inclusion and exclusion criteria for studies reviewed**

Characteristic	Inclusion	Exclusion
Population	Patients w/ any spinal pathology treated surgically and followed postoperatively (e.g. scoliosis, spinal trauma, cervical myelopathy)	Patients treated non-operatively
Intervention	Postoperative bracing	
Comparison	No postoperative bracing Standard of care	
Outcome	<p>KQ1: Efficacy and effectiveness</p> <ul style="list-style-type: none"> • Disability (e.g. NDI) • Pain (e.g. VAS) • Quality of life (e.g. SF-36) • Functional outcomes (e.g. JOA, mJOA, Nurick) <p>KQ2: Radiographic Outcomes</p> <ul style="list-style-type: none"> • Fusion rate • Sagittal alignment • ROM <p>KQ3: Safety</p> <ul style="list-style-type: none"> • Complications or adverse events (e.g. dysphagia, hardware, skin complications, reoperation, revision) 	<p>KQ1: Subjective neurological status, patient satisfaction, improvement of symptoms</p>

surgery, infection, hematoma)

KQ4: Cost-effectiveness

- Incremental cost-effectiveness ratio
(or similar)
- Cost per unit of outcome

Study design	KQ1, 2, 3: Comparative studies (e.g. RCT, prospective cohort, case-control studies) designed to compare a postoperative bracing group with a control group; n ≥ 10 per group. KQ4: Full economic studies	Case reports Nonclinical studies Animal studies
Publication	Studies published in peer-review journals and in English	Abstracts, editorials, letters Duplicate publications of the same study that do not report on different outcomes Narrative or systematic reviews

- 1 JOA = Japanese Orthopedic Association; mJOA = modified Japanese Orthopedic Association;
- 2 NDI = neck disability index; ROM = range of motion; SF-36 = short form-36; VAS = visual
- 3 analog scale

1 **Table 2. Characteristics of included studies**

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
Abbott et al., 2013 (RCT, pilot)	Brace (n = 17)	M, 53%	53.4 (NR, 13)	Cervical spondylosis (n = 8), Cervical disk herniation (n = 4), Cervical degenerative disc disease (n = 5)	ACDF with interbody cage	Philadelphia Collar daytime only for 6 weeks	KQ1: Borg CR-10 (neck and arm pain), NDI, SF-36 (PCS and MCS), FES, unipedal balance standing test	1.5, 3, 6, 12, and 24 months postoperative	55% (n = 18)
	Control (n = 16)	M, 69%	47.3 (NR, 11)	Cervical spondylosis (n = 5), Cervical disk herniation (n = 4),			KQ2: CROM, fusion rate,		

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
				Cervical degenerative disc disease (n = 7)			sagittal alignment KQ3: None KQ4: None		
Campbell et al., 2008 (Prospective cohort)‡	Brace (n = 149) Control (n = 108)	M, 43.6% M, 49.1%	44.3 (NR, 8.8) 43.3 (NR, 9.0)	Single-level radiculopathy or myelopathy	Decompression and arthrodesis using allograft and anterior cervical plate or	Cervical collar	KQ1: NDI, neck and arm pain scales, SF-36 KQ2: Fusion success KQ3: Second procedure§,	1.5, 3, 6, 12, and 24 months postoperative	NR

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
					arthoplasty		instrumentation n failure, graft extrusion KQ4: None		
Christodoulou et al., 1987 (RCT)	Brace (n = 25) Control (n = 25)	NA NA	NR NR	Adolescent idiopathic scoliosis with thoracic curves $\geq 35^\circ$	Posterior decompression and fusion with Harrington instrumentation augmented	Plaster body cast for 6 months	KQ1: None KQ2: Spinal curve KQ3: None KQ4: None	3, 6, 12, and 24 months postoperative	NR

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
					by a Cotrel				
					bar or by				
					sublaminal				
					Luque wires				
Hida et al., 2017 (RCT)	Brace (n = 45)	M, 73%	72.0 (NR, 8.7)	Cervical myelopathy secondary to	Double-door cervical laminoplasty	Philadelphia collar for 2 weeks	KQ1: VAS, JOA, SF-36 (PCS and MCS)	0.5, 3, 6, and 12 months	18% (n = 16)
	Control (n = 45)	M, 62%	71.6 (NR, 9.6)	multisegmental cervical spondylotic stenosis	without instrumentati on		KQ2: ROM (total, extension and flexion),	postoperati ve	

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
							lordotic angle C2-7 KQ3: Perioperative complications ¶ KQ4: None		
Yee et al., 2008 (RCT)	Brace (n = 46)	M, 43% **	52 (NR, 15.2) **	Spondylosis/stenosis (n = 11), degenerative spondylolisthesis (n = 13), isthmic	Posterior lumbar spinal arthrodesis	Canvas lumbar corset full-time for 8	KQ1: DPQ, SF-36 KQ2: Fusion rate KQ3: Revision	12 and 24 months postoperative	20% (n = 18)

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing weeks	Outcome Factors Assessed	Follow-up	Drop-out, n†
				spondylolisthesis (n = 8), junctional syndrome (n = 1), pseudarthrosis (n = 2), iatrogenic/post-op instability (n = 2)			surgery††, complications ‡‡ KQ4: None		
	Control (n = 44)	M, 54% **	53 (NR, 15.4) **	Spondylosis/stenosis (n = 12), degenerative spondylolisthesis					

Authors & Year	Population, n	Sex, %	Mean Age (Range, SD)*	Diagnosis	Surgery	Bracing	Outcome Factors Assessed	Follow-up	Drop-out, n†
				(n = 12), isthmic spondylolisthesis					
				(n = 5), junctional syndrome (n = 2), pseudarthrosis (n = 2), iatrogenic/post-op instability (n = 1), congenital stenosis (n = 1)					
				††					

- 1 ACDF = anterior cervical discectomy and fusion; CROM = cervical range of motion; DPQ = Dallas pain questionnaire; FES = falls efficacy scale; JOA = Japanese Orthopedic Association; MCS = mental component score; NDI= neck disability index; NR = not
- 2

- 1 reported; PCS = physical component score; ROM = range of motion; SD = standard deviation; SF-36 = short form-36; VAS = visual
2 analog scale
- 3 * Age in years
- 4 † Drop-out before end of follow-up
- 5 ‡ Postoperative care, including immobilization techniques and activity restrictions, was left to the discretion of the attending surgeon
- 6 § Included revisions, removals, reoperations, supplemental fixations, and external bone growth stimulators
- 7 ¶ Included surgical site infection, epidural hematoma, and C5 palsy
- 8 ** Based on patients who completed follow-up
- 9 †† Included later-stage revision surgery due to symptomatic nonunion or later-stage hardware removal due to prominence/bursitis
- 10 ‡‡ Included intraoperative incidental durotomy, intraoperative pedicle-screw-placement issues, new postoperative radiculopathy, early
11 postoperative pulmonary embolism, wound seroma/hematoma, deep wound infection or persistent lumbar radiculopathy
12 postoperatively

1 **Table 3. Results of statistical analysis**

Authors & Year	Statistical Analysis	Differences at Baseline	Postoperative Differences
Abbott et al., 2013	ANCOVA with adjustment for covariates, repeated measure analysis of covariance	There were no significant differences between the cervical collar group and the control group with respect to gender, age, diagnosis, level of operation, cervical range of motion, unipedal balance and baseline NDI, SF-36 (subscales, PCS and MCS), FES and Borg CR-10 (neck and arm pain) scores ($p = \text{NR}$)	ANCOVA: <i>NDI Scores at 1.5 months ($p = 0.042$)*</i> Mean difference between groups: -4.4 (95% CI: -8.6 to -0.2); Cohen's effect size: -0.77 <i>SF-36 BP scores</i> <i>6 months ($p = 0.016$)*</i> Mean difference between groups: 21.4 (95% CI: 4.4 to 38.5); Cohen's effect size: 0.73 <i>12 months ($p = 0.031$)*</i> Mean difference between groups: 17.5 (95% CI: 1.7 to 33.2); Cohen's effect size: 0.57 <i>SF-36 SF scores at 12 months ($p = 0.049$)*</i>

Mean difference between groups: 16.5 (95% CI: 0.1 to 32.9); Cohen's effect size: 0.45

SF-36 PCS

*1.5 months (p = 0.025)**

Mean difference between groups: 5.8 (95% CI: 0.8 to 10.7); Cohen's effect size: 0.84

*3 months (p = 0.038)**

Mean difference between groups: 6.8 (95% CI: 0.4 to 13.1); Cohen's effect size: 0.63

*6 months (p = 0.017)**

Mean difference between groups: 7.4 (95% CI: 1.4 to 13.4); Cohen's effect size: 0.80

*12 months (p = 0.041)**

Mean difference between groups: 7.5 (95% CI: 0.3 to 14.6); Cohen's effect size: 0.66

There were no significant differences between the cervical collar group and the control group with respect to

- 1) neck pain, arm pain, FES, SF-36 PF, SF-36 RP, SF-36 GH, SF-36 VT, SF-36 RE, SF-36 MH, SF-36 MCS at all time points assessed (1.5, 3, 6, 12, and 24 months) ($p > 0.05$)
- 2) all components of the unipedal balance test (right/ foot, right/left foot soft surface, right/left foot eyes closed) at all time points assessed (1.5, 3, 6, 12, and 24 months) ($p > 0.05$)
- 3) all components of CROM (right/left lateral flexion, flexion, extension, right/left rotation) at all time points assessed (1.5, 3, 6, 12, and 24 months) ($p > 0.05$)
- 4) fusion rates and sagittal alignment ($p = \text{NR}$)

Repeated measures analysis of covariance showed that controlling for the combined effects of all prospective measures gave significantly better outcome for the cervical collar group in neck pain ($p = 0.038$), SF-36 PCS ($p = 0.010$) and SF-36 BP ($p = 0.029$).

Campbell et al., 2008	ANOVA, ANCOVA, Fisher exact test, Student t test	SF-36 PCS was higher in the control group (31.1 ± 7.2) than in the braced group (33.4 ± 7.8) ($p = 0.019$)	<p><i>Mean improvement in NDI Scores at 1.5 months ($p = 0.008$)</i></p> <p>Braced: 21.6 ± 18.4</p> <p>Not Braced: 28.4 ± 19.0</p>
		<p>There were no significant differences between the braced group and the control group with respect to age ($p = 0.367$), gender ($p = 0.447$), worker's compensation ($p = 0.458$), litigation ($p = 1.000$), smoking status ($p = 1.000$)</p>	<p>There were no significant differences between the braced group and the control group with respect to NDI scores at 3-months ($p = 0.468$), 6-months ($p = 0.169$), 12-months ($p = 0.415$) and 24-months ($p = 0.693$), SF-36 PCS at 6-months ($p = 0.481$), 12-months ($p = 0.260$)</p>

and occupational status ($p = 0.695$), and baseline NDI ($p = 0.141$), neck pain ($p = 0.523$) and arm pain ($p = 0.710$) scores.

and 24-months ($p = 0.279$), average neck pain scores at 24-months ($p = 0.622$), and average arm pain scores at 24-months ($p = 0.260$).

There were no significant differences in fusion success at any time period between groups, though higher rates of fusion were reported in the non-braced group ($p = 0.552$ at 24-months).

There were no significant differences in rates of secondary surgeries or procedures between the braced group and the control group: revisions ($p = 0.653$), removals ($p = 0.724$), reoperations ($p = 1.000$), supplemental fixations ($p = 0.286$), any surgery ($p = 0.184$).

Christodoulou NR

Mean curves ($p = \text{NR}$)

Postoperative mean curves ($p = \text{NR}$)

et al., 1987	Braced: 58.0° Not Braced: 54.0°	Braced: 23.0° Not Braced: 22.8° There was no significant difference between the braced group and the control group with respect to the mean loss of correction at 24 months (7.0° for braced group, 6.3° for control group, p = NR).
Hida et al., 2017	Two-way repeated ANOVA, Fisher exact test, Student t test, x ² test, There were no significant differences between the collar-fixation group and the control group with respect to age (p = 0.73), gender (p = 0.26), height (p = 0.59), weight (p = 0.66), operation time (p = 0.57), intraoperative blood loss (p = 0.69), number of operated levels (p = 0.67), VAS (p = 0.33), JOA (p = 0.67), lordotic angle (p =	There were no significant differences between the collar-fixation group and the control group with respect to 1) VAS (p = 0.487), JOA recovery rates (p = 0.80), SF-36 PCS (p = 0.537), SF-36 MCS (p = 0.504), and SF-36 BP subscores (p = 0.848) at 12 months follow-up. 2) the decrease in the C2-7 lordotic angle (p = 0.82) and ROM (p = 0.61).

0.84), ROM (p = 0.88) and SF-36 PCS (p = 0.68), MCS (p = 0.80) and BP (p = 0.57).
 3) incidence of complications (p = 0.53).

<p>Yee et al., 2008</p>	<p>Mann-Whitney U test, chi-square test, Fisher exact test, two-way ANOVA</p>	<p>There were no significant differences between the brace group and the control group with respect to age (p = 0.97), gender (p = 0.35), CCI (p = 0.6), number of levels included in the arthrodesis (p = 0.42), smoking status (p = 0.89), worker's compensation or litigation (p = 0.48), revision surgery (p = 0.88), BMI (p = 0.74), diagnosis (p = 0.8), and preoperative SF-36 MCS (p = 0.9), SF-36 PCS (p = 0.19), SF-36 domain scores (p > 0.05) and DPQ category scores (p > 0.05)</p>	<p>There were no significant differences between the brace group and the control group with respect to</p> <ol style="list-style-type: none"> 1) the distribution of surgical complications or subsequent revision rates (p = 0.8) 2) postoperative DPQ category scores (p = 0.34 for the daily activity category, p = 0.67 for the work/leisure category, p = 0.17 for the anxiety-depression category and p = 0.40 for the social category) 3) postoperative SF-36 domain and component scores (p = 0.38 for PF, p = 0.28 for BP, p = 0.23 for GH, p = 0.41 for RP, p = 0.25 for VT, p = 0.79 for SF, p = 0.86 for RE, p = 0.30 for MH,
-----------------------------	---	--	--

p = 0.30 for PCS and p = 0.57 for MCS)

4) rates of fusion seen radiographically at 12 months (p = 0.8) or 24 months (p = 0.9) postoperatively.

5) Rates of revision surgery due to symptomatic non-union (p = 0.43)

- 1 ANCOVA = analysis of covariance; ANOVA = analysis of variance; BMI = body mass index; BP = bodily pain; CCI = Charlson
2 comorbidity index; CI = confidence interval; CROM = cervical range of motion; DPQ = Dallas pain questionnaire; FES = falls
3 efficacy scale; GH = general health; JOA = Japanese Orthopedic Association; MCS = mental component score; MH = mental health;
4 NDI = neck disability index; NR = not reported; OR = odds ratio; PCS = physical component score; PF = physical functioning; RE =
5 role limitations emotional; ROM = range of motion; RP = role limitations physical; SF = social functioning; SF-36 = short form-36;
6 VAS = visual analog scale; VT = vitality
- 7 * No significant differences at all other time points assessed (1.5, 3, 6, 12, and 24 months postoperative)

1 **Table 4. Summary and Strength of Evidence**

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
KQ1: What is the efficacy and effectiveness of postoperative bracing compared with no bracing based on disability, pain, quality of life and functional outcomes?			
Degenerative cervical myelopathy or radiculopathy	n = 2(Campbell, Carreon et al. 2009, Abbott, Halvorsen et al. 2013) n = 3(Campbell, Carreon et al. 2009, Abbott, Halvorsen et al. 2013, Hida, Sakai et al. 2017)	Insufficient Insufficient Low Insufficient Insufficient Insufficient	Results were inconsistent across studies: 1) a prospective cohort study reported that patients in the non-braced group had better NDI scores at 6 weeks after surgery compared to patients in the braced group,(Campbell, Carreon et al. 2009) whereas 2) a pilot randomized controlled trial indicated that patients in the braced group had better NDI scores at 6 weeks after surgery compared to patients in the non-braced group.(Abbott, Halvorsen et al. 2013) Results were inconsistent across studies: 1) a pilot randomized controlled trial indicated that patients in the non-
NDI			
SF-36 PCS			
SF-36 MCS			
SF-36 BP subscale			
SF-36 SF subscale			
SF-36 other subscales			
VAS			
JOA recovery rate			
FES			

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
Unipedal balance standing test	2013, Hida, Sakai et al. 2017)	Insufficient	braced group had better SF-36 PCS scores at 6 weeks, 3, 6,
Neck pain	n = 1(Abbott, Halvorsen et al.	Moderate	and 12 months after surgery compared to patients in the
Arm pain	2013)	Moderate	braced group,(Abbott, Halvorsen et al. 2013) whereas 2) two
	n = 1(Abbott, Halvorsen et al.		studies (a prospective cohort study and a randomized
	2013)		controlled trial) reported no significant differences in SF-36
	n = 1(Hida, Sakai et al. 2017)		PCS scores between the braced and non-braced groups at all
	n = 1(Hida, Sakai et al. 2017)		time points assessed.(Campbell, Carreon et al. 2009, Hida,
	n = 1(Abbott, Halvorsen et al.		Sakai et al. 2017)
	2013)		
	n = 1(Abbott, Halvorsen et al.		Results were inconsistent across studies: 1) a pilot
	2013)		randomized controlled trial indicated that patients in the
	n = 2(Campbell, Carreon et al.		braced group had better SF-36 BP subscale scores at 6 and
	2009, Abbott, Halvorsen et al.		12 months after surgery compared to patients in the non-
	2013)		braced group,(Abbott, Halvorsen et al. 2013) whereas 2) a
	n = 2(Campbell, Carreon et al.		randomized controlled trial reported no significant

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
Degenerative disease of the lumbar spine	2009, Abbott, Halvorsen et al. 2013)	Low	<p>differences in SF-36 BP subscale scores between the braced and non-braced groups at all time points assessed.(Hida, Sakai et al. 2017)</p> <p>Patients in the braced group had better SF-36 SF subscale scores at 12 months after surgery compared to patients in the non-braced group.(Abbott, Halvorsen et al. 2013)</p> <p>There were no significant differences in SF-36 MCS and other subscales, VAS, JOA recovery rate, FES, unipedal balance standing test, neck pain, and arm pain between the braced and non-braced groups at all time points assessed.</p>
DPQ	n = 1(Yee, Yoo et al. 2008)	Low	There were no significant differences in DPQ, SF-36 PCS,

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
SF-36 PCS	n = 1(Yee, Yoo et al. 2008)	Low	MCS and subscales between the braced and non-braced groups at all time points assessed.
SF-36 MCS	n = 1(Yee, Yoo et al. 2008)	Low	
SF-36 subscales	n = 1(Yee, Yoo et al. 2008)	Low	
KQ2: What is the impact of postoperative bracing compared with no bracing on radiographic outcomes?			
Degenerative cervical myelopathy or radiculopathy	n = 2(Abbott, Halvorsen et al. 2013, Hida, Sakai et al. 2017) n = 2(Campbell, Carreon et al. 2009, Abbott, Halvorsen et al. 2013) n = 1(Abbott, Halvorsen et al. 2013) n = 1(Hida, Sakai et al. 2017)	Low	There were no significant differences in ROM, fusion rate, sagittal alignment, and decrease in lordotic angle C2-7 between the braced and non-braced groups at all time points assessed.
ROM		Low	
Fusion rate		Moderate	
Sagittal alignment		Low	
Lordotic angle C2-7		Low	
Degenerative disease of the			There was no significant difference in fusion rate between

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
lumbar spine			the braced and non-braced groups at all time points assessed.
Fusion rate	n = 1(Yee, Yoo et al. 2008)	Low	
Adolescent idiopathic scoliosis with thoracic curves $\geq 35^\circ$			There was no significant difference in mean loss of spinal curve correction between the braced and non-braced groups
Spinal curve	n = 1(Christodoulou, Prince et al. 1987)	Low	at all time points assessed.
KQ3: What is the safety profile of postoperative bracing compared with no bracing?			
Degenerative cervical myelopathy or radiculopathy			There were no significant differences in rate of revision surgery, second procedure or complications between the
Revision surgery or second procedure	n = 1(Campbell, Carreon et al. 2009)	Low	braced and non-braced groups at all time points assessed.
Complications	n = 2 (Campbell, Carreon et al. 2009, Hida, Sakai et al. 2017)	Moderate	
Degenerative disease of the			There were no significant differences in rate of revision

Patient Population and Outcome of Interest	Studies	Strength of Evidence	Overall Effect and Conclusions
lumbar spine Revision surgery or second procedure	n = 1(Yee, Yoo et al. 2008)	Low	surgery, second procedure or complications between the braced and non-braced groups at all time points assessed.
Complications	n = 1(Yee, Yoo et al. 2008)	Low	
KQ4: What is the cost-effectiveness of postoperative bracing?			
None	n = 0	NA	None


1 CROM = cervical range of motion; DPQ = Dallas pain questionnaire; FES = falls efficacy scale; JOA = Japanese Orthopedic Association; MCS =
2 mental component score; NDI = neck disability index; NA = not applicable; PCS = physical component score; ROM = range of motion; SF-36 =
3 short form-36; VAS = visual analog scale

4

5

6

7



1

2 Figures.JPG