

Spectropolarimetry of the massive post-Red Supergiants IRC +10420 and HD 179821

M. Patel^{1,2}, R.D. Oudmaijer¹, J. S. Vink^{3,4}, J.E. Bjorkman⁵, B. Davies^{1,6,7}
M.A.T. Groenewegen⁷, [A.S. Miroshnichenko](#)⁸, J.C. Mottram¹

¹School of Physics & Astronomy, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK

²Astrophysics Group, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, UK

³Armagh Observatory, College Hill, Armagh BT61 9DG, Northern Ireland, UK

⁴Lennard-Jones Laboratory, Astrophysics, Keele University, Keele ST5 5BG, UK

⁵Ritter Observatory, M.S. 113, Department of Physics and Astronomy, University of Toledo, Toledo, OH 43606-3390, USA

⁶Chester F. Carlson Center for Imaging Science, Rochester Institute of Technology, 54 Lomb Memorial Drive, Rochester, NY 14623-5604, USA

⁷Instituut voor Sterrenkunde, K. U. Leuven, Celestijnenlaan 200 D, 3001 Leuven, Belgium

⁸Department of Physics and Astronomy, University of North Carolina at Greensboro, P.O. Box 261 70, Greensboro NC 27402-6170, USA

ABSTRACT

We present medium resolution spectropolarimetry and long term photopolarimetry of two massive post-red supergiants, IRC +10420 and HD 179821. The data provide new information on their circumstellar material as well as their evolution. In IRC +10420, the polarization of the H α line is different to that of the continuum, which indicates that the electron-scattering region is not spherically symmetric. The observed long term changes in the polarimetry can be associated with an axi-symmetric structure, along the short axis of the extended reflection nebulosity. Long term photometry reveals that the star increased in temperature until the mid-nineties, after which the photospheric flux in the optical levelled off. As the photometric changes are mostly probed in the red, they do not trace high stellar temperatures sensitively. And so, it is not obvious whether the star has halted its increase in temperature or not. For HD 179821 we find no polarization effects across any absorption or emission lines, but observe very large polarization changes of order 5% over 15 years. During the same period, the optical photometry displayed modest variability at the 0.2 magnitude level. This is unexpected, because large polarization changes are generally accompanied by strong photometric changes. Several explanations for this puzzling fact are discussed. Most of which, involving asymmetries in the circumstellar material, seem to fail as there is no evidence for the presence of hot, dusty material close to the star. A caveat is that the sparsely available near-infrared photometry could have missed periods of strong polarization activity. Alternatively, the variations can be explained by the presence of a non-radially pulsating photosphere. Changes in the photometry hint at an increase in temperature corresponding to a change through two spectral subclasses over the past ten years.

Key words: Techniques: polarimetric - Stars: evolution - Stars: circumstellar environment - Stars individual : IRC +10420, HD 179821

1 INTRODUCTION

Due to the steepness of the Initial Mass Function, massive stars ($> 8 M_{\odot}$) are extremely rare. This is exacerbated by their comparatively short lifetimes. Yet, although rare, these objects have a crucial impact on the interstellar medium due to their strong winds and high mass-loss rates, and can dominate the light output of entire galaxies.

An area of current interest is that massive evolved

stars are often surrounded by bi-polar nebulae (Weis 2003). Later in the evolution of a star, using spectropolarimetry, it is now established that the ejecta of supernovae deviate from spherical symmetry (e.g. Wang et al. 2003; Leonard et al. 2005). It is as yet unclear whether this is due to asymmetric explosions, axi-symmetric stellar winds or a pre-existing density contrast in the surrounding material (e.g. Dwarkadas & Balick 1998; Dwarkadas & Owocki 2002). Wind-axisymmetry may imply fast rotation, and be

related to the beaming of SN explosions, which may be the origin of the extremely luminous, beamed gamma-ray bursts (e.g. Mészáros 2003; Mazzali et al. 2003).

Here, we address the issue by investigating the circumstellar ejecta of two yellow hypergiants, IRC +10420 and HD 179821. These objects are thought to have evolved off the post-Red Supergiant branch and are still surrounded by mass ejected during a previous mass losing phase. Only a few yellow hypergiants are known (see the review by de Jager 1998), and the number of such hypergiants with circumstellar dust is even smaller - only IRC +10420 and HD 179821 belong to this class (see for example Oudmaijer et al. 2008). Therefore, the study of these two unique objects is important in its own right.

As these stars are distant (3-5 kpc), the direct imaging of their innermost regions is currently beyond the reaches of current technology, although interferometry is starting to resolve the winds of evolved stars (de Wit et al. 2008). Observing with spectropolarimetry allows us to probe regions much closer to the star still. Spectropolarimetry was first effectively used in the study of classical Be stars using the presence of ‘line effects’ (Poeckert 1975; Poeckert & Marlborough 1976). These are changes in polarization across spectral lines that have an emission component. They occur because emission-line photons arise over a larger volume than the stellar continuum photons. Consequently, the emission-line photons undergo fewer scatterings as they ‘see’ fewer electrons, resulting in a lower polarization than the continuum. We normally only observe a net polarization change if the geometry of the electron scattering region is aspherical. Many authors have confirmed that this technique provides evidence that envelope geometries around Be stars are indeed disk-like (Dougherty & Taylor 1992; Quirrenbach et al. 1997; Wood et al. 1997). More recently the technique has been used to investigate the geometry of circumstellar material around Herbig Ae/Be stars. Studies by Oudmaijer & Drew (1999) and Vink et al. (2002) show that most Herbig stars exhibit line effects, indicating aspherical electron-scattering regions. For a recent review, see Oudmaijer (2007). With regard to evolved stars, Davies et al. (2005) conducted a study of Luminous Blue Variables (LBVs) using spectropolarimetry. They found that 50% of the objects observed exhibited polarization changes across $H\alpha$, indicating that some asphericity lies at the base of the stellar wind. Furthermore, they found several objects for which the position angle varied randomly with time, leading them to conclude that the wind around these stars is clumpy (see also Nordsieck et al. 2001).

IRC +10420 is now well accepted as a massive, evolved object (e.g. Jones et al. 1993; Oudmaijer et al. 1996 and Humphreys et al. 2002). This is mainly based on its large distance, high outflow velocity (40 kms^{-1}) and high luminosity implied from the hypergiant spectrum. The situation for HD 179821 is less certain. The presence of non-radial pulsations coupled with comparatively modest photometric changes suggest a massive nature (Le Coroller et al. 2003). Furthermore, its circumstellar material has a large expansion velocity of 30 km s^{-1} , as measured in CO, suggesting the star is a supergiant (Kastner & Weintraub 1995). On the other hand, the overabundance of s-process elements and the low metallicity suggest HD 179821 is perhaps a lower mass

post-AGB star (Zacs et al. 1996; Reddy & Hrivnak 1999; Thévenin et al. 2000).

Although the latter’s nature is a bit more uncertain, there are some striking similarities between IRC +10420 and HD 179821. When observed as part of a larger sample of post-AGB stars, these two objects are often markedly different from the rest. In particular, their high outflow velocities (the average outflow velocity for post-AGB and AGB stars is 15 kms^{-1}) require much higher luminosities if powered by radiation pressure alone (Habing et al. 1994). They were the only objects that showed extensive reflection nebulae in a large survey by Kastner & Weintraub (1995). Jura et al. (2001) point out the enormous difference between the space velocities of IRC +10420 and HD 179821 when compared against low mass post-AGB stars. Furthermore, both objects have an exceptionally strong OI 7774 triplet absorption feature indicating a high luminosity (Humphreys et al. 1973 and Reddy & Hrivnak 1999, based on Slowik & Peterson 1995). We therefore proceed with both objects and assume they are evolved post-Red Supergiants.

Recently, both objects have been observed at arcsecond resolution in CO by Castro-Carrizo et al. (2007) who found, in accordance with previous estimates, that their mass loss rates exceeded $10^{-4} M_{\odot} \text{yr}^{-1}$ when they were in the RSG phase. The envelopes show mild deviations from spherical symmetry in their data.

This paper is organised as follows. In Section 2, we review the experimental setup and explain how the data has been reduced. We present our results for each object in Section 3, and use the new spectropolarimetric data together with past polarization measurements to investigate the nature of the circumstellar environments around each of the stars, which is discussed in Section 4. We conclude in Section 5.

2 OBSERVATIONS

We describe our most recent data in detail. Similar observations taken earlier are briefly discussed, with relevant differences highlighted. The linear spectropolarimetric data were taken on the night of 30 September 2004 using the ISIS spectrograph on the 4.2m William Herschel Telescope (WHT), La Palma. A MARCONI2 CCD detector with a R1200R grating was used. This yielded a spectral coverage of 6150-6815 Å and gave a spectral resolution of 34 kms^{-1} at $H\alpha$. The seeing was about 2'', and a slit with a width of 1 arcsec was used. The linear polarimetric component of the data was analysed using the polarization optics equipment present on the ISIS spectrograph. The object and the sky background were simultaneously observed using additional holes in the dekker mask. A calcite block was then used to split the rays into two perpendicularly polarized beams (the o and e rays). A complete data set therefore, consists of four spectra observed at 0° and 45° (to measure Stokes Q) and 22.5° and 67.5° (to measure Stokes U).

All data reduction steps used the FIGARO software maintained by Starlink and included bias subtraction, cosmic ray removal, bad pixel correction, spectrum straightening and flat fielding. Wavelength calibration was carried out using observations of a Copper-Argon lamp taken throughout the run. The data were then imported into the package CCD2POL

Object	Telescope	Date	Julian Date	%P	P.A. (Deg)	%P (H α)	P.A. (H α) (Deg)
HD 179821							
	AAT	15/09/02	2452533	1.99 \pm 0.11	40 \pm 2		
	WHT	30/09/04	2453279	2.00 \pm 0.15	36 \pm 3		
IRC +10420							
	NOT	16/01/98	2450830	1.95 \pm 0.05	174 \pm 1		
	NOT	16/05/98	2450950	1.80 \pm 0.03	174 \pm 1		
	AAT	18/09/02	2452536	2.12 \pm 0.11	173 \pm 2	1.28 \pm 0.11	10 \pm 2
	AAT	15/08/03	2452867	2.35 \pm 0.11	179 \pm 1	1.61 \pm 0.11	11 \pm 1
	WHT	30/09/04	2453279	3.40 \pm 0.15	174 \pm 2	1.93 \pm 0.15	11 \pm 2

Table 1. New polarimetric observations of HD 179821 and IRC +10420 measured in the R band. For the spectropolarimetric data (those taken at the AAT and WHT), the polarization was measured in the continuum region close to H α , while the final columns are the polarizations at the line centers. The H α line-center polarization of HD 179821 was not calculated as no line effect was observed, and therefore this measurement would be equal to the continuum polarization. The systematic error of the AAT and WHT spectropolarimetric data is estimated to be of order 0.10% and 0.15%, respectively.

Julian Date	Telescope	$B - V$	V	$V - R$	$R - I$	J	H	K
2450268	CST					5.62		
2450303	CST					5.36	4.40	3.45
2450643	CST					5.40	4.44	3.50
2450690	CST					5.40	4.45	3.49
2450707	CST					5.37	4.43	3.38
2450950	NOT	2.76	11.06	2.40	1.70			
2450985	CST					5.38	4.44	3.48
2451037	TSAO	2.58	11.12	2.42	1.60			
2451039	TSAO	2.67	11.11	2.41	1.58			
2451042	TSAO	2.58	11.13	2.42	1.54			
2451043	TSAO	2.73	11.06	2.49	1.61			
2451047	TSAO	2.51	11.01	2.41	1.58	5.63	4.55	3.73
2451048	TSAO	2.75	11.03	2.47	1.60	5.43	4.57	3.58
2451050	TSAO	2.67	11.01	2.46	1.66			
2451052	TSAO	2.75	11.03	2.47	1.59	5.35	4.44	3.53
2451057	TSAO	2.76	10.96	2.47	1.64			
2451063	TSAO	2.60	10.95	2.43	1.59			
2451075	TSAO	2.70	11.16	2.52	1.60			
2451082	TSAO	2.66	11.09	2.47	1.59			
2451083	TSAO	2.72	11.05	2.48	1.60			
2451087	CST					5.24	4.29	3.33
2451099	TSAO	2.62	11.16	2.56	1.66			
2451100	TSAO	2.71	11.16	2.54	1.66			
2451103	TSAO	2.76	11.13	2.51	1.54			
2451104	TSAO	2.72	11.03	2.48	1.59			
2451147	CST					5.35	4.42	3.46
2451245	TSAO					5.32	4.47	3.56
2451292	CST					5.40	4.45	3.57

Table 2. Photometry of IRC +10420. The data come from the Carlos Sánchez Telescope (CST), the Nordic Optical Telescope (NOT) and the Tien-Shan Astronomical Observatory (TSAO), see text for details. Typically, the photometric errors are of order 0.01-0.03 magnitude.

(also maintained by Starlink), to produce the Stokes Q and U parameters. The degree of polarization and its PA can then be found from

$$p^2 = Q^2 + U^2 \quad (1)$$

$$\theta = 0.5 \times \arctan\left(\frac{U}{Q}\right) \quad (2)$$

Finally, the data was position angle-calibrated using a set of polarization standard stars observed during the run.

Polarimetric data should only be limited by photon statistics. For example we expect an error of about 0.1% for

a detection of 10^6 photons, and significantly smaller when large parts of the spectra are binned. However, the presence of systematic errors such as scattered light and instrumental polarization often outweigh these photon-noise errors and we estimate the errors in the present data to be of order 0.15%. The observations of two zero-polarization objects show that the instrumental polarization is less than 0.15%. We have made no attempt to correct for either instrumental or interstellar polarization (ISP), as these only add a constant (Q,U) vector to the data and therefore will not affect whether we observe a line effect.




Figure 1. Spectropolarimetry of IRC +10420 around $H\alpha$ at different epochs. The upper diagrams are ‘triplots’ of the spectropolarimetric data, which were taken one year apart from one another. The bottom panel of the triplot presents the direct intensity spectrum, the data normally obtained from spectroscopy. The middle panel shows the polarization measured as a function of wavelength. The upper panel shows the corresponding polarization angle. The data are adaptively binned such that each bin has the same observational error derived from Poisson statistics. Here we have chosen errors of 0.1%, 0.1% and 0.3% respectively for the 2002, 2003, and 2004 data. Below the triplots, the Stokes Q,U vectors are plotted against each other, and binned to the same precision as the spectra. Depolarization around $H\alpha$ is seen each time and the associated excursions in Q-U space all point in the same direction.

2.1 Additional Data

2.1.1 Spectropolarimetry

Further data of IRC +10420 and HD 179821 was obtained during previous observing runs. Spectropolarimetric observations of IRC +10420 were obtained using the 3.9m Anglo-Australian Telescope (AAT) on 18/9/2002 and 15/8/2003. HD 179821 was observed on 15/9/2002 at the AAT. The 2002 data are taken with a lower spectral resolution (2.1 Å) and the 2003 data are taken at comparable resolution to the current WHT data. We refer to Davies et al. (2005) for more details on the observations taken at the AAT in 2002 and 2003.

2.1.2 Photometry and polarimetry

We also obtained additional photometric and polarimetric data of IRC +10420. Hitherto unpublished broad band polarimetry was obtained in January and May 1998 using the 2.5m Nordic Optical Telescope in La Palma, and on one of these nights *BVRI* photometry were obtained. These ob-

servations are described in Oudmajer et al. (2001). Johnson *BVRIJK* observations were obtained in 1998 on the 1 m telescope at the Tien-Shan Astronomical Observatory (Kazakhstan) with the 2d-channel photometer-polarimeter FP3U of the Pulkovo Observatory (Bergner et al. 1988). Further near-infrared photometry was obtained employing the 1.5m Carlos Sánchez Telescope in Tenerife. Details of the observing procedures can be found in Kerschbaum et al. (2006). The results are presented in Tables 1 and 2. *R* band polarimetric data and optical and near-IR photometric data taken from the literature, for both objects, is described in Tables 3 and 4 respectively.

3 RESULTS

3.1 IRC +10420

3.1.1 Spectropolarimetry

In Fig. 1 we have plotted the polarimetric data around $H\alpha$ taken in 2002, 2003 and 2004. These data show ‘triplots’

Object	Date	N	Source	Method
HD 179821	May 1989 - May 1991	4	Parthasarathy et al. (2005)	Broadband
	October 1991	1	Trammell et al. (1994)	Spectropolarimetry
	August 1993	1	HPOL database, Johnson (priv. communication)	Spectropolarimetry
	June 1997 - October 1998	30	Melikian et al. (2000)	Broadband
IRC +10420	May 1976 - June 1976	1	Craine et al. (1976)	Broadband
	May 1989	1	Johnson & Jones (1991)	Broadband
	October 1991	1	Jones et al. (1993)	Broadband
	October 1991	1	Trammell et al. (1994)	Spectropolarimetry

Table 3. *R* band polarimetric observations of HD 179821 and IRC +10420 taken from the literature. The observational period is given in the second column. The last three columns show the number of observations taken, the literature source of the data and the method. Although Craine et al. (1976) made several observations of IRC +10420, these values were averaged to a single data point in our study. We measured the continuum close to $H\alpha$ from the spectropolarimetric data in order to facilitate comparison with the *R* band data.

Object	Date	Band	N	Source
HD 179821	March 1988 - April 1988	<i>V, J, K</i>	1,2	Hrivnak et al. (1993)
	May 1990 - October 1999	<i>V</i>	169	Arkipova et al. (2001)
	May 1990 - November 2005	<i>V</i>	254	ASAS catalogue
	June 1992	<i>J, K</i>	1	Kastner & Weintraub (1995)
	October 1998	<i>J, K</i>	1	2MASS database
IRC +10420	May 1996 - June 1996	<i>J, K</i>	2,2	Humphreys et al. (1997)
	April 2001	<i>J, K</i>	4,5	Kimeswenger et al. (2004)
	April 2002 - October 2004	<i>V</i>	104	ASAS Catalogue

Table 4. A summary of recent photometric observations of IRC +10420 and HD 179821. In the fourth column we give the number of observations taken in each passband, respectively. For IRC +10420, this acts as an update to the previous tables compiled by Jones et al. (1993) and Oudmaijer et al. (1996).




Figure 2. Left : The polarization and position angle variability of IRC +10420 in the *R*-band. Both the polarization and PA show an increase with time. The bottom 2 panels show the evolution of the Stokes *Q* and *U* vectors. Right : An updated version from the photometric variations presented in Oudmaijer et al (1996). The change in the *J* band, which traces the stellar photosphere, indicates a temperature increase of the star. Also included is the evolution of the $H\alpha$ Equivalent Width which displays a marked strengthening over the years.




Figure 3. A Q-U diagram of IRC +10420. The *R*-band (Q,U) data-points come from the literature and our new broadband polarimetry. The dashed line represents the vector towards the ISP, denoted by the crossed circle, as derived by Jones et al. (1993). The continuum values from the spectropolarimetric data are represented by dots and the straight lines show the excursions to the line center polarization measured from the spectropolarimetry.

of the spectropolarimetry, along with graphs plotting the Stokes Q,U vectors against each other.

IRC +10420 displays $H\alpha$ emission with a total equivalent width (EW) of -50 \AA . The emission line is unresolved in the lower resolution 2002 data. The other, higher resolution, data indicate a double-peaked and variable emission-line profile. In 2003, the red peak is stronger than the blue peak, but in 2004 the blue peak has become the stronger, while the central absorption component of the double-peaked line has become deeper. Comparison with published data reveals that the red peak is very strong in the present spectrum. Oudmajer (1998) show data taken in 1993 of the object, where the blue peak is 2 to 3 times stronger than the red one. The EW of the line was -25 \AA , about half that observed now. This evolving blue-to-red peak ratio was also noted by Klochkova et al. (2002).

The spectropolarimetric data show large polarization changes across the line. In each case, this line effect consists of a depolarization over the central part of the line coupled with a rotation to larger angles. The associated Q-U plots (Fig. 1) also clearly show the line effects. The cluster of data points in the lower right hand corners of the graphs are due to the stellar continuum, while the excursions to the upper left hand corners trace the polarization over the $H\alpha$ line. At first glance, the excursions might seem to suggest the presence of structure. However, given that the binning is at the 0.1% and 0.3% level respectively, and that the ‘structure’ is mostly due to single points offset by less than 3σ from the global trends, we proceed under the assumption that the excursions are linear. The intrinsic polarization angle for IRC +10420 is measured by the excursion from the line-center (which has a lower contribution by the polarization due to electron scattering) to the continuum. Using Eq. 2 this is 158° , with an estimated error of 2° , and it does not seem to change with time. The magnitude of the excursion

gives an indication of the strength of the polarization change and is similar in 2002 and 2003, but almost twice as large in 2004. Strictly speaking the length of the vector represents a lower limit to the real depolarization because a finite spectral resolution can wash out the line-effect. However, the 2002, lower resolution data has a depolarization vector of similar length to the higher-resolution, 2003 data. It is therefore plausible that the depolarization (and thus the polarization of the continuum due to electron scattering) is significantly stronger in the 2004 data.

The continuum polarization towards IRC +10420 consists of three components. Firstly, there is interstellar polarization due to dichroic absorption of interstellar dust. This is constant with time and is constant over a small part of the spectrum. In data such as in Fig. 1 this would add a constant vector to the intrinsic Q,U spectrum. Importantly, the shape of the QU behaviour is not affected by the ISP. Secondly, some polarization may be due to scattered light reflected off circumstellar dust grains. The presence of scattered light is revealed by Kastner & Weintraub (1995) who show that the polarization is significant at large distances from the star. Later HST images show substantial extended emission in blue light which is interpreted as scattered light as well (Humphreys et al. 1997, 2002; see also Davies et al. 2007). When obtaining spatially unresolved data, such as here, any net polarization is due to the asymmetry of the scattering material on the sky. Thirdly, the continuum light can be scattered off free electrons that are very close to the star. Here, imaging polarimetry can not achieve the spatial resolution required to study the electron scattering region. The line-effect however readily reveals its presence.

3.1.2 Long term photo-polarimetry

Our new data are combined with literature data and are shown in Figure 2. Over the past 30 years, both the polarization and the angle have increased. To illustrate this, the Q and U Stokes vectors are also plotted. The Stokes Q parameter increases by about 2%, while the U vector essentially remains constant (see also Fig. 3). Here we note the Trammell et al. (1994) data. Their data point (3.7% at 149°) was taken in the same month as the Jones et al. (1993) data which is 2.0% at 174° . The Trammell et al. measurement strongly deviates from the general, 30-year, trend in P , Θ and U and is also significantly different from data taken in the same month. We will proceed assuming that the trend of increasing polarization with time reflects the true long term polarization behaviour of the object, and have a caveat that, perhaps, the evolution is more complex. In Fig. 3 we plot the broad-band Stokes Q and U vectors against one another. Jones et al. (1993) derive an interstellar *R* band polarization of 8 per cent towards IRC +10420 - a value that is not inconsistent with the large reddening towards the object. From their figures we measure their ISP to be $Q_{R,ISP} \approx 7.5\%$, $U_{R,ISP} \approx 2.7\%$. This value is indicated in Fig. 3. Assuming the $H\alpha$ line center is not polarized due to electron scattering, its net polarization is due to a combination of ISP and scattering off the circumstellar dust. We can thus get a (very) rough estimate of the circumstellar dust polarization by taking the difference between the $H\alpha$ line center polarization and the ISP value. Our observed line center polarizations are close each other. Remarkably, the

recent $H\alpha$ line polarization values all lie on the vector connecting the ISP with the origin. Their average distance to the ISP value is ($Q = -6.0\%$, $U = -2.15\%$), corresponding to a circumstellar dust polarization of ($P = 6.4\%$, $\Theta = 9.9^\circ$). The true value may have a larger polarization at a smaller angle, as the polarization in the center of the line is probably affected by the resolution. Note that an interesting situation has arisen where the circumstellar dust polarization effectively cancels part of the ISP resulting in smaller net polarization values. Using Eq. 2, we measure an intrinsic polarization angle for the variations of $6 \pm 2^\circ$. This is very close to the orientation found for the circumstellar dust polarization described above and pinpoints the changes in circumstellar dust polarization as a likely cause for the observed variations.

The photometric variability has been discussed extensively by Jones et al. (1993) and Oudmaijer et al. (1996). Here we give an update on the long term variability (see Table 4). The K band data show a steady fading starting in the mid-seventies. This dimming is mimicked by the data in the J band, which is known to trace the stellar photosphere (Oudmaijer et al. 1996). After around 1995, the J band magnitudes appear to reach a steady state.

Long term changes have also been observed in the hydrogen recombination lines. In 1992, Oudmaijer et al. (1994) discovered NIR recombination emission lines that were previously seen in absorption in 1984. Irvine & Herbig (1986) detected $H\alpha$ emission in 1986, while previous observations did not show emission. Therefore, the onset of hydrogen recombination emission can be traced to a period between 1984 and 1986 (Oudmaijer et al. 1994). This coincides precisely with a period of strong photometric changes.

EW data of $H\alpha$ were collected from the literature as well. It is not trivial to obtain these data because of the large width of the line. For example, EW measurements from high resolution echelle data are not possible as the free spectral range is too small to properly define a baseline and measure the equivalent width accurately. We therefore restrict ourselves to results from spectra with a large free spectral range - essentially no echelle observations. Data are taken from Oudmaijer et al. (1994), Oudmaijer (1998), Humphreys et al. (2002), this paper and Davies et al. (2007) respectively. The results are also shown in Fig. 2. $H\alpha$ has been increasing in strength, going from upper limits of the EW in the seventies to -23 \AA in the early nineties to -50 \AA in 2005. As the optical brightness has remained constant in that time, these EW changes reflect true line flux changes and imply that more ionized material is present.

3.2 HD 179821

3.2.1 Spectropolarimetry

Spectropolarimetric data of HD 179821 are shown in Fig. 4. No polarization changes are visible across the faint $H\alpha$ line. The 2002 data has a higher signal-to-noise, but a weaker $H\alpha$ emission than in 2004. There are few reports of the $H\alpha$ line in the literature, but it would appear it is variable in the sense that it varies between weak emission above the continuum (i.e. 10-20% emission above the continuum was observed by Zacs et al. 1996) and weak emission that hardly fills in the underlying absorption as observed here.

The weak emission implies that the number of free scattering electrons is rather limited, which is not inconsistent with the low surface temperature of the star (its spectral type is G4 0-Ia, Keenan & McNeil 1989). Therefore any continuum polarization due to electron scattering is bound to be low. In such a case, extremely high SNR data are required to reveal a line depolarization, if it were present at all. Alternatively, the hydrogen recombination line emission in HD 179821 could be due to photospheric shocks. The object has been found to show periodic photometric variations of 140 and 200 days (see Le Coroller et al. 2003). These variations are most likely due to pulsations, and it is common to observe $H\alpha$ emission due to shocked layers in the photosphere of cool, pulsating stars (e.g. Schmidt et al. 2004 and references therein).

3.2.2 Long term photo-polarimetry

The photometry and polarimetry from this paper and the literature of HD 179821 are plotted in Fig. 5. The longer term optical variability is discussed by Arkhipova et al. (2001). From 1989 to about 1999 the star became fainter by about 0.1 magnitude, and then returned to its original magnitude. Arkhipova et al. (2001) suggest the photometry may be cyclic on long timescales. They also find that the object is bluer in $U - B$ when the star is brightest in V , which they attribute to pulsations. This is confirmed by the detailed study on shorter term variations in the V band presented by Le Coroller et al. (2003), which are due to pulsations with periods of order hundreds of days. Here we report the hitherto unnoticed large change in the J and K bands. These became fainter by around 0.4 magnitude in the period 1988 - 2000.

Figure 5 shows that HD179821 experiences polarization changes (the star is even observed to exhibit no polarization at all for four consecutive days in November 1997), suggesting that the star is intrinsically polarized. The polarization varies more or less randomly between 0 and 2.5 per cent over the past 15 years, but we can identify two clear values for the PA, a low value around 40° and a high value of around 120° . When plotting these data in QU space (Fig. 6) we can see that the data points are distributed around what we can loosely describe as a straight line through the origin.

The move through the origin of the QU diagram is responsible for the large change in observed polarization angle. This must have happened at least five times: the first occurred between the 1989-1991 Parthasarathy et al. (2005) dataset (PA= 40°) and the 1991 Trammell et al. (1994) data point (PA= 120°) taken half a year later, when the object moved from the upper half to the lower half in the QU diagram. After this, a move occurred between 1991 and the HPOL data taken in 1993 (Johnson private communication, using HPOL, Wolff et al. 1996). The object returned back to the lower half as observed by Melikian et al. (2000), who also directly detected such an excursion in 1997-1998 (the non-detections which lie, by definition, at the origin). Finally a crossing happened after their observing run, but before our present data was taken. The move through the origin does not necessarily reflect a real 90° rotation, as the ISP may contribute significantly to the total Q, U vectors observed towards the object.

It is hard to determine the ISP towards any object,




Figure 4. As in Fig. 1, but now for HD 179821 in 2002 (left) and 2004 (right). The data has been rebinned such that the 1σ error in polarization corresponds to the value stated at the top of the triplot, i.e. 0.2% and 0.5% respectively, as calculated from photon statistics.

but we can make an educated guess. Since $A_V \sim 2$ for HD 179821 (Hrivnak et al. 1989) and one typically finds that the percentage polarization is equal to the extinction in magnitudes (e.g. Oudmaijer et al. 2001), we can expect an ISP of up to 2 per cent towards the object. A value for the interstellar polarization angle may be estimated from the polarization of surrounding field stars. To this end we investigated the compilation of polarized stars due to Heiles (2000). The local ISP appears very ordered towards this part of the sky. We selected all objects with a polarization measurement larger than 0% and within a radius of 5 degrees from the position of HD 179821. These 22 objects have an average polarization angle of 50° with a scatter of 24° . The catalogue also lists the photometric distances, and the average angle is the same for the 11 furthest ($>500\text{pc}$) objects. Whatever the magnitude of the polarization, these values indicate that the ISP can be located in the top quadrants of Fig. 6. When taking these angles at face value, and assuming the ISP is oriented in the same direction at large distances as locally, then the ISP towards HD 179821 is 50° at 2%. This is close to the present-day value of HD 179821, and, if true, would suggest that HD 179821 is presently in a relatively quiescent state, yet the polarization is still variable by up to a few per cent. The instances when the object occupies the bottom of the QU diagram can then be identified as periods where the polarization of HD 179821 is particularly strong. The optical photometry does not signal any such activity however, and continues its gradual decline to fainter magnitudes.

In short, during the last 15 years, the observed polar-

ization of HD 179821 crossed the origin in the QU -diagram several times. The changes occur along a straight line and suggest some preferred orientation, for which we derive a very rough estimate for the intrinsic angle of 38° . In addition, the polarization changes by 4-5%, while only modest changes in the optical photometry of less than 0.2 magnitude occur. Near-infrared photometry is sparse, and indicates a gradual faintening of the star. The most recent data point obtained on 12 October 1998, which was a few weeks before the end of Melikian et al. (2000)’s campaign, i.e. when HD 179821 was, if the ISP value is correct, still in a high state of polarization.

4 DISCUSSION

We have observed the only two post-Red Supergiants with large infrared excesses using multi-epoch spectropolarimetry and supplemented these data with long term photopolarimetry. We find some striking similarities between the two objects, but also large differences. We will start with the latter.

4.1 The ionized gas

The spectropolarimetry reveals a strong line-effect in the data of IRC +10420, but none for HD 179821. The latter displays weak $H\alpha$ emission and is much cooler than IRC +10420. It is therefore not surrounded by much ionized material. As electron-scattering is the prime mechanism to




Figure 5. The variability of HD 179821 with time. Plotted is the V-band photometry provided by Hrivnak et al. (1989) [squares], Arkhipova et al (2001) [triangles] and the ASAS Catalogue (Pojmanski, 2002) [circles]. The error associated with the photometry measurements provided by Arkhipova is 0.005 mag and those from the ASAS Catalogue are less than 0.06 mag. Plotted in the upper two plots are PA and polarization data taken from Parthasarathy et al. (2005) [crosses], HPOL database [triangle], Trammell et al. (1994) [star], Melikian et al. (2000) [circles] and data presented in this paper from 2002 [squares] and 2004 [rhombi]. The PA exhibits large rotations over a 15 year period, while the polarization indicates the presence of short-term variability. The lower panels show that HD 179821 has become fainter in J and K .

produce line-effects, it can then be easily understood why there is no line-effect towards HD 179821.

As described earlier, the intrinsic polarization angle due to electron scattering for IRC +10420 is measured by the excursion from the line-center, and is 158° . This angle remained constant within the errorbars during our three observing epochs, whereas the observed polarization increased by 1.5%. Since the excursion across the line in QU space became larger between 2003 and 2004, we can attribute this increase in continuum polarization to more scatterings of continuum photons by free electrons rather than dust particles.

The presence of the line-effect is strong evidence that the electron scattering region has a geometry that deviates from spherical symmetry. The constant intrinsic polarization angle indicates that the structure has remained stable. The nature of the $H\alpha$ emitting region, and especially its geometry, has been the topic of much debate, often centering on the question whether it is spherical or not (see the review of the literature by Davies et al. 2007). Davies et al. (2007) use integral field spectroscopy to study reflected light off circumstellar dust of the emission line and show that the $H\alpha$

line emitting region is not spherically symmetric. Here we confirm this result.

It is unclear however what the polarization angle corresponds to. Assuming that the scattering is optically thin, the structure responsible for the scattering is oriented perpendicular to this at an angle of 68° . A comparison with the larger scale structure observed by the Hubble Space Telescope data which traces the reflection nebulosity (Humphreys et al. 1997, 2002) is inconclusive. The angle does not correspond straightforwardly with either the long or short axis. The Hubble data reveal very clumpy structures at sub-arcsec scales and it proves impossible to link the even smaller scales probed by the electron scattering, with the clumps or larger scale structures in the imaging.

Finally, the increased ionization in IRC +10420 may be either explained by an increase of mass close to the star, due to either an infall or outflow or due to an increase in temperature.

4.2 Temperature evolution

The photometry shows some remarkable similarities between the two objects. Both stars have been subject to studies of their spectral energy distribution (SED, Oudmaijer et al. (1996) and Hrivnak et al. (1989) for IRC +10420 and HD 179821 respectively). From fits to their SEDs, it is clear that the near-infrared J band photometry samples their photospheres and not the hot dust. In both cases, this J band magnitude has become fainter over the past decades. At the same time, their V band magnitudes varied only weakly, if at all.

The photometric changes in J , in particular the decrease in $V - J$, can be explained as being due to bolometric correction effects associated with an increase in stellar temperature. For IRC +10420, this is consistent with published spectra (see Oudmaijer 1998; Klochkova et al. 1997). This trend is less obvious more recently. It is likely that the J band magnitude has reached a plateau, but it is not apparent whether this implies that the temperature increase has halted or not. The $V - J$ colour is not very sensitive to changes in spectral type earlier than A0, so an ongoing temperature increase is not ruled out based on the photometry.

J band data of HD 179821 is much more sparse, but the trend is clear. The object has become fainter in J . If the change in $V - J$ colour is due to changes in effective temperature, then the star has become hotter during this period, in the same way as for IRC +10420. For an A_V of 2 (Hrivnak et al. 1989), and combining the intrinsic optical colours for supergiants as provided by Straizys & Kuriliene (1981) and intrinsic near-infrared colours due to Koornneef (1983), we find that the star must have evolved to an earlier spectral type by about one or two spectral subtypes.

There is some ambiguity concerning the star's temperature as derived from chemical abundance modelling. For example Thévenin et al. (2000) favour a lower temperature, consistent with the G spectral type. This is 1000K less than Reddy & Hrivnak (1999) and Zacs et al. (1996) determinations using data taken earlier. Thus, it is thus hard to draw any conclusions on the evolution of the stellar temperature. The near-infrared photometric changes may be seen as independent evidence that the star appears to have been evolving towards higher temperatures. Alternatively, the temper-




Figure 6. A Q-U diagram of HD 179821. Each point corresponds to the wavelength-averaged *R*-band continuum values of Q and U taken at different epochs. There are groupings of points in two separate areas of the plot, one corresponding to low PA angles (top) and the other to high angles (bottom). The filled circle shows the position of the ISP (see text). The other symbols are as in the previous figure.

ature changes may be due to the formation of a pseudo-photosphere (Smith et al. 2004).

The main finding in this section is that both objects appear to be evolving towards the blue in the HR diagram on human timescales. Judging from the plateauing of the *J* band photometry, IRC +10420 may have slowed down its temperature evolution, or even halted altogether. It has been proposed that the object is hitting the yellow void (de Jager 1998; de Jager et al. 2001; Humphreys et al. 2002) and/or the red-edge of the bi-stability zone (Smith et al. 2004). In this case, further evolution is prevented unless a large part of its (pseudo-)photosphere is shredded. It will be interesting to follow the object to see whether a large outburst indeed will happen.

4.3 Polarization Changes

A further similarity between the two objects is the significant polarization variability over the past decades. IRC +10420 increased its polarization (as seen in the QU diagram, Fig. 3) by about 2.5%. HD 179821’s polarization changed by more than 5%, following a straight line in QU space (Fig. 6). Such changes are not uncommon in stellar objects, for example UX Ori variables and some T Tauri stars (see e.g. Oudmaijer et al. 2001; Grinin 1994). For these objects, rotating clumpy circumstellar disks can result in large polarization changes. These occur when the clumps move into the line of sight of the objects, reducing the optical brightness, often considerably by many magnitudes, and thereby increasing the contribution of polarized light. However, here, in the case of both objects, the changes are not accompanied by large changes in the optical photometry. To make matters worse, a direct consequence of the UX Ori scenario, to explain the strong polarizations variations, is that the obscuring dust is located close to the star in a compact, hot structure (Dullemond et al. 2003). This means that a strong near-infrared excess should be present and readily observable. IRC +10420 shows evidence for a

significant near-infrared excess due to hot dust. In contrast, the near-infrared photometry of HD 179821 does not show any sign of hot dust close to the star and is best explained by photospheric emission (Hrivnak et al. 1989).

4.3.1 The electron and dust scattering of IRC +10420

The, variable, continuum polarization of IRC +10420 is due to a combination of dust and electron scattering. The intrinsic PA (10°) of the circumstellar dust polarization was derived earlier from the vector connecting the ISP and the line centers of the $H\alpha$ emission. This PA corresponds to the short axis observed in the larger scale nebulosity of IRC +10420 and may signify a circumstellar dusty disk structure. The intrinsic angle of the electron scattering region is 158° and is not easily identified with the larger scale structures observed in the HST images. In the last observing epoch (2004), the line depolarization vector was much stronger than previously observed. Although resolution effects may have affected the amplitude of the excursion, the much larger continuum polarization in 2004 indicates that enhanced electron scattering is responsible for the most recent increase in polarization. This is supported by the fact that the continuum polarization moved along the line vector in the QU diagram (Fig. 3) as well. The PA measured for the QU vector between the two recentmost dates is $166 \pm 3^\circ$, very close to the line polarization vectors. Such an increase is not inconsistent with the fact that the $H\alpha$ emission EW hardly changed. Polarization due to (optically thin) electron scattering is more sensitive than (optically thick) $H\alpha$ emission for tracing ionized material (see e.g. Bjorkman & Meade 2005).

How does this compare to the longer term changes? The polarization evolves along a more or less straight line with a slope corresponding to a PA of $6 \pm 2^\circ$, which closely corresponds to the intrinsic PA of the dust scattering material. If the ISP determined value by Jones et al. (1993) is correct, then it is fair to assume that a decreasing circumstellar dust polarization is the main contributor to the long term variability. A probable cause for this is a clearing dust shell. This can be tested by producing SED model fits to photometric data taken at several epochs.

In summary, by combining the intrinsic PA of both dust and electron scattering and using the ISP, a relatively simple picture seems to emerge from the variability of the polarization of IRC +10420. At first a clearing dust shell dominates the polarization behaviour, followed by a significant increase of polarization due to electron scattering. The larger electron scattering is consistent with the observed increase in $H\alpha$ emission, while the dust clearing can be tested with modelling the circumstellar dust.

This global picture is probably not the final answer however. Even the data discussed here present more questions already. For example, it is not clear why the point taken by Trammell et al. in 1991 strongly deviates from all other values (even one taken in the same month). The $H\alpha$ depolarization they observe is consistent with what we find ten years later, and confirms that part of the polarization is due to electron scattering. The large continuum value would indicate a violent event that led to a polarization increase of several per cent within weeks, however there is no other evidence to suggest such sudden changes in other observations of IRC +10420. Similarly, it is not obvious why the

onset of H α line emission in the early nineties, and its corresponding polarization of 1-1.5%, is not detected in the long term polarimetry, which follows a more or less straight line from the seventies to the current data. A speculation is that a varying, rotating component of electron scattering was present from the onset. Given the increase in H α emission, it is possible that the electron scattering has become more optically thick. If part of the photons scatter more than once in their escape from the electron scattering region, a full or partial flip in the position angle can result (see e.g. Vink et al. 2005).

4.3.2 But what about HD 179821?

The changes observed for HD 179821 are less straightforward to interpret. The major question is how large changes over such an extended period can occur while, apart from the polarization, other observations of the star show mild variability at best. Firstly, we note that the ISP is hardly likely to vary on human timescales. This is supported by the results from a survey of polarization standard stars which reveals no variability in the observed polarization down to extremely low levels (Clarke & Naghizadeh-Khouei 1994). The H α emission is insufficiently strong to suggest the presence of a large electron scattering region, let alone one that it is variable enough to explain the observed polarization changes. Below we discuss the several possibilities that can account for the polarization and its variability observed towards HD 179821.

4.3.2.1 Asymmetric circumstellar material Having been forced to rule out electron scattering as being responsible for the variability, we now consider scattering by circumstellar dust. The presence of dust scattering was shown in the imaging polarimetry by Kastner & Weintraub (1995). The fact that our, unresolved, data show a net polarization would normally indicate that part of the scattering geometry deviates from spherical symmetry. However, this is not supported by existing imaging data of the circumstellar material of the object. The scattering dust traced by Kastner & Weintraub (1995) stretches over several arcseconds, but appears to be distributed spherically symmetrically. HST images published by Ueta et al. (2000) show small scale structure but largely a round appearance. Bujarrabal et al.'s 1992 CO rotational data maps show a slightly elongated structure at larger (~ 5 arcsec) distances from the star at a PA of 50° .

Even if the slight asymmetry would be responsible for the observed net polarization, it is hard to reconcile this with the observed variability. With a kinematic age of order thousands of years (e.g. Kastner & Weintraub 1995), one would not expect changes at a fraction of this period, i.e. 15 years, to still be traceable. A more practical objection to the extended emission being variable is that the intensity of the scattered light rapidly drops as a function of distance to the star, so there is simply not enough light available to induce variations of 4% in polarization in the total light.

A final possibility is that a compact asymmetric structure much closer to the star is responsible for the variable polarization. As mentioned earlier, the SED of the object leaves no room for a significant, hot dusty component. A caveat is

of course that the near-infrared photometry of the object is limited, so any sudden, short-lived events could have been easily missed. If the ISP value is, as the data suggest, in the upper half of the *QU* diagram, the major polarization activity occurs when the object occupies the lower half of the *QU* diagram. A prediction would then be that during such periods, excess radiation in the near-infrared due to hot dust would be present. NIR data of HD 179821 is very sparse, but the last set of *J*, *K* photometry (from 2MASS), was taken during Melikian's observing period. These data follow the gradual faintening of the star in the NIR and do not indicate any excess emission. In addition, the optical photometry appears steady throughout.

For completeness, we do note that Oudmaijer et al. (1995) found variable first-overtone CO emission at $2.3 \mu\text{m}$ towards this object. This emission typically arises in hot, dense material within the dust sublimation radius. Oudmaijer et al. interpreted this as variable mass loss. If the CO emission originates from a geometry that is not spherically symmetric, it could be associated with ejections of dense gas similar to, but shorter than, for example the asymmetric mass loss event observed towards HD 45677 (Patel et al. 2006). The resulting situation could give rise to Rayleigh scattering and a net polarization of light. However, whether this would give rise to the large polarization variations is not clear. We do point out that CO emission from the yellow hypergiant ρ Cas could also be explained by shocks due to pulsations in the stellar photosphere (Gorlova et al. 2006). This brings us to the central star as the origin of the variations.

4.3.2.2 An asymmetric star illuminating the dust shell Above, we considered the case of a central source surrounded by asymmetric scattering media. The asymmetries involved result in a net polarization of the total light. However, we tacitly assumed the central object to be either a point source or an isotropically emitting sphere.

It is well known that when a central source illuminates a spherically symmetric envelope, any asymmetries in the *central source* can induce a net polarization. As for example reviewed by Boyle et al. (1986), who find that a photosphere with a nonuniform surface brightness can result in net polarization when its radiation is scattered off circumstellar dust. Other examples are changes found in the polarization spectra of evolved stars where it has been inferred that different molecular bands cover different parts of the stellar photosphere (Biegging et al. 2006). The polarization changes over molecular bands can be explained by both Rayleigh scattering by particles in clouds high in the atmosphere and scattering due to circumstellar dust (Raveendran 1991). In all these cases the break from symmetry is not due to the circumstellar material, but due to the asymmetric illuminating source.

The extensive photometric observations and period analysis of Le Coroller et al. (2003) indicate that HD 179821 is a non-radial pulsator (NRP). It is not yet clear whether the shape of the star changes altogether, as we lack the spectral data (Le Coroller et al. quote unpublished data implying that the velocity variations are modest), or due to changes in surface temperature, the situation Le Coroller et al. prefer. In either case, the circumstellar material will be irradiated by an asymmetric source. Then, the polarization changes can be expected to occur around a preferred axis, as, apart from

rotation, the pulsation nodes are not expected to stochastically change position on the star.

In order to investigate whether it is plausible that a star with an aspherical shape can yield observable polarization, we performed some test calculations of a star covered by large spots, surrounded by a spherically symmetric dust shell. The model set-up consists of a star with a spot. In the situation where one (bright) spot is present, the polarization will increase with viewing angle. That is to say, if the spot is located in the line of sight, we will see its light directly and not much net polarization will be measured. If on the other hand the system is inclined, less direct light will be seen and the polarized light contribution increases, hence a larger polarization is observed. We should also note that unless the scattering occurs very close to the star (at a few stellar radii), the polarization percentage is independent of the radius of the shell. This is due to the constant geometry of the situation.

For the simple case where all the stellar flux is radiated from one spot covering 10 per cent of the stellar surface, we find that polarizations up to 3 per cent are readily produced when the star is mildly inclined. For smaller surface areas, the polarization can be even larger, exceeding 10 per cent. This result confirms earlier studies which used less extreme parameters (Raveendran 1991; Al-Malki et al. 1999). Of course this is an idealized situation, but it demonstrates that net polarizations of order per cent are certainly possible. Indeed, from an observational point of view, polarization changes of around 1 per cent have been observed towards Betelgeuse (Hayes 1984). These changes were due to variable hotspots on the stellar surface. Therefore, polarization changes can indeed be of order few per cent if the star itself is anisotropic.

One could argue why the polarization variability is not modulated by the 140 or 200 day period. Perhaps it is on the shorter term, but we lack the data to properly investigate this. However, where we could match photometric and polarimetric data, on shorter timescales it appears that the polarization values closely follow the photometry. It is therefore possible that the polarization is variable at similar timescales to the pulsations. A further question is why such long term changes are observed. We note that the mean photometry shows long term changes, probably tracing the global evolution of the photosphere, which would have an effect on the non-radial pulsations. Significantly, the preferred plane would not necessarily change, and this is found.

A critical test of the scenarios discussed above is simultaneous photo-polarimetric monitoring of HD 179821. If non-radial pulsations are responsible for the observed polarization, both observables should be strongly correlated. If, on the other hand, episodic mass ejections are the main cause of the variations, we would expect enhanced near-infrared emission due to hot dust during significant polarization changes.

5 CONCLUSION

We have presented spectropolarimetry and long term photopolarimetry for two post-Red Supergiants. A strong depolarization across the H α emission line is found for IRC +10420, suggesting an electron-scattering region that is not circularly

symmetric, confirming the results of Davies et al. (2007), who found such evidence from their integral field spectroscopy of the object. The time evolution indicates that the source has increased in temperature until at least 1995, after which the *J* band photometry indicates this may have levelled off. If the temperature increase of the object has indeed halted, an increase in mass loss or mass infall rate can be responsible for the observed H α emission and polarization.

HD 179821, a cooler object, has less H α emission, and no depolarization across the H α line could be detected. The photometry implies that this evolved object is also undergoing a change in temperature. Hitherto, temperature determinations of the object were inconclusive, but if the star is a G supergiant, then the observed change in photometry suggests it has become earlier by one or two subclasses. Strong changes at the 5 per cent level in polarization over the past 15 years have been detected. During the same time, the optical photometry has only varied by at most 0.2 magnitudes. The most obvious explanations for this observation such as changes in either electron or dust scattering can be ruled out. A complication is that the polarization is not correlated with any other observable. The possibility that the star undergoes irregular, asymmetric, mass ejections is discussed. However, there is little evidence for the occurrence of such events. Instead, it is proposed that the star itself is asymmetric and that its anisotropic radiation, which is scattered off the circumstellar dust results in the net polarization observed.

Acknowledgments

We wish to thank the referee, Prof. Kenneth H. Nord-sieck, for his detailed reading of the paper and constructive comments. We thank Willem-Jan de Wit for reading the manuscript. Based on observations conducted on the WHT, the NOT and the CST on the Canary islands, the AAT in Australia, and the TSAO in Kazakhstan. RDO is grateful for the support from the Leverhulme Trust for awarding a Research Fellowship. JCM and MP acknowledge PhD student funding from PPARC/STFC and the University of London respectively. We are grateful to Joni Johnson for allowing us to use her HPOL data on HD 179821.

REFERENCES

- Al-Malki M. B., Simmons J. F. L., Ignace R., Brown J. C., Clarke D., 1999, *A&A*, 347, 919
- Arhipova V. P., Ikonnikova N. P., Noskova R. I., Sokol G. V., Shugarov S. Y., 2001, *Astron. Lett.*, 27, 156
- Bergner Y. K., Bondarenko S. L., Miroschnichenko A. S., Yudin R. V., Yu. Y. N., Moralev Y. D., Schumacher A. V., 1988, *Izvestia Glavn. Astron. Obs. v Pulkove*, 205, 142
- Biegging J. H., Schmidt G. D., Smith P. S., Oppenheimer B. D., 2006, *ApJ*, 639, 1053
- Bjorkman K. S., Meade M. R., 2005 Vol. 343 of *ASP Conference Series*, p. 406
- Boyle R. P., Aspin C., McLean I. S., Coyne G. V., 1986, *A&A*, 164, 310
- Bujarrabal V., Alcolea J., Planesas P., 1992, *A&A*, 257, 701

- Castro-Carrizo A., Quintana-Lacaci G., Bujarrabal V., Neri R., Alcolea J., 2007, *A&A*, 465, 457
- Clarke D., Naghizadeh-Khouei J., 1994, *AJ*, 108, 687
- Davies B., Oudmaijer R. D., Sahu K. C., 2007, *ApJ*, 671, 2059
- Davies B., Oudmaijer R. D., Vink J. S., 2005, *A&A*, 439, 1107
- de Jager C., 1998, *A&AR*, 8, 145
- de Jager C., Lobel A., Nieuwenhuijzen H., Stothers R., 2001, *MNRAS*, 327, 452
- de Wit W., Davies B., Oudmaijer R. D., Groenewegen M., Hoare M., Malbet F., 2008, *A&A* accepted, arXiv:0711.4975
- Dougherty S. M., Taylor A. R., 1992, *Nature*, 359, 808
- Dullemond C. P., van den Ancker M. E., Acke B., van Boekel R., 2003, *ApJL*, 594, L47
- Dwarkadas V. V., Balick B., 1998, *AJ*, 116, 829
- Dwarkadas V. V., Owocki S. P., 2002, *ApJ*, 581, 1337
- Gorlova N., Lobel A., Burgasser A. J., Rieke G. H., Ilyin I., Stauffer J. R., 2006, *ApJ*, 651, 1130
- Grinin V. P., 1994, in The P. S., Perez M. R., van den Heuvel E. P. J., eds, *ASP Conf. Series Vol. 62. The Nature and Evolutionary Status of Herbig Ae/Be Stars*. Astron.Soc. Pac., San Francisco, p. 63
- Habing H. J., Tignon J., Tielens A. G. G. M., 1994, *A&A*, 286, 523
- Hayes D. P., 1984, *ApJS*, 55, 179
- Heiles C., 2000, *AJ*, 119, 923
- Hrivnak B. J., Kwok S., Volk K. M., 1989, *ApJ*, 346, 265
- Humphreys R. M., Davidson K., Smith N., 2002, *AJ*, 124, 1026
- Humphreys R. M., Smith N., Davidson K., et al. 1997, *AJ*, 114, 2778
- Humphreys R. M., Strecker D. W., Murdock T. L., Low F. J., 1973, *ApJ*, 179, L49
- Irvine C. E., Herbig G. H., 1986, *IAU Circ*, 4286, 1
- Jones T. J., Humphreys R. M., Gehrz R. D., et al. 1993, *ApJ*, 411, 323
- Jura M., Velusamy T., Werner M. W., 2001, *ApJ*, 556, 408
- Kastner J. H., Weintraub D. A., 1995, *ApJ*, 452, 833
- Keenan P. C., McNeil R. C., 1989, *ApJS*, 71, 245
- Kerschbaum F., Groenewegen M. A. T., Lazaro C., 2006, *A&A*, 460, 539
- Klochkova V. G., Chentsov E. L., Panchuk V. E., 1997, *MNRAS*, 292, 19
- Klochkova V. G., Yushkin M. V., Chentsov E. L., Panchuk V. E., 2002, *Astronomy Reports*, 46, 139
- Koornneef J., 1983, *A&A*, 128, 84
- Le Coroller H., Lèbre A., Gillet D., Chapellier E., 2003, *A&A*, 400, 613
- Leonard D. C., Li W., Filippenko A. V., Foley R. J., Chornock R., 2005, *ApJ*, 632, 450
- Mazzali P. A., Deng J., Tominaga N., Maeda K., Nomoto K., Matheson T., Kawabata K. S., Stanek K. Z., Garnavich P. M., 2003, *ApJL*, 599, L95
- Melikian N. D., Magnan C., Eritsian M. A., Karapetian A. A., 2000, *Astrophysics*, 43, 430
- Mészáros P., 2003, *Nature*, 423, 809
- Nordsieck K. H., Wisniewski J., Babler B. L., et al. 2001 Vol. 233 of *ASP Conference Series*, . Astron. Soc. Pac., San Francisco, p. 261
- Oudmaijer R. D., 1998, *A&AS*, 129, 541
- Oudmaijer R. D., 2007, in *Diffuse Matter From Star Forming Regions to Active Galaxies*, Editors T.W. Hartquist, S.A.E.G. Falle, J.M. Pittard, Springer, astro-ph/0610515
- Oudmaijer R. D., Davies B., de Wit W., Patel M., 2008, in D. Luttermoser B. Smith . R. S., ed., *ASP Conf. Series in press. Biggest, Baddest, Coolest stars*. Astron.Soc. Pac., San Francisco, pp –
- Oudmaijer R. D., Drew J. E., 1999, *MNRAS*, 305, 166
- Oudmaijer R. D., Geballe T. R., Waters L. B. F. M., Sahu K. C., 1994, *A&A*, 281, L33
- Oudmaijer R. D., Groenewegen M. A. T., Matthews H. E., Blommaert J. A. D. L., Sahu K. C., 1996, *MNRAS*, 280, 1062
- Oudmaijer R. D., Palacios J., Eiroa C., Davies J. K., de Winter D., Ferlet R. e. a., 2001, *A&AS*, 379, 564
- Oudmaijer R. D., Waters L. B. F. M., van der Veen W. E. C. J., Geballe T. R., 1995, *A&A*, 299, 69
- Parthasarathy M., Jain S. K., Sarkar G., 2005, *AJ*, 129, 2451
- Patel M., Oudmaijer R. D., Vink J. S., Mottram J. C., Davies B., 2006, *MNRAS*, 373, 1641
- Poeckert R., 1975, *ApJ*, 196, 777
- Poeckert R., Marlborough J. M., 1976, *ApJ*, 206, 182
- Quirrenbach A., Bjorkman K. S., Bjorkman J. E., et al. 1997, *ApJ*, 479, 477
- Raveendran A. V., 1991, *A&A*, 243, 445
- Reddy B. E., Hrivnak B. J., 1999, *AJ*, 117, 1834
- Schmidt E. G., Johnston D., Lee K. M., Langan S., Newman P. R., Snedden S. A., 2004, *AJ*, 128, 2988
- Slowik D. J., Peterson D. M., 1995, *AJ*, 109, 2193
- Smith N., Vink J. S., de Koter A., 2004, *ApJ*, 615, 475
- Straizys V., Kuriliene G., 1981, *Ap&SS*, 80, 353
- Thévenin F., Parthasarathy M., Jasniewicz G., 2000, *A&A*, 359, 138
- Trammell S. R., Dinerstein H. L., Goodrich R. W., 1994, *AJ*, 108, 984
- Ueta T., Meixner M., Bobrowsky M., 2000, *ApJ*, 528, 861
- Vink J. S., Drew J. E., Harries T. J., Oudmaijer R. D., 2002, *MNRAS*, 337, 356
- Vink J. S., Drew J. E., Harries T. J., Oudmaijer R. D., Unruh Y., 2005, *MNRAS*, 359, 1049
- Wang L., Baade D., Höflich P., Wheeler J. C., 2003, *ApJ*, 592, 457
- Weis K., 2003, *A&A*, 408, 205
- Wolff M. J., Nordsieck K. H., Nook M. A., 1996, *AJ*, 111, 856
- Wood K., Bjorkman K. S., Bjorkman J. E., 1997, *ApJ*, 477, 926
- Zacs L., Klochkova V. G., Panchuk V. E., Spelmanis R., 1996, *MNRAS*, 282, 1171

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.