

COMPARISON OF FOUR CLONES OF THE ICHTHYOTOXIC FLAGELLATE
PRYMNESIUM

Melissa A. Clouse

A Thesis Submitted to
University of North Carolina Wilmington in Partial Fulfillment
Of the Requirements for the Degree of
Master of Science

Department of Biology and Marine Biology

University of North Carolina Wilmington

2005

Approved by

Advisory Committee

Richard M. Dillaman

Thomas E. Lankford, Jr.

Carmelo R. Tomas
Chair

Accepted by

Robert Roer
Dean, Graduate School

TABLE OF CONTENTS

ABSTRACT	v
ACKNOWLEDGMENTS	vi
LIST OF TABLES	vii
LIST OF FIGURES	viii
INTRODUCTION	1
MATERIALS AND METHODS.....	8
Culturing	8
Nutrient Concentration Studies	9
Growth Studies	9
Erythrocyte Lysis Assay	10
Fish Bioassay	14
Statistical Analysis	15
RESULTS.....	16
Growth: Nutrient-Replete Cultures	16
Growth: Nitrogen-Deficient Cultures.....	16
Growth: Phosphorus-Deficient Cultures.....	23
Hemolytic Activity: Nutrient-Replete Cultures	34
Hemolytic Activity: Nitrogen-Deficient Cultures.....	34
Hemolytic Activity: Phosphorus-Deficient Cultures.....	37
Hemolytic Activity: Effect of Nutrient Concentration.....	40
Ichthyotoxicity	53

DISCUSSION.....	57
Growth.....	57
Hemolytic Activity.....	57
Hemolytic Activity: Effect of Nutrient Concentration.....	58
Ichthyotoxicity.....	59
Comparison between <i>P. parvum</i> and <i>P. calathiferum</i>	62
Hemolytic Activity: Supernatant versus Pellet.....	63
Hemolytic Activity: Growth Phase.....	64
Summary.....	66
CONCLUSIONS.....	67
LITERATURE CITED.....	69
APPENDICES.....	74

ABSTRACT

Since the mid 1980s blooms of the ichthyotoxic flagellate *Prymnesium parvum* have resulted in recurrent fish kills in Texas lakes, rivers, and reservoirs. South Carolina experienced a bloom of *P. parvum* in a brackish golf course pond in summer 2001. No dead fish were reported since the pond had no resident fish. The following year at Artesian Aquafarms in N.C., all hybrid striped bass perished to blooms of *P. parvum*. In the present study, clonal cultures from each of these blooms were grown in laboratory studies to determine response variation with nutrient-stressed (N-limited, P-limited) and replete cells for growth, hemolytic activity, and ichthyotoxicity. A congener, *P. calathiferum* originally isolated from a New Zealand bloom, was used for comparison. Of the *P. parvum* clones, the TX clone overall grew slower (0.21-0.31 div d^{-1}), had lower hemolytic activity (40-7164 units), but had the highest ichthyotoxicity (1 hr to kill fish in P-limited, 3 hrs in Replete and N-limited). This clone was the most sensitive to nutrient stress and conditioning was reduced to 1 week. In contrast, overall growth and hemolytic activity were greater in the NC (0.21-0.56 div d^{-1} , 77-21399 units) and SC clones (0.20-0.70 div d^{-1} , 45-20795 units) with lower ichthyotoxicity for both (≥ 4 hrs in N-&P-limited). *Prymnesium calathiferum* showed substantially lower hemolytic activity (8-779 units), but grew faster (0.30-0.73 div d^{-1}) than the *P. parvum* clones. Nitrogen-deficient cultures were similar to or more hemolytic than P-deficient cultures for *P. parvum*, but the P-deficient cultures were the most ichthyotoxic. Under nutrient-replete conditions, *P. calathiferum* was the most ichthyotoxic of the clones with fish mortality occurring in one hour as compared to three hours for the TX clone. Toxicity in *P. parvum* is a complex interaction of hemolytic and ichthyotoxic components.

ACKNOWLEDGMENTS

I wish to thank my advisor, Dr. Carmelo Tomas, who graciously took me on as a graduate student. Thanks for the support and patience throughout this learning process.

I also wish to thank my committee members, Dr. Richard Dillaman and Dr. Thomas Lankford, for their insight and involvement in this project. Thanks for the helpful comments that guided my thinking. Also thanks to Dr. Blum who guided my statistical analyses.

This study would not have been possible without the *P. parvum* clones themselves, so thanks to Artesian Aquafarms for the NC bloom samples, Dr. Alan Lewitus from the USC Belle Baruch Institute for the SC bloom samples, Ms. Joan Glass and Dave Buzan from Texas Parks and Wildlife Department for the TX bloom samples, and the NC Department of Environment and Natural Resources for the New River samples.

Financial support for this study was provided by NC SeaGrant (# 5-50470) and the NC Department of Health and Human Services (CDC Grant to NC #015-0414). Also thanks to the UNCW Graduate School for the Travel Award to attend the Baltimore GEOHAB conference.

LIST OF TABLES

Table	Page
1. ELA Buffer for Hemolysis Assay	11
2. Initial cell densities of three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i>	17
3. Growth rates (k) of three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i>	19
4. Time (in hours) required to kill two fish in each dilution for the three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i>	54

LIST OF FIGURES

Figure	Page
1. Distribution of <i>P. parvum</i> blooms in the United States.....	3
2. Structure of prymnesin	5
3. Growth of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> under nutrient-replete conditions.....	18
4. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under nutrient replete conditions for Trial 1	20
5. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under nutrient replete conditions for Trial 2	21
6. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under nutrient replete conditions for Trial 3	22
7. Growth of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> under N-deficient conditions.....	24
8. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under N-deficient conditions for Trial 1.....	25
9. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under N-deficient conditions for Trial 2.....	26
10. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under N-deficient conditions for Trial 3.....	27
11. Growth of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> under P-deficient conditions	29
12. Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under P-deficient conditions for Trial 1.....	30

13.	Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under P-deficient conditions for Trial 2.....	31
14.	Regression analysis for log phase for three geographically-distinct clones of <i>Prymnesium parvum</i> and <i>P. calathiferum</i> under P-deficient conditions for Trial 3.....	32
15.	Mean terminal densities of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	33
16.	Hemolytic activity by the supernatant of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under nutrient-replete conditions	35
17.	Hemolytic activity by the pellet of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under nutrient-replete conditions	36
18.	Hemolytic activity by the supernatant of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under N-deficient conditions.....	38
19.	Hemolytic activity by the pellet of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under N-deficient conditions.....	39
20.	Hemolytic activity by the supernatant of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under P-deficient conditions	41
21.	Hemolytic activity by the pellet of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> grown under P-deficient conditions	42
22.	Hemolytic activity by the supernatant of the North Carolina <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	44
23.	Hemolytic activity by the pellet of the North Carolina <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	45

24.	Hemolytic activity by the supernatant of the South Carolina <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	46
25.	Hemolytic activity by the pellet of the South Carolina <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient)	48
26.	Hemolytic activity by the supernatant of the Texas <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	49
27.	Hemolytic activity by the pellet of the Texas <i>Prymnesium parvum</i> clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	50
28.	Hemolytic activity by the supernatant of <i>Prymnesium calathiferum</i> grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	51
29.	Hemolytic activity by the pellet of <i>Prymnesium calathiferum</i> grown under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	52
30.	Cell densities for the fish bioassays of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	55
31.	Hemolytic activity for the fish bioassays of three geographically-distinct clones of <i>Prymnesium parvum</i> (NC=North Carolina; SC=South Carolina; TX=Texas) and <i>P. calathiferum</i> under different nutrient treatments (Replete, N-Deficient, P-Deficient).....	56

INTRODUCTION

Prymnesium parvum N. Carter (1937), a toxic phytoplankton usually <10 µm in size, was first discovered in the 1920s in a brackish tide pool on the Isle of Wight, England (Carter 1937). Its distinguishing features include two chloroplasts, two flagella (12-15 µm) and a single stiff haptonema (3-5 µm) (Green *et al.* 1982). Species identification involves viewing its body scales using electron microscopy since all species of the genus *Prymnesium* have an outer covering of organic scales. *Prymnesium parvum* has two layers of scales with those on the inner layer having wide inflexed rims, while those of the outer layer have narrow rims (Green *et al.* 1982). The scales of its distal face have a pattern of concentric circles, while those of its proximal face have a pattern of radial ridges (Manton and Leedale 1963). Scale morphology (as well as studies on physiology, toxicity, and genetics) led to the conclusion that a once separately identified species (*P. patelliferum*) is only a different life stage of *P. parvum* (Larsen and Medlin 1997, Larsen and Edvardsen 1998, Larsen 1999).

The devastating fish kills caused by *P. parvum* led to detailed studies of its blooms. Carp mortalities associated with *P. parvum* blooms occurred in Israel in 1945 (Reich and Aschner 1947, Yariv and Hestrin 1961) and in China since 1963 (Guo *et al.* 1996). In November-December 1969, the Yuzhnyy Fish Farms in Ukraine experienced fish-killing blooms (Krasnoshchek and Abramovich 1971). That same year, the Norfolk Thurne Broads, United Kingdom, began experiencing kills of perch, roach, bream, and eels (Holdway *et al.* 1978, Wortley and Phillips 1987) during similar blooms. Annual blooms of *P. parvum* from 1989–1996 in the Sandsfjord, Norway damaged fisheries of Atlantic salmon and rainbow trout (Kaarstvedt *et al.* 1991, Larsen and Edvardsen 1998). In Morocco (Sabour *et al.* 2000), *P. parvum* blooms killed carp, barbells, eels, sunpoles, gambusia, shrimp, bivalves, and other

invertebrates during 1998-1999. Lake Koronia experienced the first reported *P. parvum* bloom in Greece in August-September 2004. The blooms, ranging in densities from 120-1450 x10⁶ cells/L in waters having N:P ratios of 10:1, resulted in the death of thousands of birds (30 species, including the endangered *Pelecanus crispus*) and hundreds of fish (Moustaka-Gouni *et al.* 2004).

In the United States, *Prymnesium* blooms were first reported during the mid-1980s in Texas (Figure 1). Affected waters in Texas include Baylor Lake, Brazos River, Buffalo Springs Lake, California Creek, Colorado City Lake, Colorado River, E.V. Spence Reservoir, Lake Diversion, Lake Granbury, Lake Kemp, Lake Sweetwater, Lake Whitney, Lubbock Lake, Moss Creek Lake, Paint Creek, Pecos River, Possum Kingdom Lake/Reservoir, and Red Bluff Reservoir (James and de la Cruz 1989, Rhoades and Hubbs 1992, Texas Parks and Wildlife Department website). More recently, South Carolina experienced a bloom on May 22, 2001 at a brackish golf course pond on Kiawah Island (Lewitus *et al.* 2003). There were no fish kills reported because there were no fish in the pond. Artesian Aquafarms in Elizabeth City, North Carolina experienced *P. parvum* blooms in 2002, which started in March and persisted through October. This aquaculture facility grows hybrid striped bass. When the water source was changed from fresh to brackish to optimize fish growth, *Prymnesium* blooms occurred, causing large fish kills. *Prymnesium parvum* was also observed during a two-year study of the New River, NC where unexplained fish mortalities were commonly reported since 1980, though fish mortality directly linked to *P. parvum* was not established (Tomas *et al.* 2004). Other areas in the U.S. with *P. parvum* blooms have included Colorado, New Mexico, Arizona, and Florida (Websites for Texas Parks and Wildlife Department, Colorado Department of Natural Resources, Arizona Game and Fish Department).



Figure 1. Distribution of *P. parvum* blooms in the United States.

Yariv first isolated the toxin of *P. parvum* in 1958 (Yariv and Hestrin 1961). Properties of prymnesins include a ninety-carbon skeleton with a methyl group as the only carbon branching, five ether rings, possessing a hydrophilic end and a hydrophobic end, and functional groups including conjugated double and triple bonds, chlorine atoms, an amino group and glycosidic residues (Morohashi *et al.* 2001) (Figure 2). Presently, there are two known structures of prymnesins with four known effects - ichthyotoxicity, hemolysis, cytolysis, and neurotoxicity (Igarashi *et al.* 1998). These effects appear to be related to the ability of the prymnesins to change the permeability of cell membranes (Johansson and Granéli 1999).

The ichthyotoxic component affects fish, tadpoles, and invertebrates by disrupting the permeability of the gills (Yariv and Hestrin 1961, James and de la Cruz 1989). Gill damage can occur after only ten to fifteen minutes of exposure. Uptake of the toxin is considered to occur in two stages, 1) a reversible damage to the gills results in a loss of selective permeability and 2) the fish become more sensitive to other toxins due to breakdown in the gill membranes (Ulitzur and Shilo 1966). Cofactors are required to activate the ichthyotoxin including monovalent and divalent cations such as calcium and magnesium (Parnas *et al.* 1962, Parnas 1963). Using streptomycin, neomycin, and spermine have also increased ichthyotoxicity in laboratory experiments (Yariv and Hestrin 1961, Ulitzur and Shilo 1964).

The hemolytic portion of prymnesin was found to contain at least six compounds with two similar glycolipids as the dominating compound of the mixture (Meldahl *et al.* 1994). A specific receptor site on the surface of the blood cells is suggested for the hemolysin, and an aggregation of toxin molecules may be involved (Igarashi *et al.* 1998). The hemolysins become inactivated with high pH, monovalent cations, and possibly by some lipids (Padilla 1970, Ulitzur and Shilo 1970, Igarashi *et al.* 1998).




Figure 2. Structure of prymnesin. (from Igarashi *et al.* 1999)

Cytolytic abilities are also reported for the prymnesins. Cells swell due to the free flux of ions (sodium and potassium), amino acids, and nucleotides. As with the ichthyotoxic effect, two stages are involved – water uptake and swelling, followed by cell lysis (Dafni and Giberman 1972).

Neurotoxic effects of prymnesins caused respiratory failure in frogs, mice and cats in laboratory experiments (Parnas 1963). This toxin attacks the central nervous system by blocking postsynaptic membranes at the neuromuscular junction as well as contracting smooth muscle (Parnas and Abbott 1965, Meldahl *et al.* 1994).

One factor implicated in toxin production is nutrient limitation. Shilo (1967) found low levels of phosphorus in the medium with highly toxic *P. parvum* cells. Dafni *et al.* (1972) proposed that phosphate limitation might lead to increased toxin production by disrupting membrane phospholipids, which would act to excrete the toxin. Similarly, blooms in the United Kingdom (Holdway and Watson 1978, Holdway *et al.* 1978, Wortley and Phillips 1987), Norway (Aure and Rey 1992), and Morocco (Sabour *et al.* 2000) occurred in areas with low phosphate concentrations. Johansson and Granéli (1999) studied hemolytic activity of *P. parvum* under both nitrogen and phosphorus limitation and found hemolytic activity to increase under nutrient limitation compared to non-limiting conditions. From these results, they suggested that cellular physiological stress might cause increased toxin production. They argued that this is somewhat perplexing in that the linkage between nutrient limitation and toxicity may not be direct, since nitrogen and phosphorus are not main components of the toxin. In studies for allelopathy, defined as “the release of chemical substances by individuals of a population that have an effect on the individuals of another population” (Hulot and Huisman 2004), toxin production by nutrient-stressed *P. parvum* cultures was associated with inhibiting various

phytoplankton (*Thalassiosira weissflogii*, *Prorocentrum minimum*, *Rhodomonas cf. baltica*) and decreased survival rates of the ciliate *Euplotes affinis* (Granéli and Johansson 2003a, 2003b).

While several laboratory studies suggest phosphorus limitation increases toxin production and release, the study of Johansson and Granéli (1999) emphasizes that either nitrogen or phosphorus limitation may lead to increased toxicity. Adding to this confusion are the reports that natural blooms are associated with phosphorus limitation. The recent blooms in the United States have occurred in areas of high eutrophication, but there was some variation among the bloom sites. Two blooms occurred in brackish man-made facilities (NC and SC), while the other occurred in alkaline natural lakes (TX). The blooms in Texas and North Carolina caused fish kills, while those in South Carolina did not due to a lack of fish in the waterbody.

The purpose of this study was to test the relationship between nutrient concentration, growth phase, and toxicity in clones of *P. parvum* from the United States blooms (North Carolina, South Carolina, and Texas) and compare them to a clone of *P. calathiferum* Chang and Ryan (1985), a toxic species from New Zealand.

The research objectives were:

- to determine toxicity as measured by hemolytic activity of the four clones of *Prymnesium*.
- to examine toxicity to *Gambusia holbrooki* of the four clones of *Prymnesium*.
- to examine the effects of nutrient concentration (nitrogen and phosphorus) on hemolytic activity.
- to examine how growth phase (lag, log, stationary) affects hemolytic activity.

The null hypotheses were that:

- the four clones do not differ in their hemolytic activity and are not capable of killing fish.

- neither nutrient concentration nor growth phase affect the hemolytic activity of the four clones.

MATERIALS AND METHODS

Culturing

The Tomas lab (UNCW Center for Marine Science) has clonal cultures of *P. parvum* established from blooms in the United States. One of the fourteen clones from the North Carolina bloom, picked at random, was used in this study (CMS TAC PP7). *Prymnesium parvum* clones from South Carolina (CMS TAC PP22) and Texas (CMS TAC PP18) were also used. Another *Prymnesium* species from New Zealand (*P. calathiferum* (CCMP Strain CHANG1) was used to compare with *P. parvum* from the United States.

Clonal cultures were grown with f/2 media (Guillard and Ryther 1962) at salinities of 4 for *P. parvum* and 36 for *P. calathiferum*. To make the media of different salinities, full strength seawater (at salinity of 38-39), collected 20-50 miles offshore, was filtered through a 47 mm Whatman GF/F glass microfibre filter, and diluted with pyrogen-free deionized water equivalent to Milli-Q ultrapure water. Salinity was checked with a refractometer. Water of the appropriate salinity was then sterilized by autoclaving at 121° C for 15 minutes in Teflon bottles. Sterile nutrients were added aseptically once the salinities were cooled to room temperature. The cultures were maintained in Erlenmeyer (150 mL) and Fernbach (1.5 L) flasks and kept in a walk-in growth chamber of constant temperature (22 ± 1° C) with a 16: 8 hour light:dark photoperiod under cool fluorescent 40-Watt light. For the experiments, only cultures grown in Fernbach flasks were used, with transfers done every 2-3 weeks.

Nutrient Concentration Studies

For the nutrient studies, *f/2* media was used to make nutrient-replete, nitrogen-limited, and phosphorus-limited conditions. The nitrogen-limited media was modified to N:P = 4:1, using 16 μM nitrate and 4 μM phosphate. The phosphorus-limited media was modified to N:P = 80:1, using 80 μM nitrate and 1 μM phosphate. Approximately 30 mL of culture grown in nutrient-replete media were transferred into 150 mL of nutrient-deficient media in Erlenmeyer flasks. After one week of growth, the cultures were transferred again to nutrient-deficient media. After another week of growth, the cultures were inoculated into Fernbach flasks and the experiments began. The Texas *P. parvum* clone (for nitrogen and phosphorus limitation) and *P. calathiferum* (for only nitrogen limitation) did not exhibit growth after inoculation into the Fernbach flasks after two weeks of nutrient deprivation. To allow studies of these cultures, they were conditioned for only one week in nutrient-deficient media before inoculation into Fernbach flasks.

Growth Studies

Prymnesium cultures were counted daily using a Beckman Coulter Multisizer 2E Particle Counter with an aperture of 100 μm to generate growth curves as well as to determine the number of cells used in each hemolysis assay. For counting, full strength seawater was diluted with pyrogen-free deionized water to obtain the correct salinity. This water was then filtered through two filters, 47 mm Whatman GF/F glass microfibre filter and a 0.2 μm GTTP Millipore Isopore Membrane Filter, to minimize particle interference. Cultures were diluted with their respective salinities of 4 for *P. parvum* and 36 for *P. calathiferum* to optimize the counter capabilities. Very dense cultures gave a higher percentage of coincidence, defined as a greater

chance that more than one organism was passing through the orifice at the same time. The percentage of coincidence was kept below 5% by diluting the samples before counting. Four replicate counts were taken daily from one sample of each culture. Each sample was continuously stirred and 500 μ L of each culture was counted. To obtain the number of cells in one mL, the raw count was multiplied by 2 and corrected for the dilution factor. The means of the four replicate counts were used to generate growth curves using SigmaPlot 2001. Using the growth curves, log phase was determined by eye. The mean cells/mL were converted to Log_2 and plotted against the days in log phase. A linear regression was performed on the cell densities in log phase using SigmaPlot 2001. The slope of the line gave the growth rate k .

$$k = \log_2 (N_1/N_0)/(t_1-t_0) \text{ (Guillard 1973).}$$

Erythrocyte Lysis Assay

To test hemolytic activity, the erythrocyte lysis assay (ELA, Eschbach *et al.* 2001) was performed. Outdated human red blood cells, obtained from and screened for pathogens by the American Red Cross, were used for this assay due to easier availability and convenience. Other sources, such as fish blood, were not readily available and in insufficient quantities to guarantee a reliable and stable supply. The blood was stored in a 4° C refrigerator and kept on ice while used in the assay. At the time of analysis, the test blood was placed into a 15mL centrifuge tube to which approximately 10 mL of cold ELA buffer were added (Table 1).

The buffer was kept cold so as not to lyse blood cells due to heat shock. The centrifuge tube of blood and buffer was inverted five times and spun at 300 rpm at 4° C for five minutes with a Hermle (Model #Z383K) refrigerated centrifuge. The supernatant was discarded and the cells were resuspended with 10 mL of cold buffer. This washing procedure was repeated until the supernatant was clear (approximately 3-6 times). After the final centrifugation, the

Table 1. ELA Buffer for Hemolysis Assay*

Reagent	Concentration (mM)	Molarity (g/mol)	Grams for 2L
NaCl	150	58.44	17.54
KCl	3.2	74.56	0.4772
MgSO ₄	1.25	246.48	0.6162
CaCl ₂	3.75	110.99	0.8234
Trizma pre-set crystals, pH 7.0	12.2	154.8	3.772

* pH adjusted to 7.4 at 4° C.

supernatant was removed and a volume of cold buffer to match the volume of blood was added. A 1:40 through 1:60 dilution of erythrocytes with buffer was made to give an optimal concentration of erythrocytes. This was done to maximize the capability of the microplate reader, giving a full positive control optical density reading of 3.00 OD.

Prymnesium cultures were selected on days representing lag, log, and stationary phases of growth. Fifty mL aliquots of the whole culture were centrifuged at 3200 rpm for 15 minutes at 4° C. The first 50 mL aliquot centrifuged had the supernatant poured off and used in the assay. Due to the small nature of *Prymnesium* cells, obtaining a visible pellet usually took many rounds of centrifuging 50 mL aliquots of culture, removing the majority of the supernatant, and adding more culture to centrifuge. The amount of culture added to obtain a visible pellet was recorded to use in later calculations to correct for the different volumes centrifuged. When a visible pellet of a known cell number was obtained, 3 mL of cold buffer was added and the mixture was sonicated continuously for 30 seconds at an amplitude of 45 on a 20 kHz Ultrasonic Processor (Model #GE 130 PB, Sonics and Materials, Inc.). Cold buffer was used to yield 1:2 and 1:10 dilutions of the supernatant and 1:10 and 1:100 dilutions for the sonicated pellet. The 100% concentrations of the supernatant and pellet were also used in the assay. Throughout the assay, all culture samples were kept on ice.

Using a Transferpette-8 pipette, 125 µl each of erythrocytes and the *P. parvum* dilutions were placed into a Costar 96-well microtitre plate with V-shaped bottoms. Throughout the experiment, the plates were kept on ice. Negative controls consisted of erythrocytes incubated with buffer only; positive controls were erythrocytes incubated with saponin (Sigma S4521) which lysed all erythrocytes. The saponin reagent was made by dissolving 0.008 g of saponin with 50 mL of pyrogen-free DIW to give a concentration of 20 µg/125µL/well. Once made, the

saponin was placed into cryovials, kept in a -80° C freezer, and then thawed before use in the assay. Both the negative and positive controls consisted of eight replicate wells and the culture samples consisted of 4-8 replicates. The plates were sealed and incubated at 4° C for 24 hours. After incubation, the plates were centrifuged at 1250 rpm for 10 minutes at 4° C and 250 µl aliquots of the supernatant were then transferred to a 96-well Costar microtitre plate with flat bottoms. The optical densities of the samples (representing released hemoglobin) were read with on a Bio-Tek Powerwave X microplate reader at 415 nm equipped with the K.C. Junior program. The mean of the negative control replicates was subtracted from the rest of the values to correct for any hemolysis by the buffer. Each corrected value was divided by the mean of the saponin replicates and multiplied by 100 to obtain a percentage of lysis. The mean of the replicates were calculated for each culture concentration. This gave percent hemolysis by the supernatant and the pellet.

Toxicity was then normalized on a per cell basis to compare the same number of cells in each clone. As mentioned before, centrifuging the exact number of cells in each culture was not possible due to different cell densities and problems obtaining a pellet. The Coulter count (in cells/mL) was multiplied by the volume of sample centrifuged (50 mL for the supernatant, usually a larger volume for the pellet) as well as the volume of cells in each well (always 125 µL). This number (N) was then divided into the percent hemolysis (PH) value, giving toxicity on a per cell basis. The number was then multiplied by a constant (1×10^7) to obtain an easier to read value. This gave normalized hemolysis by the supernatant and pellet.

$$N = (\# \text{ cells/mL from Coulter counter}) * (\text{Volume centrifuged}) * (0.125 \text{ mL})$$

$$(\text{PH}/N) * (1 \times 10^7) = \text{Normalized Lysis Value}$$

Toxicity per growth phase (lag, log, stationary) was examined in all four clones. Three replicate trials were performed for each nutrient treatment.

Fish Bioassay

The toxicity of the four clones was also examined with a fish bioassay. Assays using the eastern mosquitofish *Gambusia holbrooki* are commonly used to determine the presence of *Prymnesium* toxicity. For this study, results from the bioassay were compared with those from a hemolysis assay. The procedure outlined by Shilo and Aschner (1953) was used. *Gambusia* were captured in local brackish ponds and maintained in an aquarium at the UNCW Center for Marine Science (IACUC 01-008).

Before the fish bioassay, the cultures were counted every few days with a Coulter Counter to verify that they were in stationary phase. Stationary phase was chosen because the cultures would be the most dense and easier to work with (i.e. centrifuging 40 mL of culture would result in a visible pellet). An aliquot of 200 mL of each clone (NC, SC, TX, Pcal) were placed into sterile beakers. Aliquots of 40 mL of each culture were centrifuged at 3200 rpm for 15 minutes at 4°C. The supernatant was then separated from the pellet. Both fractions, to be used later in hemolysis assays, were placed in a -80° C freezer.

The fish bioassay took place in 25 Pyrex 50-mL beakers.

- 1 beaker – negative control
- 2 beakers each – NC [100%]; NC [50%]; NC [10%]
- 2 beakers each – SC [100%]; SC [50%]; SC [10%]
- 2 beakers each – TX [100%]; TX [50%]; TX [10%]
- 2 beakers each – Pcal [100%]; Pcal [50%]; Pcal [10%]

The cultures were diluted with their respective sterile media. The 100% concentration consisted of 40 mL of culture. The 50% and 10% concentrations were made by diluting the culture with media. The negative control consisted of 40 mL of aquarium water. One mL of the cofactor DADPA (0.003 M DADPA (3,3' – diaminodipropylamine, Sigma I 7006) was added to every beaker, including the negative control, to make the fish more sensitive to the toxin (Ulitzur and Shilo 1966). Timing began after one *Gambusia* was added to all the beakers. The assay was monitored closely for up to six hours. Observations were recorded every fifteen minutes for the first two hours and every thirty minutes thereafter. Symptoms of toxicity that were looked for included erratic movements, loss of equilibrium, release of blood into the medium, and subsurface bleeding at the snout, gills, and pectoral fins. Death was determined by immobility of the body and gills. At the end of the experiment, dead fish were discarded. The survivors were removed from the experimental beakers and then released into a pond far from the normal collection site.

Statistical Analysis

Growth rates during log phase were examined by linear regression analysis with SigmaPlot 2001, after Log₂-transforming the data. Hemolytic activity differences were examined using two-way and three-way analysis of variance (ANOVA) and Tukey HSD with SigmaStat and JMP, after Log₁₀-transforming the hemolytic activity data. A significant level of $\alpha=0.05$ was used in all tests.

RESULTS

Growth: Nutrient-Replete Cultures

The initial cell densities varied significantly ($P < 0.001$) for every trial and clone (Table 2). All four clones exhibited a very short lag phase in all three trials (Figure 3). Throughout the trials, the *P. parvum* clones had similar log phase lengths, but differed in terminal densities. For Trial 1, the *P. parvum* clones reached similar ($0.477 < P < 0.979$) terminal cell densities. The NC clone reached the highest terminal density ($P < 0.001$), while the similar ($P = 0.326$) SC and TX clones reached slightly lower terminal densities in Trial 2. For Trial 3, the NC reached a significantly ($P < 0.001$) higher terminal density than the SC clone, while the TX clone had a smaller ($P < 0.001$) terminal density than both the NC and SC clones. *Prymnesium calathiferum* had a shorter log phase and reached a smaller terminal density ($P < 0.001$) for Trials 1 and 2 compared with the *P. parvum* clones. In Trial 3, *P. calathiferum* exhibited no true stationary phase, but rather went from log phase to declining phase.

The three *P. parvum* clones exhibited similar growth rates during log phase for all three trials in the nutrient-replete treatment (Table 3, Figures 4-6). The growth rates for *P. calathiferum* were not significantly different ($0.052 < P < 0.079$) from those for the three *P. parvum* clones.

Growth: Nitrogen-Deficient Cultures

The initial cell densities varied significantly ($P < 0.001$) for every trial and clone (Table 2). All four clones exhibited a very short lag phase in all trials. In Trial 1, the three *P. parvum*

Table 2. Initial cell densities of three geographically-distinct clones of *Prymnesium parvum** and *P. calathiferum*.

Nutrient Treatment	Trial	NC	SC	TX	Pcal
Replete	1	20058	12469	22621	21298
	2	17815	10855	15775	3645
	3	9255	11830	14330	11790
N-deficient	1	938	1252	7555	5505
	2	2535	924	13845	3700
	3	4660	3140	3905	3675
P-deficient	1	5760	5212	3755	4780
	2	8185	10470	15550	10415
	3	1360	2135	7525	2910

* NC=North Carolina; SC=South Carolina; TX=Texas




Figure 3. Growth of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* under nutrient-replete conditions. Error bars represent one standard deviation about the mean of four daily counts on a Coulter Counter. (A) Trial 1. (B) Trial 2. (C) Trial 3. (D) Terminal densities of the clones for each trial.

Table 3. Growth rates (k) of three geographically-distinct clones of *Prymnesium parvum** and *P. calathiferum*. Parentheses indicate days in log phase.

Nutrient Treatment	Trial	NC	SC	TX	Pcal
Replete					
	1	0.27 (5-10)	0.48 (4-8)	0.26 (5-12)	0.61 (5-19)
	2	0.27 (5-14)	0.30 (6-14)	0.31 (9-16)	0.42 (7-14)
	3	0.21 (9-17)	0.20 (8-18)	0.24 (5-18)	0.30 (5-14)
N-deficient					
	1	0.38 (4-9)	0.45 (2-6)	0.23 (0-3)	0.58 (1-4)
	2	0.22 (2-10)	0.33 (4-8)	0.21 (0-3)	0.46 (1-4)
	3	0.32 (2-5)	0.43 (2-5)	0.30 (1-3)	0.66 (1-3)
P-deficient					
	1	0.44 (5-10)	0.51 (5-10)	0.24 (2-7)	0.73 (3-8)
	2	0.54 (2-7)	0.49 (2-7)	0.22 (1-7)	0.69 (2-6)
	3	0.56 (9-15)	0.70 (4-13)	0.23 (1-5)	0.46 (6-11)

* NC=North Carolina; SC=South Carolina; TX=Texas




Figure 4. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under nutrient-replete conditions for Trial 1. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 5. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under nutrient-replete conditions for Trial 2. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 6. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under nutrient-replete conditions for Trial 3. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*

clones all reached similar terminal cell densities of approximately $1.2-1.3 \times 10^5$ cells/mL, though the TX clone had a shorter log phase than the other clones (Figure 7). *Prymnesium calathiferum* reached a higher terminal density ($P < 0.001$) than the *P. parvum* clones. In Trial 2, the SC clone had a slightly shorter log phase, but reached a statistically similar ($P = 0.720$) terminal density to the NC clone. The TX clone differed from both the NC and SC clones by having a shorter log phase and a larger ($P < 0.001$) terminal density. *Prymnesium calathiferum* also had a very short log phase, but reached the smallest terminal density ($P < 0.001$) of the four clones. In Trial 3, NC and SC clones grew to similar terminal densities ($P = 0.998$), while the terminal density of the TX clone was significantly lower ($P < 0.001$) than those of the other *P. parvum* clones, but similar ($P = 0.754$) to that of *P. calathiferum*.

The SC clone had a high growth rate for the *P. parvum* clones for all three trials, but was only significantly different from the TX clone ($P = 0.016$) and not the NC clone ($P = 0.113$) (Table 3, Figures 8-10). In Trial 1, the NC and SC clones had similar growth rates, while the TX clone had a slightly lower growth rate. In Trials 2 and 3, the NC and TX had similar ($P = 0.113$) growth rates. *Prymnesium calathiferum* had the significantly ($P \leq 0.013$) highest growth rate of the four clones for all three trials.

Growth: Phosphorus-Deficient Cultures

The initial cell densities varied significantly ($P < 0.001$) for every trial and clone (Table 2). All four clones had short lag and log phases. Lag phase was longer in Trial 3. The SC clone had significantly higher ($P < 0.001$) terminal densities than the NC clone for all three trials (Figure 11). The TX clone reached the smallest terminal density ($P < 0.001$) of the four clones for all




Figure 7. Growth of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* under N-deficient conditions. Error bars represent one standard deviation about the mean from four daily counts on a Coulter Counter. (A) Trial 1. (B) Trial 2. (C) Trial 3. (D) Terminal densities of the clones for each trial.




Figure 8. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under N-deficient conditions for Trial 1. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 9. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under N-deficient conditions for Trial 2. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 10. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under N-deficient conditions for Trial 3. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*

three trials. *Prymnesium calathiferum* reached a terminal density higher than the TX clone, but less than the NC and SC clones for all three trials ($P < 0.001$).

For Trials 1 and 2, the NC and SC clones had similar growth rates. For Trial 3, the SC clone had a much higher growth rate than the other three clones. The growth rate of the TX clone was much lower than the other clones for all trials, but was only significantly different from the SC clone ($P = 0.042$) and *P. calathiferum* ($P = 0.021$). *Prymnesium calathiferum* had the highest growth rate of the four clones for Trials 1 and 2 (Table 3, Figures 12-14), but the growth rates of the NC, SC, and *P. calathiferum* clones were not significantly different ($P \geq 0.638$).

Overall, when averaging the trials and comparing nutrient treatments, all clones reached their maximum growth under nutrient-replete conditions (Figure 15). Nutrient-deficient conditions produced limited growth. The P-deficient conditions produced larger terminal densities than the N-deficient conditions for all four clones. *Prymnesium calathiferum* had the significantly ($P \leq 0.019$) highest growth rate throughout the nutrient treatments (Table 3). The NC and SC clones had their highest growth rates in the nutrient-deficient treatments, particularly in the P-deficient treatment, which was significantly different ($P \leq 0.035$) from the nutrient-replete treatment. The TX and *P. calathiferum* clones had similar growth rates throughout the nutrient treatments.




Figure 11. Growth of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* under P-deficient conditions. Error bars represent one standard deviation about the mean from four daily counts on a Coulter Counter. (A) Trial 1. (B) Trial 2. (C) Trial 3. (D) Terminal densities of the clones for each trial.




Figure 12. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under P-deficient conditions for Trial 1. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 13. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under P-deficient conditions for Trial 2. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 14. Regression analysis for log phase for three geographically-distinct clones of *Prymnesium parvum* and *P. calathiferum* under P-deficient conditions for Trial 3. (A) North Carolina (B) South Carolina (C) Texas (D) *P. calathiferum*




Figure 15. Mean terminal densities of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean of three trials.

Hemolytic Activity: Nutrient-Replete Cultures

There were no significant differences in mean hemolytic activity among the three *P. parvum* clones for either the supernatant ($P > 0.300$) or the pellet ($P > 0.600$) for any of the growth phases (Figures 16 and 17). For the supernatant, the NC and SC clones were significantly more hemolytic ($P < 0.040$) than *P. calathiferum* in lag and stationary phases, but the TX clone (lag, $P = 0.077$; stationary, $P = 0.065$) was not statistically different from *P. calathiferum*. In log phase, all three *P. parvum* clones were significantly more hemolytic ($P \leq 0.004$) than *P. calathiferum*. For the pellet, the three *P. parvum* clones were more hemolytic ($P < 0.040$) than *P. calathiferum* in lag and log phases. However, in stationary phases, there were no differences in mean hemolytic activity ($0.999 < P < 0.5$) among the four clones.

Lag phase was the most hemolytic ($P < 0.030$) phase for the supernatant in all four clones. Lag phase was also the most hemolytic ($P \leq 0.007$) for the pellet for the NC and SC clones. For the TX clone, lag phase was the most hemolytic for the pellet, but was only statistically different from stationary phase ($P = 0.003$) and not log phase ($P = 0.059$). For *P. calathiferum*, there were no differences in hemolytic activity ($P > 0.200$) among the growth phases for the pellet. There was no difference in hemolytic activity between the supernatant and the pellet for the NC, SC, and *P. calathiferum* clones. For the TX clone, the pellet was significantly more hemolytic ($P < 0.001$) than the supernatant.

Hemolytic Activity: Nitrogen-Deficient Cultures

There were clonal differences in mean hemolytic activity for the supernatant (Figure 18A). The NC and SC clones were similar in hemolytic activity, but more




Figure 16. Hemolytic activity by the supernatant of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under nutrient-replete conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 17. Hemolytic activity by the pellet of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under nutrient-replete conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3

hemolytic ($P < 0.001$) than both the TX and *P. calathiferum* clones in all growth phases. This trend was supported by all three of the individual trials (Figure 18B-D). Overall, the TX clone was more hemolytic ($P < 0.001$) than *P. calathiferum*, but differences in growth phase were seen. In lag phase, there was no significant difference ($P = 0.743$) between the TX and *P. calathiferum* clones, but the TX clone was more hemolytic ($P \leq 0.003$) than *P. calathiferum* in log and stationary phases. There were no significant differences in hemolytic activity ($P > 0.100$) among the growth phases for any of the *P. parvum* clones. For *P. calathiferum*, lag phase was more hemolytic than log ($P = 0.016$) and stationary ($P = 0.010$) phases.

For the pellet, there were no significant differences ($P > 0.400$) in mean hemolytic activity among the three *P. parvum* clones (Figure 19A). In addition, all *P. parvum* clones were significantly more hemolytic ($P < 0.001$) than *P. calathiferum* for all growth phases. For the NC and SC clones, lag phase was more hemolytic ($P < 0.050$) than stationary phase. For the TX clone, there was no difference in hemolytic activity among the growth phases. For *P. calathiferum*, lag phase was the most hemolytic ($P \leq 0.037$). The supernatant was the most hemolytic for the NC, SC, and *P. calathiferum* clones ($P \leq 0.011$). For the TX clone, the pellet was the most hemolytic ($P < 0.001$).

Hemolytic Activity: Phosphorus-Deficient Cultures

For the supernatant, clonal differences in mean hemolytic activity were seen in the *P. parvum* clones for individual growth phases (Figure 20A). The NC clone was significantly more hemolytic ($P = 0.016$) than the TX clone during lag phase, while the TX clone was significantly more hemolytic ($P = 0.022$) than the SC clone during stationary




Figure 18. Hemolytic activity by the supernatant of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under N-deficient conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 19. Hemolytic activity by the pellet of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under N-deficient conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.

phase. These two differences were supported by all three of the individual trials (Figure 20B-D). The NC and SC clones were significantly more hemolytic ($P < 0.001$) than *P. calathiferum* for all growth phases. Overall, the TX clone was also more hemolytic ($P < 0.001$) than *P. calathiferum*, but differences were seen in growth phases. The TX clone was only significantly more hemolytic ($P < 0.001$) than *P. calathiferum* for log and stationary phases, but there was no significant difference ($P = 0.059$) during lag phase. Lag phase was the most hemolytic phase ($P \leq 0.022$) for the NC, SC, and *P. calathiferum* clones, but all phases of the TX clone had similar hemolytic activity ($P \geq 0.755$).

For the pellet, there were no differences in mean hemolytic activity among the *P. parvum* clones for lag and log phases, but the TX clone was statistically more hemolytic than the NC ($P = 0.027$) and SC ($P = 0.008$) clones in stationary phase (Figure 21A). This difference was supported in all three of the individual trials (Figure 21B-D). Overall, the *P. parvum* clones were more hemolytic ($P < 0.001$) than *P. calathiferum*, but this was not seen in the individual growth phases for the NC and SC clones. *Prymnesium calathiferum* was significantly less hemolytic than the NC and SC clones only for lag ($P < 0.001$) and log ($P \leq 0.022$) phases, but there was no difference among the three clones in stationary phase. For the NC and SC clones, lag phase was the most hemolytic phase ($P \leq 0.012$). For the TX and *P. calathiferum* clones, there were no significant differences among growth phases. There was no difference among the supernatant and pellet for the four clones ($0.194 \leq P \leq 0.395$).

Hemolytic Activity: Effect of Nutrient Concentration

For the supernatant of the NC clone, the nutrient-replete culture was significantly less hemolytic ($P < 0.001$) than the nutrient-deficient cultures for all growth phases (Figure




Figure 20. Hemolytic activity by the supernatant of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under P-deficient conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 21. Hemolytic activity by the pellet of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown under P-deficient conditions. Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.

22A). In lag phase, there was no significant difference in hemolytic activity for the nutrient-deficient cultures ($P=0.622$). This was supported in two of the three individual trials (Figure 22B-D). In the first trial, however, the N-deficient culture was significantly more hemolytic ($P<0.001$) than the P-deficient culture. In log and stationary phases, the N-deficient culture was more hemolytic ($P\leq 0.001$) than the P-deficient culture.

For the pellet of the NC clone, the nutrient-replete culture was significantly less hemolytic than the nutrient-deficient cultures ($P\leq 0.012$) for both lag and log phases (Figure 23A). In stationary phase, however, the nutrient-replete culture was less hemolytic ($P<0.001$) than the N-deficient culture, but there was no difference ($P=0.132$) between the nutrient-replete and P-deficient cultures. The N-deficient culture was only significant more hemolytic than the P-deficient culture in stationary phase ($P=0.006$) and not in log phase ($P=0.121$). For Trials 1 and 2, the N-deficient culture was more hemolytic than the P-deficient culture for all growth phases (Figure 23B-C). However, in Trial 3, the P-deficient culture had a much higher normalized hemolysis value in lag phase than in the other trials and was more hemolytic than the N-deficient culture (Figure 23D).

For the supernatant of the SC clone, the nutrient-replete culture was significantly less hemolytic than the nutrient-deficient cultures ($P\leq 0.05$) in all growth phases (Figure 24A). In lag phase, there was no difference between the nutrient-deficient cultures. The P-deficient culture was the most hemolytic for Trial 1 (Figure 24B), but the N-deficient culture was the most hemolytic Trials 2 and 3 in lag phase (Figure 24C-D). However, the N-deficient culture was significantly more hemolytic ($P\leq 0.008$) than the P-deficient culture in log and stationary phases.




Figure 22. Hemolytic activity by the supernatant of the North Carolina *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 23. Hemolytic activity by the pellet of the North Carolina *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 24. Hemolytic activity by the supernatant of the South Carolina *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.

For the pellet of the SC clone, the nutrient-replete culture was significantly less hemolytic ($P=0.010$) than the nutrient-deficient cultures in lag phase (Figure 25A). There was no overall difference in hemolytic activity between the nutrient-deficient cultures in lag phase. The P-deficient culture was the most hemolytic in two trials (Figure 25B, 25D), while the N-deficient culture was the most hemolytic in one trial (Figure 25C). In log and stationary phases, the N-deficient culture was more hemolytic than the other cultures ($P<0.040$), but the P-deficient and nutrient-replete cultures were not significantly different ($P>0.130$).

For the TX clone, the nutrient-replete culture was significantly less hemolytic than the nutrient-deficient cultures for only log and stationary phases for hemolysis by both the supernatant ($P\leq 0.041$) and pellet ($P<0.080$) (Figures 26 and 27). There were no significant differences between the nutrient-deficient cultures for any of the growth phases.

For the supernatant of *P. calathiferum*, the nutrient-replete culture was less hemolytic ($P=0.027$) than the N-deficient culture in lag phase (Figure 28). There was no significant difference between the nutrient-replete and P-deficient cultures in lag phase. The nutrient-replete culture was significantly less hemolytic than the both nutrient-deficient cultures in log and stationary phases ($P\leq 0.010$). The N-deficient culture was more hemolytic than the P-deficient culture in log ($P=0.069$) and stationary ($P=0.007$) phases, but there was no difference between the two nutrient-deficient cultures in lag phase. For the pellet of *P. calathiferum*, there were no effects of nutrient concentration on mean hemolytic activity (Figure 29).




Figure 25. Hemolytic activity by the pellet of the South Carolina *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 26. Hemolytic activity by the supernatant of the Texas *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 27. Hemolytic activity by the pellet of the Texas *Prymnesium parvum* clone grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 28. Hemolytic activity by the supernatant of *Prymnesium calathiferum* grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.




Figure 29. Hemolytic activity by the pellet of *Prymnesium calathiferum* grown under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean. (A) Means of the three trials. (B) Trial 1. (C) Trial 2. (D) Trial 3.

Ichthyotoxicity

The fish in the control beakers survived in each trial and did not show signs of ichthyotoxicity. None of the test fish displayed visible symptoms of the hemolytic part of the toxin. There was no visible bleeding from the snout or the gills, and there was no release of blood into the surrounding medium.

The nutrient-replete *P. calathiferum* culture was more ichthyotoxic than the *P. parvum* cultures. Two fish were killed in one hour in the 100% concentration of *P. calathiferum*, while only one fish was killed in three hours in the 100% concentration of the TX clone (Table 4). Fish were not killed in any of the concentrations for the other *P. parvum* clones. *Prymnesium calathiferum* was also the least dense and least hemolytic of the four clones for this nutrient treatment (Figures 30-31).

The P-deficient *P. parvum* cultures were the most ichthyotoxic of the nutrient treatments. For the NC clone, fish were killed at all concentrations of the P-deficient culture. Only one fish was killed with the N-deficient NC culture and no fish were killed with the nutrient-replete NC culture. For the SC clone, only the P-deficient culture caused fish kills. For the TX clone, all six fish were killed at all culture concentrations of the P-deficient cultures. Three fish were killed with the N-deficient TX culture and one was killed in the nutrient-replete TX culture.

The TX clone was the most ichthyotoxic, most hemolytic, and least dense of the *P. parvum* clones for all nutrient treatments (Table 4, Figures 30-31). The TX clone killed fish in all nutrient treatments, while the SC clone only killed fish under P-deficient conditions and the NC clone only killed under nutrient-deficient conditions. In addition, it took less time to kill fish exposed to the TX clone compared with the NC and SC clones. It took only one hour to kill fish

Table 4. Time (in hours) required to kill two fish in each concentration for three geographically-distinct clones of *Prymnesium parvum** and *P. calathiferum*.

Nutrient Treatment	Concentration	NC	SC	TX	Pcal
Replete					
	10%	—	—	—	—
	50%	—	—	—	—
	100%	—	—	3**	1.5
N-deficient					
	10%	—	—	—	—
	50%	—	—	3**	—
	100%	4**	—	3	—
P-deficient					
	10%	6**	—	2	—
	50%	5.5	—	1	5.5**
	100%	4	5.5	1	—

* NC=North Carolina; SC=South Carolina; TX=Texas

**=only 1 fish killed; — = no fish killed




Figure 30. Cell densities for the fish bioassays of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* under different nutrient treatments (Replete, N-Deficient, P-Deficient).




Figure 31. Hemolytic activity for the fish bioassays of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* under different nutrient treatments (Replete, N-Deficient, P-Deficient). Error bars represent one standard deviation about the mean (n=8). (A) Hemolytic activity by the supernatant. (B) Hemolytic activity by the pellet.

with the TX clone, while it took over three hours for the NC and SC clones to kill fish. For the 100% concentration of the phosphorus-deficient cultures, it took the NC clone 4 hours to kill fish, while TX killed fish in 2.5 hours.

All *P. parvum* clones were more hemolytic in nitrogen-deficient conditions, yet more ichthyotoxic under phosphorus-deficient conditions (Table 4, Figures 30-31).

DISCUSSION

Growth

The *P. parvum* clones had similar growth rates for the nutrient-replete treatments, but had differences when grown in nutrient-deficient media. All four clones experienced smaller terminal densities in the nutrient-deficient treatments, indicating that nutrient limitation caused limited growth. The TX clone, however, had lower growth rates than the SC clone for the N-deficient treatment and both the NC and SC clones in the P-deficient treatment. As mentioned previously, the TX clone showed no growth after two weeks of pre-conditioning in nutrient-deficient media. Even with only one week of pre-conditioning, growth was severely limited with smaller terminal densities.

Hemolytic Activity

There were more similarities than differences in hemolytic activity among the *P. parvum* clones. In the nutrient-replete treatment, there were no differences among the clones. In the N-deficient treatment, clonal differences occurred in the supernatant only with the NC and SC clones having similar hemolytic activity and the TX clone being less hemolytic. For the P-deficient treatment, clonal differences were seen only in the individual growth phases. For the

supernatant, the TX clone was less hemolytic than the NC clone in lag phase, but more hemolytic than the SC clone in stationary phase. For the pellet, the TX clone was more hemolytic than the NC and SC clones in stationary phase.

Overall, the NC and SC clones were more similar to each other than to the TX clone. The NC and SC clones had similar hemolytic activity, but each deviated from the TX clone on various occasions.

Hemolytic Activity: Effect of Nutrient Concentration

The effect of nutrient concentration on hemolytic activity varied with clone. For all *P. parvum* clones, the N-deficient cultures were more hemolytic than the nutrient-replete cultures.

For the NC and SC clones, both nutrient-deficient cultures were more hemolytic than the nutrient-replete cultures for the supernatant, while the P-deficient cultures were similar to or more hemolytic than the nutrient-replete cultures for the pellet. This result agrees with the study of Hagstrom and Granéli (2005), which found hemolytic activity in nutrient-sufficient and P-deficient cultures to be similar. Also for the NC and SC clones, the N-deficient cultures were similar to or more hemolytic than the P-deficient cultures. This finding agrees with Johansson and Granéli (1999), who found N-deficient cultures to be more hemolytic on a per cell basis than P-deficient cultures, though their finding was not statistically significant.

For the TX clone, both nutrient-deficient cultures were more hemolytic than the nutrient-replete cultures, and there was no difference in hemolytic activity between the nutrient-deficient cultures. This was similar to the finding by Johansson and Granéli (1999), which looked at total hemolytic activity.

Based on the findings of this study, nitrogen deficiency caused greater hemolytic activity than nitrogen sufficiency for *P. parvum*. Phosphorus deficiency (compared with phosphorus sufficiency) caused greater hemolytic activity in the supernatant and had greater or similar hemolytic activity in the pellet depending on clone and growth phase. Nitrogen deficiency caused similar or greater hemolytic activity than phosphorus deficiency depending on clone and growth phase.

Ichthyotoxicity

For the fish bioassay, the TX clone was the most ichthyotoxic of the *P. parvum* clones. The TX clone was the most stressed by nutrient deficiencies in terms of growth, and while this did not correlate well with hemolytic activity, more nutrient-stressed in this case correlated with being more ichthyotoxic. The TX clone was the most hemolytic on a per cell basis.

If a water sample containing *P. parvum* was hemolytic, there might be a higher chance of having a fish kill. More studies are needed, however. Previous reports found no relationship between hemolytic activity and ichthyotoxicity (Simonsen and Moestrup 1997). Kim and Padilla (1977) found three prymnesin fractions that were hemolytic and only one fraction that was both ichthyotoxic and hemolytic. Both the hemolysis assay and fish bioassay relate information about the toxicity of *Prymnesium*, but it remains to be seen whether the two toxin properties are related.

The nutrient-replete *P. parvum* cultures showed minimal ichthyotoxicity. Previous studies found *P. parvum* to be ichthyotoxic even under nutrient-replete conditions (Yariv and Hestrin 1961, Reich and Parnas 1962, Ulitzur and Shilo 1964, Kim and Padilla 1977). Inactivation of ichthyotoxicity under laboratory settings can occur with constant illumination,

heating, high pH, and high salinity (Reich and Parnas 1962, Parnas *et al.* 1962, Parnas 1963, Ulitzur and Shilo 1964, Shilo 1981). None of these factors occurred during the experiment.

Nutrient-deficient cultures were the most ichthyotoxic, particularly the P-deficient cultures. Fish kills have mostly occurred in high nitrogen P-limited systems, although Moustaka-Gouni *et al.* (2004) did report a fish kill in a N-limited waterbody. Since all the clones came from regions associated with P-limited conditions at the time of the bloom (Lewitus *et al.* 2003, Tomas *et al.* 2004), finding that the clones were most ichthyotoxic during P-deficient conditions was consistent with field observations. The N:P ratios in this study were consistent with natural conditions.

Extrapolating this bioassay into natural environments, if a eutrophic waterbody experienced a *P. parvum* bloom, fish kills would likely occur. If a non-eutrophic waterbody experienced a *P. parvum* bloom, fish kills may or may not occur. Since only one trial was done for each treatment, solid conclusions cannot be drawn from the fish bioassays. It should also be noted that the *P. parvum* densities used in this fish bioassay, approximately 10^2 cells/L, were well below documented densities for fish kills. While the number of *P. parvum* cells is never an indicator of whether or not a fish kill will occur, studies have shown that at least 10^4 cells/L are required (Reich and Aschner 1947). Most fish kills occur with blooms between 10^6 and 10^9 cells/L (Edwardsen and Paasche 1998). In this study, the TX was the most ichthyotoxic clone while being the least dense.

In this study, the clones that came from samples with fish kills (NC and TX clones) were the most ichthyotoxic. In this case, it seems that once a bloom has killed fish, it may be more likely to kill fish again in the future. Most areas have recurrent fish kills associated with *P. parvum* blooms. *Prymnesium* may produce resting stages in natural environments. These resting

stages that exist in other harmful algal bloom species including *Gymnodinium catenatum*, *Pyrodinium bahamense*, and species of *Alexandrium* are a period of dormancy brought on by adverse environmental conditions (Anderson 1998, Usup and Azanza 1998, Amorim *et al.* 2002). Despite the lack of information on resting stages in *Prymnesium*, this by no means suggests that the cysts do not exist. *Prymnesium* cysts would be very small and difficult to detect, especially in a small population size. The next step in future *Prymnesium* research is to confirm the presence of these cysts and their ability to form motile populations.

What might be an explanation for the differences in the *P. parvum* clones? The basic assumption is that the three clones should behave similarly (similar growth rates, hemolytic activity, and ichthyotoxicity) because they are *P. parvum*. However, the clones did show differences. These differences may be attributed to different environmental conditions, including geographical origin, type of waterbody, and whether fish kills occurred. Another explanation may be the genetic makeup of the different clones. Further studies should examine the genetic differences between the NC, SC, and TX clones. While this study focused on physiological ecology, other studies should compare the toxins from each clone. Six hemolytic compounds have recently been isolated for the NC clone (Wright *et al.* 2005). Further experiments should also compare clones from the same geographic area, but one from a natural waterbody and one from a manmade waterbody.

Another question is whether clones in the same bloom vary in characteristics. This is a fundamental problem in the study of harmful algal blooms – if a bloom occurs, do all the clones respond in the same manner? This study examined geographical differences. One solution to this problem would be to isolate many different clones from the same bloom and test genetic

differences. The development of genetic fingerprinting for HAB species, including *P. parvum*, is critical.

Variation in harmful algal bloom species is not rare. Larsen published several studies looking at three geographically-distinct clones of *P. parvum* from Norway, Denmark, and England (Larsen and Medlin 1997, Larsen and Edvardsen 1998, Larsen and Bryant 1998). The studies found that the clones from the same area were more physiologically similar than clones from different areas of the same species. The three *P. parvum* clones did display differences in genetics, growth rate, and toxicity (Larsen and Bryant 1998). Intraspecific variation has also been found in non-*Prymnesium* species. Strom and Bright (2003) found intraspecific variation in organic and inorganic nitrogen requirements for the haptophyte *Emiliana huxleyi*. Doblin *et al.* (2000) found intraspecific variation in selenium requirements for the dinoflagellate *Gymnodinium catenatum*.

Comparison between *P. parvum* and *P. calathiferum*

There were notable differences between the *P. parvum* and *P. calathiferum* cultures. When *P. calathiferum* reached stationary phase, clumps of particles appeared along with an unpleasant odor. Neither characteristic was ever associated with the *P. parvum* clones. *Prymnesium calathiferum* reached stationary phase more rapidly, normally having lower terminal densities, and had the highest growth rates. So, why does *P. calathiferum* grow faster than *P. parvum*? These are the first growth rates on *P. calathiferum* and there are no known published growth rates for comparison. In the short term, *P. calathiferum* may have some competitive advantage in being able to grow faster.

Relative to hemolytic activity, *P. parvum* was highly hemolytic, while *P. calathiferum* showed limited hemolytic activity. *Prymnesium calathiferum* was used as a negative control in this study. Again, these are the first observations on *P. calathiferum*.

However, *P. calathiferum* was reported as being ichthyotoxic. In a previous study, the supernatant from a nutrient-replete *P. calathiferum* culture killed fish in three hours (Chang 1985). In the fish bioassays performed for this study, the nutrient-replete *P. calathiferum* culture killed fish in 1.5 hours while the nutrient-deficient cultures killed no fish. The fish killed at the 50% concentration of the P-deficient culture was probably not due to the culture since the fish at the 100% concentration were not killed. From these results, it is unlikely that fish would be killed in a eutrophic waterbody where a *P. calathiferum* bloom formed.

More work needs to be done on *P. calathiferum*. The initial finding of *P. calathiferum* was from a bloom where it was not even the dominant phytoplankton (Chang and Ryan 1985). From this study, there is a distinct difference in toxicity between *P. parvum* and *P. calathiferum*. Since *P. parvum* produces prymnesin, it is assumed that *P. calathiferum* also produces the same toxin. However, there are no studies confirming that *P. calathiferum* produces prymnesin.

Hemolytic Activity: Supernatant versus Pellet

This study also examined where hemolytic activity was the greatest – in the cells (represented by the pellet) or released into the surrounding medium (represented by the supernatant). Prymnesin is an unusual toxin in that it is extracellular and released into the medium (Shilo and Aschner 1953). This should mean that the extracellular component (the supernatant in this study) should be the most toxic. Studies have found the extracellular component to be highly hemolytic (Simonsen and Moestrup 1997, Fistarol *et al.* 2003).

The results from this study vary according to clone and nutrient treatment. For the P-deficient cultures of the four clones, hemolytic activity in the cells equaled that in the surrounding medium. For the N-deficient cultures, the hemolytic activity in the extracellular component was greater in the NC, SC, and *P. calathiferum* clones, while the hemolytic activity in the cells were greater in the TX clone. For the nutrient-replete cultures, hemolytic activity was greater in the cells than was released for the TX clone, while hemolytic activity was equally distributed between the intracellular and extracellular components of the NC, SC, and *P. calathiferum* clones.

Why is the media more hemolytic than the cells if the toxin is extracellular? Possibly there was no trigger to release toxin. Nitrogen deficiency was the only nutrient treatment to show the supernatant being most hemolytic. One suggestion is that the membranes are less intact and more leaky due to imbalanced metabolism from nitrogen deficiency. Being stressed by limited nutrients is thought to lead to an imbalance in metabolism, such that the toxin is expelled through leaks in the membrane (Dafni *et al.* 1972). The TX clone was the only organism to show the pellet being the most hemolytic. Perhaps membrane leakage is less likely in nitrogen deficiency for the TX clone.

Hemolytic Activity: Growth Phase

Hemolytic activity was also assessed at different growth phases – lag phase, log phase, and stationary phase. As a secondary metabolite, toxins should accumulate when the organism is in active growth and be at their highest levels in stationary phase (Calvo *et al.* 2002).

This study found that there was either no difference among the growth phases or that lag phase was the most hemolytic. Both findings were very inconsistent with previous studies. A

few studies on *P. parvum* have found hemolysis to be highest in stationary phase (Shilo and Rosenberger 1960, Igarashi *et al.* 1995, Rosetta *et al.* 2003). Padilla (1970) found that the hemolysin accumulated in log phase. Simonsen and Moestrup (1997) found that hemolytic activity was greatest in log phase, and that lag and stationary phases were similar to each other and less hemolytic. One study that looked at the dinoflagellate *Alexandrium* found that total hemolysis increased with growth, but that hemolysis on a per cell basis decreased from log to stationary phases (Arzul *et al.* 1999).

In addition, finding lag phase to be the most hemolytic or similar to later phases in the supernatant was also inconsistent with previous studies. Simonsen and Moestrup (1997) found hemolytic activity in the medium to be highest in stationary phase, but not present in lag or log phases. Shilo (1967) found that in lag phase, only the intracellular toxin was present. The extracellular toxin was not present until later in the growth cycle.

The best explanation for lag phase being the most hemolytic would be carryover from inoculation. For the nutrient-replete cultures, they were inoculated from cultures in stationary phase where hemolysin production is known to occur – not only accumulating in the cells but also released into the surrounding medium. When cultures are inoculated, they receive the cells and the medium, which have both presumably accumulated the toxin. Lag phase being more hemolytic than stationary phase could be due to toxin release by stressed organisms. Inoculation into a new surrounding would seem to be stressful – if not, lag phase would not exist, and the cells would start multiplying rapidly. Possibly this stress may cause the cells to rupture and release prymnesin. Nonetheless, it would be impossible for hemolysin production to occur in lag phase. Enzyme function is thought to be low in lag phase. Most enzymes are inactivated during

stationary phase, and during lag phase, the organisms are replenishing their nutrient supply in order to re-activate the enzymes and rapidly reproduce (Fogg and Thake 1987).

Summary

As this study found, intraspecific variation in growth rate, hemolytic activity, and ichthyotoxicity did occur in geographically-distinct strains of *P. parvum*. The next step would be to examine genetic differences. The question as to whether nitrogen or phosphorus limitation greater affects toxicity in *P. parvum* still does not have a firm answer. Experiments using different forms of nitrogen and using different concentrations of nitrate and phosphate should be done to look at the effects on toxicity, particularly hemolytic activity.

Observations of low hemolytic activity and ichthyotoxicity limited to nutrient-replete conditions suggest *P. calathiferum* should pose a minimal problem. However, observations of high hemolytic activity and ichthyotoxicity support *P. parvum* being a major problem. The question exists why has the United States recently started experiencing *P. parvum* blooms? Has it always been there and just not noticed? Alternatively, has it been introduced from other areas? The larger question is how do you get rid of the *P. parvum* blooms? Two mitigation and control problems exist – just the bloom and the toxicity of the bloom. Eliminating the organism all together poses many problems – in a manmade waterbody, such as the NC and SC ponds, elimination would be an easier problem than in a natural system such as TX. There are many management methods, including ammonium sulfate, copper sulfate, potassium permanganate, hydrogen peroxide, and clay. Each has its set of problems (Barkoh and Fries 2005, Hagstrom and Graneli 2005).

This study suggests that eutrophication may increase the ichthyotoxicity and possibly the hemolytic activity of *P. parvum* blooms. Waterbodies affected by *P. parvum* should be monitored for nitrate, ammonia, and phosphate concentrations. When the waters become severely N- or P-limited, decreasing the nutrient source may be beneficial to decrease the potential of fish kills.

CONCLUSIONS

- Intraspecific variation in growth rate, hemolytic activity, and ichthyotoxicity occurred in the three clones of *P. parvum* used in this study.
- *Prymnesium parvum* and *P. calathiferum* showed discernible differences in growth rate, hemolytic activity, and ichthyotoxicity.
- Nitrogen deficiency caused greater hemolytic activity than nitrogen replete conditions for *P. parvum*.
- Phosphorus deficiency (compared with phosphorus replete conditions) caused greater hemolytic activity in the supernatant and had greater or similar hemolytic activity in the pellet depending on clone and growth phase.
- Nitrogen deficiency caused similar or greater hemolytic activity than phosphorus deficiency depending on clone and growth phase.
- *Prymnesium calathiferum* cultures grown under nutrient-replete conditions were more ichthyotoxic than those under nutrient-deficient conditions.
- Phosphorus deficiency caused the greatest ichthyotoxicity in the *P. parvum* clones.
- There was no difference in hemolytic activity between the supernatant and the pellet in the phosphorus-deficient treatment for the four clones. For the nutrient-replete treatment,

there was no difference between the supernatant and pellet for the NC, SC, and *P. calathiferum* clones.

- For the nitrogen-deficient treatment, the supernatant was the most hemolytic for the NC, SC, and *P. calathiferum* clones.
- For the nutrient-replete and nitrogen-deficient treatments, the pellet was the most hemolytic for the TX clone.
- There was either no difference in hemolytic activity among the growth phases or lag phase was the most hemolytic.

LITERATURE CITED

- Amorin, A., B. Dale, R. Godinho, and V. Brotas. 2002. *Gymnodinium catenatum*-like cysts (Dinophyceae) in recent sediments from the coast of Portugal. *Phycologia*. 40(6): 572-582.
- Anderson, D.M. 1998. Physiology and bloom dynamics of toxic *Alexandrium* species, with emphasis on life cycle transitions. NATO ASI Series. V G 431. In: *Physiological Ecology of Harmful Algal Blooms*. Edited by D.M. Anderson, A.D. Cembella, and G.M. Hallegraeff. 29-48.
- Arizona Game and Fish Department website. Golden alga frequently asked questions. http://www.azgfd.gov/nrm/temp/golden_alga_faqs.shtml#3. Accessed 13 November 2005
- Aure, J. and F. Rey. 1992. Oceanographic conditions in the Sansfjord system, western Norway, after a bloom of the toxic prymnesiophyte *Prymnesium parvum* Carter in August 1990. *Sarsia*. 76: 247-254.
- Arzul, G., P. Gentien, and M. Crassous Ifremer. 1994. A haemolytic test to assay toxin excreted by the marine dinoflagellate *Gyrodinium cf. aureolum*. *Wat. Res.* 28(4): 961-965.
- Barkoh, A. and L.T. Fries. 2005. Management of *Prymnesium parvum* at Texas state fish hatcheries. TPWD Management Data Series. No. 236. 104 pp. http://www.tpwd.state.tx.us/landwater/water/environconcerns/hab/habwork/papers/pp_mangement.phtml. Accessed 8 November 2005.
- Calvo, A.M., R.A. Wilson, J.W. Bok, and N.P. Keller. 2002. Relationship between secondary metabolism and fungal development. *Microbiol. Mol. Biol. Rev.* 66(3): 447—489.
- Carter, N. 1937. New or interesting algae from brackish water. *Arch. Protistenk.* 90: 1-68
- Chang F.H. 1985. Preliminary toxicity test of *Prymnesium calathiferum* n. sp. isolated from New Zealand. In: *Toxic Dinoflagellates*. Ed. by D.M. Anderson, A.W. White and D.G. Baden., pp. 109-112. Elsevier, New York.
- Chang F.H. and K.G. Ryan. 1985. *Prymnesium calathiferum* sp. nov. (Prymnesiophyceae), a new species isolated from Northland, New Zealand. *Phycologia*. 24: 191-198.
- Colorado Department of Natural Resources Website. Golden Algae suspected in fish kill. <http://www.dnr.state.co.us/news/press.asp?pressid=1922> Accessed 13 November 2005.
- Dafni, Z. and E. Giberman. 1972. Nature of initial damage to Ehrlich ascites cells caused by *Prymnesium parvum* toxin. *Biochim. Biophys Acta.* 255: 380-385.
- Dafni, Z., S. Ulitzur, and M. Shilo. 1972. Influence of light and phosphate on toxin production and growth of *Prymnesium parvum*. *J General Microbiology.* 70: 199-207.
- Doblin, M.A., S.I. Blackburn, and G.M. Hallegraeff. 2000. Intraspecific variation in the selenium requirement of different geographic strains of the toxic dinoflagellate *Gymnodinium catenatum*. *J. Plankton Research.* 22: 421-432.
- Edvardsen, B. and E. Paasche. 1998. Bloom dynamics and physiology of *Prymnesium* and *Chrysochromulina*. NATO ASI Series. V G 431. In: *Physiological Ecology of Harmful Algal Blooms*. Edited by D.M. Anderson, A.D. Cembella, and G.M. Hallegraeff. 193-208.

- Eschbach, E., J.P. Scharsack, U. John, and L.K. Medlin. 2001. Improved erythrocyte lysis assay in microtitre plates for sensitive detection and efficient measurement of haemolytic compounds from ichthyotoxic algae. *J. Applied Toxicology*. 21: 513-519.
- Fistarol G.O., C. Legrand, and E. Granéli. 2003. Allelopathic effect of *Prymnesium parvum* on a natural plankton community. *Marine Ecology Progress Series*. 255: 115-125
- Fogg, G.E. and B. Thake 1987. *Algal cultures and phytoplankton ecology*. 3rd edition. Madison: U. of Wisconsin Press, 269 pp.
- Granéli, E. and N. Johansson. 2003a. Increase in production of allelopathic substances by *Prymnesium parvum* cells grown under N- or P-deficient conditions. *Harmful Algae*. 2: 135-145.
- Granéli, E. and N. Johansson. 2003b. Effects of the toxic haptophyte *Prymnesium parvum* on the survival and feeding of a ciliate: the influence of different nutrient conditions. *Marine Ecology Progress Series*. 254: 49-56.
- Green, J.C., D.F. Hibberd, and R.N. Pienaar. 1982. The taxonomy of *Prymnesium* (Prymnesiophyceae) including a description of a new cosmopolitan species, *P. patellifera* sp. Nov., and further observations on *P. parvum* N. Carter. *British Phycological Journal*. 17: 363-382.
- Guillard, K. 1973. Division Rates. In: Stein, J.R. *Handbook of Phycological Methods Culture Methods and Growth Measurements*. Cambridge University Press, New York, pp. 289-31. Guillard 1973
- Guillard R.R.L. and J.H. Ryther 1962. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* Cleve. *Can. J. Microbiol.* 8: 229-239
- Guo, M, P.J. Harrison, and F.J.R Taylor. 1996. Fish kills related to *Prymnesium parvum* N. Carter (Haptophyta) in the People's Republic of China. *J. Applied Phycology*. 8: 111-117.
- Hagstrom, J.A., and E Granéli. 2005. Removal of *Prymnesium parvum* (Haptophyceae) cells under different nutrient conditions by clay. *Harmful Algae*. 4: 249-260.
- Holdway, P.A. and R.A. Watson. 1978. Growth and ichthyotoxicity in two strains of *Prymnesium parvum* Carter (Haptophyta) as affected by phosphorous concentration. *British Phycological J.* 13: 201-202.
- Holdway, P.A., R.A. Watson, and B. Moss. 1978. Aspects of the Ecology of *Prymnesium parvum* (Haptophyta) and water chemistry in the Norfolk Broads, England. *Freshwater Biology*. 8: 295-311.
- Hulot, F.D. and J. Huisman. 2004. Allelopathic interactions between phytoplankton species: the roles of heterotrophic bacteria and mixing intensity. *Limnology and Oceanography*. 49(4): 1424-1434.
- Igarashi, T., M. Satake, and T. Yasumoto. 1999. Structures and partial stereochemical assignments for prymnesin-1 and prymnesin-2: potent hemolytic and ichthyotoxic glycosides isolated from the red tide alga *Prymnesium parvum*. *J. American Chemical Society*. 121 (37): 8499-8511.
- Igarashi, T, S. Aritake, and T. Yasumoto. 1998. Biological activities of Prymnesin-2 isolated from a red tide alga *Prymnesium parvum*. *Nat. Toxins*. 6: 35-41.

- Igarashi, R., Y. Oshima, M. Murata, and T. Yasumoto. 1995. Chemical studies on prymnesins isolated from *Prymnesium parvum*. Harmful marine algal blooms, proceedings of the Sixth International Conference on Toxic Marine Phytoplankton, 303-308.
- James, T.L. and A. de la Cruz. 1989. *Prymnesium parvum* Carter (Chrysophyceae) as a suspect of mass mortalities of fish and shellfish communities in western Texas. *Texas J. of Science*. 41: 429-430.
- Johansson, N. and E.Granéli. 1999. Influence of different nutrient conditions on cell density, chemical composition and toxicity of *Prymnesium parvum* (Haptophyta) in semi-continuous cultures. *J. Experimental Marine Biology and Ecology*. 239: 243-258.
- Kaartvedt, S., T.M. Johnsen, D.L. Aksnes, U. Lie, and H. Svendsen. 1991. Occurrence of the toxic phytoflagellate *Prymnesium parvum* and associated fish mortality in a Norwegian fjord system. *Canadian J. of Fisheries and Aquatic Sciences*. 48: 2316-2323.
- Kim, Y.S. and G.M. Padilla. 1977. Hemolytically active components from *P. parvum* and *G. breve* toxins. *Life Sciences*. 21: 1287-1292.
- Krasnoshchek, G.P. and L.S. Abramovich. 1971. Mass development of *Prymnesium parvum* Cart. in fish-breeding ponds. *Hydrobiol. J.* 7:54-55.
- Larsen, A. and L.K. Medlin. 1997. Inter- and intraspecific genetic variation in twelve *Prymnesium* (Haptophyceae) clones. *J. Phycol.* 33: 1007-1015.
- Larsen, A. and B. Edvardsen. 1998. Relative ploidy levels in *Prymnesium parvum* and *P. patelliferum* (Haptophyta) analyzed by flow cytometry. *Phycologia*. 37(6): 412-424.
- Larsen, A. and S. Bryant. 1998. Growth rate and toxicity of *Prymnesium parvum* and *Prymnesium patelliferum* (Haptophyta) in response to changes in salinity, light and temperature. *Sarsia*. 83(5): 409-418.
- Larsen, A. 1999. *Prymnesium parvum* and *P. patelliferum* (Haptophyta) – one species. *Phycologia*. 38(6): 541-543.
- Lewitus, A. J., L.B. Schmidt, L.J. Mason, J.W. Kempton, S.B. Wilde, J.L. Wolny, B. J. Williams, K. C. Hayes, S. N. Hymel, C. J. Keppler, A. H. Ringwood. 2003. Harmful algal blooms in South Carolina residential and golf course ponds. *Population and Environment*. 24 (5):387-413.
- Manton, I and G.F. Leedale. 1963. Observations on the fine structure of *Prymnesium parvum* Carter. *Arch. Mikrobiol.* 45: 285-303.
- Meldahl, A.S., B. Edvardsen, and F. Fonnum. 1994. Toxicity of four potentially ichthyotoxic marine phytoflagellates determined by four different test methods. *J. Toxicology and Environmental Health*. 42: 289-301.
- Morohashi, A., M. Satake, Y. Oshima, T. Igarashi, and T. Yasumoto. 2001. Absolute configuration at C14 and C85 in Prymnesin-2, a potent hemolytic and ichthyotoxic glycoside isolated from the red tide alga *Prymnesium parvum*. *Chirality*. 13: 601-605.
- Moustaka-Gouni, M., C.M. Cook, S. Gkelis, E. Michaloudi, K. Pantlelidakis, M. Pyrovetsi, and T. Lanaras. 2004. The coincidence of a *Prymnesium parvum* bloom and the mass kill of birds and fish in Lake Koronia. *Harmful Algae News*. 26: 1-2.
- Padilla, G.M. 1970. Growth and toxigenesis of the chrysomonad *Prymnesium parvum* as a function of salinity. *J Protozool*. 17:456-462.
- Parnas I., K. Reich, and F. Bergmann. 1962. Photoactivation of ichthyotoxin from axenic cultures of *Prymnesium parvum* Carter. *Appl. Microbiol.* 10: 237-239.
- Parnas, I. 1963. The toxicity of *Prymnesium parvum*. *Israel J. Zool.* 12: 15-23.

- Parnas, I. and B.C. Abbott. 1965. Physiological activity of the ichthyotoxin from *Prymnesium parvum*. *Toxicon*. 3: 133-145.
- Reich, K and M. Aschner. 1947. Mass development and control of the phytoflagellate *Prymnesium parvum* in fish ponds in Palestine. *Palestine J. of Botany*. 4:14-23.
- Reich, K. and I. Parnas 1962. Effect of illumination on ichthyotoxin in an axenic culture of *Prymnesium parvum* Carter. *J. Protozool*. 9:38-40.
- Rhoades, K. and C. Hubbs. 1992. Recovery of Pecos River fishes from a red tide fish kill. *The Southwestern Naturalist*. 37(2):178-187.
- Rosetta, C.H. and G.B. McManus. 2003. Feeding by ciliates on two harmful algal bloom species, *Prymnesium parvum* and *Prorocentrum minimum*. *Harmful Algae*. 2:109-126.
- Sabour, B, M. Loudiki, B. Oudra, S. Oubraim, B. Fawzi, S. Fadlaoui, M. Chlaida and V. Vasconcelos. 2000. Blooms of *Prymnesium parvum* associated with fish mortalities in a hypereutrophic brackish lake in Morocco. *Harmful Algae News*. 21: 8-9.
- Shilo, M. and M. Aschner. 1953. Factors governing the toxicity of cultures containing the phytoflagellate *Prymnesium parvum* Carter. *J. General Microbiology*. 8: 333-343.
- Shilo, M. 1967. Formation and mode of action of algal toxins. *Bacteriological Reviews*. 31: 180-193.
- Shilo, M and R.F. Rosenberger. 1960. Studies on the toxic principles formed by the chryomonad *Prymnesium parvum* Carter. *Annals of New York Academy of Sciences*. 90:866-876.
- Shilo, M. 1981. The toxic principles of *Prymnesium parvum*. *Environmental Science Research*. 20: 37-47.
- Simonsen, S. and O. Moestrup. 1997. Toxicity tests in eight species of *Chrysochromulina* (Haptophyta). *Can J Bot*. 75: 129-136.
- Strom, S.L. and K. Bright. 2003. Effects of planktonic food webs of organic nitrogen use by bloom-forming phytoplankton. Abstract in Proceedings of the Biennial Meeting of the Estuarine Research Federation. Seattle, WA, USA, ERF, Port Republic, MD, USA Texas Parks and Wildlife Department website. Golden Alga in Texas. <http://www.tpwd.state.tx.us/hab/ga> Accessed 10 November 2005.
- Tomas, C., J. Glass, J. Ralph, and A. Lewitus. 2004. Blooms of the ichthyotoxic flagellate *Prymnesium parvum* in U.S. waters: an emerging or a perennial problem? In: Steidinger, K.A., Lansberg, J.H., Tomas, C.R., and G.A. Vargo. (Eds). *Harmful Algae 2002*. Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of UNESCO.
- Ulitzur, S. and M. Shilo. 1964. A sensitive assay system for determination of the ichthyotoxicity of *Prymnesium parvum*. *J. General Microbiol*. 36:161-169.
- Ulitzur, S. and M. Shilo. 1966. Mode of action of *Prymnesium parvum* ichthyotoxin. *J. Protozool*. 13:322-336.
- Ulitzur, S and M. Shilo. 1970. Procedure for purification and separation of *Prymnesium parvum* toxins. *Biochim. Biophys Acta*. 201: 350-363.
- Usup, G., and R.V. Azanza. 1998. Physiology and bloom dynamics of the tropical dinoflagellate *Pyrodinium bahamense*. NATO ASI Series. V G 431. In: *Physiological Ecology of Harmful Algal Blooms* Edited by D.M. Anderson, A.D. Cembella, and G.M. Hallegraeff. 81-94.
- Wortley, J.S. and G.L. Phillips. 1987. Fish mortalities and *Prymnesium* in the Norfolk Broads. *Institute of Fisheries Management*. 18: 152-162.

- Wright, J.L.C., C. Tomas, M. Satake, N. Alam, T. Shuster. 2005. Structure of several new hemolytic toxins from strains of *Prymnesium parvum* isolated from fish ponds in North Carolina. *Proceedings of the 3rd Symposium on Harmful Algae in the U.S.* Pacific Grove, CA., October 2-7.
- Yariv, S. and S. Hestrin. 1961. Toxicity of the extracellular phase of *Prymnesium parvum* cultures. *J. Gen. Microbiol.* 24:165-175.

APPENDICES

Appendix A. Daily Counts of Cultures.

North Carolina *P. parvum* clone – Trial 1 of Nutrient-Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	968	19360	20510	12	7646	611680	626900
	976	19520			8173	653840	
	1121	22420			7323	585840	
	1037	20740			8203	656240	
1	1781	35620	33830	13	8342	667360	669580
	1729	34580			8383	670640	
	1618	32360			8421	673680	
	1638	32760			8333	666640	
2	2717	54340	54470	14	8544	683520	676160
	2729	54580			8392	671360	
	2685	53700			8284	662720	
	2763	55260			8588	687040	
3	4296	85920	76984	15	9376	750080	756080
	4128	82560			9523	761840	
	4031	80620			9687	774960	
	4028	80560			9218	737440	
4	5768	115360	118715	16	9790	783200	783460
	6093	121860			9690	775200	
	5834	116680			9896	791680	
	6048	120960			9797	783760	
5	8692	173840	173725	17	10357	828560	847600
	8769	175380			10510	840800	
	8603	172060			10798	863840	
	8681	173620			10715	857200	
6	11508	230160	224930	18	11035	882800	883800
	11321	226420			11079	886320	
	10867	217340			11153	892240	
	11290	225800			10923	873840	
7	4042	323360	303660	19	12075	966000	981300
	3887	310960			12426	994080	
	3749	299920			12266	981280	
	3505	280400			12298	983840	
8	5026	402080	379640	20	12929	1034320	1020080
	4581	366480			12478	998240	
	4745	379600			12976	1038080	
	4630	370400			12621	1009680	
9	5364	429120	401520	21	11277	902160	933780
	4608	368640			11936	954880	
	5174	413920			11510	920800	
	4930	394400			11966	957280	
10	6153	492240	483100	22	13149	1051920	1040140
	6004	480320			12916	1033280	
	5881	470480			12908	1032640	
	6117	489360			13034	1042720	
11	6719	537520	537480				
	6767	541360					
	6630	530400					
	6758	540640					

South Carolina *P. parvum* clone – Trial 1 of Nutrient-Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	622	12440	12465	12	6589	527120	501140
	572	11440			5579	446320	
	672	13440			6864	549120	
	627	12540			6025	482000	
1	1057	21120	20225	13	6844	547520	563320
	995	19880			7091	567280	
	899	17980			7069	565520	
	1097	21920			7162	572960	
2	1787	35740	36865	14	7776	622080	628620
	1755	35100			7843	627440	
	1813	36260			7866	629280	
	2018	40360			7946	635680	
3	3074	61480	66305	15	8437	674960	688200
	3543	70860			8343	667440	
	3145	62900			8622	689760	
	3499	69980			9008	720640	
4	4972	99440	101535	16	9942	795360	793360
	4977	99540			9872	789760	
	5092	101840			9758	780640	
	5266	105320			10096	807680	
5	7224	144480	145995	17	9584	766720	782300
	7228	144560			9673	773840	
	7326	146520			9814	785120	
	7421	148420			10044	803520	
6	7224	144480	224460	18	9996	799680	801120
	7228	144560			10030	802400	
	7326	146520			9940	795200	
	7421	148420			10090	807200	
7	10480	209600	292540	19	12323	985840	980140
	11806	236120			12301	984080	
	11109	222180			12198	975840	
	11497	229940			12185	974800	
8	3499	279920	373973	20	11712	936960	943060
	3742	299360			11685	934800	
	3636	290880			11902	952160	
	4178	334240			11854	948320	
9	3805	304400	323360	21	12402	992160	988280
	3914	313120			12507	1000560	
	4271	341680			12249	979920	
	4373	349840			12256	980480	
10	4715	377200	370587	22	13197	1055760	1051900
	4809	384720			13070	1045600	
	5217	417360			13230	1058400	
	5314	425120			13098	1047840	
11	5198	415840	422260				
	5384	430720					

Texas *P. parvum* clone – Trial 1 of Nutrient-Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	1124	22480	22620	12	7909	632720	640900
	1137	22740			8220	657600	
	1173	23460			7942	635360	
	1090	21800			7974	637920	
1	2757	55140	56920	13	8348	667840	659260
	2831	56600			8124	649920	
	2849	56980			8163	653040	
	2948	58960			8328	666240	
2	4523	90460	89985	14	8843	707440	712500
	4427	88540			8903	712240	
	4568	91360			8868	709440	
	4479	89580			9011	720880	
3	6804	136080	136880	15	9487	758960	756420
	6889	137780			9469	757520	
	6690	133800			9218	737440	
	6993	139860			9647	771760	
4	8460	169200	168790	16	10007	800560	815380
	8357	167140			10310	824800	
	8475	169500			10101	808080	
	8466	169320			10351	828080	
5	8778	175560	176340	17	10625	850000	871300
	8980	179600			10679	854320	
	8673	173460			10970	877600	
	8837	176740			11291	903280	
6	10758	215160	217210	18	11100	888000	902260
	10795	215900			11427	914160	
	10991	219820			11059	884720	
	10898	217960			11527	922160	
7	3396	271680	275040	19	12614	1009120	1009120
	3455	276400					
	3466	277280					
	3435	274800					
8	3827	306160	301640	20	12344	987520	1004140
	3686	294880			12477	998160	
	3814	305120			12567	1005360	
	3755	300400			12819	1025520	
9	4124	329920	339480	21	12322	985760	992380
	4358	348640			12192	975360	
	4359	348720			12554	1004320	
	4133	330640			12551	1004080	
10	5661	452880	453240	22	13590	1087200	1084320
	5776	462080			13407	1072560	
	5588	447040			13617	1089360	
	5637	450960			13602	1088160	
11	6383	510640	518000				
	6501	520080					
	6385	510800					
	6631	530480					

Prymnesium calathiferum – Trial 1 of Nutrient-Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	1037	20740	19830	12	6858	548640	571800
	1110	22220			7077	566160	
	872	17440			7532	602560	
	946	18920			7123	569840	
1	989	19780	14280	13	7142	571360	548784
	593	11860			6517	521360	
	603	12060			6443	515440	
	671	13420			6966	557280	
2	1126	22520	22696	14	7231	578480	476180
	1058	21160			5820	465600	
	1267	25340			5862	468960	
	1034	20680			5979	478320	
3	1189	23780	34970	15	6148	491840	548784
	1772	35440			4417	353360	
	1651	33020			4752	380160	
	1804	36080			4627	370160	
4	1767	35340	63780	16	4981	398480	407960
	3196	63920			4956	396480	
	3124	62480			5085	406800	
	3282	65640			5019	401520	
5	3154	63080	144460	17	5338	427040	412240
	7323	146460			5048	403840	
	7057	141140			5152	412160	
	7311	146220			5067	405360	
6	6934	138680	145495	18	5345	427600	392640
	7490	149800			4860	388800	
	7322	146440			4751	380080	
	7121	142420			4877	390160	
7	7551	151020	249520	19	5144	411520	617600
	7105	142100			6836	546880	
	2702	216160			6921	553680	
	3237	258960			8559	684720	
8	3356	268480	359420	20	8564	685120	501380
	3181	254480			5943	475440	
	4536	362880			5931	474480	
	4455	356400			6468	517440	
9	4468	357440	645620	21	6727	538160	508200
	4512	360960			5757	460560	
	8105	648400			6067	485360	
	8081	646480			6475	518000	
10	8059	644720	572200	22	7111	568880	672380
	8036	642880			8450	676000	
	7031	562480			8398	671840	
	6876	550080			8584	686720	
11	7282	582560	800840		8187	654960	
	7421	593680					
	9827	786160					
	9941	795280					
	10170	813600					
	10353	828240					

North Carolina *P. parvum* clone – Trial 2 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	877	17540	17815	12	8641	691280	696900
	871	17420			8730	698400	
	922	18440			8656	692480	
	893	17860			8818	705440	
1	1360	27200	28145	13	9662	772960	779980
	1418	28360			9835	786800	
	1461	29220			9417	753360	
	1390	27800			10085	806800	
2	1904	38080	38585	14	11749	939920	944600
	1863	37260			11730	938400	
	1892	37840			11819	945520	
	2058	41160			11932	954560	
3	3373	67460	75005	15	10931	874480	876520
	3582	71640			10868	869440	
	3921	78420			10963	877040	
	4125	82500			11064	885120	
4	4980	99600	108865	16	12473	997840	1003940
	5341	106820			12334	986720	
	5706	114120			12678	1014240	
	5746	114920			12712	1016960	
5	8166	163320	164995	17	12974	1037920	1058220
	7951	159020			13284	1062720	
	8438	168760			13695	1095600	
	8444	168880			12958	1036640	
6	11238	224760	219835	18	13551	1084080	1111780
	10665	213300			13750	1100000	
	10985	219700			14199	1135920	
	11079	221580			14089	1127120	
7	14602	292040	289560	19	14314	1145120	1157580
	14198	283960			14292	1143360	
	14429	288580			14446	1155680	
	14683	293660			14827	1186160	
8	17924	358480	358265	20	15642	1251360	1282760
	17626	352520			15733	1258640	
	17977	359540			15985	1278800	
	18126	362520			16778	1342240	
9	21635	432700	430620				
	20941	418820					
	21819	436380					
	21729	434580					
10	7168	573440	574000				
	7197	575760					
	7184	574720					
	7151	572080					
11	4608	368640	557560				
	7637	610960					
	7728	618240					
	7905	632400					

South Carolina *P. parvum* clone – Trial 2 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	517	10340	10885	12	6760	540800	548340
	528	10560			6812	544960	
	569	11380			6811	544880	
	563	11260			7034	562720	
1	947	18940	19265	13	6904	552320	599300
	919	18380			7460	596800	
	1011	20220			7463	597040	
	976	19520			8138	651040	
2	1312	26240	27380	14	8542	683360	753180
	1209	24180			9538	763040	
	1408	28160			8841	707280	
	1547	30940			10738	859040	
3	2204	44080	45665	15	8076	646080	664840
	2145	42900			8132	650560	
	2307	46140			8470	677600	
	2477	49540			8564	685120	
4	3052	61040	65685	16	8753	700240	705720
	3165	63300			8598	687840	
	3376	67520			8911	712880	
	3544	70880			9024	721920	
5	4822	96440	97570	17	8755	700400	720140
	4672	93440			8825	706000	
	4889	97780			8896	711680	
	5131	102620			9531	762480	
6	5939	118780	121660	18	9136	730880	737820
	6072	121440			8805	704400	
	5966	119320			9189	735120	
	6355	127100			9761	780880	
7	9056	181120	189060	19	9811	784880	793400
	8962	179240			9762	780960	
	9283	185660			9939	795120	
	10511	210220			10158	812640	
8	13685	273700	284630	20	9990	799200	840540
	13564	271280			10389	831120	
	14347	286940			10311	824880	
	15330	306600			11337	906960	
9	14955	299100	298380				
	14870	297400					
	13895	277900					
	15956	319120					
10	4681	374480	421600				
	5209	416720					
	5265	421200					
	5925	474000					
11	5409	432720	457060				
	5514	441120					
	5821	465680					
	6109	488720					

Texas *P. parvum* clone – Trial 2 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	799	15980	15775	12	4394	351520	352080
	777	15540			4429	354320	
	814	16280			4281	342480	
	765	15300			4500	360000	
1	962	19240	19505	13	5647	451760	466520
	949	18980			5781	462480	
	986	19720			5874	469920	
	1004	20080			6024	481920	
2	1498	29960	29430	14	6893	551440	573080
	1430	28600			7430	594400	
	1468	29360			6853	548240	
	1490	29800			7478	598240	
3	1710	34200	33915	15	7623	609840	613080
	1635	32700			7510	600800	
	1734	34680			7654	612320	
	1704	34080			7867	629360	
4	2147	42940	43490	16	8812	704960	694860
	2156	43120			8674	693920	
	2211	44220			8637	690960	
	2184	43680			8620	689600	
5	3530	70600	70160	17	8644	691520	697520
	3519	70380			8826	706080	
	3485	69700			8631	690480	
	3498	69960			8775	702000	
6	4619	92380	93480	18	9094	727520	729140
	4665	93300			9008	720640	
	4717	94340			9186	734880	
	4695	93900			9169	733520	
7	5842	116840	117530	19	9999	799920	799120
	5913	118260			9864	789120	
	5895	117900			9962	796960	
	5856	117120			10131	810480	
8	7502	150040	150990	20	9781	782480	798120
	7588	151760			10001	800080	
	7598	151960			10052	804160	
	7510	150200			10072	805760	
9	7628	152560	155340				
	7780	155600					
	7818	156360					
	7842	156840					
10	2841	227280	232240				
	2877	230160					
	2969	237520					
	2925	234000					
11	3115	249200	250580				
	3111	248880					
	3163	253040					
	3140	251200					

Prymnesium calathiferum – Trial 2 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	182	3640	3675	12	3870	309600	349740
	190	3800			4289	343120	
	192	3840			4496	359680	
	171	3420			4832	386560	
1	182	3640	3685	13	4194	335520	387280
	181	3620			4467	357360	
	185	3700			4954	396320	
	189	3780			5749	459920	
2	323	6460	6635	14	5770	461600	507420
	310	6200			5956	476480	
	348	6960			6633	530640	
	346	6920			7012	560960	
3	707	14140	14500	15	4864	389120	441320
	674	13480			5432	434560	
	742	14840			5738	459040	
	777	15540			6032	482560	
4	1077	21540	21765	16	5130	410400	456600
	1070	21400			5551	444080	
	1060	21200			5814	465120	
	1146	22920			6335	506800	
5	1861	37220	37305	17	4371	349680	390800
	1877	37540			4801	384080	
	1837	36740			5088	407040	
	1886	37720			5280	422400	
6	2602	52040	53365	18	4822	385760	403440
	2637	52740			4982	398560	
	2730	54600			5031	402480	
	2704	54080			5337	426960	
7	3156	63120	64620	19	3257	260560	298020
	3163	63260			3635	290800	
	3275	65500			3952	316160	
	3330	66600			4057	324560	
8	4497	89940	97105	20	3870	309600	349740
	4693	93860			4289	343120	
	4924	98480			4496	359680	
	5307	106140			4832	386560	
9	6626	132520	147860				
	7348	146960					
	7504	150080					
	8094	161880					
10	2105	168400	191100				
	2295	183600					
	2499	199920					
	2656	212480					
11	2797	223760	266680				
	3239	259120					
	3492	279360					
	3806	304480					

North Carolina *P. parvum* clone – Trial 3 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	468	9360	9255	14	7506	600480	748920
	437	8740			7374	589920	
	467	9340			7405	592400	
	479	9580			7625	610000	
1	855	17100	17310	15	9285	742800	720100
	870	17400			9374	749920	
	864	17280			9407	752560	
	873	17460			9380	750400	
2	1225	24500	25970	16	8818	705440	835200
	1315	26300			8901	712080	
	1365	27300			8999	719920	
	1289	25780			9287	742960	
3	1947	38940	38745	17	10231	818480	800020
	1895	37900			10257	820560	
	1981	39620			10477	838160	
	1926	38520			10795	863600	
4	2537	50740	53955	18	9827	786160	808680
	2589	51780			9833	786640	
	2750	55000			10016	801280	
	2915	58300			10325	826000	
5	3634	72680	75045	19	9996	799680	972120
	3599	71980			10034	802720	
	3842	76840			10045	803600	
	3934	78680			10359	828720	
6	6215	124300	128090	20	11867	949360	1006000
	6530	130600			12098	967840	
	6405	128100			12281	982480	
	6468	129360			12360	988800	
7	9700	194000	198575	21	12178	974240	926740
	9826	196520			12166	973280	
	10049	200980			12697	1015760	
	10140	202800			13259	1060720	
8	14283	285660	286670	22	11233	898640	1139140
	14364	287280			11520	921600	
	14113	282260			11770	941600	
	14574	291480			11814	945120	
9	15047	300940	298795	23	14013	1121040	1044040
	14023	280460			13960	1116800	
	15753	315060			14423	1153840	
	14936	298720			14561	1164880	
10	17875	357500	351645	24	15310	1224800	1260560
	16855	337100			15424	1233920	
	18059	361180			16077	1286160	
	17540	350800			16217	1297360	
11	21668	433360	459350	25	13993	1119440	1190960
	22720	454400			14518	1161440	
	23198	463960			15083	1206640	
	24284	485680			15954	1276320	
12	7044	563520	579020	26	15417	1233360	1266080
	7205	576400			15698	1255840	
	7170	573600			15681	1254480	
	7532	602560			16508	1320640	
13	7044	563520	598200	27	14413	1153040	720100
	7205	576400			14601	1168080	
	7170	573600			14930	1194400	
	7532	602560			16024	1281920	

South Carolina *P. parvum* clone – Trial 3 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	554	11080	11830	14	6078	486240	578840
	547	10940			5954	476320	
	599	11980			6238	499040	
	666	13320			6292	503360	
1	1109	22180	24225	15	5669	453520	565500
	1222	24440			5781	462480	
	1204	24080			5907	472560	
	1310	26200			6244	499520	
2	1405	28100	28555	16	6877	550160	711720
	1331	26620			7105	568400	
	1534	30680			7296	583680	
	1441	28820			7664	613120	
3	1889	37780	40945	17	6906	552480	752020
	1961	39220			6915	553200	
	2103	42060			6913	553040	
	2236	44720			7541	603280	
4	2689	53780	57585	18	8493	679440	977500
	2911	58220			8763	701040	
	2895	57900			8764	701120	
	3022	60440			9566	765280	
5	3608	72160	75765	19	8900	712000	845760
	3804	76080			9073	725840	
	3763	75260			9451	756080	
	3978	79560			10177	814160	
6	4896	97920	105280	20	11722	937760	932940
	5148	102960			11999	959920	
	5313	106260			12393	991440	
	5699	113980			12761	1020880	
7	6835	136700	146285	21	10332	826560	911380
	7206	144120			10424	833920	
	7463	149260			10384	830720	
	7753	155060			11148	891840	
8	10714	214280	214590	22	11099	887920	1028520
	10600	212000			11568	925440	
	10170	203400			11679	934320	
	11434	228680			12301	984080	
9	13754	275080	273040	23	10675	854000	1142120
	13043	260860			11046	883680	
	13947	278940			11619	929520	
	13864	277280			12229	978320	
10	15543	310860	315595	24	12027	962160	1071020
	14760	295200			12347	987760	
	16585	331700			13279	1062320	
	16231	324620			13773	1101840	
11	17560	351200	23464.91	25	14519	1161520	1064860
	18174	363480			14624	1169920	
	18300	366000			12620	1009600	
	20267	405340			15343	1227440	
12	6078	486240	491240	26	12869	1029520	1007100
	5954	476320			13072	1045760	
	6238	499040			13630	1090400	
	6292	503360			13980	1118400	
13	5669	453520	472020	27	12376	990080	1143900
	5781	462480			12927	1034160	
	5907	472560			13565	1085200	
	6244	499520			14375	1150000	

Texas *P. parvum* clone – Trial 3 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	728	14560	14330	14	5314	425120	311600
	697	13940			5240	419200	
	700	14000			5475	438000	
	741	14820			5638	451040	
1	1054	21080	21195	15	4102	328160	330320
	1016	20320			4277	342160	
	1106	22120			4287	342960	
	1063	21260			4424	353920	
2	1299	25980	26210	16	5685	454800	433340
	1317	26340			5665	453200	
	1318	26360			5722	457760	
	1308	26160			5682	454560	
3	1678	33560	34490	17	6224	497920	341800
	1680	33600			6254	500320	
	1798	35960			6406	512480	
	1742	34840			6563	525040	
4	2449	48980	48995	18	6883	550640	455080
	2405	48100			6890	551200	
	2514	50280			7111	568880	
	2431	48620			6993	559440	
5	3044	60880	60685	19	6735	538800	508940
	2999	59980			6657	532560	
	3015	60300			6789	543120	
	3079	61580			6948	555840	
6	4174	83480	83785	20	6828	546240	557540
	4106	82120			6920	553600	
	4122	82440			6874	549920	
	4355	87100			6950	556000	
7	5423	108460	109075	21	6901	552080	542580
	5457	109140			6821	545680	
	5428	108560			7069	565520	
	5507	110140			6991	559280	
8	7568	151360	147730	22	7395	591600	551440
	7283	145660			7456	596480	
	7474	149480			7665	613200	
	7221	144420			7590	607200	
9	7180	143600	142585	23	6938	555040	555640
	6911	138220			7065	565200	
	7250	145000			7188	575040	
	7176	143520			7172	573760	
10	9407	188140	194415	24	8127	650160	602120
	9832	196640			8251	660080	
	9653	193060			8433	674640	
	9991	199820			8418	673440	
11	9407	188140	194415	25	7509	600720	567260
	9832	196640			7659	612720	
	9653	193060			7693	615440	
	9991	199820			7887	630960	
12	3866	309280	311600	26	7918	633440	664580
	3920	313600			7783	622640	
	3735	298800			7890	631200	
	4059	324720			7991	639280	
13	4061	324880	330320	27	7735	618800	614960
	4139	331120			7769	621520	
	4117	329360			7790	623200	
	4199	335920			7841	627280	

Prymnesium calathiferum – Trial 3 of Nutrient Replete Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	614	12280	11790	12	7527	602160	645260
	614	12280			7749	619920	
	563	11260			8515	681200	
	567	11340			8472	677760	
1	434	8680	9095	13	8613	689040	705940
	464	9280			8708	696640	
	476	9520			8910	712800	
	445	8900			9066	725280	
2	635	12700	13610	14	9617	769360	792460
	718	14360			9658	772640	
	630	12600			10027	802160	
	739	14780			10321	825680	
3	1056	21120	22090	15	8550	684000	722580
	1124	22480			9042	723360	
	1121	22420			9289	743120	
	1117	22340			9248	739840	
4	1889	37780	38700	16	7223	577840	590120
	1928	38560			7359	588720	
	2014	40280			7404	592320	
	1909	38180			7520	601600	
5	5264	105280	108090	17	4808	384640	409300
	5362	107240			5148	411840	
	5421	108420			5353	428240	
	5571	111420			5156	412480	
6	9319	186380	186085				
	9236	184720					
	9431	188620					
	9231	184620					
7	11052	221040	228145				
	11329	226580					
	11460	229200					
	11788	235760					
8	15615	312300	323030				
	16210	324200					
	16210	324200					
	16571	331420					
9	23394	467880	475205				
	23616	472320					
	24358	487160					
	23673	473460					
10	5528	442240	453720				
	5622	449760					
	5851	468080					
	5685	454800					
11	7177	574160	588560				
	7214	577120					
	7716	617280					
	7321	585680					

North Carolina *P. parvum* clone – Trial 1 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	122	976	938	12	578	11560	11270
	128	1024			554	11080	
	114	912			557	11140	
	105	840			565	11300	
1	161	1288	1116	13	575	11500	11445
	135	1080			549	10980	
	126	1008			585	11700	
	136	1088			580	11600	
2	197	1576	1630	14	580	11600	11480
	212	1696			572	11440	
	207	1656			568	11360	
	199	1592			576	11520	
3	277	2216	2278	15	604	12080	11485
	291	2328			549	10980	
	295	2360			581	11620	
	276	2208			563	11260	
4	418	3344	3244	16	568	11360	11355
	404	3232			522	10440	
	410	3280			586	11720	
	390	3120			595	11900	
5	591	4728	4420	17	510	10200	10410
	579	4632			511	10220	
	528	4224			513	10260	
	512	4096			548	10960	
6	838	6704	6782				
	828	6624					
	818	6544					
	907	7256					
7	991	7928	8510				
	1107	8856					
	1030	8240					
	1127	9016					
8	1278	10224	10020				
	1315	10520					
	1288	10304					
	1129	9032					
9	619	12380	12095				
	586	11720					
	633	12660					
	581	11620					
10	549	10980	11415				
	584	11680					
	604	12080					
	546	10920					
11	525	10500	11055				
	565	11300					
	534	10680					
	587	11740					

South Carolina *P. parvum* clone – Trial 1 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	161	1288	1252	12	546	10920	11535
	146	1168			607	12140	
	162	1296			542	10840	
	157	1256			612	12240	
1	262	2096	2106	13	613	12260	12400
	282	2256			611	12220	
	256	2048			608	12160	
	253	2024			648	12960	
2	418	3344	3314	14	615	12300	12180
	396	3168			618	12360	
	402	3216			591	11820	
	441	3528			612	12240	
3	766	6128	5990	15	626	12520	11995
	705	5640			613	12260	
	797	6376			623	12460	
	727	5816			537	10740	
4	1112	8896	8948	16	557	11140	12135
	1095	8760			609	12180	
	1120	8960			625	12500	
	1147	9176			636	12720	
5	1294	10352	10754	17	595	11900	12550
	1372	10976			615	12300	
	1332	10656			649	12980	
	1379	11032			651	13020	
6	1472	11776	11890				
	1451	11608					
	1532	12256					
	1490	11920					
7	1279	10232	11122				
	1416	11328					
	1307	10456					
	1559	12472					
8	1423	11384	11880				
	1444	11552					
	1567	12536					
	1506	12048					
9	581	11620	12095				
	623	12460					
	612	12240					
	603	12060					
10	614	12280	12340				
	614	12280					
	619	12380					
	621	12420					
11	594	11880	12605				
	675	13500					
	612	12240					
	640	12800					

Texas *P. parvum* clone – Trial 1 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	413	8260	7555	12	569	11380	11460
	382	7640			609	12180	
	363	7260			556	11120	
	353	7060			558	11160	
1	508	10160	9970				
	491	9820					
	517	10340					
	478	9560					
2	544	10880	11320				
	546	10920					
	541	10820					
	633	12660					
3	602	12040	12425				
	614	12280					
	630	12600					
	639	12780					
4	564	11280	11460				
	568	11360					
	584	11680					
	576	11520					
5	574	11480	12160				
	596	11920					
	621	12420					
	641	12820					
6	613	12260	12145				
	607	12140					
	633	12660					
	576	11520					
7	579	11580	12335				
	650	13000					
	577	11540					
	661	13220					
8	673	13460	13480				
	675	13500					
	698	13960					
	650	13000					
9	648	12960	13135				
	673	13460					
	631	12620					
	675	13500					
10	664	13280	12830				
	652	13040					
	632	12640					
	618	12360					
11	565	11300	12030				
	625	12500					
	624	12480					
	592	11840					

Prymnesium calathiferum – Trial 1 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	299	5980	5505				
	288	5760					
	262	5240					
	252	5040					
1	309	6180	6105				
	307	6140					
	310	6200					
	295	5900					
2	586	11720	11260				
	550	11000					
	583	11660					
	533	10660					
3	983	19660	19590				
	1011	20220					
	974	19480					
	950	19000					
4	934	18680	19375				
	934	18680					
	997	19940					
	1010	20200					
5	1149	22980	22945				
	1118	22360					
	1139	22780					
	1183	23660					
6	1287	25740	26515				
	1360	27200					
	1358	27160					
	1298	25960					
7	1432	28640	28335				
	1386	27720					
	1409	28180					
	1440	28800					
8	1299	25980	25735				
	1303	26060					
	1281	25620					
	1264	25280					

North Carolina *P. parvum* clone – Trial 2 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	144	2880	2535	12	732	14640	14600
	123	2460			719	14380	
	123	2460			744	14880	
	117	2340			725	14500	
1	164	3280	2805	13	1004	20080	19570
	147	2940			922	18440	
	125	2500			966	19320	
	125	2500			1022	20440	
2	219	4380	4175	14	723	14460	13565
	215	4300			699	13980	
	212	4240			615	12300	
	189	3780			676	13520	
3	266	5320	4775	15	771	15420	15010
	237	4740			700	14000	
	231	4620			790	15800	
	221	4420			741	14820	
4	332	6640	6240	16	680	13600	13865
	282	5640			728	14560	
	318	6360			676	13520	
	316	6320			689	13780	
5	323	6460	6570	17	698	13960	12955
	323	6460			636	12720	
	327	6540			604	12080	
	341	6820			653	13060	
6	407	8140	7855	18	713	14260	13635
	391	7820			682	13640	
	366	7320			658	13160	
	407	8140			674	13480	
7	413	8260	8335				
	449	8980					
	394	7880					
	411	8220					
8	559	11180	10985				
	584	11680					
	518	10360					
	536	10720					
9	681	13620	12775				
	625	12500					
	600	12000					
	649	12980					
10	653	13060	13435				
	672	13440					
	679	13580					
	683	13660					
11	731	14620	14285				
	669	13380					
	723	14460					
	734	14680					

South Carolina *P. parvum* clone – Trial 2 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	129	1032	924	12	877	17540	17635
	118	944			923	18460	
	101	808			892	17840	
	114	912			835	16700	
1	276	2208	2078	13	862	17240	16955
	263	2104			814	16280	
	253	2024			851	17020	
	247	1976			864	17280	
2	344	2752	2830	14	952	19040	18975
	327	2616			923	18460	
	370	2960			924	18480	
	374	2992			996	19920	
3	494	3952	4014	15	843	16860	16670
	492	3936			829	16580	
	486	3888			773	15460	
	535	4280			889	17780	
4	366	7320	7125	16	776	15520	16185
	338	6760			786	15720	
	350	7000			814	16280	
	371	7420			861	17220	
5	464	9280	10290	17	795	15900	15925
	521	10420			754	15080	
	478	9560			823	16460	
	595	11900			813	16260	
6	785	15700	15865	18	723	14460	15865
	794	15880			799	15980	
	844	16880			805	16100	
	750	15000			846	16920	
7	769	15380	16060	19	823	16460	16830
	911	18220			860	17200	
	674	13480			794	15880	
	858	17160			889	17780	
8	921	18420	18020	20	833	16660	17045
	878	17560			820	16400	
	899	17980			851	17020	
	906	18120			905	18100	
9	848	16960	17110	21	793	15860	16785
	869	17380			819	16380	
	772	15440			846	16920	
	933	18660			899	17980	
10	852	17040	16060				
	789	15780					
	845	16900					
	726	14520					
11	800	16000	16445				
	799	15980					
	854	17080					
	836	16720					

Texas *P. parvum* clone – Trial 2 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	711	14220	13895	12	939	18780	19110
	738	14760			959	19180	
	659	13180			958	19160	
	671	13420			966	19320	
1	1026	20520	20660				
	1030	20600					
	1060	21200					
2	1016	20320	21770				
	1086	21720					
	1105	22100					
	1098	21960					
3	1065	21300	22190				
	1163	23260					
	1121	22420					
4	1061	21220	21280				
	1093	21860					
	1064	21280					
	1046	20920					
5	1080	21600	21165				
	1066	21320					
	1038	20760					
	1050	21000					
6	1071	21420	20500				
	1074	21480					
	1065	21300					
	1045	20900					
7	1007	20140	22000				
	983	19660					
	1051	21020					
	1105	22100					
8	1103	22060	22250				
	1141	22820					
	1062	21240					
	1167	23340					
9	1115	22300	21500				
	1106	22120					
	1083	21660					
	1064	21280					
10	1118	22360	21500				
	1035	20700					
	1083	21660					
	1064	21280					
11	1118	22360	19875				
	1035	20700					
	969	19380					
	994	19880					
	993	19860					
	1019	20380					

Prymnesium calathiferum – Trial 2 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	206	4120	3700				
	184	3680					
	185	3700					
	165	3300					
1	141	2820	2920				
	143	2860					
	147	2940					
	153	3060					
2	308	6160	6230				
	312	6240					
	306	6120					
	320	6400					
3	366	7320	6820				
	328	6560					
	317	6340					
	353	7060					
4	423	8460	8205				
	378	7560					
	425	8500					
	415	8300					
5	333	6660	6880				
	355	7100					
	336	6720					
	352	7040					
6	348	6960	6930				
	351	7020					
	340	6800					
	347	6940					
7	343	6860	6685				
	330	6600					
	345	6900					
	319	6380					
8	262	5240	5425				
	272	5440					
	290	5800					
	261	5220					

North Carolina *P. parvum* clone – Trial 3 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	247	4940	4660				
	249	4980					
	206	4120					
	230	4600					
1	256	5120	5285				
	269	5380					
	285	5700					
	247	4940					
2	323	6460	6450				
	315	6300					
	312	6240					
	340	6800					
3	382	7640	7750				
	419	8380					
	369	7380					
	380	7600					
4	551	11020	10755				
	516	10320					
	545	10900					
	539	10780					
5	645	12900	12145				
	623	12460					
	570	11400					
	591	11820					
6	642	12840	12140				
	596	11920					
	597	11940					
	593	11860					
7	613	12260	11635				
	568	11360					
	581	11620					
	565	11300					
8	642	12840	12585				
	619	12380					
	651	13020					
	605	12100					
9	676	13520	13040				
	674	13480					
	642	12840					
	616	12320					
10	700	14000	13205				
	673	13460					
	645	12900					
	623	12460					

South Carolina *P. parvum* clone – Trial 3 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	160	3200	3140				
	141	2820					
	157	3140					
	170	3400					
1	196	3920	3700				
	179	3580					
	166	3320					
	199	3980					
2	242	4840	4905				
	227	4540					
	236	4720					
	276	5520					
3	341	6820	7100				
	370	7400					
	356	7120					
	353	7060					
4	434	8680	9320				
	451	9020					
	471	9420					
	508	10160					
5	606	12120	12140				
	571	11420					
	615	12300					
	636	12720					
6	554	11080	11760				
	579	11580					
	575	11500					
	644	12880					
7	641	12820	12460				
	620	12400					
	654	13080					
	577	11540					
8	567	11340	12255				
	608	12160					
	642	12840					
	634	12680					
9	657	13140	13140				
	633	12660					
	661	13220					
	677	13540					
10	660	13200	13260				
	670	13400					
	677	13540					
	645	12900					

Texas *P. parvum* clone – Trial 3 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	208	4160	3905				
	190	3800					
	201	4020					
	182	3640					
1	228	4560	4820				
	234	4680					
	246	4920					
	256	5120					
2	364	7280	7225				
	373	7460					
	366	7320					
	342	6840					
3	382	7640	7305				
	341	6820					
	363	7260					
	375	7500					
4	414	8280	8005				
	404	8080					
	388	7760					
	395	7900					
5	438	8760	8370				
	423	8460					
	412	8240					
	401	8020					
6	445	8900	8775				
	426	8520					
	425	8500					
	459	9180					
7	455	9100	9015				
	443	8860					
	450	9000					
	455	9100					
8	411	8220	8475				
	439	8780					
	420	8400					
	425	8500					

Prymnesium calathiferum – Trial 3 of Nitrogen-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	193	3860	3675				
	179	3580					
	194	3880					
	169	3380					
1	174	3480	3545				
	181	3620					
	188	3760					
	166	3320					
2	397	7940	8130				
	410	8200					
	397	7940					
	422	8440					
3	443	8860	8795				
	475	9500					
	420	8400					
	421	8420					
4	424	8480	8840				
	455	9100					
	440	8800					
	449	8980					
5	444	8880	9290				
	451	9020					
	459	9180					
	504	10080					
6	433	8660	9095				
	446	8920					
	460	9200					
	480	9600					
7	386	7720	7645				
	361	7220					
	380	7600					
	402	8040					
8	342	6840	6780				
	321	6420					
	348	6960					
	345	6900					

North Carolina *P. parvum* clone – Trial 1 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	252	5040	5760	12	10774	215480	219400
	340	6800			10784	215680	
	249	4980			11033	220660	
	311	6220			11289	225780	
1	249	4980	5970	13	11335	226700	232115
	326	6520			11607	232140	
	226	4520			11616	232320	
	393	7860			11865	237300	
2	692	13840	13630	14	11082	221640	223870
	665	13300			10930	218600	
	743	14860			11322	226440	
	626	12520			11440	228800	
3	904	18080	18180	15	11074	221480	223870
	883	17660			11088	221760	
	905	18100			11670	233400	
	944	18880			11394	227880	
4	1295	25900	26995	16	11007	220140	226130
	1308	26160			11164	223280	
	1359	27180			11316	226320	
	1437	28740			11375	227500	
5	1911	38220	39765	17	10676	213520	224310
	1999	39980			10902	218040	
	1957	39140			11073	221460	
	2086	41720			10982	219640	
6	2725	54500	55890	18	9292	185840	218165
	2774	55480			9309	186180	
	2863	57260			9360	187200	
	2816	56320			9495	189900	
7	3928	78560	79865				
	4032	80640					
	3951	79020					
	4062	81240					
8	5327	106540	109380				
	5384	107680					
	5486	109720					
	5679	113580					
9	7073	141460	142080				
	7000	140000					
	7104	142080					
	7239	144780					
10	9097	181940	183165				
	9155	183100					
	8996	179920					
	9385	187700					
11	9960	199200	203180				
	10082	201640					
	10060	201200					
	10534	210680					

South Carolina *P. parvum* clone – Trial 1 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	641	5128	5212	12	12604	252080	257580
	622	4976			12890	257800	
	649	5192			12894	257880	
	694	5552			13128	262560	
1	994	7952	8246	13	13077	261540	264885
	1023	8184			13211	264220	
	1125	9000			13316	266320	
	981	7848			13373	267460	
2	594	11880	11570	14	13272	265440	269180
	566	11320			13226	264520	
	590	11800			13424	268480	
	564	11280			13914	278280	
3	992	19840	19115	15	13115	262300	266030
	917	18340			13183	263660	
	948	18960			13266	265320	
	966	19320			13642	272840	
4	1316	26320	26720	16	13000	260000	261270
	1362	27240			13050	261000	
	1335	26700			13145	262900	
	1331	26620			13059	261180	
5	1975	39500	38645	17	13140	262800	267425
	1917	38340			13282	265640	
	1890	37800			13427	268540	
	1947	38940			13636	272720	
6	2934	58680	59895	18	13099	261980	266350
	3018	60360			13239	264780	
	3058	61160			13336	266720	
	2969	59380			13596	271920	
7	4657	93140	92555	19	12732	254640	258035
	4624	92480			12850	257000	
	4566	91320			12823	256460	
	4664	93280			13202	264040	
8	5614	112280	120915	20	12810	256200	257925
	6443	128860			12821	256420	
	5314	106280			12983	259660	
	6812	136240			12971	259420	
9	8697	173940	185395				
	9304	186080					
	9393	187860					
	9685	193700					
10	10830	216600	224040				
	11198	223960					
	11189	223780					
	11591	231820					
11	12466	249320	252440				
	12489	249780					
	12569	251380					
	12964	259280					

Texas *P. parvum* clone – Trial 1 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	173	3460	3755				
	189	3780					
	188	3760					
	201	4020					
1	272	5440	5725				
	285	5700					
	280	5600					
	308	6160					
2	363	7260	6985				
	341	6820					
	336	6720					
	357	7140					
3	449	8980	8765				
	444	8880					
	425	8500					
	435	8700					
4	453	9060	9245				
	490	9800					
	436	8720					
	470	9400					
5	579	11580	11335				
	555	11100					
	539	10780					
	594	11880					
6	640	12800	12405				
	621	12420					
	606	12120					
	614	12280					
7	867	17340	17755				
	921	18420					
	917	18340					
	846	16920					
8	1070	21400	22380				
	1146	22920					
	1010	20200					
	1250	25000					
9	1190	23800	24395				
	1214	24280					
	1263	25260					
	1212	24240					

Prymnesium calathiferum – Trial 1 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	596	4768	4780	12	6715	134300	127610
	588	4704			6039	120780	
	607	4856			6757	135140	
	599	4792			6011	120220	
1	492	3936	3860	13	6850	137000	140610
	519	4152			7055	141100	
	477	3816			7010	140200	
	442	3536			7207	144140	
2	316	6320	6180	14	6666	133320	135235
	301	6020			6695	133900	
	319	6380			6792	135840	
	300	6000			6894	137880	
3	556	11120	9870	15	6400	128000	130045
	473	9460			6453	129060	
	445	8900			6622	132440	
	500	10000			6534	130680	
4	645	12900	13885	16	5490	109800	111075
	732	14640			5432	108640	
	673	13460			5619	112380	
	727	14540			5674	113480	
5	755	15100	15535	17	4756	95120	95225
	755	15100			4734	94680	
	835	16700			4804	96080	
	762	15240			4751	95020	
6	1873	37460	40390				
	2266	45320					
	1698	33960					
	2241	44820					
7	3698	73960	74420				
	3586	71720					
	3733	74660					
	3867	77340					
8	4857	97140	101300				
	5124	102480					
	5155	103100					
	5124	102480					
9	5677	113540	116860				
	5582	111640					
	6430	128600					
	5683	113660					
10	6912	138240	142695				
	7182	143640					
	7195	143900					
	7250	145000					
11	6784	135680	136930				
	7024	140480					
	6401	128020					
	7177	143540					

North Carolina *P. parvum* clone – Trial 2 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	440	8800	8185	12	10040	200800	204145
	403	8060			10138	202760	
	391	7820			10093	201860	
	403	8060			10558	211160	
1	536	10720	9540	13	10098	201960	215930
	464	9280			10751	215020	
	449	8980			10947	218940	
	459	9180			11390	227800	
2	563	11260	12200	14	8992	179840	184605
	599	11980			8911	178220	
	615	12300			9433	188660	
	663	13260			9585	191700	
3	1035	20700	21145	15	10037	200740	207890
	1031	20620			10270	205400	
	1063	21260			10604	212080	
	1100	22000			10667	213340	
4	1451	29020	33155				
	1526	30520					
	1756	35120					
	1898	37960					
5	1791	35820	36890				
	1795	35900					
	1833	36660					
	1959	39180					
6	2422	48440	49855				
	2428	48560					
	2514	50280					
	2607	52140					
7	4896	97920	98235				
	4908	98160					
	4923	98460					
	4920	98400					
8	6247	124940	125835				
	6235	124700					
	6282	125640					
	6403	128060					
9	8327	166540	169670				
	8559	171180					
	8438	168760					
	8610	172200					
10	9546	190920	202470				
	9820	196400					
	10440	208800					
	10688	213760					
11	10010	200200	204725				
	10141	202820					
	10104	202080					
	10690	213800					

South Carolina *P. parvum* clone – Trial 2 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	500	10000	10470	12	13646	272920	275980
	533	10660			13616	272320	
	518	10360			13688	273760	
	543	10860			14246	284920	
1	771	15420	15955	13	13640	272800	280170
	779	15580			13937	278740	
	810	16200			13941	278820	
	831	16620			14516	290320	
2	1049	20980	21835	14	13870	277400	283125
	1031	20620			14082	281640	
	1077	21540			14113	282260	
	1210	24200			14560	291200	
3	1654	33080	33420	15	14556	291120	296730
	1652	33040			14493	289860	
	1643	32860			14916	298320	
	1735	34700			15381	307620	
4	2117	42340	45245				
	2166	43320					
	2297	45940					
	2469	49380					
5	2956	59120	61510				
	3058	61160					
	3100	62000					
	3188	63760					
6	3894	77880	80605				
	3941	78820					
	4088	81760					
	4198	83960					
7	6362	127240	129410				
	6445	128900					
	6495	129900					
	6580	131600					
8	8227	164540	166200				
	8250	165000					
	8272	165440					
	8491	169820					
9	10271	205420	208125				
	10324	206480					
	10407	208140					
	10623	212460					
10	12013	240260	243970				
	11906	238120					
	12239	244780					
	12636	252720					
11	13115	262300	266885				
	13184	263680					
	13453	269060					
	13625	272500					

Texas *P. parvum* clone – Trial 2 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	783	15660	15550	12	3330	66600	67700
	732	14640			3440	68800	
	801	16020			3370	67400	
	794	15880			3463	69260	
1	1325	26500	26480	13	3298	65960	66750
	1337	26740			3377	67540	
	1300	26000			3338	66760	
	1334	26680			3370	67400	
2	1708	34160	34435				
	1718	34360					
	1718	34360					
	1743	34860					
3	2063	41260	41945				
	2055	41100					
	2168	43360					
	2103	42060					
4	2250	45000	44515				
	2213	44260					
	2255	45100					
	2185	43700					
5	2608	52160	51025				
	2521	50420					
	2606	52120					
	2470	49400					
6	3103	62060	62060				
7	3432	68640	68695				
	3467	69340					
	3428	68560					
	3412	68240					
8	3168	63360	64390				
	3271	65420					
	3369	67380					
	3266	65320					
9	3175	63500	65550				
	3235	64700					
	3320	66400					
	3380	67600					
10	3375	67500	68060				
	3427	68540					
	3407	68140					
	3419	68380					
11	3235	64700	67495				
	3362	67240					
	3468	69360					
	3434	68680					

Prymnesium calathiferum – Trial 2 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	575	11500	10415				
	504	10080					
	508	10160					
	496	9920					
1	366	7320	6685				
	328	6560					
	339	6780					
	304	6080					
2	508	10160	10770				
	599	11980					
	497	9940					
3	550	11000	12635				
	631	12620					
	632	12640					
	633	12660					
4	631	12620	28520				
	1300	26000					
	1465	29300					
	1494	29880					
5	1445	28900	44665				
	2310	46200					
	2229	44580					
	2187	43740					
6	2207	44140	61675				
	3126	62520					
	3046	60920					
	3122	62440					
7	3041	60820	79385				
	3920	78400					
	4011	80220					
	3967	79340					
8	3979	79580	108845				
	5414	108280					
	5477	109540					
	5341	106820					
9	5537	110740	117215				
	5948	118960					
	5754	115080					
	5793	115860					
10	5948	118960	103300				
	5093	101860					
	5124	102480					
	5292	105840					
	5151	103020					

North Carolina *P. parvum* clone – Trial 3 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	77	1540	1360	12	3472	69440	70645
	64	1280			3616	72320	
	71	1420			3454	69080	
	60	1200			3587	71740	
1	170	1360	1462	13	4473	89460	91840
	182	1456			4678	93560	
	197	1576			4676	93520	
	182	1456			4541	90820	
2	242	1936	1928	14	6558	131160	130090
	240	1920			6311	126220	
	249	1992			6490	129800	
	233	1864			6659	133180	
3	299	2392	2468	15	8918	178360	177005
	308	2464			8663	173260	
	311	2488			8847	176940	
	316	2528			8973	179460	
4	469	3752	3758	16	8840	176800	180945
	472	3776			9199	183980	
	450	3600			8667	173340	
	488	3904			9483	189660	
5	630	5040	5040	17	9881	197620	202370
	612	4896			10145	202900	
	621	4968			10046	200920	
	657	5256			10402	208040	
6	848	6784	6664	18	9948	198960	205070
	794	6352			10068	201360	
	825	6600			10456	209120	
	865	6920			10542	210840	
7	1076	8608	8756	19	10496	209920	212955
	1063	8504			10296	205920	
	1111	8888			10656	213120	
	1128	9024			11143	222860	
8	1601	12808	12786	20	10201	204020	210570
	1572	12576			10343	206860	
	1590	12720			10530	210600	
	1630	13040			11040	220800	
9	2027	16216	17296				
	2228	17824					
	2108	16864					
	2285	18280					
10	3581	28648	28106				
	3394	27152					
	3630	29040					
	3448	27584					
11	2270	45400	44305				
	2225	44500					
	2177	43540					
	2189	43780					

South Carolina *P. parvum* clone – Trial 3 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	110	2200	2135	12	10452	418080	416590
	101	2020			10240	409600	
	110	2200			10540	421600	
	106	2120			10427	417080	
1	350	2800	2756	13	11209	448360	450230
	323	2584			11059	442360	
	346	2768			11207	448280	
	359	2872			11548	461920	
2	428	3424	3492	14	11823	472920	483280
	443	3544			11866	474640	
	447	3576			12051	482040	
	428	3424			12588	503520	
3	719	5752	5638	15	12368	494720	494500
	707	5656			12198	487920	
	701	5608			12357	494280	
	692	5536			12527	501080	
4	1001	8008	7764	16	11238	449520	464660
	917	7336			11790	471600	
	990	7920			11256	450240	
	974	7792			12182	487280	
5	1470	11760	11026	17	12308	492320	492940
	1317	10536			12297	491880	
	1360	10880			12147	485880	
	1366	10928			12542	501680	
6	2305	18440	18494	18	12504	500160	503220
	2332	18656			12493	499720	
	2334	18672			12341	493640	
	2276	18208			12984	519360	
7	3494	27952	28584	19	12605	504200	508620
	3617	28936			12460	498400	
	3605	28840			12835	513400	
	3576	28608			12962	518480	
8	5769	46152	46442	20	11681	467240	474020
	5530	44240			11552	462080	
	6087	48696			11820	472800	
	5835	46680			12349	493960	
9	7981	63848	65214				
	8336	66688					
	7783	62264					
	8507	68056					
10	13199	105592	104592				
	12690	101520					
	13268	106144					
	13139	105112					
11	7480	299200	292410				
	7207	288280					
	7375	295000					
	7179	287160					

Texas *P. parvum* clone – Trial 3 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	340	6800	7525				
	410	8200					
	359	7180					
	396	7920					
1	523	10460	10095				
	500	10000					
	521	10420					
	475	9500					
2	832	16640	16460				
	795	15900					
	866	17320					
	799	15980					
3	824	16480	16935				
	845	16900					
	864	17280					
	854	17080					
4	884	17680	17530				
	905	18100					
	908	18160					
	809	16180					
5	1085	21700	22045				
	1083	21660					
	1156	23120					
	1085	21700					
6	1069	21380	21185				
	1074	21480					
	1029	20580					
	1065	21300					
7	1140	22800	23295				
	1201	24020					
	1122	22440					
	1196	23920					
8	1190	23800	24425				
	1139	22780					
	1296	25920					
	1260	25200					
9	1114	22280	23705				
	1061	21220					
	1294	25880					
	1272	25440					
10	1062	21240	22220				
	1361	27220					
	1015	20300					
	1006	20120					

Prymnesium calathiferum – Trial 3 of Phosphorus-Deficient Treatment

Day	Cells/0.5 mL	Cells/mL	Mean cells/mL	Day	Cells/0.5 mL	Cells/mL	Mean cells/mL
0	154	3080	2910	12	5025	100500	102910
	132	2640			5008	100160	
	150	3000			5268	105360	
	146	2920			5281	105620	
1	107	2140	2405	13	4803	96060	97200
	139	2780			4751	95020	
	125	2500			5053	101060	
	110	2200			4833	96660	
2	182	3640	3905	14	4696	93920	96145
	195	3900			4767	95340	
	213	4260			4914	98280	
	191	3820			4852	97040	
3	267	5340	5465	15	4820	96400	96025
	292	5840			4752	95040	
	262	5240			4761	95220	
	272	5440			4872	97440	
4	314	6280	6470	16	4137	82740	84045
	309	6180			4175	83500	
	338	6760			4177	83540	
	333	6660			4320	86400	
5	368	7360	7060	17	3764	75280	74540
	341	6820			3695	73900	
	386	7720			3719	74380	
	317	6340			3730	74600	
6	990	19800	20050				
	1002	20040					
	998	19960					
	1020	20400					
7	1851	37020	38850				
	1896	37920					
	1989	39780					
	2034	40680					
8	2861	57220	59040				
	3017	60340					
	2966	59320					
	2964	59280					
9	2742	54840	53105				
	2527	50540					
	2714	54280					
	2638	52760					
10	3980	79600	81860				
	4064	81280					
	4287	85740					
	4041	80820					
11	5870	117400	122725				
	6111	122220					
	6263	125260					
	6301	126020					

Appendix B: Erythrocyte Lysis Assay Data

Replete: Trial Averages

Hemolytic activity of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)
	Lag	70.877 \pm 33.219	3543.172 \pm 973.283	48.645 \pm 36.153	3639.586 \pm 1371.491	20.116 \pm 11.471	1958.295 \pm 1701.731	1.114 \pm 0.739	187.553 \pm 133.058
	Log	77.973 \pm 10.283	259.223 \pm 111.171	33.886 \pm 11.621	143.808 \pm 79.522	30.255 \pm 11.339	203.530 \pm 144.659	3.766 \pm 2.385	8.662 \pm 5.333
	Stationary	51.658 \pm 4.143	77.711 \pm 10.591	27.845 \pm 17.538	45.837 \pm 30.097	18.363 \pm 11.003	40.742 \pm 30.975	3.907 \pm 4.034	28.323 \pm 31.049

Hemolytic activity of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)
	Lag	89.000 \pm 8.742	1440.031 \pm 765.165	97.592 \pm 8.031	2713.572 \pm 2415.977	91.475 \pm 20.171	2395.659 \pm 1824.391	1.054 \pm 0.514	64.847 \pm 61.544
	Log	98.496 \pm 2.214	261.677 \pm 124.744	97.841 \pm 4.381	326.539 \pm 190.61	101.594 \pm 5.484	456.98 \pm 263.425	13.522 \pm 3.972	45.182 \pm 10.9
	Stationary	91.596 \pm 3.684	113.368 \pm 36.977	83.426 \pm 29.258	126.509 \pm 80.509	97.253 \pm 9.442	174.982 \pm 59.814	38.472 \pm 9.628	97.390 \pm 44.456

Replete: Trial 1

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on February 7, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis (Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis (Cell)
2	Lag	97.46 ±1.849	2862.794 ±54.226	90.041 ±2.018	3907.191 ±87.592	8.73 ±0.32	155.230 ±5.685	1.966 ±0.716	138.626 ±50.452
15	Log	69.763 ±3.763	147.636 ±7.963	22.541 ±2.033	52.408 ±4.727	17.273 ±0.741	36.534 ±1.567	3.416 ±0.387 ^a	21.904 ±2.481 ^a
22	Stationary	53.797 ±4.834	82.739 ±7.436	42.817 ±4.815	65.116 ±7.324	19.288 ±2.517	28.456 ±3.714	8.397 ±1.026	19.979 ±2.441

*-n=4; a=Log phase on Day 7

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on February 7, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
2	Lag	98.179 ±5.615*	576.677 ±32.989*	106.395 ±6.712*	923.373 ±58.259*	108.266 ±1.16*	384.869 ±4.124*	1.645 ±0.099*	23.197 ±1.393*
15	Log	96.610 ±7.160	204.452 ±15.152	100.65 ±5.698	234.010 ±13.248	107.607 ±3.754	227.594 ±7.940	2.545 ±0.239* ^a	16.322 ±1.536* ^a
22	Stationary	95.814 ±5.998	147.360 ±9.226	100.605 ±4.045	152.998 ±6.153	107.926 ±5.542	159.225 ±8.178	48.429 ±4.782	115.221 ±11.379

*-n=4; a=Log phase on Day 7

Replete: Trial 2

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on March 14, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis (Cell)
2	Lag	53.530 ±23.923	2219.527 ±992.023	23.277 ±2.780	1360.109 ±162.430	22.983 ±2.152	1249.307 ±117.001	1.379 ±0.353	334.794 ±85.737
9	Log	35.520 ±2.988	131.877 ±11.104	33.352 ±2.256	181.877 ±12.302	21.794 ±1.040	224.448 ±10.712	0.707 ±0.304	7.649 ±3.285
20	Stationary	53.952 ±2.907	67.304 ±3.622	32.167 ±1.555	61.240 ±2.959	26.153 ±1.348	52.435 ±2.702	0.530 ±0.324 ^a	2.846 ±1.738 ^a

a - Stationary phase on Day 19

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on March 14, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis (Cell)
2	Lag	88.699 ±7.789	3677.734 ±322.965	90.666 ±7.063	5297.750 ±412.731	92.556 ±7.837	5031.126 ±426.052	0.576 ±0.548	69.433 ±66.112
9	Log	93.061 ±5.638	345.746 ±20.949	100.079 ±4.255	545.751 ±23.204	89.473 ±2.035	921.451 ±20.965	3.595 ±0.458	38.895 ±4.959
20	Stationary	96.067 ±3.702	119.841 ±4.618	100.030 ±5.432	190.437 ±10.340	101.628 ±4.455	203.760 ±8.931	9.190 ±0.961 ^a	49.332 ±5.159 ^a

a - Stationary phase on Day 19

Replete: Trial 3

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on April 25, 2005. (n=8)

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	33.637 ±3.801 ^a	3108.704 ±351.310 ^a	32.618 ±2.513 ^a	2154.015 ±165.964 ^a	31.670 ±2.120	3536.292 ±236.691	0.633 ±0.510	85.886 ±69.160
11	Log	74.649 ±7.610 [*]	260.059 ±26.506 [*]	45.766 ±9.681 [*]	197.140 ±41.693 [*]	35.262 ±6.140 [*]	290.246 ±50.527 [*]	6.318 ±1.092 ^{*b}	44.297 ±7.655 ^{*b}
27	Stationary	46.882 ±3.246	65.542 ±4.330	8.550 ±1.229 ^d	11.156 ±1.604 ^d	6.927 ±0.513	17.795 ±1.317	0.588 ±0.007 ^c	2.298 ±2.744 ^c

* - n=4; a - Lag phase on Day 1; b- Log phase on Day 7, c-Stationary phase on Day 17; d-Stationary phase on Day 29

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nutrient replete conditions. All clones were inoculated on April 25, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)
0	Lag	88.049 ±7.647 ^a	2034.317 ±176.708 ^a	95.714 ±10.011 ^a	1755.746 ±183.672 ^a	69.085 ±9.496	2857.046 ±392.690	0.712 ±0.337	35.805 ±16.947
11	Log	100.933 ±25.938	175.813 ±45.173	92.793 ±22.836 [*]	199.855 ±49.175 [*]	96.869 ±21.334	398.672 ±87.788	9.624 ±4.649 ^b	35.514 ±17.160 ^b
27	Stationary	89.013 ±7.504	118.748 ±10.011	49.664 ±7.372 ^d	36.094 ±4.968 ^d	93.848 ±9.674	241.097 ±24.857	37.777 ±0.055 ^c	147.685 ±6.85 ^c

* - n=4; a - Lag phase on Day 1; b- Log phase on Day 7, c-Stationary phase on Day 17; d-Stationary phase on Day 29

N-Deficient: Trial Averages

Hemolytic activity of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
	Lag	33.662 \pm 10.573	17487 \pm 1601.087	28.369 \pm 9.451	20795.93 \pm 14476.8	5.259 \pm 1.680	1456.904 \pm 1309.345	2.226 \pm 1.504	779.944 \pm 352.989
	Log	63.162 \pm 36.237	12291.53 \pm 8079.995	57.482 \pm 28.469	7850.901 \pm 837.061	8.108 \pm 1.062	1133.417 \pm 472.738	1.151 \pm 0.151	214.897 \pm 59.021
	Stationary	76.324 \pm 22.766	9599.747 \pm 3307.954	84.901 \pm 9.164	9815.413 \pm 1873.03	19.587 \pm 7.686	2423.249 \pm 690.949	1.504 \pm 1.355	190.649 \pm 12.719

Hemolytic activity of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
	Lag	92.521 \pm 7.958	13228.492 \pm 5806.117	90.382 \pm 5.191	14670.09 \pm 11760.71	82.794 \pm 6.939	7164.427 \pm 5545.626	1.785 \pm 1.322	177.615 \pm 139.822
	Log	91.208 \pm 1.756	5224.072 \pm 1682.887	93.794 \pm 3.869	3469.88 \pm 561.342	92.024 \pm 4.961	3484.36 \pm 1315.627	0.839 \pm 0.384	35.194 \pm 21.286
	Stationary	87.965 \pm 7.376	3107.485 \pm 283.225	91.067 \pm 2.425	3235.977 \pm 111.060	92.008 \pm 6.426	3123.968 \pm 1442.671	0.749 \pm 0.431	26.451 \pm 6.237

N-Deficient: Trial 1

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). The NC and SC clones were inoculated on March 14, 2005, TX was inoculated on May 16, 2005, and Pcal was inoculated on May 22, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
3	Lag	26.878 ±3.801	18,882.043 ±2669.789	25.568 ±6.709	6830.685 ±1791.935	4.543 ±0.481 ^a	962.087 ±101.913 ^a	3.899 ±1.271 ^a	1133.365 ±369.273 ^a
7	Log	21.347 ±1.202	4013.765 ±225.960	51.381 ±6.902	7398.104 ±646.164	7.597 ±0.764 ^c	1060.522 ±106.638 ^c	1.037 ±0.509 ^b	147.373 ±72.330 ^b
14	Stationary	87.611 ±5.588	12,211.162 ±778.779	91.055 ±6.382	11,961.745 ±838.374	13.664 ±2.509 ^c	1817.574 ±333.708 ^c	3.068 ±1.221 ^d	190.731 ±75.919 ^d

a – Lag Phase on Day 0; b- Log phase on Day 2; c-Log phase on Day 4; d-Stationary phase on Day 8; e-Stationary phase on Day 11

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). The NC and SC clones were inoculated on March 14, 2005, TX was inoculated on May 16, 2005, and Pcal was inoculated on May 22, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
3	Lag	97.582 ±9.243	17,577.219 ±1664.694	94.370 ±9.621	6464.586 ±658.947	75.179 ±13.090 ^a	2793.115 ±486.370 ^a	2.806 ±0.643 ^a	302.081 ±69.203 ^a
7	Log	90.645 ±3.109	4869.523 ±167.000	95.057 ±2.924	3907.284 ±120.173	97.700 ±13.290 ^c	4011.579 ±545.716 ^c	0.405 ±0.223 ^b	13.079 ±7.212 ^b
14	Stationary	93.405 ±6.075	3425.974 ±222.801	90.236 ±7.957	3152.211 ±277.942	86.481 ±10.128 ^c	2875.958 ±336.774 ^c	1.246 ±0.681 ^d	19.363 ±10.577 ^d

a – Lag Phase on Day 0; b- Log phase on Day 2; c-Log phase on Day 4; d-Stationary phase on Day 8; e-Stationary phase on Day 11

N-Deficient: Trial 2

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). NC was inoculated on April 11, 2005, SC was inoculated on April 1, 2005, TX was inoculated on May 16, 2005, and Pcal was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	28.264 ±3.505	17,840.126 ±2212.11	20.636 ±1.132	35,735.106 ±1960.151	4.056 ±0.476	467.076 ±54.835	0.988 ±0.553	427.387 ±238.959
3	Log	82.760 ±2.172 ^b	20,158.048 ±529.033 ^b	88.506 ±4.589 ^c	8816.828 ±457.214 ^c	9.329 ±1.005 ^a	701.361 ±75.535 ^a	1.094 ±0.318	256.648 ±74.623
8	Stationary	91.241 ±12.035 ^e	10,708.154 ±1412.216 ^e	89.280 ±6.774 ^f	8511.660 ±645.708 ^f	28.272 ±6.541 ^d	2276.333 ±526.546 ^d	0.689 ±0.601	203.326 ±177.137

a-Log phase on Day 4; b- Log phase on Day 5; c-Log phase on Day 7; d- Stationary phase on Day 11; e- Stationary phase on Day 18; f- Stationary phase on Day 21

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). NC was inoculated on April 11, 2005, SC was inoculated on April 1, 2005, TX was inoculated on May 16, 2005, and Pcal was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	83.348 ±6.186	15,473.113 ±1148.420	84.512 ±4.329	28,143.889 ±1441.445	84.441 ±11.868	1705.773 ±239.761	0.292 ±0.236	26.280 ±21.278
3	Log	89.803 ±5.137 ^b	7055.985 ±403.562 ^b	96.875 ±5.589 ^c	2836.916 ±163.760 ^c	89.855 ±14.102 ^a	1986.893 ±311.843 ^a	1.137 ±0.388	55.541 ±18.986
8	Stationary	90.920 ±13.405 ^e	2883.898 ±425.147 ^e	93.799 ±6.175 ^f	3193.766 ±210.211 ^f	90.486 ±7.021 ^d	1821.380 ±141.299 ^d	0.464 ±0.571	31.101 ±38.260

a-Log phase on Day 4; b- Log phase on Day 5; c-Log phase on Day 7; d- Stationary phase on Day 11; e- Stationary phase on Day 18; f- Stationary phase on Day 21

N-Deficient: Trial 3

Hemolytic activity ($n=8 \pm$ one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). NC and SC were inoculated on May 4, 2005, TX was inoculated on May 22, 2005, and Pcal was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis (Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	45.844 ± 6.390	15,738.825 ± 2193.979	38.905 ± 3.211	19,821.984 ± 1636.208	7.179 ± 0.755	2941.548 ± 309.398	1.790 ± 0.670	779.079 ± 291.914
3	Log	85.380 $\pm 8.473^b$	12,702.783 $\pm 1260.536^b$	32.558 $\pm 6.608^*$	7337.772 $\pm 1489.092^*$	7.399 $\pm 1.092^a$	1638.369 $\pm 241.887^a$	1.323 ± 0.357	240.670 ± 64.974
10	Stationary	50.120 ± 8.856	5,879.926 ± 1039.101	74.370 ± 10.606	8972.830 ± 1279.726	16.825 $\pm 1.988^c$	3175.841 $\pm 375.304^c$	0.754 $\pm 0.652^d$	177.889 $\pm 153.802^d$

* - $n=4$; a-Log phase on Day 2; b-Log phase on Day 4; c-Stationary phase on Day 8

Hemolytic activity ($n=8 \pm$ one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in nitrogen-deficient conditions (N:P=4:1). NC and SC were inoculated on May 4, 2005, TX was inoculated on May 22, 2005, and Pcal was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	96.634 ± 7.112	6635.144 ± 488.400	92.264 ± 10.293	9401.804 ± 1048.959	88.762 ± 8.664	6994.394 ± 682.639	2.256 ± 1.494	204.563 ± 135.550
3	Log	93.177 $\pm 6.724^b$	3746.708 $\pm 270.339^b$	89.451 $\pm 18.002^*$	3665.439 $\pm 737.596^*$	88.516 $\pm 13.879^a$	4454.609 $\pm 698.533^a$	0.975 ± 0.354	36.962 ± 13.417
10	Stationary	79.570 ± 7.723	3012.582 ± 292.433	89.168 ± 6.618	3361.954 ± 249.544	99.059 $\pm 9.075^c$	4674.566 $\pm 428.318^c$	0.539 $\pm 0.472^c$	28.889 $\pm 25.889^c$

* - $n=4$; a-Log phase on Day 2; b-Log phase on Day 4; c-Stationary phase on Day 8

P-Deficient: Trial Averages

Hemolytic activity of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N:P=80:1). Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
	Lag	43.472 \pm 31.767	13755.502 \pm 4880.402	29.767 \pm 9.941	8581.064 \pm 3782.827	12.659 \pm 8.903	3406.254 \pm 2602.884	2.243 \pm 1.731	457.391 \pm 407.998
	Log	85.607 \pm 4.026	3084.742 \pm 347.592	81.398 \pm 7.211	2039.98 \pm 382.495	25.611 \pm 9.827	2081.198 \pm 1357.01	1.968 \pm 0.892	62.139 \pm 25.679
	Stationary	91.756 \pm 2.4481	691.985 \pm 15.783	92.751 \pm 4.173	462.845 \pm 139.360	61.078 \pm 35.632	2849.406 \pm 1723.339	3.144 \pm 3.767	40.949 \pm 41.264

Hemolytic activity of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N:P=80:1). Values for means of three individual trials \pm one standard deviation.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
	Lag	89.540 \pm 6.881	21339.202 \pm 28043.698	72.926 \pm 14.657	10430.26 \pm 9753.698	93.006 \pm 9.359	4758.803 \pm 3284.013	2.284 \pm 0.723	168.536 \pm 117.662
	Log	88.774 \pm 1.456	1085.792 \pm 113.142	94.589 \pm 1.934	809.984 \pm 79.376	94.769 \pm 1.411	3456.237 \pm 782.519	9.712 \pm 5.746	128.736 \pm 44.239
	Stationary	94.243 \pm 5.381	448.771 \pm 250.089	94.165 \pm 7.07	298.831 \pm 48.01	99.335 \pm 5.971	2595.053 \pm 697.868	13.084 \pm 9.523	149.204 \pm 124.195

P-Deficient: Trial 1

Hemolytic activity (n=8 ± one standard deviation) by the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). NC was inoculated on April 11, 2005, SC and Pcal were inoculated on April 3, 2005, and TX was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	29.593	8218.893	39.526	12,133.848	12.115	5162.640	2.345	784.908
		±5.713	±1586.902	±3.406	±1045.656	±1.583	±674.380	±0.937	±313.779
5	Log	82.760	3330.526	86.497	2310.445	18.831	2215.244	2.649	56.967
		±2.172	±87.407	±6.937 ^a	±185.311 ^a	±2.525	±297.063	±0.584 ^b	±12.545 ^b
	Stationary	92.877	681.241	96.034	595.829	23.021	1509.645	7.481	88.516
		±11.889 ^c	±87.195 ^c	±4.491 ^f	±27.859 ^f	±2.207 ^c	±144.743 ^c	±2.181 ^d	±25.810 ^d

a- Log phase on Day 6; b – Log phase on Day 7; c- Stationary phase on Day 9; d-Stationary phase on Day 14; e- Stationary Phase on Day 18; f- Stationary phase on Day 20

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). NC was inoculated on April 11, 2005, SC and Pcal were inoculated on April 3, 2005, and TX was inoculated on May 31, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	94.459	7716.049	87.995	7944.989	92.630	8223.559	3.025	297.773
		±5.885	±480.824	±4.092	±369.493	±15.259	±1354.554	±0.931	±91.618
5	Log	89.803	1165.795	95.956	753.857	96.266	4355.501	15.589	176.408
		±5.137	±66.677	±3.243 ^a	±25.477 ^a	±19.211	±869.135	±3.156 ^b	±35.713 ^b
	Stationary	98.297	257.499	94.675	309.156	101.162	1951.100	18.664	116.227
		±14.068 ^e	±36.849 ^e	±4.706 ^f	±15.366 ^f	±16.604 ^c	±320.289 ^c	±1.556 ^d	±9.691 ^d

a- Log phase on Day 6; b – Log phase on Day 7; c- Stationary phase on Day 9; d-Stationary phase on Day 14; e- Stationary Phase on Day 18; f- Stationary phase on Day 20

P-Deficient: Trial 2

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). All clones were inoculated on May 2, 2005, except TX, which was inoculated on May 17, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	79.816 ±9.483	15,613.867 ±1854.853	30.123 ±3.303	4603.973 ±504.832	4.041 ±0.943	415.873 ±97.056	3.821 ±1.493	586.911 ±229.334
5	Log	88.453 ±11494 ^a	2838.957 ±368.890 ^a	76.299 ±14.844*	1769.515 ±586.286*	21.120 ±2.973	662.139 ±93.234	2.296 ±0.787*	90.011 ±29.955*
15	Stationary	88.948 ±12.007	684.609 ±92.412	88.055 ±12.382	474.825 ±66.767	93.649 ±8.362 ^c	2245.051 ±200.443 ^c	1.263 ±0.672 ^b	19.567 ±10.416 ^b

* - n=4; a-Log phase on Day 6; b- Stationary phase on Day 10; c-Stationary Phase on Day 13

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). All clones were inoculated on May 2, 2005, except TX, which was inoculated on May 17, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	83.154 ±12.497	2709.547 ±407.155	72.063 ±11.050	2159.623 ±331.103	83.840 ±11.012	1691.755 ±222.176	2.245 ±1.133	67.616 ±34.138
5	Log	87.744 ±9.586 ^a	1005.788 ±109.872 ^a	93.221 ±6.706*	866.111 ±62.303*	93.463 ±9.609	2930.251 ±301.325	9.441 ±4.049*	120.795 ±51.799*
15	Stationary	88.138 ±11.172	357.042 ±45.253	86.853 ±11.294	246.497 ±32.052	104.179 ±9.464 ^c	2497.490 ±226.854 ^c	18.500 ±2.988 ^b	286.560 ±46.281 ^b

* - n=4; a-Log phase on Day 6; b- Stationary phase on Day 10; c-Stationary Phase on Day 13

P-Deficient: Trial 3

Hemolytic activity (n=8 ± one standard deviation) of the supernatant for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). NC and SC were inoculated on March 14, 2005, TX was inoculated on June 9, 2005, and Pcal was inoculated on July 13, 2005.

Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	21.006 ±1.309	17,433.747 ±1086.058	19.653 ±1.848	9005.370 ±846.678	21.882 ±1.930 ^a	4640.248 ±410.315 ^a	0.463 ±0.355 ^a	254.445 ±195.218 ^a
9	Log	21.918 ±1.403	2027.403 ±129.803	28.548 ±3.811	700.353 ±93.503	36.881 ±5.100 ^b	3366.210 ±465.487 ^b	0.958 ±0.314 ^c	39.439 ±12.947 ^c
20	Stationary	93.443 ±4.186	710.106 ±31.807	94.165 ±4.803	317.882 ±16.211	66.563 ±10.217 ^d	4793.522 ±735.687 ^d	0.688 ±0.381 ^e	14.763 ±8.178 ^e




a- Lag phase on Day 2; b- Log phase on Day 4; c-Log phase on Day 7; d- Stationary phase on Day 10; e-Stationary phase on Day 17

Hemolytic activity (n=8 ± one standard deviation) of the pellet for lag, log, and stationary phases of three geographically-distinct clones of *Prymnesium parvum* (NC=North Carolina; SC=South Carolina; TX=Texas) and *P. calathiferum* grown in phosphorus-deficient conditions (N: P = 80:1). NC and SC were inoculated on March 14, 2005, TX was inoculated on June 9, 2005, and Pcal was inoculated on July 13, 2005.




Day	Growth Stage	NC		SC		TX		Pcal	
		% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)	% Lysis	Lysis(Cell)
0	Lag	82.008 ±4.688	52,355.271 ±2992.825	58.719 ±4.962	20,697.243 ±1748.742	102.547 ±15.667 ^a	4361.096 ±666.249 ^a	1.581 ±0.690 ^a	140.219 ±85.173 ^a
9	Log	92.055 ±2.157	3153.710 ±73.894	90.261 ±2.653	820.128 ±24.105	94.578 ±6.279 ^b	3082.960 ±204.665 ^b	4.107 ±0.288 ^c	89.005 ±6.243 ^c
20	Stationary	96.294 ±4.450	731.772 ±33.814	100.966 ±5.223	340.841 ±17.629	92.663 ±9.692 ^d	3336.569 ±348.953 ^d	2.088 ±0.772 ^e	44.826 ±16.565 ^e

a- Lag phase on Day 2; b- Log phase on Day 4; c-Log phase on Day 7; d- Stationary phase on Day 10; e-Stationary phase on Day 17




Appendix C: Comparison of Initial Cell Density and Hemolytic Activity






Comparison of initial cell density and hemolytic activity of North Carolina *P. parvum* clone under nutrient-replete conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of South Carolina *P. parvum* clone under nutrient-replete conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of Texas *P. parvum* clone under nutrient-replete conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of *P. calathiferum* under nutrient-replete conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of North Carolina *P. parvum* clone under nitrogen-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of South Carolina *P. parvum* clone under nitrogen-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.




Comparison of initial cell density and hemolytic activity of Texas *P. parvum* clone under nitrogen-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of *P. calathiferum* under nitrogen-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.






Comparison of initial cell density and hemolytic activity of North Carolina *P. parvum* clone under phosphorus-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.



Comparison of initial cell density and hemolytic activity of South Carolina *P. parvum* clone under phosphorus-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.



Comparison of initial cell density and hemolytic activity of Texas *P. parvum* clone under phosphorus-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.



Comparison of initial cell density and hemolytic activity of *P. calathiferum* under phosphorus-deficient conditions for three trials. (A) Lag phase. (B) Log phase. (C) Stationary phase.