



Approximation of maps into spheres by regulous maps

MACIEJ ZIELIŃSKI

Abstract. Let X be a compact real algebraic set of dimension n . We prove that every Euclidean continuous map from X into the unit n -sphere can be approximated by a regulous map. This strengthens and generalizes previously known results.

Mathematics Subject Classification. 14P05, 14P25.

Keywords. Real algebraic variety, Regulous map, Sphere, Approximation.

1. Introduction. A recent direction of research in real algebraic geometry is to study intermediate classes of maps between continuous and regular maps. Such classes as continuous rational maps, stratified-regular maps, and regulous maps (which often coincide) have been studied in a series of papers [3, 7, 8, 11–27]. The aim of this note is to strengthen a certain related result of [21] and [15].

We begin by fixing some terminology. A *real algebraic variety* is a locally ringed space isomorphic to some algebraic subset of \mathbb{R}^n , for some positive integer n , endowed with the Zariski topology and the sheaf of regular functions. It is worth recalling that this class is identical with the class of quasi-projective real varieties, for more detail and information see [4]. A morphism of real algebraic varieties is called a *regular map*. We will be also interested in the Euclidean topology of such varieties and this is the topology we will mean, unless explicitly stated otherwise, when using topological notions. By a smooth map we understand a map of class C^∞ .

Let X be a real algebraic variety. A *stratification* \mathcal{S} of X is a finite collection of pairwise disjoint Zariski locally closed subvarieties whose union is equal to X . A map $f: X \rightarrow Y$ of real algebraic varieties is said to be *regulous* if it is continuous and if there exists some stratification \mathcal{S} of X such that $f|_S$ is a regular map for every $S \in \mathcal{S}$. We denote the set of all regulous maps between X and Y by $\mathcal{R}^0(X, Y)$. We shall treat $\mathcal{R}^0(X, Y)$ as a subspace of the space

$\mathcal{C}(X, Y)$ of all continuous maps endowed with the compact-open topology. Note that regulous maps in the sense of our definition were called stratified-regular in [21] and the follow-up papers [20, 22]. This definition is different but equivalent to that of [7] where the terminology was introduced, see [21, Remark 2.3] or [24].

Each regulous map is also *continuous rational*—i.e. f is continuous and $f|_{X^0}$ is regular for some Zariski open dense subset $X^0 \subset X$. While the converse is false in general, it is true if X is nonsingular, see [11].

Let us first recall the following result contained in [15].

Theorem 1.1. *Let X be a compact nonsingular real algebraic variety of dimension $p \geq 1$. Then the set $\mathcal{R}^0(X, \mathbb{S}^p)$ is dense in $\mathcal{C}(X, \mathbb{S}^p)$.*

A related weaker result allowing for a singular X is contained in [21]:

Theorem 1.2. *Let X be a compact real algebraic variety of dimension $p \geq 1$. Then any continuous map $X \rightarrow \mathbb{S}^p$ is homotopic to a regulous map.*

Our aim is to strenghten Theorem 1.2 by showing that the nonsingularity assumption of Theorem 1.1 is unnecessary.

Theorem 1.3. *Let X be a compact real algebraic variety of dimension $p \geq 1$. Then the set $\mathcal{R}^0(X, \mathbb{S}^p)$ is dense in $\mathcal{C}(X, \mathbb{S}^p)$.*

It is well known that analogous results do not hold if regulous maps in Theorems 1.1, 1.2, and 1.3 are replaced with regular maps. For example, a continuous map $\mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{S}^2$ is homotopic to a regular map if and only if it is null-homotopic, cf. [5].

2. Proof of the main theorem. We shall use the concept of the algebraic cohomology classes of a real algebraic variety which we recall now. Let X be a compact nonsingular real algebraic variety. A class in $H^*(X; \mathbb{Z}/2)$ is said to be algebraic if it is Poincaré dual to a homology class in $H_*(X, \mathbb{Z}/2)$ represented by an algebraic subset. The set $H_{\text{alg}}^*(X; \mathbb{Z}/2)$ of all algebraic cohomology classes is a subring of $H^*(X; \mathbb{Z}/2)$ and if $f: X \rightarrow Y$ is a regular map, then the induced map f^* in cohomology maps $H_{\text{alg}}^*(Y; \mathbb{Z}/2)$ into $H_{\text{alg}}^*(X, \mathbb{Z}/2)$, cf. [2, 4, 6].

An important tool we need is [15, Lemma 2.2], which allows controlled approximation of continuous maps into projective space by regular maps. We restate it here for convenience.

Lemma 2.1. *Let X be a compact nonsingular real algebraic variety, and let A be a Zariski closed subvariety of X . Let $f: X \rightarrow \mathbb{P}^n(\mathbb{R})$ be a continuous map whose restriction $f|_A: A \rightarrow \mathbb{P}^n(\mathbb{R})$ is a regular map. Assume that*

$$f^*(H^1(\mathbb{P}^n(\mathbb{R}); \mathbb{Z}/2)) \subset H_{\text{alg}}^1(X; \mathbb{Z}/2).$$

Then one can find a regular $g: X \rightarrow \mathbb{P}^n(\mathbb{R})$ arbitrarily close to f and satisfying $g|_A = f|_A$ (i.e. every neighborhood of f in $\mathcal{C}(X, \mathbb{P}^n(\mathbb{R}))$ contains such a map).

We are now ready to prove Theorem 1.3

Proof. Let f be any map in $\mathcal{C}(X, \mathbb{S}^p)$. Treating X as a closed subset of \mathbb{R}^m for some $m \in \mathbb{N}$, one can find a smooth map $f_0: U \rightarrow \mathbb{S}^p$ defined on some neighborhood U of X in \mathbb{R}^m such that $f_0|_X$ is arbitrarily close to f . Let Σ denote the singular locus of X . Then $f_0(\Sigma) \subsetneq \mathbb{S}^p$, since $\dim \Sigma < p$. This allows us to approximate $f_0|_\Sigma$ by regular maps using the stereographic projection and Weierstrass approximation theorem. We can therefore reduce the proof (by replacing f with suitably modified f_0) to the case where f is a restriction of a smooth map defined on a neighborhood of X in \mathbb{R}^m and $f|_\Sigma$ is regular with $f(\Sigma) \subsetneq \mathbb{S}^p$. Then, by Sard's theorem, there exists an $s_0 \in \mathbb{S}^p \setminus f(\Sigma)$ which is a regular value of the smooth map $f|_{X \setminus \Sigma}$.

By Hironaka's resolution of singularities theorem [9, 10], there exists a finite composition of blowups $\pi: Y \rightarrow X$ over Σ with Y nonsingular. The restriction $\bar{\pi}: Y \setminus \pi^{-1}(\Sigma) \rightarrow X \setminus \Sigma$ of π is then a biregular isomorphism and s_0 is a regular value of the smooth map $f \circ \bar{\pi}$. Letting $F = (f \circ \pi)^{-1}(s_0) = (f \circ \bar{\pi})^{-1}(s_0)$, consider the blowup of Y with center F , which we will denote $\sigma: B(Y, F) \rightarrow Y$, and the blowup $\tau: B(\mathbb{S}^p, s_0) \rightarrow \mathbb{S}^p$ of \mathbb{S}^p over s_0 . Since $\dim X = p$, the set F is finite as a fiber of the smooth map $f \circ \bar{\pi}$ over its regular value s_0 , hence $B(Y, F)$ is a real algebraic variety. Finiteness of F and the fact that $f \circ \pi$ is also a smooth map allow us to apply [1, Lemma 2.5.9] in order to construct a smooth lifting g of $f \circ \pi$ to a map between $B(Y, F)$ and $B(\mathbb{S}^p, s_0)$ making the following diagram commute:

$$\begin{array}{ccc}
 B(Y, F) & \xrightarrow{g} & B(\mathbb{S}^p, s_0) \\
 \sigma \downarrow & & \tau \downarrow \\
 Y & \xrightarrow{f \circ \pi} & \mathbb{S}^p \\
 \pi \downarrow & & \parallel \\
 X & \xrightarrow{f} & \mathbb{S}^p
 \end{array}$$

Our aim for now is to find a regular map $H: B(Y, F) \rightarrow B(\mathbb{S}^p, s_0)$ arbitrarily close to g in $\mathcal{C}(B(Y, F), B(\mathbb{S}^p, s_0))$ in such a way that the map $\tilde{f}: X \rightarrow \mathbb{S}^p$ making the following diagram commute will be regulous and close to f :

$$\begin{array}{ccc}
 B(Y, F) & \xrightarrow{H} & B(\mathbb{S}^p, s_0) \\
 \pi \circ \sigma \downarrow & & \tau \downarrow \\
 X & \xrightarrow{\tilde{f}} & \mathbb{S}^p
 \end{array}$$

Let $D = \sigma^{-1}(F)$ and $E = \tau^{-1}(s_0)$. Then D and E are nonsingular algebraic hypersurfaces in the respective blowups. Let $u \in H^1(B(Y, F); \mathbb{Z}/2)$ be the cohomology class Poincaré dual to the homology class in $H_*(B(Y, F); \mathbb{Z}/2)$ represented by D and let v be the class in $H^1(B(\mathbb{S}^p, s_0); \mathbb{Z}/2)$ dual to the class in $H_*(B(\mathbb{S}^p, s_0); \mathbb{Z}/2)$ represented by E . Recall that there exists a biregular isomorphism $\varphi: B(\mathbb{S}^p, s_0) \rightarrow \mathbb{P}^p(\mathbb{R})$ such that

$$\varphi(E) = \mathbb{P}^{p-1}(\mathbb{R}) \subset \mathbb{P}^p(\mathbb{R}). \tag{1}$$

Hence $H^1(B(\mathbb{S}^p, s_0); \mathbb{Z}/2) = \varphi^*(H^1(\mathbb{P}^p(\mathbb{R}); \mathbb{Z}/2)) \cong \mathbb{Z}/2$ with v the generator. Since g is transverse to E and $D = g^{-1}(E)$ we have $u = g^*(v)$ (cf. [6, Proposition 2.15]) and it follows that

$$g^*(H^1(B(\mathbb{S}^p, s_0); \mathbb{Z}/2)) \subset H_{\text{alg}}^1(B(Y; F); \mathbb{Z}/2). \quad (2)$$

Let $i: D \rightarrow B(Y, F)$ and $j: E \rightarrow B(\mathbb{S}^p, s_0)$ be the inclusion maps and let $\bar{g}: D \rightarrow E$ be the map induced by the restriction of g to D . Then, since $g \circ i = j \circ \bar{g}$ we have $\bar{g}^*(j^*(v)) = i^*(g^*(v)) = i^*(u)$. This allows us to apply Lemma 2.1 to approximate \bar{g} by regular maps. Indeed, since by (1) $H^1(E; \mathbb{Z}/2)$ is generated by $j^*(v)$ and by definition $i^*(u)$ is in $H_{\text{alg}}^1(D; \mathbb{Z}/2)$, we have

$$\bar{g}^*(H^1(E; \mathbb{Z}/2)) \subset H_{\text{alg}}^1(D; \mathbb{Z}/2).$$

Therefore, there exists a regular map $r: D \rightarrow E$ arbitrarily close to \bar{g} in $\mathcal{C}(D, E)$.

Let $L = (\pi \circ \sigma)^{-1}(\Sigma)$ and let $h: \Sigma \rightarrow B(\mathbb{S}^p, s_0)$ be the regular map given by $(\tau|_{\tau^{-1}(\mathbb{S}^p \setminus \{s_0\})})^{-1} \circ (f|_{\Sigma})$. We now extend $j \circ r$ to a smooth map $G: B(Y, F) \rightarrow B(\mathbb{S}^p, s_0)$ close to g such that $G|_L = (h \circ \pi \circ \sigma)|_L$ (this is possible since the sets E and L are disjoint). If G is close enough to g , we have $g^* = G^*$ and hence by (2) we can apply Lemma 2.1 to G with $A = D \cup L$. This gives a regular map $H: B(Y, F) \rightarrow B(\mathbb{S}^p, s_0)$ which is close to g and satisfies $H|_L = (h \circ \pi \circ \sigma)|_L$ and $H(D) \subset E$. The latter shows that the map $\bar{H}: Y \rightarrow \mathbb{S}^p$ given by

$$\bar{H}(y) = \begin{cases} s_0, & \text{for } y \in F, \\ \tau(H(\sigma^{-1}(y))), & \text{for } y \notin F, \end{cases}$$

is continuous and well-defined. Moreover $\bar{H}(y) = f \circ \pi(y)$ for $y \in \pi^{-1}(\Sigma)$. Also note that \bar{H} is close to $f \circ \pi$, since $(\tau \circ H)|_{Y \setminus F}$ is close to $\tau \circ f \circ \pi$. Therefore, the map $\tilde{f}: X \rightarrow \mathbb{S}^p$ given by

$$\tilde{f}(x) = \begin{cases} f(x), & \text{for } x \in \Sigma, \\ \bar{H}(\pi^{-1}(x)), & \text{for } x \notin \Sigma, \end{cases}$$

is well-defined, continuous, and can be chosen arbitrarily close to f .

It remains to check that \tilde{f} is regulous. Since $\tilde{f}|_{\Sigma}$ is regular as is \tilde{f} in restriction to the finite set $f^{-1}(s_0)$, it is enough to show $\tilde{f}|_{X \setminus A}$ is regular where $A = \Sigma \cup f^{-1}(s_0)$. Since we have $\tilde{f}|_{X \setminus A} = \tau \circ H \circ (\sigma|_{Y \setminus F})^{-1} \circ (\pi|_{X \setminus \Sigma})^{-1}$ with all the maps on the right-hand side regular, the proof is finished. \square

Acknowledgements. The author was partially supported by the National Science Centre (Poland) under Grant Number 2014/15/B/ST1/00046.

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- [1] S. AKBULUT AND H. KING, *Topology of Real Algebraic Sets*, Mathematical Sciences Research Institute Publications, 25, Springer, New York, 1992.
- [2] R. BENEDETTI AND A. TOGNOLI, Remarks and counterexamples in the theory of real algebraic vector bundles and cycles, In: *Real Algebraic Geometry and Quadratic Forms* (Rennes, 1981), 198–211, Lecture Notes in Math., 959, Springer, Berlin, 1982.
- [3] M. BILSKI, W. KUCHARZ, A. VALETTE, AND G. VALETTE, Vector bundles and regulous maps, *Math. Z.* **275** (2013), 403–418.
- [4] J. BOCHNAK, M. COSTE, AND M.-F. ROY, *Real Algebraic Geometry*, *Ergebnisse der Mathematik und ihrer Grenzgebiete* (3), 36, Springer, Berlin, 1998.
- [5] J. BOCHNAK AND W. KUCHARZ, Realization of homotopy classes by algebraic mappings, *J. Reine Angew. Math.* **377** (1987), 159–169.
- [6] A. BOREL AND A. HAEFLIGER, La classe d’homologie fondamentale d’un espace analytique, *Bull. Soc. Math. France* **89** (1961), 461–513.
- [7] G. FICHO, J. HUISMAN, F. MANGOLTE, AND J.-Ph. MONNIER, Fonctions régulières, *J. Reine Angew. Math.* **718** (2016), 103–151.
- [8] G. FICHO, J.-P. MONNIER, AND R. QUAREZ, Continuous functions in the plane regular after one blowing up, *Math. Z.* **285** (2017), 287–323.
- [9] H. HIRONAKA, Resolution of singularities of an algebraic variety over a field of characteristic zero, *Ann. of Math.* **79** (1964), 109–326.
- [10] J. KOLLÁR, *Lectures on Resolution of Singularities*, *Annals of Mathematics Studies*, 166, Princeton University Press, Princeton, NJ, 2007.
- [11] J. KOLLÁR AND K. NOWAK, Continuous rational functions on real and p-adic varieties, *Math. Z.* **279** (2015), 85–97.
- [12] J. KOLLÁR, W. KUCHARZ, AND K. KURDYKA, Curve-rational functions, to appear in *Math. Ann.*
- [13] W. KUCHARZ, Rational maps in real algebraic geometry, *Adv. Geom.* **9** (2009), 517–539.
- [14] W. KUCHARZ, Regular versus continuous rational maps, *Topology Appl.* **160** (2013), 1086–1090.
- [15] W. KUCHARZ, Approximation by continuous rational maps into spheres, *J. Eur. Math. Soc.* **16** (2014), 1555–1569.
- [16] W. KUCHARZ, Continuous rational maps into the unit 2-sphere, *Arch. Math.* **102** (2014), 257–261.
- [17] W. KUCHARZ, Some conjectures on stratified-algebraic vector bundles, *J. Singul.* **12** (2015), 92–104.
- [18] W. KUCHARZ, Continuous rational maps into spheres, *Math. Z.* **283** (2016), 1201–1215.
- [19] W. KUCHARZ, Stratified-algebraic vector bundles of small rank, *Arch. Math.* **107** (2016), 239–249.

- [20] W. KUCHARZ AND K. KURDYKA, Some conjectures on continuous rational maps into spheres, *Topology Appl.* **208** (2016), 17–29.
- [21] W. KUCHARZ AND K. KURDYKA, Stratified-algebraic vector bundles, to appear in *J. Reine Angew. Math.*
- [22] W. KUCHARZ AND K. KURDYKA, Comparison of stratified-algebraic and topological K-theory, [arXiv:1511.04238](https://arxiv.org/abs/1511.04238) [math.AG].
- [23] W. KUCHARZ AND K. KURDYKA, Linear equations on real algebraic surfaces, [arXiv:1602.01986](https://arxiv.org/abs/1602.01986) [math.AG].
- [24] W. KUCHARZ AND M. ZIELIŃSKI, Regulous vector bundles, [arXiv:1703.05566](https://arxiv.org/abs/1703.05566) [math.AG].
- [25] J.-P. MONNIER, Semi-algebraic geometry with rational continuous functions, [arXiv:1603.04193](https://arxiv.org/abs/1603.04193) [math.AG].
- [26] K.J. NOWAK, Some results of algebraic geometry over Henselian rank one valued fields, *Sel. Math. New Ser.* **28** (2017), 455–495.
- [27] M. ZIELIŃSKI, Homotopy properties of some real algebraic maps, *Homology Homotopy Appl.* **18** (2016), 287–294.

MACIEJ ZIELIŃSKI

Institute of Mathematics, Faculty of Mathematics and Computer Science

Jagiellonian University

ul. Łojasiewicza 6

30-348 Kraków

Poland

e-mail: Maciej.Zielinski@im.uj.edu.pl

Received: 26 May 2017