

## ROLE OF NITRIC OXIDE IN SJÖGREN'S SYNDROME

YRJÖ T. KONTTINEN, LOUISE A. M. PLATTS, SUSANNE TUOMINEN, KARI K. EKLUND,  
NINA SANTAVIRTA, JYRKI TÖRNWALL, TIMO SORSA, MIKA HUKKANEN, and JULIA M. POLAK

**Objective.** To measure levels of salivary nitrite ( $\text{NO}_2^-$ ) and to localize nitric oxide synthases (NOS) in the labial salivary glands (LSGs) of patients with Sjögren's syndrome (SS).

**Methods.**  $\text{NO}_2^-$  was measured by the Griess reaction. LSGs were analyzed using NADPH-diaphorase histochemical and immunohistochemical studies to determine the constitutive NOS (neuronal [ncNOS] and endothelial [ecNOS]) and inducible NOS (iNOS) isoforms.

**Results.** The  $\text{NO}_2^-$  concentration (mean  $\pm$  SEM  $307 \pm 51 \mu\text{M}$  versus  $97 \pm 16 \mu\text{M}$ ;  $P < 0.05$ ) and output ( $166 \pm 46$  nmoles/minute versus  $37 \pm 7$  nmoles/minute) were increased in SS patients compared with healthy control subjects. NADPH-diaphorase was found in some nerve fibers and endothelial cells, and, in SS, was found in myoepithelial, acinar, and ductal epithelial cells, but in only a few inflammatory cells. In SS, ncNOS-immunoreactive nerve fibers were sparse and ecNOS was found in a minority of the CD31-positive vascular endothelial cells and acinar cells, whereas iNOS was localized in myoepithelial, acinar, and ductal epithelial cells, often together with tumor necrosis factor  $\alpha$ .

**Conclusion.** Nitrite was found in normal human saliva. NO produced by ncNOS probably acts as a non-adrenergic, noncholinergic neurotransmitter, whereas that produced by ecNOS exerts a vasodilatory effect. SS

patients had increased  $\text{NO}_2^-$  concentrations, with most of the superfluous salivary NO being produced not by the immigrant inflammatory cells, but rather, by the resident salivary gland cells. NO may contribute to inflammatory damage and acinar cell atrophy in SS.

Nitric oxide (NO) is produced by vascular endothelium, nerve terminals, and inflammatory cells in a reaction catalyzed by NO synthase (NOS), which converts L-arginine to NO and L-citrulline (1). NOS occurs in at least 3 different isoforms, endothelial constitutive NOS (ecNOS), neuronal constitutive NOS (ncNOS), and inducible NOS (iNOS) (2,3). Both ecNOS and ncNOS are constitutively expressed, but are regulated as to their level of activity by various soluble ligands and/or shear forces through calcium/calmodulin-dependent second messenger pathways. NO concentrations produced by this route are usually relatively low (in the picomolar range). In contrast, regulation of iNOS occurs at the level of gene transcription and is relatively slow, but leads to long-lasting and significant increases in NO production (giving rise to nanomolar local concentrations of NO) (1,3,4).

NO has a short half-life and is rapidly converted to nitrites and nitrates. Therefore, its range of action is limited. This is important from the point of view of regulation, because NO itself is a small, lipid-soluble compound, which easily penetrates the cell membrane of nearby cells and can therefore affect them in a receptor-independent autocrine and/or paracrine manner. At the same time, NO is a highly reactive compound, due to an unpaired electron in its molecular orbital. At low concentrations, it preferentially activates guanylate cyclase, which thus produces cGMP. At high concentrations, NO inhibits various iron-containing DNA synthetases and mitochondrial enzymes, and thus inhibits cell growth and division, finally leading to cell death (5).

Due to its potential dual role in normal salivary gland function and damage in pathologic conditions, it was considered important to characterize the eventual

Supported by the Signe and Ane Gyllenberg Foundation, the Sigrid Juselius Foundation, the Finnish Academy, Helsinki University Central Hospital (Evo Grant), Finland, and the Great Charity of Freemasons, UK.

Yrjö T. Konttinen, MD, PhD: Helsinki University Central Hospital, Helsinki, Finland; Louise A. M. Platts, BSc, Mika Hukkanen, PhD, Julia M. Polak: Royal Postgraduate Medical School, University of London, London, UK; Susanne Tuominen, DDS, Kari K. Eklund, MD, Nina Santavirta, MSc, Jyrki Törnwall, DDS, Timo Sorsa, DDS: University of Helsinki, Helsinki, Finland.

Address reprint requests to Yrjö T. Konttinen, MD, PhD, Institute of Biomedicine, Department of Anatomy, PO Box 9 (Siltavuorenpenger 20 A), FIN-00014 University of Helsinki, Helsinki, Finland.

Submitted for publication February 26, 1996; accepted in revised form December 6, 1996.

presence of NO in saliva and to localize the cells/NOS isoforms responsible for NO production in patients with Sjögren's syndrome (SS).

## PATIENTS AND METHODS

**Patients and samples.** All patients and healthy volunteers gave their informed consent. The Declaration of Helsinki (6) was followed throughout the study, which was approved by the local ethics committee.

Both patients and controls provided a stimulated salivary sample, as described elsewhere (7). Briefly, stimulation of saliva was achieved by the chewing of paraffin for 5 minutes, and the stimulated whole saliva was then collected. Immediately after collecting the saliva, the samples for  $\text{NO}_2^-$  measurement were centrifuged at 1,000g for 5 minutes, and the supernatants were frozen at  $-70^\circ\text{C}$  until analyzed. For vasoactive intestinal peptide (VIP) measurement, the saliva was collected into chilled tubes containing 50,000 IU/ml aprotinin and 10 mM EDTA to minimize artifactual neuropeptide degradation *in vitro* (8).

The mean age of the 12 primary SS patients, who were all women, was 51.6 years (range 21–74). The duration of the SS was a mean of 9 years (range 1–23). All patients were nonsmokers. The mean age of the 10 healthy control women was 47.7 years (range 39–56).

The dental/periodontal features of all patients were evaluated for decayed, missing, or filled surfaces using the DMFS index (9), bleeding on probing using the BOP index (10), presence of visible plaque using the visible plaque index (11), retentive calculus (subgingival calculus) using the RC index (12), items on the community periodontal index of treatment needs (13), and pocket probing depth (10). Eventual correlations were sought between the dental and periodontal status and salivary  $\text{NO}_2^-$  concentration.

Between 6 and 8 labial salivary glands (LSGs) were obtained from each patient by biopsy. Samples were taken by blunt dissection under infiltration anesthesia and fixed by immersion in freshly made 1% paraformaldehyde in 0.01M phosphate buffered 0.15M saline (PBS; pH 7.1) at  $4^\circ\text{C}$  for 16–18 hours. After fixation, cryoprotection was performed in 15% (weight/volume) sucrose and 0.01% (w/v) sodium azide in PBS for several days at  $4^\circ\text{C}$  before the samples were embedded in Tissue-Tek OCT compound (Lab-Tek Products, Elkhart, IN), snap frozen in liquid nitrogen, and stored at  $-70^\circ\text{C}$  until staining. LSGs were obtained from 4 patients with SS who fulfilled the European criteria for SS (14), and from 2 healthy controls who did not have SS (14) and who had a focus score of  $<1$  (15).

**Jenkins activity survey (JAS).** The JAS scale consists of 84 items, which are divided into 4 subscales. For the present study, we recorded the values of the actual A-B factor, which involves 21 items on the JAS that best reflect the stress behavior of a person (16). The reliability of the stress factor was confirmed with the use of the Cronbach  $\alpha$  test (17).

**Nitrite analysis.**  $\text{NO}_2^-$ , a stable end-product of NO oxidation, was measured using the Griess reagent, consisting of equal volumes of 0.1% *N*-(1-naphthyl)ethylenediamine HCl and 1% sulfanilamide plus 5%  $\text{H}_3\text{PO}_4$  (18). Briefly, 50- $\mu\text{l}$

aliquots of the saliva sample, in duplicate, from each donor were incubated with an equal volume of the Griess reagent in a 96-well U-bottomed microtiter plate for 5 minutes at  $22^\circ\text{C}$ . With each study, a standard curve was established using  $\text{NaNO}_2$  in a range of 10–1,000  $\mu\text{M}$  for each assay. After a 10-minute reaction, the microplate was read on a spectrophotometer at 545 nm. The data were derived from the slope of the standard curve and were presented as the mean value of the duplicates. Results of the salivary  $\text{NO}_2^-$  measurements were expressed as a concentration (in  $\mu\text{M}$ ), but also as  $\text{NO}_2^-$  output (output = concentration  $\times$  volume; in nmoles/minute).

**NADPH-diaphorase histochemical analysis.** For histochemical demonstration of NADPH-diaphorase activity, which is an indicator of NOS activity, freshly cut and PBS-rinsed 10- $\mu\text{m}$  cryostat sections from the midsagittal plane of LSGs were mounted on formol-gelatin-coated slides and incubated in 1 mM  $\beta$ -NADPH, 0.5 mM nitroblue tetrazolium, 1.5 mM L-malic acid, and 0.1% Triton X-100 in 50 mM Tris HCl, pH 7.4, at  $37^\circ\text{C}$  for 1 hour. Finally, slides were rinsed in distilled water and mounted (19). This procedure resulted in an intense deep-blue nitroblue tetrazolium formazan formation, which was not observed when NADPH was not added to the reaction mixture.

**Acetylcholinesterase histochemical analysis.** Acetylcholinesterase activity in LSG tissue was visualized as previously described (20). Briefly, the cryostat sections were first immersed in an incubation medium for 45 minutes at  $37^\circ\text{C}$ . The acetylcholinesterase activity was then visualized by immersion of the sections for 5 minutes in 50 mM Tris HCl buffer, pH 7.6, containing 0.04% 3,3'-diaminobenzidine and 0.3% nickel ammonium sulfate. Sections were incubated in this solution for a further 5 minutes after addition of hydrogen peroxide, for a final concentration of 0.003%.

**Immunocytochemical analysis for NOS.** Polyclonal antibodies for the cNOS and iNOS isoforms used in this study have been characterized in detail elsewhere. Briefly, antibodies for ncNOS, kindly provided by Drs. S. Moncada and V. Riveros-Moreno (Wellcome Research Laboratories, Beckenham, Kent, UK), were raised against the whole protein purified from rat brain and against synthetic peptide fragment 53 of rat brain ncNOS; both of these antisera cross-react with the ecNOS isoform at high concentrations (21). Monoclonal antibodies for ecNOS were raised against a purified whole protein of particulate bovine aortic endothelial NOS isoform (22) and were kindly provided by Dr. J. S. Pollock (Abbott Laboratories, Abbott Park, IL). Antibodies for iNOS were raised against synthetic peptides corresponding to amino acid residues 47–71 of murine macrophage iNOS and 54–76 of human hepatocyte iNOS (23–25). Murine macrophage iNOS antibodies were kindly provided by Drs. S. Moncada and V. Riveros-Moreno (Wellcome Research Laboratories) and human hepatocyte iNOS antibodies by Dr. T. J. Evans (Royal Postgraduate Medical School).

Antibodies against the synthetic 47–71 peptide of macrophage iNOS have been characterized for specificity in detail elsewhere, with the use of an antigen preabsorption test and Western blotting (25). Similarly, the specificity of the human hepatocyte iNOS antibodies have been confirmed by antigen reabsorption test with the use of a synthetic 54–76 peptide, by Western blotting, and by comparison with *in situ*

hybridization (26). There was no cross-reactivity between eNOS and iNOS antibodies.

Antibodies against tumor necrosis factor  $\alpha$  (TNF $\alpha$ ; Dr. W. A. Buurman, University of Limburg, Maastricht, The Netherlands) and CD68 (Dako, Glostrup, Denmark) were used to demonstrate a possible induction mechanism of iNOS and its localization in macrophages. Antibodies against CD31 (platelet endothelial cell adhesion molecule; Dr. A. V. Mazurov, Institute of Experimental Cardiology, Cardiology Research Centre, Moscow, Russia) and against nitrotyrosine (TCS Biological, Bucks, UK) were used, in some cases, to demonstrate endothelial cells and the possible site of action of oxidative NO-induced damage in inflammatory LSG tissue, respectively.

Fifteen micrometer-thick cryostat sections were cut through the midsagittal planes of LSGs, transferred onto formol-gelatin-coated slides, and allowed to dry for 1 hour before any further procedures were carried out. Endogenous peroxidase was inhibited by soaking the sections in 0.3% (v/v) hydrogen peroxide in methanol at 4°C for 20 minutes. The sequential sections were then placed in a humid chamber and incubated 1) overnight at 4°C with primary antibodies as described above; 2) for 30 minutes at 22°C with biotinylated goat anti-rabbit (for polyclonal antibodies) or horse anti-mouse (for monoclonal antibodies) IgG (Vector Laboratories, Peterborough, UK), diluted 1:100 in PBS with 0.1% w/v bovine serum albumin; and 3) for 60 minutes at 22°C with avidin-biotin-peroxidase complex (ABC; avidin and biotinylated horseradish peroxidase diluted 1:200 in 0.1M PBS, pH 7.4 [27], for 5 minutes at 22°C with 3,3'-diaminobenzidine and nickel ammonium sulfate containing glucose and glucose oxidase). After each step, the slides were rinsed 3 times in PBS for 5 minutes. Finally, the slides were dehydrated in ethanol, cleared in Inhibisol (Kalon Chemicals, Cramlington, UK), and mounted in Pertex (Histolab Products, Gothenburg, Sweden).

Omission of the primary antibodies and use of irrelevant immunoglobulin from the same species and of the same class, isotype, and concentration were used as staining controls. Conventional hematoxylin and eosin staining of sequential sections was used for comparative histopathologic analysis.

**Other immunohistochemical analyses.** VIP immunolocalization was performed using specific antibodies and the ABC technique, as described in detail elsewhere (28,29). All samples were also stained with anti-CD31, anti-CD68, and antimyeloperoxidase antibodies as markers for endothelial cells, macrophages, and neutrophils, respectively. Furthermore, all samples were similarly assessed for the presence, localization, and extent of expression of TNF $\alpha$ .

An indirect immunofluorescence method was used for colocalization of iNOS and CD68-positive macrophages. Cryostat sections, 7–10- $\mu$ m thick, mounted on formol-gelatin-coated slides, were washed in PBS and incubated with 1) diluted normal goat serum for 30 minutes at 22°C before blotting it away; 2) monoclonal anti-CD68 antibodies for 16 hours at 4°C; 3) tetramethylrhodamine isothiocyanate (TRITC)-labeled goat anti-mouse IgG antiserum (1:100) for 60 minutes at 22°C in the dark; 4) polyclonal iNOS antibodies (1:300) for 16 hours at 4°C in the dark; and 5) fluorescein isothiocyanate (FITC)-labeled goat anti-rabbit IgG (1:100) for 60 minutes at 22°C in the dark before mounting with Vectashield (Vector Laboratories) to reduce quenching of the

**Table 1.** Correlations between the salivary nitrite (NO $_2^-$ ) concentration and dental and periodontal status of 10 patients with Sjögren's syndrome\*

Dental/periodontal status	Spearman's rank correlation coefficient	P
DMFS	0.2492	0.4874
BOP	0.5294	0.1155
VPI	-0.2339	0.5155
RC	-0.4786	0.1617
PPD (4–5 mm)	-0.3556	0.3133
PPD (>6 mm)	-0.0580	0.8735


\* DMFS = decayed, missing, or filled surfaces; BOP = bleeding on probing; VPI = visible plaque index (yes/no); RC = retentive calculus (subgingival calculus); PPD = pocket probing depth. The community periodontal index of treatment need values were zero for all patients studied. None of the correlations were statistically significant.

fluorescent signal. Between the steps, slides were washed in PBS, if not stated otherwise. Immunostained preparations were examined using an Olympus BX60 (Olympus Optical, Tokyo, Japan) research microscope with selective filters for specific emission wavelengths of TRITC and FITC.

**Cell counting.** The total number of CD68-positive macrophages and cells immunoreactive for the murine and human iNOS antibodies were counted manually using a graticule and a 20 $\times$  objective magnification. Four randomly selected fields which contained at least 1 inflammatory cell focus with more than 50 inflammatory mononuclear cells were counted in each study subject. The cells that were counted included inflammatory cells, acinar cells, and some ductal epithelial cells. Cells with typical mast cell morphology were excluded from cell counts.

**Radioimmunoassay for VIP.** The neuropeptide concentration in saliva was measured using a standard competitive  $^{125}$ I-radioimmunoassay kit (Peninsula, Belmont, CA) as described in detail elsewhere (8). Briefly, 1) 100  $\mu$ l of sample or standard was incubated in polypropylene tubes for 24 hours at 4°C with 100  $\mu$ l of rabbit anti-human VIP IgG, 2) 100  $\mu$ l of  $^{125}$ I-labeled peptide (10,000–15,000 counts per minute/100  $\mu$ l) was added and incubated for 24 hours at 4°C, 3) 100  $\mu$ l of goat anti-rabbit IgG and 100  $\mu$ l of normal rabbit serum were added and incubated for 90 minutes at room temperature before 4) 500  $\mu$ l of radioimmunoassay buffer was added, followed by centrifugation at 1,700g for 20 minutes at 4°C. The supernatant was discarded and the radioactivity of the pellet was measured with a programmed gamma scintillation counter. All samples were analyzed in duplicate, with individual controls for apparent binding in the absence of antibody (i.e., total counts, nonspecific binding, total binding).

**Statistical analysis.** Results are presented as the mean  $\pm$  SEM. Comparisons between the 2 groups were performed with the Student's *t*-test for normally distributed variables, and with the Mann-Whitney U test for non-normally distributed variables. For analysis of the linear relationship between 2 variables, Pearson's correlation coefficient was used for normally distributed variables and Spearman's rank correlation coefficient for non-normally distributed variables. BMDP-PC, version 7.01 (BMDP Statistical Software, Cork, Ireland), statistical software was used for all calculations



**Figure 1.** Histochemical and immunohistochemical localization of NADPH-diaphorase activity and the 3 different isoforms of nitric oxide synthase (NOS) in labial salivary glands from patients with Sjögren's syndrome. **A,** Histochemical demonstration of NADPH-diaphorase activity within myoepithelial cells (**arrows**) and acinar cells of acini (**open arrow**) and excretory duct epithelia (**arrowheads**). **B,** Neuronal NOS immunoreactivity in nerve fibers (**arrows**) terminating near an acinus. Neuronal NOS-containing nerve fibers can also be seen in connective tissue surrounding the acinus. **C,** CD31-positive vascular endothelial cells (**arrows**) within a small focal inflammatory cell infiltrate surrounding an excretory duct. **D,** Endothelial NOS (**arrows**) was found to colocalize with CD31-positive cells, as evidenced by immunostaining of serial sections. **E,** Immunoreactivity for endothelial NOS in several acini localized mainly in acinar cells (**arrows**). **F,** Antibodies for human hepatocyte inducible NOS (iNOS) isoform reveal numerous iNOS-immunoreactive mononuclear cells (**open arrows**) within an inflammatory cell infiltrate. (Original magnification  $\times 400$ .)

except for those in the JAS, which were analyzed using the SPSS software package (SPSS Inc., Chicago, IL).

## RESULTS

**Relationship of salivary  $\text{NO}_2^-$  to SS, stress, and VIP.** The  $\text{NO}_2^-$  concentration was higher in patients with SS (mean  $\pm$  SEM  $307 \pm 51 \mu\text{M}$ ) than in healthy control individuals ( $97 \pm 16 \mu\text{M}$ ;  $P < 0.05$ ). There was no correlation between the salivary  $\text{NO}_2^-$  concentration and the dental or periodontal status of the SS patients (Table 1). The  $\text{NO}_2^-$  output was  $37.2 \pm 6.6$  nmoles/minute in healthy controls compared with  $166.3 \pm 46.2$  nmoles/minute in patients with SS ( $P < 0.01$ ). The salivary VIP concentration was  $68.6 \pm 24.2$  pg/ml in the patient group compared with  $40.8 \pm 23.6$  pg/ml in the healthy control group ( $P < 0.05$ ). However, there were no significant correlations between salivary  $\text{NO}_2^-$  and VIP concentrations or between salivary  $\text{NO}_2^-$  concentration and stress in the patients or in the controls.

**NADPH-diaphorase activity in LSGs in SS.** The most conspicuous finding was that, in the LSGs, intense NADPH-diaphorase activity was localized in myoepithelial cells of the glandular acini and less frequently in acinar end-piece cells (Figure 1A). Epithelial cells of intercalated, striated, and excretory ducts also showed intense NADPH-diaphorase activity (Figure 1A). It was particularly interesting that such epithelial-cell NADPH-diaphorase expression was also seen in the LSGs obtained from healthy controls. NADPH-diaphorase activity was found in some nerve trunks, fibers, and terminals located around the acini and salivary ducts. Such NADPH-diaphorase-containing nerve fibers and terminals were relatively evenly distributed, but seemed to be absent from lymphocyte-infiltrated areas (i.e., lymphocyte foci) in SS patients.

NADPH-diaphorase activity was less frequently localized in endothelial cells of arterioles, capillaries, or venules. NADPH-diaphorase-positive endothelial cells were, in particular, observed in lymphocyte infiltrates in SS patients (for more details, see below). NADPH-diaphorase activity was seen in only a few of the inflammatory cells of the characteristic focal lymphocyte-rich foci (for more details, see below).

**Neuronal cNOS in LSGs in SS.** Antibodies raised against the whole cNOS protein purified from rat brain and the synthetic peptide fragment 53 of rat brain cNOS showed a very sparse distribution of cNOS-containing nerve fibers, mainly localized around the acini and along the salivary ducts (Figure 1B). However, cNOS immunoreactive nerve fibers were not seen in

inflammatory cell foci in SS. In contrast, acetylcholinesterase histochemical and immunocytochemical localization of VIP-containing nerve fibers revealed a very dense distribution of cholinergic and VIPergic (parasympathetic) nerve fibers coiling around the acini and salivary ducts in noninflamed areas of LSG, although such fibers appeared to be mostly absent from the lymphocyte foci in SS (data not shown).


**Endothelial cNOS in LSGs in SS.** Monoclonal antibodies raised against the bovine aortic eNOS isoform revealed a very sparse endothelial cell eNOS immunoreactivity compared with immunoreactivity for CD31. However, it was of interest to note that in SS, CD31-positive endothelial cells in focal inflammatory cell infiltrates showed a strong immunoreactivity for eNOS (Figures 1C and D). Some ductal epithelial cells showed dense granular immunoreactivity and many acini were also immunoreactive for this antiserum (Figure 1E).

**Inducible NOS in LSGs in SS.** Antibodies raised against both the murine macrophage and human hepatocyte iNOS peptide sequences were used in the present study. The 2 antibodies gave somewhat different profiles of iNOS-immunoreactive cells in inflammatory LSG tissue. Murine macrophage iNOS antiserum showed immunoreactivity for acinar structures and some inflammatory cells, whereas human hepatocyte iNOS antibodies showed more frequent immunoreactivity for mononuclear inflammatory cells (Figure 1F) as well as for some ductal epithelial cells (data not shown).

In SS patients, granular  $\text{TNF}\alpha$  staining was seen in some acini, vascular endothelial cells, and, most notably, ductal epithelial cells. Immunostaining of serial sections for  $\text{TNF}\alpha$  and human hepatocyte iNOS peptide sequence showed colocalization of  $\text{TNF}\alpha$  with iNOS in ductal epithelial cells, but not in acinar or vascular endothelial cells (Figures 2A and B).

The chemical reaction of NO with superoxide generates peroxynitrate, which has been shown to cause nitration of protein tyrosine groups. Antibodies against nitrosylated tyrosine (nitrotyrosine) residues labeled some ductal epithelial cells, but mainly detected nitrotyrosine in periductal connective tissue. Periacinar connective tissue was also, in some cases, nitrotyrosine positive, but nitrotyrosine immunoreactivity could not be detected within the acinar cells (data not shown). Nitrotyrosine-staining controls were negative.

Cell counting showed human hepatocyte iNOS in some immigrant inflammatory cells in the lymphocyte-rich foci in patients with SS. Approximately  $16 \pm 3$  (mean  $\pm$  SEM) such cells were seen per  $1 \text{ mm}^2$  of




**Figure 2.** Immunoreactivity for human inducible nitric oxide synthase (iNOS) antibodies (A) (arrows) was found to colocalize with that for tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) (B) (arrows) in ductal epithelial cells in labial salivary glands from patients with Sjögren's syndrome. The 2 sections immunostained for iNOS and TNF $\alpha$  were taken 40  $\mu$ m apart, but several ducts can still be traced through the tissues. Note the absence of iNOS and TNF $\alpha$  immunoreactivity in acini. (Original magnification  $\times$  400.)

inflammatory cells, compared with  $21 \pm 2$  CD68+ macrophages. However, colocalization studies with double labeling suggested that there was only a minor overlap between these 2 cell subpopulations (data not shown), although some CD68-positive macrophages were also immunoreactive for human hepatocyte iNOS (Figures 3A and B).

### DISCUSSION

SS is characterized by a focal adenitis, autoimmunity, and sicca symptoms. In salivary glands, this is

expressed in the form of focal T lymphocyte-rich infiltrates and xerostomia. The simplistic and prevailing view has been that sialopenia and xerostomia are due to structural destruction of the secretory acinar cells, which are responsible for the secretion of the primary saliva. However, there is a clear discrepancy between the often focal and local salivary gland involvement and the greatly diminished salivary flow seen in SS. Significantly, many patients draw advantage from the use of sialogogues, such as xylitol gum or pilocarpine. Therefore, an alternative hypothesis has been that, before permanent



**Figure 3.** Indirect immunofluorescence technique was used for colocalization of inducible nitric oxide synthase (iNOS) immunoreactivity (A) (arrows) with CD68-positive macrophages (B) (arrows). In these photomicrographs, only 2 CD68-positive macrophages (B) also express iNOS immunoreactivity, although some additional inflammatory and/or resident cells were immunoreactive for the iNOS isoform (A). (Original magnification  $\times$  450.)

salivary gland parenchymal damage has occurred in a long-lasting and chronic disease, the focal perivascular and periductal inflammation alters the vasoneural regulation of the various vascular, motor, and secretory components of the normal salivary flow.

Using VIP and neuropeptide Y as markers for postganglionic parasympathetic and sympathetic nerve fibers, respectively, a clearly decreased innervation of the salivary glands in SS has been shown (28,29). In addition to adrenergic and cholinergic nerves, the non-adrenergic, noncholinergic system plays a role in normal and abnormal neuronal regulation of the salivary glands. An important part of the nonadrenergic, noncholinergic system is mediated by VIP, which, via its low-affinity binding sites, is linked to cAMP formation and, via its high-affinity binding sites, to  $\text{Ca}^{2+}$  mobilization and promotion of neuronal survival (30–32). More recently, NO has been measured in human saliva (33). It was also confirmed that NO is released from salivary glands rather than being produced from the activity of oral bacteria (33). Furthermore, there were no significant correlations between the salivary  $\text{NO}_2^-$  concentration and the periodontal status of the SS patients, which rules out any contribution of inflammatory gingival crevicular fluid to the salivary  $\text{NO}_2^-$ . This is consistent with recent studies on the presence of NADPH-diaphorase activity and NOS isoforms in feline submandibular glands (34) and with the current findings on the presence of NADPH-diaphorase activity and of all 3 different NOS isotypes in human salivary glands.

Formation of primary saliva involves vasodilation and acinar cell secretion. Changes in the regulatory vasoneural interactions may lead to sialopenia and dry mouth, such as is seen in psychogenic dry mouth in depression and anxiety or as an anticholinergic side effect of many widely used drugs. Changes in the L-arginine/NO-generating pathway may contribute to these crucial events. NO may influence parasympathetic vasodilation and salivary flow by regulating peptide release and second messenger systems for both VIP and acetylcholine, as has been shown in pigs and rats (35–37). As described in the present study, the elements of the L-arginine/NO-generating pathways are also found in human salivary glands and saliva.

There were some interesting differences between the SS patients and controls. The increase in  $\text{NO}_2^-$  seen in SS may have been related to an inflammation-induced increase in the expression of the iNOS rather than to the increased activity of the cNOS isoforms. The increased salivary  $\text{NO}_2^-$  concentration is not likely to be a reflection of the decreased “water” content of the saliva, because the salivary  $\text{NO}_2^-$  output was also significantly

higher in the SS patients than in the controls. Furthermore, a decrease in the salivary gland tissue mass, e.g., by replacement with inflammatory cell infiltrates, fibrosis, and fat atrophy, would lead to an apparently low salivary  $\text{NO}_2^-$  concentration, so that a 50% decrease in the mass would decrease the salivary  $\text{NO}_2^-$  concentration/output by half. Perhaps even more importantly, deranged architecture/function of the diseased glandular tissue would also contribute to an apparently low salivary  $\text{NO}_2^-$  concentration/output, so that a 50% decrease in the function would also decrease the salivary  $\text{NO}_2^-$  concentration/output by half. Therefore, the observed differences between patients and controls may reflect quite significant differences in the salivary gland tissues.

Studies in pigs have demonstrated that release of VIP from the salivary gland, elicited by parasympathetic stimulation (10 Hz, 30 seconds), was markedly reduced by a NOS inhibitor,  $N^G$ -nitro-L-arginine (36,37). Interestingly, in the present study, no significant correlations were found between the salivary  $\text{NO}_2^-$  and VIP concentrations, suggesting that this NO-mediated regulation of VIP release may not be the only system controlling VIP release from the peripheral VIPergic nerve fibers in human salivary glands.

In addition to the constitutively expressed and rapidly regulated endothelial and neuronal NOS, iNOS has been described in numerous cell types. However, the presence of iNOS in human macrophages is still a matter of controversy (38,39). The iNOS gene is located in chromosome 7, and its level of expression is regulated by various cytokines and lipopolysaccharides. In particular, certain cytokine combinations, such as interferon- $\gamma$  (IFN $\gamma$ ), interleukin-1 (IL-1), and TNF $\alpha$ , are very effective and synergistic in the up-regulation of iNOS gene expression (40,41). Inducible NOS produces high concentrations of NO for extended time periods, and, often, high cytotoxic and bactericidal concentrations of NO are produced.

Increased NO production hardly represents attempts to compensate for diminished cholinergic stimulation. It seems that the increased NO production is caused by local inflammatory changes and by cytokines, and, as such, appears to be an effect of inflammation rather than of a physiologic process intended to compensate for diminished cholinergic stimulation. Moreover, increased inflammatory NO production mediated by iNOS might not be able to compensate for the highly regulated neuronal events that are involved in the coordinated vascular, motor, and secretory components of the reflex salivary flow. Inflammatory NO production would be very unlikely to have that type of effect,

coordinated in time and at site, that is characteristic of a well-functioning normal neurovascular system of the salivary gland.

It was of particular interest to note that NADPH-diaphorase activity and, to a lesser extent, iNOS expression, were seen in the acinar and ductal epithelial cells of LSGs from SS patients, but also to some extent in those from healthy controls. In thyroid glands, NADPH-diaphorase activity and NOS immunoreactivity have been described in nerve fibers and vascular endothelium, but not in the glandular parenchymal cells (42). Lohinai and coworkers, in a study of feline submandibular glands, noticed that the epithelial layer of the salivary ductal branches was NOS immunonegative, but NADPH-diaphorase positive (34). Inducible NOS/NADPH-diaphorase in salivary glands might be induced by bacterial lipopolysaccharides (40). However, considering the cytokine-mediated regulation of iNOS, it seems more likely that in SS, iNOS immunoreactivity and NADPH-diaphorase activity are induced by cytokines, such as TNF $\alpha$ , IL-1, and IFN $\gamma$  (40,41), which are produced by the salivary gland epithelial cells and/or activated T lymphocytes (43–45).

One interesting feature of the minor salivary glands is the occurrence of focal accumulations of lymphocytes around the duct walls (46–48), which are believed to have a role in the immune surveillance of the mouth (48). Although the control LSGs were from subjects who did not have SS and who had a focus score (10) of <1, they did have occasional lymphocyte foci. These foci might produce enough cytokines (43–45) to up-regulate the iNOS immunoreactivity/NADPH-diaphorase activity even in normal minor salivary glands, although not to the same extent as in SS, as suggested by the present findings on salivary NO $_2^-$  concentration.

The salivary concentration of NO $_2^-$  may not quite reflect the extent of local NO production, because at least part of the locally produced NO could be disposed via pathways other than that leading to saliva. It is therefore possible that the increased, inflammation-associated, and probably cytokine-induced local NO production in SS interferes with the normal function of the cNOS isoforms and may, by itself, lead to cytopathic changes and acinar cell atrophy.

Thus, NO is produced in physiologic amounts in normal salivary glands mainly by constitutively expressed ncNOS and ecNOS isoforms, and may contribute to various events that regulate vascular reactivity, salivary flow, and neuropeptide release. In focal adenitis in SS, cytokine-induced iNOS is expressed by salivary gland acinar and ductal epithelial cells together with TNF $\alpha$ ,

suggesting a role for these resident cells in local pathomechanisms, which may contribute to various cytopathic and atrophic cell changes.

## REFERENCES

1. Moncada S, Higgs A: The L-arginine-nitric oxide pathway. *N Engl J Med* 329:2002–2012, 1993
2. Marletta MA: Nitric oxide synthase: aspects concerning structure and catalysis. *Cell* 23:927–930, 1994
3. Nathan C, Xie Q: Nitric oxide synthases: roles, tolls, and controls. *Cell* 78:915–918, 1994
4. Schmidt HH, Walter U: NO at work. *Cell* 78:919–925, 1994
5. Stamler JS: Redox signaling: nitrosylation and related target interactions of nitric oxide. *Cell* 78:931–936, 1994
6. Declaration of Helsinki, Hong Kong amendment, September 1989
7. Malmström MJ, Segerberg-Kontinen M, Tuominen TS, Hietanen JH, Wolf JE, Sane JI, Kontinen YT: Xerostomia due to Sjögren's syndrome: diagnostic criteria, treatment and outlines for a continuous dental care programme and an open trial with Sulfarlem®. *Scand J Rheumatol* 17:77–86, 1988
8. Törnwall J, Kontinen YT, Hietanen J, Sorsa T, Hukkanen M, Uusitalo H: VIP in salivary glands in Sjögren's syndrome. *Br J Rheumatol* 34:891–893, 1995
9. Kariologiska principer: Nordisk Lärobok i Kariologi. Fifth edition. Edited by Y Ericsson. Stockholm, Tandläkarförbundet, 1980
10. Lindhe J, editor: Textbook of Clinical Periodontology. Second edition. Copenhagen, Munksgaard, 1989
11. Ainamo J, Bay I: Problems and proposals for recording gingivitis and plaque. *Int Dent J* 25:229–235, 1975
12. Björby A, Löe H: The relative significance of different local factors in the initiation and development of periodontal inflammation. *J Periodontol Res* 1:76–77, 1967
13. Ainamo J, Barmes D, Beagrie G, Cutress T, Martin J, Sardo-Infirri J: Development of the World Health Organization (WHO) community periodontal index of treatment needs (CPITN). *Int Dent J* 32:281–291, 1982
14. Vitali C, Bombardieri S, Moutsopoulos HM, Balestrieri G, Ben-civelli W, Bernstein RM, Bjerrum KB, Braga S, Coll J, de Vita S, Drosos AA, Ehrenfeld M, Hatron PY, Hay EM, Isenberg DA, Janin A, Kalden JR, Kater L, Kontinen YT, Maddison PJ, Maini RN, Manthorpe R, Meyer O, Ostuni P, Pennec Y, Prause JU, Richards A, Sauvezie B, Schiødt M, Sciuto M, Scully C, Shoenfeld Y, Skopouli FN, Smolen JS, Snaith ML, Tishler M, Todesco S, Valesini G, Venables PJW, Wattiaux MJ, Youinou P: Preliminary criteria for the classification of Sjögren's syndrome: results of a prospective concerted action supported by the European Community. *Arthritis Rheum* 36:340–347, 1993
15. Greenspan JS, Daniels TE, Talal N, Sylvester RA: The histopathology of Sjögren's syndrome in labial salivary gland biopsies. *Oral Surg Oral Med Oral Pathol* 37:217–229, 1974
16. Jenkins CD, Zyzanski SJ, Rosenman RH: Progress toward validation of a computer-scored test for the type A coronary-prone behavior pattern. *Psychosom Med* 33:193–202, 1971
17. Snedecor GW, Cochran WG: Statistical Methods. Seventh edition. Ames, The Iowa State University Press, 1980
18. Ding AH, Nathan CF, Stuehr DJ: Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages: comparison of activation cytokines and evidence for independent production. *J Immunol* 141:2407–2412, 1988
19. Hope BT, Michael GJ, Knigge KM, Vincent SR: Neuronal NADPH diaphorase is a nitric oxide synthase. *Proc Natl Acad Sci U S A* 88:2811–2814, 1991
20. Wharton J, Polak JM, Gordon L, Banner NR, Springall DR, Rose



- M, Khagani A, Wallwork J, Yacoub MH: Immunohistochemical demonstration of human cardiac innervation before and after transplantation. *Circ Res* 66:900-912, 1990
21. Springall DR, Riveros-Moreno V, Buttery L, Suburo A, Bishop AE, Merrett M, Moncada S, Polak JM: Immunological detection of nitric oxide synthase(s) in human tissues using heterologous antibodies suggesting different isoforms. *Histochemistry* 98:259-266, 1992
  22. Pollock JS, Nakane M, Buttery LDK, Martinez A, Springall DR, Polak JM, Förstermann U, Murad F: Characterization and localization of endothelial nitric oxide synthase using specific monoclonal antibodies. *Am J Physiol* 265:C1379-C1387, 1993
  23. Hevel JM, White KA, Marletta MA: Purification of the inducible murine macrophage nitric oxide synthase: identification as a flavoprotein. *J Biol Chem* 266:22789-22791, 1991
  24. Hamid Q, Springall DR, Riveros-Moreno V, Chanez P, Howarth P, Redington A, Bousquet J, Godard P, Holgate S, Polak JM: Induction of nitric oxide synthase in asthma. *Lancet* 342:1510-1513, 1993
  25. Buttery LDK, Evans TJ, Springall DR, Carpenter A, Cohen J, Polak JM: Immunohistochemical localization of inducible nitric oxide synthase in endotoxin-treated rats. *Lab Invest* 71:755-764, 1994
  26. Buttery LDK, Springall DR, Chester AH, Evans TJ, Standfield N, Parums DV, Yacoub MH, Polak JM: Inducible NO synthase is present within human atherosclerotic lesions and promotes the formation and activity of peroxynitrate. *Lab Invest* 75:77-85, 1996
  27. Hsu SM, Raine L, Fanger H: Use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques: a comparison between ABC and unlabeled antibody (PAP) procedures. *J Histochem Cytochem* 29:577-580, 1981
  28. Konttinen YT, Hukkanen M, Kempainen P, Segerberg M, Sorsa T, Malmström M, Rose S, Itescu S, Polak JM: Peptide-containing nerves in labial salivary glands in Sjögren's syndrome. *Arthritis Rheum* 35:815-820, 1992
  29. Konttinen YT, Sorsa T, Hukkanen M, Segerberg M, Kuhlefelt-Sundström M, Malmström M, Polak JM: Topology of innervation of labial salivary glands by protein gene product 9.5 and synaptophysin immunoreactive nerves in patients with Sjögren's syndrome. *J Rheumatol* 19:30-37, 1992
  30. Brenneman DE, Eiden LE: Vasoactive intestinal peptide and electrical activity influence neuronal survival. *Proc Natl Acad Sci U S A* 83:1159-1162, 1986
  31. Fatatis A, Holtzclaw LA, Avidor R, Brenneman DE, Russell JT: Vasoactive intestinal peptide increases intracellular calcium in astroglia: synergism with  $\alpha$ -adrenergic receptors. *Proc Natl Acad Sci U S A* 91:2036-2040, 1994
  32. Konttinen YT, Kempainen P, Segerberg M, Hukkanen M, Rees R, Santavirta S, Sorsa T, Pertovaara A, Polak JM: Peripheral and spinal neural mechanisms in arthritis, with particular reference to treatment of inflammation and pain. *Arthritis Rheum* 37:965-982, 1994
  33. Bodis S, Haregewoin A: Evidence for the release and possible neural regulation of nitric oxide in human saliva. *Biochem Biophys Res Commun* 194:347-350, 1993
  34. Lohinai Z, Szekely AD, Soos L, Feher E: Distribution of nitric oxide synthase containing elements in the feline submandibular gland. *Neurosci Lett* 192:9-12, 1995
  35. Damas J: Pilocarpine-induced salivary secretion, kinin system and nitric oxide in rats. *Arch Int Physiol Biochim Biophys* 102:103-105, 1994
  36. Modin A: Non-adrenergic, non-cholinergic vascular control with reference to neuropeptide Y, vasoactive intestinal polypeptide and nitric oxide. *Acta Physiol Scand* 622 (Suppl):1-74, 1994
  37. Modin A, Weitzberg E, Lundberg JM: Nitric oxide regulates peptide release from parasympathetic nerves and vascular reactivity to vasoactive intestinal polypeptide in vivo. *Eur J Pharmacol* 261:185-197, 1994
  38. Schneemann M, Schoedon G, Hofer S, Blau N, Guerrero L, Schaffner A: Nitric oxide synthase is not a constituent of the antimicrobial armature of human mononuclear phagocytes. *J Infect Dis* 167:1358-1363, 1993
  39. Padgett EL, Pruett SB: Evaluation of nitrite production by human macrophages. *Biochem Biophys Res Commun* 186:775-781, 1992
  40. Curran RD, Billiar TR, Stuehr DJ, Ochoa JB, Harbrecht BG, Flint SG, Simmons RL: Multiple cytokines are required to induce hepatocyte nitric oxide production and inhibit total protein synthesis. *Ann Surg* 212:462-471, 1990
  41. Liew FY, Li Y, Millott S: Tumor necrosis factor-alpha synergizes with IFN-gamma in mediating killing of *Leishmania major* through the induction of nitric oxide. *J Immunol* 145:4306-4310, 1990
  42. Syed MA, Leong SK, Chan AS: Localization of NADPH-diaphorase reactivity in the chick and mouse thyroid gland. *Thyroid* 4:475-478, 1994
  43. Fox RI, Kang HI, Ando D, Abrams J, Pisa E: Cytokine mRNA expression in salivary gland biopsies of Sjögren's syndrome. *J Immunol* 152:5532-5539, 1994
  44. Ogawa N, Dang H, Talal N: PCR analysis of cytokines produced in salivary glands of Sjögren's syndrome patients. In: *Sjögren's Syndrome: State of the Art*. Edited by M Homma, S Sugai, T Tojo, N Miyasaka, M Akizuki. Amsterdam, Kugler Publications, 1994
  45. Boumba D, Skopouli FN, Moutsopoulos HM: Cytokine mRNA expression in the labial salivary gland tissues from patients with primary Sjögren's syndrome. *Br J Rheumatol* 34:326-333, 1995
  46. Segerberg-Konttinen M, Konttinen YT, Bergroth V: Focus score in the diagnosis of Sjögren's syndrome. *Scand J Rheumatol Suppl* 61:47-51, 1986
  47. Segerberg-Konttinen M: Focal adenitis in lacrimal and salivary glands: a post-mortem study. *Scand J Rheumatol* 17:379-385, 1988
  48. Dale AC: Salivary glands. In: *Oral Histology: Development, Structure, and Function*. Fourth edition. Edited by AR Ten Cate. St. Louis, Mosby, 1994