

Fluid Flow Induced Calcium Response in Bone Cell Network

BO HUO,^{1,2} XIN L. LU,¹ CLARK T. HUNG,³ KEVIN D. COSTA,⁴ QIAOBING XU,⁵
GEORGE M. WHITESIDES,⁵ and X. EDWARD GUO¹

¹Bone Bioengineering Laboratory, Department of Biomedical Engineering, Columbia University, 351 Engineering Terrace, 1210 Amsterdam Avenue, Mail Code 8904, New York, NY 10027, USA; ²Institute of Mechanics, Chinese Academy of Sciences, Beijing 100080, P.R. China; ³Cellular Engineering Laboratory, Department of Biomedical Engineering, Columbia University, New York, NY 10027, USA; ⁴Cardiac Cell Mechanics Laboratory, Department of Biomedical Engineering, Columbia University, New York, NY 10027, USA; and ⁵Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA

(Received 15 January 2008; accepted 19 February 2008; published online 14 March 2008)

Abstract—In our previous work, bone cell networks with controlled spacing and functional intercellular gap junctions had been successfully established by using microcontact printing and self assembled monolayers technologies [Guo, X. E., E. Takai, X. Jiang, Q. Xu, G. M. Whitesides, J. T. Yardley, C. T. Hung, E. M. Chow, T. Hantschel, and K. D. Costa. *Mol. Cell. Biomech.* 3:95–107, 2006]. The present study investigated the calcium response and the underlying signaling pathways in patterned bone cell networks exposed to a steady fluid flow. The glass slides with cell networks were separated into eight groups for treatment with specific pharmacological agents that inhibit pathways significant in bone cell calcium signaling. The calcium transients of the network were recorded and quantitatively evaluated with a set of network parameters. The results showed that 18 α -GA (gap junction blocker), suramin (ATP inhibitor), and thapsigargin (depleting intracellular calcium stores) significantly reduced the occurrence of multiple calcium peaks, which were visually obvious in the untreated group. The number of responsive peaks also decreased slightly yet significantly when either the COX-2/PGE₂ or the NOS/nitric oxide pathway was disrupted. Different from all other groups, cells treated with 18 α -GA maintained a high concentration of intracellular calcium following the first peak. In the absence of calcium in the culture medium, the intracellular calcium concentration decreased slowly with fluid flow without any calcium transients observed. These findings have identified important factors in the flow mediated calcium signaling of bone cells within a patterned network.

Keywords—Bone cell, Fluid shear, Gap junction, PGE₂, Nitric oxide, ATP, Calcium store.

INTRODUCTION

Osteocytes are interconnected through numerous intercellular processes, forming extensive cell networks throughout the bone tissue.^{6,36,39} It has been shown that osteocyte density is an important physiological parameter, which decreases with age and microdamage accumulation.³⁴ However, most previous studies on mechanotransduction in osteocytes and osteoblasts were performed on confluent or sub-confluent uncontrolled monolayers of cells.^{38,40} In order to examine the roles of the cellular connectivity between bone cells in mechanotransduction, a controlled bone cell network is necessary for quantitative studies. In our previous study,¹¹ a two-dimensional patterned bone cell network was successfully established to mimic the *in vivo* osteocyte network by using microcontact printing and self assembled monolayers (SAM) technologies. Each individual bone cell in the network was connected with four neighboring cells via functional gap junctions. Under applied single cell nano-indentation with an atomic force microscope probe, calcium wave propagation within the bone cell network was observed. It was also demonstrated that multiple calcium peaks observed within an individual bone cell resulted from the self-perpetuating signal propagation within the circuit of the bone cell network. One of the major objectives of the present study was to examine the spatiotemporal characteristics of the intracellular calcium response in a well-controlled bone cell network under a steady fluid flow, a more physiologically relevant stimulation mode than the nano-indentation on a single bone cell in the network.

Calcium signaling is essential for numerous bone cell functions, such as proliferation and differentiation.^{3,15,29,41,42} It has been demonstrated that mechanical stimuli such as fluid flow can induce a

Address correspondence to X. Edward Guo, Bone Bioengineering Laboratory, Department of Biomedical Engineering, Columbia University, 351 Engineering Terrace, 1210 Amsterdam Avenue, Mail Code 8904, New York, NY 10027, USA. Electronic mail: ed.guo@columbia.edu

consistent and robust intracellular calcium ($[Ca^{2+}]_i$) response and intercellular calcium signaling in bone cells.^{21,28} The fluid flow induced elevation of cytosolic calcium comes mainly from two sources: intracellular stores (e.g., endoplasmic reticulum, ER) and the extracellular environment.^{4,14} $[Ca^{2+}]_i$ signaling is mediated by various molecular pathways, e.g., inositol triphosphate (IP_3), adenosine 5'-triphosphate (ATP), prostaglandin E_2 (PGE_2), and nitric oxide (NO) (Fig. 1). IP_3 can lead to a rapid release of calcium stored in the ER via binding to the ER membrane receptor.^{3,15} After the cytosolic calcium concentration is elevated to a critical level by intra/extracellular sources, the depleted intracellular calcium stores tend to recover their calcium reservation to original level and become ready for a next release of calcium.^{3,24} When a bone cell is under fluid flow induced shear stress, the activation of gap junction hemichannels (connexin 43) induces ATP efflux from the cytosol to the pericellular environment.^{8,9} Extracellular ATP can elicit a significant $[Ca^{2+}]_i$ response by binding with the P_2 purinergic membrane receptor.¹³ Fluid shear stress can also prompt the induction of COX-2 protein and further PGE_2 release in MC3T3-E1 osteoblast-like cells.²⁵ It was previously reported that fluid shear stress can elicit nitric oxide production in osteocytes and osteoblasts accompanied by an increased expression of nitric oxide synthase (NOS).^{23,40} Nitric oxide modulates $[Ca^{2+}]_i$ signaling via a cyclic guanosine monophosphate (cGMP) dependent pathway²² or nitrosylation of proteins.³⁵ Moreover, nitric oxide could directly contribute to the $[Ca^{2+}]_i$ release via triggering of an influx pathway that is, in part, responsible for the refilling of internal calcium stores.²⁰

Gap junctions are membrane-spanning channels, where each pair of connexons (i.e. hemichannels) forms a cylinder with a pore in the center through which small molecules (<1 kDa) can pass from one cell to another cell.³¹ It is widely accepted that this intercellular connection plays a significant role in coordinating bone cell network activities. Messenger molecules mentioned previously, such as IP_3 and calcium, may be able to transfer between the neighboring cells through gap junctions and thereby mediate propagation of $[Ca^{2+}]_i$ signaling in bone cells.^{17,38} The quantitative description of $[Ca^{2+}]_i$ signaling within a controlled bone cell network connected by gap junctions under fluid shear has yet to be obtained experimentally.

In the present study, using microcontact printing and SAM technologies developed previously,¹¹ bone cell networks with controlled spacing and functional intercellular gap junctions were built with MC3T3-E1 osteoblast-like bone cells. The cell networks were divided into 8 groups to test the effects of treatment with a battery of pharmacological agents to interrupt or inhibit $[Ca^{2+}]_i$ signaling. The $[Ca^{2+}]_i$ responses of the networks under steady fluid flow were recorded with fluorescence images, and the spatiotemporal characteristics of the $[Ca^{2+}]_i$ responses were quantitatively characterized with a set of network parameters, such as number of calcium spikes, time to reach first peak, and time to 50% relaxation. The inter/intra cellular calcium signaling mechanisms and the roles of different molecular pathways in $[Ca^{2+}]_i$ signaling were further analyzed.

MATERIALS AND METHODS

Bone Cell Network

Microcontact printing and SAM surface chemistry technologies were used to prepare the bone cell networks as described previously.¹¹ In brief, a chromium mask containing the desired square grid of connected circles was fabricated. The pattern was replicated to a master made of Shipley 1818 positive photoresist (MicroChem Corp, Newton, MA) by exposing it to UV light through the mask. The geometric parameters of the pattern adopted here ($2\ \mu\text{m}$ width connection line, $20\ \mu\text{m}$ diameter circle, and $70\ \mu\text{m}$ separation distance) were optimized for osteoblast-like cells.¹¹ A polydimethylsiloxane (PDMS, *Sylgard 184*, Dow Corning, Midland, MI) stamp with the desired micro-pattern was then generated by curing PDMS elastomer on the master. On the day before the fluid flow experiment, the stamp was coated with an adhesive SAM (octadecanethiol; Sigma-Aldrich Co., St. Louis,




FIGURE 1. A schematic drawing of calcium signaling pathways in bone cells and corresponding inhibitors employed in the current study. Red arrow: influx or upregulation activity, green arrow: efflux from cytosol, dash arrow: inhibition or blocking with a specific inhibitory intervention noted and underlined.




FIGURE 2. Pseudo-color images of typical intracellular calcium responses of a bone cell network from untreated group. Fluid flow was applied at 60 s. Arrowheads highlight those cells responding. Some cells were able to respond multiple times.

MO), and pressed onto a gold-coated glass slide (150 Å Au using an e-beam evaporator, SC2000, SEMICORE Inc., Livermore, CA) for 60 s. The stamped glass slide was immediately immersed in a non-adhesive ethylene glycol terminated SAM solution (HS-C11-EG3; Prochimia, Sopot, Poland) for 3 h to cover areas that were not patterned with the adhesive SAM. To facilitate cell adhesion, the patterned slide was further incubated in 10 $\mu\text{g}/\text{mL}$ fibronectin (Sigma–Aldrich Co., St. Louis, MO) in PBS for 1 h, which can only be absorbed by the adhesive SAM coated region. Osteoblast-like MC3T3-E1 cells with controlled density (1.0×10^4 cell/ cm^2) were seeded on the patterned slide and cultured in α -MEM medium supplemented with 2% charcoal-stripped fetal bovine serum (CS-FBS, Hyclone Laboratories Inc., Logan, UT) for 24 h before further testing. CS-FBS was used here to control the growth of cells and to avoid double-cell-occupancy on a single circular spot on the pattern. Figure 2 shows the snapshots of a typical bone cell network in the untreated group with a 70 μm separation distance.

Steady Fluid Flow and $[\text{Ca}^{2+}]_i$ Response

To indicate $[\text{Ca}^{2+}]_i$, patterned bone cells on the glass slide were loaded with 5 μM Fluo-4 AM (Molecular Probes, Eugene, OR) in α -MEM medium supplemented with 0.5% CS-FBS for 1 h at 37 $^\circ\text{C}$. After calcium dye loading, the slide was rinsed with

dye-free medium twice and then mounted on a custom-built parallel plate flow chamber, which was attached to an inverted fluorescence microscope (Olympus IX71, Melville, NY) with a 10 \times objective.¹⁴ The entire operation was performed with extreme care to minimize the response of cells to early agitations. Furthermore, a 20-min resting period preceded fluid flow stimulation, which has been shown to be sufficient for bone cells to recover and to generate repetitive $[\text{Ca}^{2+}]_i$ responses.¹⁰ A magnetic gear pump was connected to the chamber to generate a steady fluid flow with a constant shear stress of 40 dyne/ cm^2 on the cell surface. The $[\text{Ca}^{2+}]_i$ response of the bone cell network under fluid flow stimulation was recorded with a high-speed CCD camera (ORCA-ER-1394, Hamamatsu Photonics K.K., Hamamatsu City, Japan) for a total 10-min period, 1 min for baseline and 9 min after the onset of fluid flow. The intensity of $[\text{Ca}^{2+}]_i$ for each cell was later normalized by its corresponding baseline.

Experimental Groups

Besides the untreated group (12 slides, 130 cells analyzed), seven specific biochemical blocking agents were employed to identify the sources and the contributions of molecular pathways to the network characteristics of $[\text{Ca}^{2+}]_i$ response. (1) *Extracellular calcium depletion*: In this group, calcium-free Dulbecco's Modified Eagle Medium (DMEM, Invitrogen Corporation, Carlsbad, CA) and calcium-free Hank's Balanced Salt Solution (HBSS, Invitrogen Corporation) were introduced to replace the regular medium after Fluo-4 AM loading (8 slides, 99 cells analyzed). (2) *ER calcium store depletion*: After Fluo-4 staining, the cell-seeded slides (6 slides, 90 cells analyzed) were incubated in 1 μM Thapsigargin (TG) (Sigma–Aldrich Co.) medium for 30 min prior to flow.¹³ TG depletes calcium from the ER store. (3) *PGE₂ blocking*: The patterned cells were pretreated with 10 μM *N*-(2-cyclohexyloxy-4-nitrophenyl)methanesulfonamide (NS-398, EMD Chemicals Inc., San Diego, CA) for 24 h before flow study. NS-398 selectively inhibits the COX-2 enzyme activity, which in turn blocks the PGE₂ release.²⁵ One hundred and twenty-three cells on 11 slides were analyzed in this group. (4) *NO blocking*: In this group, 100 μM NG-monomethyl-L-arginine (L-NMMA; EMD Chemicals Inc.) was introduced into the cell culture medium 1 day before seeding the cells on slides and continuously presented in mediums afterward. L-NMMA inhibits the production of nitric oxide via competitively inhibiting all three isoforms of nitric oxide synthase (NOS).²⁷ In this group, 189 cells on 16 slides were analyzed. (5) *Gap junction blocking*: To study the role of intercellular gap junctions on calcium responses of the bone cell network, 75 μM

18 α -Glycyrrhetic acid (18 α -GA, Sigma–Aldrich Co.), a reversible gap junction blocker which binds to membrane proteins and causes disassembly of gap junction plaques,¹² was supplied in Fluo-4 AM loading medium and fluid flow medium.³⁸ In this group, 61 cells on 6 slides were studied. (6) *ATP blocking*: Suramin (EMD Chemicals Inc.), a P2 purinergic receptor blocker, was employed to investigate the effect of ATP release on the $[Ca^{2+}]_i$ response. 100 μ M suramin was applied to the cells 30 min before exposing to shear flow.³⁷ Fifty cells from 3 slides were analyzed. (7) *Vehicle DMSO control*: Among all the previously employed chemicals, NS-398, 18 α -GA, and TG were dissolved into DMSO. Therefore vehicle control testing was performed with 0.3% (v/v) DMSO presented in medium (3 slides, 51 cells analyzed). The concentrations of above agents were chosen from the relevant literature in bone cells where they have been demonstrated to be effective in inhibiting their respective pathways.

Data Analysis

To quantitatively analyze the network characteristics of the $[Ca^{2+}]_i$ responses, a set of network parameters were defined from the time-lapse of $[Ca^{2+}]_i$ response (Fig. 3). For each individual cell, the total number of $[Ca^{2+}]_i$ responsive peaks during the flow stimulation period was counted. t_1 denoted the time between the onset point of fluid flow and the maximum value of the first responsive spike, while t_2 was the time




FIGURE 3. A typical calcium response of bone cells in an untreated network under applied steady fluid flow. The characteristic temporal parameters are also illustrated. t_1 denotes the time from flow onset to the maximum value of the first responsive peak; t_2 is the time from peak value to the 50% relaxation; t_3 , t_4 , and t_5 are the time intervals between different peaks. m_1 represents the normalized magnitude of the first peak.

from first peak to its 50% relaxation point. t_3 , t_4 , and t_5 were defined as the time intervals in between successive peaks, respectively, and m_1 is the normalized magnitude of the first peak. One-way analysis of variance (ANOVA) with Bonferroni's *post hoc* analysis was performed to determine statistical differences between mean values of different treatments, $p < 0.05$. The condition of equal population variances of treatments was tested by Levene's method.

RESULTS

A typical set of intracellular calcium images during the testing period are shown in Fig. 2, clearly demonstrating multiple responses of bone cells within the network under fluid flow. A set of typical $[Ca^{2+}]_i$ traces of individual cells from the untreated group and those with different treatments are shown in Fig. 4. In each group, $[Ca^{2+}]_i$ profiles of three cells from a single glass slide were plotted. For untreated, vehicle control, NS-398 (PGE₂ blocking), and L-NMMA (NO blocking) groups, multiple $[Ca^{2+}]_i$ responsive peaks were quite obvious even by visual inspection. However, 18 α -GA (gap junction blocking) and TG (ER calcium depletion) groups tended to exhibit only a single $[Ca^{2+}]_i$ peak. In the 18 α -GA group, it was interesting to note that the $[Ca^{2+}]_i$ intensity was sustained at a relatively high level after the first calcium peak. In the absence of extracellular calcium, unlike the $[Ca^{2+}]_i$ response in any other group, $[Ca^{2+}]_i$ decreased slowly after the onset of fluid flow without any $[Ca^{2+}]_i$ peak.

To quantitatively and statistically analyze the calcium waves, the frequencies of multiple responding peaks are shown in Fig. 5a, while the mean value of the number of responding peaks are plotted in Fig. 5b. In the untreated group, over 55% of cells were able to respond three or more times with an overall mean value as 2.89 ± 0.16 times of baseline. In the vehicle control group, this value was 2.33 ± 0.19 but no significant difference was detected when compared with the untreated group. Both NS-398 (PGE₂ blocking) and L-NMMA (NO blocking) resulted in fewer multiple responses, only 50% of cells exhibited two or more calcium transients. Furthermore, 15 and 22% of cells had no response to fluid flow in these two groups, respectively, which was significantly higher than those of the untreated or vehicle control group. 18 α -GA (gap junction blocking) and suramin (ATP blocking) significantly reduced the number of calcium peaks, while cells with a single response comprised approximately 60% of the total population. ANOVA testing revealed no significant difference between the mean values of these two groups, but they were both significantly different from untreated and vehicle control groups.




FIGURE 4. A set of typical intracellular calcium traces of individual cells with or without pharmacological blocker treatment. The calcium images were recorded for 10 min (1 min for baseline and 9 min for flow stimulation). The intensity of intracellular calcium concentration shown here was normalized to the baseline.

The TG-treated group (ER depleted) generated the least number of responding calcium peaks, which was significantly different from any other group. The removal of extracellular Ca^{2+} completely abolished the $[\text{Ca}^{2+}]_i$ response.

As shown in Fig. 6, cells in the gap junction blocked group took the longest time to reach the first $[\text{Ca}^{2+}]_i$ peak ($t_1 = 61 \pm 7$ s). Moreover, the relaxation time of this group was nearly 5 times that of all other

groups ($t_2 = 222 \pm 23$ s). The untreated and TG groups shared a similar t_1 value (40 ± 4 vs. 34 ± 2 s, $p > 0.05$). The vehicle control, NS-398, L-NMMA, and suramin groups all had significantly shorter first peak response times when compared with the untreated group, but there was no significant difference among these four groups. Aside from the 18 α -GA group, all the other six groups had no significant difference in the 50% relaxation time. The 18 α -GA




FIGURE 5. (a) The frequency distribution of the number of responsive peaks of cells in different groups and (b) the average of the number of responsive peaks (the error bars are standard errors of the means (SEM) while * and ** indicate statistical significance of $p < 0.05$). The group of Ca^{2+} free medium had no response to fluid flow and was not included.

group was further characterized with the highest first peak magnitude ($m_1 = 267 \pm 22\%$ of baseline). The NS-398 and L-NMMA groups also had slightly yet significantly higher m_1 values than those of the other three groups. When comparing the time intervals between calcium peaks, i.e. t_3 , t_4 , and t_5 , it was found that only the 18 α -GA group had significantly higher t_3 value than the untreated or L-NMMA group. The mean values of these time intervals fell within the range of 100–200 s.

DISCUSSION

The network characteristics of $[\text{Ca}^{2+}]_i$ response of bone cells in a controlled cell network have been quantified, and several biochemical pathways related

to $[\text{Ca}^{2+}]_i$ signaling have been examined. Notably, multiple $[\text{Ca}^{2+}]_i$ peaks, a signature of network circuits, have been observed in bone cell networks under applied steady fluid flow. Intracellular calcium signaling in bone cells involves both internal and external sources of calcium.³ The internal calcium stores are mainly held within the membrane systems of the ER.^{2,5} Current experimental results from calcium-free media and ER depletion groups have identified the calcium source underlying the $[\text{Ca}^{2+}]_i$ responses in the bone cell networks. The cells in calcium-free media showed absolutely no $[\text{Ca}^{2+}]_i$ increase due to fluid flow, indicating the critical role of extracellular calcium. A few researchers have noted that the influx of extracellular calcium is required to evoke the opening of the calcium release channels in the ER.^{18,24} In fact, it has been shown that the gap junction dependent propagation of intercellular calcium signals in rat osteoblast cells requires influx of external calcium through L-type calcium channels.¹⁸ Moreover, the cytosolic calcium concentration was significantly reduced in the absence of extracellular calcium, which was consistent with a previous report.¹³ On the other hand when calcium was depleted from ER stores, more than 60% of cells exhibited a calcium peak under the stimulation of fluid flow, similar with those observed in randomly cultured 80–90% confluent MC3T3-E1 cells under an oscillatory fluid flow.¹ The magnitude of the first responsive peak was not significantly altered. This confirmed that extracellular calcium influx was responsible for the initial calcium peak. However, the average number of responsive $[\text{Ca}^{2+}]_i$ peaks in the cell network was significantly reduced to barely more than one. Thus ER stores appears to be the major calcium source responsible for the lower-magnitude $[\text{Ca}^{2+}]_i$ peaks following the initial calcium transient in the network of bone cells.^{1,24}

The $[\text{Ca}^{2+}]_i$ responses of the networked bone cells consist of a large initial peak followed subsequently by multiple lower-magnitude peaks, which is similar to those observed in an 80–90% confluent bone cell-culture study.⁷ In the untreated group, over 50% of the cells were able to generate three or more calcium peaks within 9 min. In the ER calcium depleted group, the ATP activation blocked group, and the gap junction blocked group, the average number of calcium peaks during 9 min was significantly reduced, and the number for the first two groups was very close to one (TG and suramin treated, respectively). The importance of ER calcium stores in the multiple responsive $[\text{Ca}^{2+}]_i$ peaks^{1,24} has been confirmed in the current networked bone cells. This new bone cell network study suggests the critical importance of the ATP pathway in mediating multiple $[\text{Ca}^{2+}]_i$ peaks of bone cells under steady fluid flow. This observation is consistent with literature




FIGURE 6. Network characteristic parameters of calcium response in the bone cell network under steady fluid flow from different treatment groups (the error bars are SEM while * and ** indicate statistical significance of $p < 0.05$): (a) time to the first peak t_1 ; (b) time to 50% relaxation after the first peak t_2 ; (c) magnitude of the first peak m_1 ; and (d) time between peaks t_3 , t_4 , and t_5 . The group of Ca^{2+} free medium had no response to fluid flow and was not included.

where calcium wave propagation in bone cells induced by cell poking was influenced by the ATP pathway.^{16–18} We also noted that 18 α -GA (gap junction blocking) did not completely extinguish the multiple transients of bone cells, as 32% of cells responded more than once. This implied that a different pathway other than gap junction signal communication also plays a significant role in the multiple calcium peaks in the bone cell network. Indeed, inhibiting COX-2/PGE₂ or nitric oxide pathways also slightly reduced the multiple calcium peaks of individual osteoblasts, which was similar to previous findings in other cell types.^{19,22,32,35} A previous study by Li *et al.* has shown that blocking the nitric oxide pathway can significantly reduce the refilling capability of intracellular calcium store,²⁰ thereby decreasing the occurrence of multiple transients.

Calcium signaling in bone cells can be propagated and amplified via intercellular gap junctions.^{11,16–18,30} An interesting result from the gap-junction blocked group is the observation of a sustained high $[\text{Ca}^{2+}]_i$ evoked by the steady fluid flow following the initial calcium peak (Fig. 4). The exact mechanism for this calcium plateau will require further investigation. In bone cells, once the $[\text{Ca}^{2+}]_i$ is elevated rapidly by both extracellular entry and ER release, the high concentration of intracellular calcium activates various plasma membrane calcium pumps to enhance the

extrusion rate to terminate this toxic-high $[\text{Ca}^{2+}]_i$ level.²⁶ It seems that the 18 α -GA dramatically impeded or retarded the function of these calcium pumps. For a large fraction of the cells in this group, the $[\text{Ca}^{2+}]_i$ never recovered to the initial resting level during the 9-min loading duration. Cells in this group showed the lowest frequency of responders with two or more calcium spikes, and the cells that exhibited multiple transients were characterized by a significantly longer time between the first and the second spikes (t_3 value). Qualitatively, the intercellular propagation of a $[\text{Ca}^{2+}]_i$ transient from within a bone cell to its neighboring cells in the network was observed in untreated group, and this was blocked by 18 α -GA. Therefore, the elevated $[\text{Ca}^{2+}]_i$ could be partially attributable to the retention of $[\text{Ca}^{2+}]_i$.

In summary, a thorough investigation of the spatiotemporal characteristics of $[\text{Ca}^{2+}]_i$ signaling of bone cells in a controlled cell network under steady fluid flow was conducted. The effects of several candidate pathways potentially involved in inter/intra cellular calcium signaling were also studied. We confirmed that the major source for the rapid initial calcium transient under mechanical stimulation was attributable to the influx of extracellular calcium. The ER calcium stores and ATP pathways are responsible for the subsequent multiple calcium peaks. The intercellular gap junctions

in the cell network demonstrated a significant influence on the $[Ca^{2+}]_i$ response. Bone cells in a gap junction-blocked network exhibited a sustained high calcium concentration after the triggering from fluid flow. It was also shown that the PGE₂ and NO pathways were both involved to a lesser extent in the calcium signaling of the bone cell network.

For future studies, it would be interesting to repeat similar experiments on an osteocyte network. It will also be important to re-examine the calcium propagation waves in a bone cell network subjected to oscillatory fluid flow, or steady flow with magnitudes closer to physiological values (5–32 dynes/cm² instead of 40 dynes/cm²). A relatively high magnitude of shear stress was used in the current study to evoke more responding cells and more multiple calcium peaks. With the techniques presented in this study, it is also possible to examine the intercellular signaling between osteocytes and osteoblasts in a controlled cell network, or to build a 3D microfluidic system to mimic the *in vivo* fluid flow stimulation on osteocytes and their processes.³³ To better understand the exact roles of individual pathways on calcium signaling, techniques that employ greater specificity such as the use of primary cells from gene knockout mice can be incorporated. Currently a theoretical model to predict the calcium responses in a gap junction connected/blocked cell network is under development in our laboratory.

ACKNOWLEDGMENTS

This work was supported by NIH grant R21 AR052417 (X. Edward Guo). The authors wish to thank Mr. Andrew D. Baik for editing of this paper.

REFERENCES

- ¹Batra, N. N., Y. J. Li, C. E. Yellowley, L. You, A. M. Malone, C. H. Kim, and C. R. Jacobs. Effects of short-term recovery periods on fluid-induced signaling in osteoblastic cells. *J. Biomech.* 38:1909–1917, 2005.
- ²Berridge, M. J. Inositol trisphosphate and calcium signaling. *Nature* 361:315–325, 1993.
- ³Berridge, M. J., P. Lipp, and M. D. Bootman. The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* 1:11–21, 2000.
- ⁴Chen, N. X., K. D. Ryder, F. M. Pavalko, C. H. Turner, D. B. Burr, J. Qiu, and R. L. Duncan. Ca²⁺ regulates fluid shear-induced cytoskeletal reorganization and gene expression in osteoblasts. *Am. J. Physiol. Cell Physiol.* 278:C989–997, 2000.
- ⁵Clapham, D. E. Calcium signaling. *Cell* 80:259–268, 1995.
- ⁶Cowin, S. C. On mechanosensation in bone under microgravity. *Bone* 22:119S–125S, 1998.
- ⁷Donahue, S. W., H. J. Donahue, and C. R. Jacobs. Osteoblastic cells have refractory periods for fluid-flow-induced intracellular calcium oscillations for short bouts of flow and display multiple low-magnitude oscillations during long-term flow. *J. Biomech.* 36:35–43, 2003.
- ⁸Genetos, D. C., D. J. Geist, D. Liu, H. J. Donahue, and R. L. Duncan. Fluid shear-induced ATP secretion mediates prostaglandin release in MC3T3-E1 osteoblasts. *J. Bone Miner. Res.* 20:41–49, 2005.
- ⁹Genetos, D. C., C. J. Kephart, Y. Zhang, C. E. Yellowley, and H. J. Donahue. Oscillating fluid flow activation of gap junction hemichannels induces ATP release from MLO-Y4 osteocytes. *J. Cell Physiol.* 212:207–214, 2007.
- ¹⁰Godin, L. M., S. Suzuki, C. R. Jacobs, H. J. Donahue, and S. W. Donahue. Mechanically induced intracellular calcium waves in osteoblasts demonstrate calcium fingerprints in bone cell mechanotransduction. *Biomech. Model. Mechanobiol.* 6:391–398, 2007.
- ¹¹Guo, X. E., E. Takai, X. Jiang, Q. Xu, G. M. Whitesides, J. T. Yardley, C. T. Hung, E. M. Chow, T. Hantschel, and K. D. Costa. Intracellular calcium waves in bone cell networks under single cell nanoindentation. *Mol. Cell. Biomech.* 3:95–107, 2006.
- ¹²Guo, Y., C. Martinez-Williams, K. A. Gilbert, and D. E. Rannels. Inhibition of gap junction communication in alveolar epithelial cells by 18alpha-glycyrrhetic acid. *Am. J. Physiol.* 276:L1018–1026, 1999.
- ¹³Hung, C. T., F. D. Allen, K. D. Mansfield, and I. M. Shapiro. Extracellular ATP modulates $[Ca^{2+}]_i$ in retinoic acid-treated embryonic chondrocytes. *Am. J. Physiol.* 272:C1611–1617, 1997.
- ¹⁴Hung, C. T., F. D. Allen, S. R. Pollack, and C. T. Brighton. Intracellular Ca²⁺ stores and extracellular Ca²⁺ are required in the real-time Ca²⁺ response of bone cells experiencing fluid flow. *J. Biomech.* 29:1411–1417, 1996.
- ¹⁵Iqbal, J., and M. Zaidi. Molecular regulation of mechanotransduction. *Biochem. Biophys. Res. Commun.* 328:751–755, 2005.
- ¹⁶Jorgensen, N. R., S. T. Geist, R. Civitelli, and T. H. Steinberg. ATP- and gap junction-dependent intercellular calcium signaling in osteoblastic cells. *J. Cell Biol.* 139:497–506, 1997.
- ¹⁷Jorgensen, N. R., Z. Henriksen, C. Brot, E. F. Eriksen, O. H. Sorensen, R. Civitelli, and T. H. Steinberg. Human osteoblastic cells propagate intercellular calcium signals by two different mechanisms. *J. Bone Miner. Res.* 15:1024–1032, 2000.
- ¹⁸Jorgensen, N. R., S. C. Teilmann, Z. Henriksen, R. Civitelli, O. H. Sorensen, and T. H. Steinberg. Activation of L-type calcium channels is required for gap junction-mediated intercellular calcium signaling in osteoblastic cells. *J. Biol. Chem.* 278:4082–4086, 2003.
- ¹⁹Krieger, N. S., K. K. Frick, K. LaPlante Strutz, A. Michalenka, and D. A. Bushinsky. Regulation of COX-2 mediates acid-induced bone calcium efflux *in vitro*. *J. Bone Miner. Res.* 22:907–917, 2007.
- ²⁰Li, N., J. Y. Sul, and P. G. Haydon. A calcium-induced calcium influx factor, nitric oxide, modulates the refilling of calcium stores in astrocytes. *J. Neurosci.* 23:10302–10310, 2003.
- ²¹Liedert, A., D. Kaspar, R. Blakytyn, L. Claes, and A. Ignatius. Signal transduction pathways involved in mechanotransduction in bone cells. *Biochem. Biophys. Res. Commun.* 349:1–5, 2006.

- ²²Looms, D. K., K. Tritsarlis, B. Nauntofte, and S. Dissing. Nitric oxide and cGMP activate Ca^{2+} -release processes in rat parotid acinar cells. *Biochem. J.* 355:87–95, 2001.
- ²³McGarry, J. G., J. Klein-Nulend, and P. J. Prendergast. The effect of cytoskeletal disruption on pulsatile fluid flow-induced nitric oxide and prostaglandin E₂ release in osteocytes and osteoblasts. *Biochem. Biophys. Res. Commun.* 330:341–348, 2005.
- ²⁴Mogami, H., A. V. Tepikin, and O. H. Petersen. Termination of cytosolic Ca^{2+} signals: Ca^{2+} reuptake into intracellular stores is regulated by the free Ca^{2+} concentration in the store lumen. *EMBO J.* 17:435–442, 1998.
- ²⁵Norvell, S. M., S. M. Ponik, D. K. Bowen, R. Gerard, and F. M. Pavalko. Fluid shear stress induction of COX-2 protein and prostaglandin release in cultured MC3T3-E1 osteoblasts does not require intact microfilaments or microtubules. *J. Appl. Physiol.* 96:957–966, 2004.
- ²⁶Pozzan, T., R. Rizzuto, P. Volpe, and J. Meldolesi. Molecular and cellular physiology of intracellular calcium stores. *Physiol. Rev.* 74:595–636, 1994.
- ²⁷Ralston, S. H., D. Todd, M. Helfrich, N. Benjamin, and P. S. Grabowski. Human osteoblast-like cells produce nitric oxide and express inducible nitric oxide synthase. *Endocrinology* 135:330–336, 1994.
- ²⁸Rubin, J., C. Rubin, and C. R. Jacobs. Molecular pathways mediating mechanical signaling in bone. *Gene* 367:1–16, 2006.
- ²⁹Sato, K., A. Suematsu, T. Nakashima, S. Takemoto-Kimura, K. Aoki, Y. Morishita, H. Asahara, K. Ohya, A. Yamaguchi, T. Takai, T. Kodama, T. A. Chatila, H. Bito, and H. Takayanagi. Regulation of osteoclast differentiation and function by the CaMK-CREB pathway. *Nat. Med.* 12:1410–1416, 2006.
- ³⁰Saunders, M. M., J. You, J. E. Trosko, H. Yamasaki, Z. Li, H. J. Donahue, and C. R. Jacobs. Gap junctions and fluid flow response in MC3T3-E1 cells. *Am. J. Physiol. Cell Physiol.* 281:C1917–C1925, 2001.
- ³¹Shapiro, F. Variable conformation of GAP junctions linking bone cells: a transmission electron microscopic study of linear, stacked linear, curvilinear, oval, and annular junctions. *Calcif. Tissue Int.* 61:285–293, 1997.
- ³²Soldati, L., A. Terranegra, B. Baggio, R. Biasion, T. Arcidiacono, G. Priante, D. Cusi, and G. Vezzoli. Arachidonic acid influences intracellular calcium handling in human osteoblasts. *Prostaglandins Leukot. Essent. Fatty Acids* 75:91–96, 2006.
- ³³Takai, E. Modulation of bone cell mechanotransduction. In: Department of Biomedical Engineering. New York: Columbia University, p. 209, 2005.
- ³⁴Vashishth, D., O. Verborgt, G. Divine, M. B. Schaffler, and D. P. Fyhrie. Decline in osteocyte lacunar density in human cortical bone is associated with accumulation of microcracks with age. *Bone* 26:375–380, 2000.
- ³⁵Volk, T., K. Mading, M. Hensel, and W. J. Kox. Nitric oxide induces transient Ca^{2+} changes in endothelial cells independent of cGMP. *J. Cell Physiol.* 172:296–305, 1997.
- ³⁶Weinbaum, S., S. C. Cowin, and Y. Zeng. A model for the excitation of osteocytes by mechanical loading-induced bone fluid shear stresses. *J. Biomech.* 27:339–360, 1994.
- ³⁷Yellowley, C. E., C. R. Jacobs, and H. J. Donahue. Mechanisms contributing to fluid-flow-induced Ca^{2+} mobilization in articular chondrocytes. *J. Cell Physiol.* 180:402–408, 1999.
- ³⁸Yellowley, C. E., Z. Li, Z. Zhou, C. R. Jacobs, and H. J. Donahue. Functional gap junctions between osteocytic and osteoblastic cells. *J. Bone Miner. Res.* 15:209–217, 2000.
- ³⁹You, L. D., S. Weinbaum, S. C. Cowin, and M. B. Schaffler. Ultrastructure of the osteocyte process and its pericellular matrix. *Anat. Rec. A Discov. Mol. Cell Evol. Biol.* 278:505–513, 2004.
- ⁴⁰Zaman, G., A. A. Pitsillides, S. C. Rawlinson, R. F. Suswillo, J. R. Mosley, M. Z. Cheng, L. A. Platts, M. Hukkanen, J. M. Polak, and L. E. Lanyon. Mechanical strain stimulates nitric oxide production by rapid activation of endothelial nitric oxide synthase in osteocytes. *J. Bone Miner. Res.* 14:1123–1131, 1999.
- ⁴¹Zayzafoon, M. Calcium/calmodulin signaling controls osteoblast growth and differentiation. *J. Cell Biochem.* 97:56–70, 2006.
- ⁴²Zayzafoon, M., K. Fulzele, and J. M. McDonald. Calmodulin and calmodulin-dependent kinase II α regulate osteoblast differentiation by controlling c-fos expression. *J. Biol. Chem.* 280:7049–7059, 2005.