Archived at the Flinders Academic Commons

http://dspace.flinders.edu.au/dspace/

This is the publisher's copyrighted version of this article.

The original can be found at: http://www.trialsjournal.com/content/8/1/5

Trials

Letter to the Editor

Handheld computers for data entry: high tech has its problems too

Tania M Shelby-James*[1,2], Amy P Abernethy[1,2,3,4], Andrew McAlindon[5] and David C Currow[1,6]

Address: [1]Department of Palliative and Supportive Services, Flinders University, Bedford Park, South Australia, Australia, [2]Southern Adelaide Palliative Services, Repatriation General Hospital, Daw Park, South Australia, Australia, [3]Division of Medical Oncology, Department of Medicine, Duke University Medical Centre, Durham, North Carolina, USA, [4]Department of Medicine, Flinders University, Adelaide, South Australia, Australia, [5]HealthConnect SA, Department of Health, South Australia, Australia and [6]Cancer Australia, Canberra, Australia

Email: Tania M Shelby-James* - tania.shelby-james@rgh.sa.gov.au; Amy P Abernethy - amy.abernethy@duke.edu; Andrew McAlindon - andrew.mcalindon@health.sa.gov.au; David C Currow - david.currow@rgh.sa.gov.au

* Corresponding author

Abstract

Background: The use of handheld computers in medicine has increased in the last decade, they are now used in a variety of clinical settings. There is an underlying assumption that electronic data capture is more accurate that paper-based data methods have been rarely tested. This report documents a study to compare the accuracy of hand held computer data capture versus more traditional paper-based methods.

Methods: Clinical nurses involved in a randomised controlled trial collected patient information on a hand held computer in parallel with a paper-based data form. Both sets of data were entered into an access database and the hand held computer data compared to the paper-based data for discrepancies.

Results: Error rates from the handheld computers were 67.5 error per 1000 fields, compared to the accepted error rate of 10 per 10,000 field for paper-based double data entry. Error rates were highest in field containing a default value.

Conclusion: While popular with staff, unacceptable high error rates occurred with hand held computers. Training and ongoing monitoring are needed if hand held computers are to be used for clinical data collection.

Background

The use of handheld computers has been increasing steadily in medicine over the last decade. As a data entry tool, handheld computers have a number of advantages over paper-based data collection including reduced paperwork, transcription errors, time and cost [1,2]. In some clinical settings it has been estimated that the use of a handheld computer can save nurses nearly two hours per day [3]. Handheld computers have been adapted into a variety of research and clinical settings and available software allows integration into many database types. However, studies have not documented the accuracy of data collection using handheld computers. There is an underlying assumption they are better than a paper-based system.

We recently completed a longitudinal prospective randomised controlled trial of 461 palliative patients [4] fol-

lowed from referral to a specialised palliative care service until death. This trial generated more than 2 million paper-based data entry points. Prior to instituting hand-held computers for data collection and entry, we assessed user accuracy. Our aim was to use handheld computers to reduce trial nurse's workload.

Participants, Methods, and Results

An Ethics Committee-approved comparison study was undertaken with 6 trial forms. Staff undertook training on Compaq Pocket PCs model iPAQ H3950. Research nurses completed a traditional paper-based form that was entered into a database by administrative staff in parallel with the same nurse completing an electronic form on the handheld computer. Data were transferred from the hand-held computer using a manual cradle. Results of these two methods were compared for discrepancies; the paper form was considered the gold standard.

A total of 2001 data elements from 29 consecutive trial participants were entered using both electronic and paper-based methods. Error rates for data entry using the hand-held computers were 67.5 errors per 1000 fields, much higher than the accepted 10 per 10,000 fields for paper-based double data entry[5]. Error rates were highest in fields containing a default value such as a date.

Staff found the handheld computers easy to use and liked using them as a reference tool however they reported dis-comfort using them for data collection. As reported else-where, [6] we experienced a number of technical problems with uploading and downloading data that caused frustration and dissatisfaction. This may have been reduced using a wireless system.

Comment

Rarely have evaluations of user accuracy been undertaken when introducing new technologies for data collection. In contrast to the belief that handheld computers would decrease error rates, we found unacceptably high error rates. This finding has implications for the introduction of electronic data collection for research or clinical purposes without an adequate testing phase.

Error rates need to be monitored if handheld computers are to be used both at the time they are introduced, and in an ongoing way.

Competing interests

The author(s) declare that they have no competing inter-ests.

Authors' contributions

TSJ carried out conception and design, obtaining funding, acquisition of data; analysis and interpretation of data

and drafting of the manuscript. AA carried out conception and design; analysis and interpretation of data and critical revision of manuscript. AM provided technical advice and design of the data base and critical revision of the manu-script. DC participated in the design of the study, interpre-tation of the data, critical revision of the manuscript and supervision. All authors read and approved the final man-uscript.

References

1. Saleh KJ, Radosevich DM, Kassim RA, Moussa M, Dykes D, Bottolfson H, Gioe TJ, Robinson H: **Comparison of commonly used ortho-paedic outcome measures using palm-top computers and paper surveys.** *Journal of Orthopaedic Research* 1920:1146-1151.
2. Shaw S, May V: **An Australian agency revolutionizes home care with the palm pilot.** *Caring* 2004, **23**:30-32.
3. **Organizational learning: could it be the key to higher quality and increased revenues?** *Healthcare Benchmarks & Quality Improve-ment* 2002, **9**:37-40.
4. Abernethy AP, Currow DC, Hunt R, Williams H, Roder-Allen G, Rowett D, Shelby-James T, Esterman A, May F, Phillips PA: **A prag-matic 2 × 2 × 2 factorial cluster randomized controlled trial of educational outreach visiting and case conferencing in pal-liative care-methodology of the Palliative Care Trial [ISRCTN 81117481].** *Contemporary Clinical Trials 27(1):83-100,* 2006.
5. Day S, Fayers P, Harvey D: **Double data entry: What value, what price?** *Controlled Clinical Trials* 1998, **19**:15-24.
6. Chan SS, Chu CP, Cheng BC, Chen PP: **Data management using the personal digital assistant in an acute pain service.** *Anaes-thesia & Intensive Care* 2004, **32**:81-6.