

SPECIAL REPORT

STROKE, COMPLEX REGIONAL PAIN SYNDROME AND PHANTOM LIMB PAIN: CAN COMMONALITIES DIRECT FUTURE MANAGEMENT?

Nicole E. Acerra, BScPT^{1,2}, Tina Souvlis, PhD¹ and G. Lorimer Moseley, PhD³

From the ¹Division of Physiotherapy, The University of Queensland, and ²Physiotherapy Department, Royal Brisbane and Women's Hospital, Brisbane, Australia, and ³Pain Imaging Neuroscience Group, Department of Human Physiology, Anatomy & Genetics & fMRIB Centre, The University of Oxford, Oxford, UK

Despite being different conditions, complex regional pain syndrome type 1, phantom limb pain and stroke share some potentially important similarities. This report examines experimental and clinical findings from each patient population. It identifies common aspects of symptomatic presentation, sensory phenomena and patterns of cortical reorganization. Based on these common findings, we argue that established principles of stroke rehabilitation are also applicable to rehabilitation of complex regional pain syndrome type 1 and phantom limb pain. In addition, we contend that promising treatment approaches for complex regional pain syndrome type 1 and phantom limb pain may be helpful in stroke rehabilitation. Examples of emerging supportive evidence for these hypotheses are provided and discussed.

Key words: stroke, reflex sympathetic dystrophy, physiotherapy, sensorimotor cortex, cortical reorganization, neurorehabilitation.

J Rehabil Med 2007; 39: 109–114

Correspondence address: Tina Souvlis, Division of Physiotherapy, School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, 4072, Australia. E-mail: t.souvlis@shrs.uq.edu.au

Submitted March 16, 2006; accepted October 11, 2006.

INTRODUCTION

The brain retains the ability to change throughout one's lifespan. This ability is demonstrated by the continually updating nature of neural representations held throughout the brain, including the somatotopic representation of the body in the primary somatosensory cortex (1). Sensorimotor plasticity may be particularly important following nervous system injury, for example during stroke recovery, which depends on coordinated cortical activation of the affected and surrounding areas (2). That principle of recovery may be relevant to conditions such as complex regional pain syndrome type 1 (CRPS1) and phantom limb pain (PLP), which are characterized by pain but also by abnormal organization of the sensorimotor cortex contralateral to the affected side (3, 4). Post-stroke cortical reorganization can be influenced by therapeutic interventions, but may be limited by areas of permanent neuronal damage (2). However, with CRPS1 and PLP, no such permanent neu-

ronal damage is likely. CRPS1, PLP and stroke originate from distinct mechanisms (peripheral trauma, deafferentation and cortical damage); however, each can be investigated with the same clinical, electrophysiological and imaging techniques to determine their clinical presentation and cortical response to treatment. Therefore, although they are different medical conditions, common features can be compared. From this perspective, CRPS1, PLP and stroke share many features, including the following: similar changes in sensorimotor activation patterns; sensibility; and response to attention-based repetitive-training regimes. Here we argue that, on the basis of clinical and cortical similarities, treatments effective in one patient population might be effective in the others. This is important because stroke, CRPS1 and PLP are all debilitating conditions for which successful rehabilitation can be elusive (5, 6). Therefore, we argue that CRPS1 and PLP may respond to interventions that restore normal cortical activity during post-stroke recovery and that effective stroke treatments may benefit patients with CRPS1 and PLP.

CLINICAL PRESENTATION OF STROKE, CRPS1 AND PLP

Each year approximately 0.5% of the population will sustain a cerebrovascular accident, or stroke (7). In each case, signs and symptoms reflect the damaged areas of the brain and may include motor, sensory, language, perceptual and cognitive deficits (7). Some patients subsequently develop shoulder or hand pain, swelling, hypersensitivity and reduced regional bone density in the affected limb; findings for which there is usually no identifiable peripheral cause (8, 9). The pathophysiology of these findings, sometimes referred to as shoulder-hand syndrome or CRPS1, is unclear, but they are not rare; reported frequencies range from 1.5% to 61% (8, 9).

CRPS1 is most common after minor trauma (10, 11), for example in 8–37% of uncomplicated wrist fractures (12). CRPS1 usually involves a single limb, but it has also been reported following minor trauma to the head (13), neck (13) and chest (14), and via psychophysiological mechanisms such as with post-traumatic stress disorder (15). Signs and symptoms include: intense, non-dermatomal pain; altered sweating and blood flow; peripheral oedema; abnormal sensation, including hyperalgesia and allodynia; restricted active and passive move-

ment; reduced regional bone density; tissue hypoxia; reduced muscle strength; and muscle tremor (see diagnostic criteria (10) and review (11)). There is growing evidence that CRPS1 is a centrally-mediated neurological condition (11) involving the central nervous system (CNS) at several integrated levels, including the somatosensory, sympathetic and somatomotor systems (8, 11).

PLP occurs in as many as 80% of amputees. It is characterized by: intense, burning, non-dermatomal pain; warmth or cold; perceived motor disturbances, including cramping and tremor; perceived limitations in active range of motion; and perceived swelling within the phantom limb (see review (16)). The phantom limb can also be perceptually distorted, for example, reports of the phantom limb in sustained uncomfortable positions (17). The pathophysiology of PLP is complex; however, like CRPS1, it appears to be mediated within or by the CNS (4, 16).

CORTICAL CHANGES IN CRPS1 AND PLP – RELEVANT TO STROKE?

The CNS is highly adaptive and dynamic – functional organization and neuronal response profiles change according to use (1). Neuroimaging techniques investigate brain activity in association with tasks and stimuli, which in turn allow evaluation of the relationship between cortical changes and symptoms (2). Alterations in the size, shape, location and activation pattern of individual body-part representations in the sensorimotor cortex can effectively be visualized (18) and these images can suggest possible mechanisms underlying clinical conditions and their recovery. The neuroimaging findings in stroke, CRPS1 and PLP are presented and interpreted from this point of view.

The nature of cortical reorganization in stroke, CRPS1 and PLP

Recovery after stroke is thought to depend on cortical reorganization (2). Treatment generally employs repetitive verbal and manual cues and task-specific component training (5, 19). A number of studies have investigated the pattern of cortical changes post-stroke; one unifying conclusion is that the best motor recovery is associated with the greatest return to a normal state of brain function (2, 20, 21). Specifically, imaging studies performed before and after successful stroke rehabilitation show a variety of altered brain activation patterns including: a change in the location and the size of the affected areas; less bilateral brain activation; more activation along the rim of the damaged area; and some degree of diaschisis, where brain areas spatially distant take over the function of stroke-affected areas (see reviews (2, 21)). Several studies show that, with treatment, motor recovery occurs in parallel with increased motor cortex activity during movement; including possible changes in sensorimotor, pre-motor and supplementary motor cortices (2, 21) and a shift in the sensorimotor cortex representation (2).

In CRPS1 the sensorimotor cortex of the affected body-part changes in activation pattern, and in the size and location of

the body-part's cortical representation (3, 22, 23). It appears that the cortical representation effectively shrinks (e.g. in unilateral upper limb CRPS1, the distance between the centre of the representation for the hand and lip and between digits one and five of the affected limb are smaller than those of the unaffected limb) (23). Interestingly, the extent of shift in representation is correlated with the pain intensity (3, 23), a relationship also reported in people with PLP (4, 24). In addition, following treatment the greatest pain relief is positively correlated with normalization of the body-part representations in the somatosensory cortex (3).

Cortical reorganization is also a feature of PLP; like CRPS1, the shift in cortical reorganization is associated with pain intensity (4, 24). Altered affected body-part cortical representations are not surprising because of the missing limb's lack of sensory input and motor ability. The somatosensory cortex changes to reflect this with an expansion of the adjacent cortical representations into the amputated body-part representation (4, 24). Notably, cortical reorganization is related to PLP, but it seems to be unrelated to phantom limb pain intensity, stump pain, referred sensations and reports of telescoping (24). Further similarities in cortical activation patterns are noted with CRPS1 and PLP treatment. As with CRPS1 (3), reduced PLP in response to treatment correlates with normalization of cortical reorganization (4).

Clinical findings in stroke, CRPS1 and PLP

Synchiria. Synchiria is a clinical phenomenon in which a cutaneous stimulus that is applied to one limb evokes sensation simultaneously in both limbs (25). Synchiria occurs following stroke (26) and in amputees with PLP (27). Synchiria and dysynchiria have been observed in people with unilateral CRPS1 (28). In dysynchiria touch to the asymptomatic limb evokes the sensation of touch in that limb and pain and dy-saesthesia at the corresponding site at a mirror-image site on the affected limb (28). Notably, the experience evoked on the affected limb matched that which would be evoked if that limb was actually touched. Synchiria and dysynchiria are assessed while the patient watches the unaffected limb stimulation with a mirror placed between their limbs such that patients can see the unaffected limb and its mirror image. Neither synchiria nor dysynchiria have been reported in healthy subjects and we have not been able to produce it in people with acute localized or radicular pain (28). However, synchiria has been reported in pain-free post-stroke patients (26), which suggests that the phenomena are not evoked solely in association with pain. The mechanisms underlying synchiria and dysynchiria remain to be elucidated, but could include any of the following working alone or in combination: (i) changes in spinal dorsal horn function (including central sensitization (29), bilateral sensory interneurons or ganglia (30), spinal cord or brainstem commissural interneurons (31), or glial cell activation (32, 33)); (ii) changes in subcortical structures (including changes in thalamic function (34, 35), associative somatosensory cortices, the insula, frontal cortices or the anterior cingulate cortex, each of which are known to change with CRPS1 (34,

36)) or altered communication between both hemisphere via the corpus callosum (37).

Referred sensations. In this type of sensory referral, a cutaneous stimulus is experienced both at the stimulated area (e.g. the face) and at another site that is anatomically remote but adjacent to the stimulated site in either the primary somatosensory cortex homunculus (S1) (e.g. the hand) (17, 38, 39), the secondary somatosensory cortex (SII), the thalamus, the posterior parietal cortex or the right dorsolateral prefrontal cortex (40). For example, touching the cheek can evoke the feeling of touch in the hand in patients with CRPS1 (39) and stroke (38) and in the phantom limb of amputees (17, 24, 40). This type of sensory referral may reflect changes in the response properties of S1 (37), SII, thalamic, posterior parietal or right dorsolateral prefrontal neurons (40) similar to that proposed to explain synaesthesia. In synaesthesia, stimulation of one sensory modality automatically triggers perception in a second modality (such as coloured numbers) and may be related to neuronal firing (37). However, the mechanisms underlying referred sensations may be more complex than a simple shift in neighbouring cortical representations (38). Possible mechanisms include mediation of extensive and interconnecting neural networks with variable synaptic strength (41), or loss of sensory input which may remove tonic inhibition from the affected zone (“disinhibition”) such that sensory input from adjacent zones is now sufficient to activate neurons normally responsible for the anatomically remote area (41). That proposal is similar to the one offered for observations in primates, in which cortical synapses normally suppressed by simultaneous input from two connected neurons are thought to become disinhibited when sensory input from one area is removed (42).

IMPLICATIONS FOR TREATMENT

We propose that these neuroimaging and clinical data raise implications for the management of stroke, CRPS1 and PLP. First, it is accepted that stroke involves direct insult to brain tissue. Therefore one goal of stroke rehabilitation is to maximize the return of normal brain and limb function. Secondly, effective treatments that target cortical changes in stroke may be applicable across conditions characterized by cortical reorganization, and vice versa. Viewing the current literature from the perspective that stroke, CRPS1 and PLP may depend on similar cortical networks to attain maximal functional recovery, novel clinical pathways may be elucidated and investigated to foster the greatest cortical reorganization and recovery independent of the inciting condition (e.g. a peripheral injury in the case of CRPS1 and PLP and central brain damage in the case of stroke). This approach advocates an expansion of current management practices to include interventions which are associated with cortical reorganization and improved function. Interventions based on this model have demonstrated positive results (2–4) and raise potential avenues for treatment in other groups, in which cortical mechanisms may be important. Here we discuss 3 types of treatment to illustrate this point:

(i) mirror therapy (43, 44) and motor imagery (45–47), (ii) constraint-induced movement therapy (CIMT) (48), and (iii) sensory discrimination training (4).

Mirror therapy and motor imagery

Mirror therapy involves bilateral limb movement while simultaneously viewing the unaffected hand and its reflected image performing the movement. During mirror therapy, the affected hand remains hidden from view. Mirror therapy has been evaluated for two different purposes: pain relief with PLP and CRPS1, and motor recovery post-stroke. There is evidence that mirror therapy reduces pain in patients with CRPS1 (44) and anecdotal data suggest likewise for amputees with PLP (27, 37). There is also evidence that mirror therapy enhances motor ability over 6 months post-stroke (43, 49). Mirror therapy has been shown to increase ipsilateral primary motor excitability in healthy controls when compared with sham therapy (50), which may account for the improvement in motor function (43, 49). Mirror therapy is currently being evaluated for its ability to enhance motor recovery early post-stroke (51) when cortical reorganization is at its peak (19).

Motor imagery, which includes imagined movements of the affected limb, is effective in motor recovery in chronic (>6 months) post-stroke populations (45, 52). Motor imagery may be effective with both simple (e.g. wrist movement (45)) and complex (e.g. walking (52)) movement patterns. Similarly, graded motor imagery, which involves combining limb laterality recognition (determining limb images as right or left), imagined movements and mirror therapy (46), imparts improvement in pain and disability and functional gains with patients with CRPS1 (46). Because the effect is lost when the components are reordered, it is unlikely to be simply due to increased attention to the limb, but may depend on sequential activation of cortical sensory and motor networks (47).

The mechanisms underlying motor imagery remain unclear. Real and imagined movements activate similar cortical networks (53), and imagined performance is an established strategy in sports and performance psychology (54). Proposed mechanisms for improved motor recovery post-stroke with mirror therapy and motor imagery include: reconciliation of motor output and sensory feedback (44); activation of so-called pre-motor “mirror neurones”, which have intimate connections with visual processing areas (55), are thought to prime the primary motor cortex (46) and to be important in imitating motor action (56); and graded activation of cortical motor networks (57). Although empirical data relating to those theories are lacking, mirror therapy and graded motor imagery programmes may be useful treatment with each of CRPS1, PLP and post-stroke patients.

Constraint-induced movement therapy (CIMT)

CIMT aims to effect cortical changes by forcing use of the stroke-affected limb through impeding use of the unaffected limb, for example by immobilizing the unaffected hand by placing it in an oven mitt (48). CIMT has shown promising results in the chronic stroke population, in whom most studies

have demonstrated expansion of the contralateral cortical motor areas corresponding to improved motor performance both after 2 weeks of treatment and at 6 months post-treatment (48). A proportional change in cortical representational and sensory and motor performance has also been demonstrated in healthy subjects given repeated peripheral input (58, 59) and in patients after stroke participation in repetitive-training regimes (60). One goal for CRPS1 treatment is to increase use of the limb through exercise, desensitization (applying gradual and variable sensory stimuli) and a gradual return to activities (61); however, to our knowledge CIMT has not been investigated formally in this population. CRPS1 has been associated with generalized disuse (62); therefore CIMT, used gradually and as pain allows, could produce good results in the treatment of CRPS1 and warrants further investigation.

Sensory discrimination training

Sensory discrimination training involves discrimination of the type and location of stimuli applied to the skin and has been investigated in amputees (4). In that study, patients localized short-duration electrical stimuli (50 Hz) on the affected limb (90 minutes daily for 10 days) and the results demonstrated three important findings: (i) patients had reduced PLP intensity; (ii) there was normalization of S1 organization; and (iii) both of the above correlated with improved performance on the sensory discrimination task (4). The mechanism underlying the effect is not clear. It is possible that active discrimination of the passive electrical stimulus influenced cortical reorganization and pain intensity as per repetitive learning regimes, or that it had a transcutaneous electrical nerve stimulation-like effect in reducing pain (63); or that it had an effect similar to sensory desensitization, which is often advocated for tissue hypersensitivity (8, 64). The neural mechanisms underlying the effects of sensory discrimination training on hypersensitivity are unclear. They may impact symptoms by influencing cortical reorganization of the somatosensory cortex in patients with CRPS1, PLP and stroke in a similar manner to repetitive learning-regimes in healthy individuals (58, 59) via normalization of somatosensory representation (4).

A common element of mirror therapy and motor imagery, CIMT and sensory discrimination training, is that they might target symptomatic and functional improvement by gradually providing adequate and appropriate input to influence cortical reorganization. In that sense, these approaches are akin to tasks such as learning to read Braille or to play a musical instrument, which are associated with altered sensory representation of the area trained in association with improved performance (58, 59).

DISCUSSION

We have proposed that it may be possible to adapt and apply treatments across post-stroke, CRPS1 and PLP patient populations because: (i) they share common neuroimaging and clinical findings; and (ii) the best recovery seems to be associated with the greatest return to normal brain function. Stroke

rehabilitation is based on the principle that functional recovery is contingent upon appropriate and maximal cortical activation and use-dependent learning. Similar use-dependent strategies might promote CRPS1 and PLP recovery. The proposed model assumes that improved cortical activation patterns underlie recovery in each of stroke, CRPS1 and PLP, and that interventions that promote maximal cortical reorganization may be applicable across conditions. The current review highlights similar neuroimaging, clinical and treatment-responses in the patient populations of stroke, CRPS1 and PLP. Based on these similarities mechanism-driven treatments that are effective in enhancing recovery in one patient population should be assessed for their effectiveness in the other patient populations. We have used the examples of mirror therapy and motor imagery, CIMT and sensory discrimination training to demonstrate the possibility that treatments associated with changes in cortical activation patterns may be effective across the conditions of stroke, CRPS1 and PLP.

ACKNOWLEDGEMENTS

NEA is supported by NHMRC Grant ID 409919 and The Royal Brisbane & Women's Hospital, Brisbane, Australia. GLM is supported by a Nuffield Oxford Medical Fellowship and is on leave from the School of Physiotherapy, The University of Sydney, Australia.

REFERENCES

- Buonomano D, Merzenich M. Cortical plasticity: from synapses to maps. *Annu Rev Neurosci* 1998; 21: 149–186.
- Cramer SC. Functional imaging in stroke recovery. *Stroke* 2004; 35: 2695–2698.
- Maihofner C, Handwerker HO, Neundorfer B, Birklein F. Cortical reorganization during recovery from complex regional pain syndrome. *Neurology* 2004; 63: 693–701.
- Flor H, Denke C, Schaefer M, Grusser S. Effect of sensory discrimination training on cortical reorganisation and phantom limb pain. *Lancet* 2001; 357: 1763–1764.
- Lincoln NB, Parry RH, Vass CD. Randomized, controlled trial to evaluate increased intensity of physiotherapy treatment of arm function after stroke. *Stroke* 1999; 30: 573–579.
- Schwartzman RJ. Reflex sympathetic dystrophy. *Curr Opin Neurol Neurosurg* 1993; 6: 531–536.
- Feigin VL, Lawes CMM, Bennett DA, Anderson CS. Stroke epidemiology: a review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurol* 2003; 2: 43–53.
- Wasner G, Schattschneider J, Binder A, Baron R. Complex regional pain syndrome – diagnostic, mechanisms, CNS involvement and therapy. *Spinal Cord* 2003; 41: 61–75.
- Daviet JC, Preux PM, Salle JY, Lebreton F, Munoz M, Dudognon P, et al. Clinical factors in the prognosis of complex regional pain syndrome type I after stroke – a prospective study. *Am J Phys Med Rehab* 2002; 81: 34–39.
- Merskey H, Bogduk N. Classification of chronic pain: descriptions of chronic pain syndromes and definitions of pain terms. Seattle: IASP Press; 1994.
- Janig W, Baron R. Complex regional pain syndrome is a disease of the central nervous system. *Clin Auton Res* 2002; 12: 150–164.
- Zollinger PE, Tuinebreijer WE, Kreis RW, Breederveld RS. Effect of vitamin C on frequency of reflex sympathetic dystrophy in wrist

- fractures: a randomised trial. *Lancet* 1999; 354: 2025–2028.
13. Melis M, Zawawi K, Al-Badawi E, Lobo SL, Mehta N. Complex regional pain syndrome in the head and neck: a review of the literature. *J Orofac Pain* 2002; 16: 93–104.
 14. Papay FA, Verghese A, StantonHicks M, Zins J. Complex regional pain syndrome of the breast in a patient after breast reduction. *Ann Plast Surg* 1997; 39: 347–352.
 15. Grande LA, Loeser JD, Ozuna J, Ashleigh A, Samii A. Complex regional pain syndrome as a stress response. *Pain* 2004; 110: 495–498.
 16. Flor H. Phantom-limb pain: characteristics, causes, and treatment. *Lancet Neurol* 2002; 1: 182–189.
 17. Ramachandran VS, Hirstein W. The perception of phantom limbs – The DO Hebb lecture. *Brain* 1998; 121: 1603–1630.
 18. Dykes RW. Mechanisms controlling neuronal plasticity in somatosensory cortex. *Can J Physiol Pharmacol* 1997; 75: 535–545.
 19. Carr JH, Shepherd RB. Stroke rehabilitation: guidelines for exercise and training to optimize motor skill. London: Butterworth-Heinemann; 2003.
 20. Nudo RJ, Friel KM. Cortical plasticity after stroke: implications for rehabilitation. *Rev Neurol (Paris)* 1999; 155: 713–717.
 21. Ward NS, Cohen LG. Mechanisms underlying recovery of motor function after stroke. *Arch Neurol* 2004; 61: 1844–1848.
 22. Juottonen K, Gockel M, Silen T, Hurri H, Hari R, Forss N. Altered central sensorimotor processing in patients with complex regional pain syndrome. *Pain* 2002; 98: 315–323.
 23. Maihofner C, Handwerker HO, Neundorfer B, Birklein F. Patterns of cortical reorganization in complex regional pain syndrome. *Neurology* 2003; 61: 1707–1715.
 24. Flor H, Elbert T, Knecht S, Wienbruch C, Pantev C, Birbaumer N, et al. Phantom-limb pain as a perceptual correlate of cortical reorganization following arm amputation. *Nature* 1995; 375: 482–484.
 25. Janet P. Un cas d’allochirie. In: *Névroses et idées fixes*. Paris; 1898, p. 234–262.
 26. Sathian K. Interannual referral of sensation to anesthetic hands. *Neurology* 2000; 54: 1866–1868.
 27. Ramachandran VS, Rogers-Ramachandran D, Cobb S. Touching the phantom limb. *Nature* 1995; 377: 489–490.
 28. Acerra NE, Moseley GL. Dysynchiria: watching the mirror image of the unaffected limb elicits pain on the affected side. *Neurology* 2005; 65: 751–753.
 29. Bausbaum A, Bushnell MC, Devor M. Pain: basic mechanisms. In: Justins DM, ed. *Pain 2005 – an updated review*. Seattle: IASP Press; 2005, p. 3–9.
 30. McLachlan EM, Janig W, Devor M, Michaelis M. Peripheral nerve injury triggers noradrenergic sprouting within dorsal root ganglia. *Nature* 1993; 363: 543–546.
 31. Koltzenburg M, Wall PD, McMahon SB. Does the right side know what the left is doing? *Trends Neurosci* 1999; 22: 122–127.
 32. Milligan ED, Twining C, Chacur M, Biedenkapp J, O’Connor K, Poole S, et al. Spinal glia and proinflammatory cytokines mediate mirror-image neuropathic pain in rats. *J Neurosci* 2003; 23: 1026–1040.
 33. Watkins LR, Maier SF. Beyond neurons: Evidence that immune and glial cells contribute to pathological pain states. *Physiol Rev* 2002; 82: 981–1011.
 34. Fukumoto M, Ushida T, Zinchuk VS, Yamamoto H, Yoshida S. Contralateral thalamic perfusion in patients with reflex sympathetic dystrophy syndrome. *Lancet* 1999; 354: 1790–1791.
 35. Dostrovsky JO. Ascending spinal and trigeminal nociceptive pathways. In: Justins DM, editor. *Pain 2005 – an updated review: refresher course syllabus*. Seattle: IASP Press; 2005, p. 43–48.
 36. Maihofner C, Forster C, Birklein F, Neundorfer B, Handwerker HO. Brain processing during mechanical hyperalgesia in complex regional pain syndrome: a functional MRI study. *Pain* 2005; 114: 93–103.
 37. Ramachandran VS. A brief tour of human consciousness: from impostor poodles to purple numbers. New York: PI Press; 2004.
 38. Turton AJ, Butler SR. Referred sensations following stroke. *Neurocase* 2001; 7: 397–405.
 39. McCabe CS, Haigh RC, Halligan PW, Blake DR. Referred sensations in patients with complex regional pain syndrome type 1. *Rheumatology* 2003; 42: 1067–1073.
 40. Grusser SM, Muhlnickel W, Schaefer M, Villringer K, Christmann C, Koeppel C, et al. Remote activation of referred phantom sensation and cortical reorganization in human upper extremity amputees. *Exp Brain Res* 2004; 154: 97–102.
 41. Knecht S, Henningsen H, Elbert T, Flor H, Hohling C, Pantev C, et al. Reorganizational and perceptual changes after amputation. *Brain* 1996; 119: 1213–1219.
 42. Gandevia S, Phegan C. Perceptual distortions of the human body image produced by local anaesthesia, pain and cutaneous stimulation. *J Phys (Lond)* 1999; 514: 609–616.
 43. Altschuler EL, Wisdom SB, Stone L, Foster C, Galasko D, Llewellyn DME, et al. Rehabilitation of hemiparesis after stroke with a mirror. *Lancet* 1999; 353: 2035–2036.
 44. McCabe CS, Haigh RC, Ring EFJ, Halligan PW, Wall PD, Blake DR. A controlled pilot study of the utility of mirror visual feedback in the treatment of complex regional pain syndrome (type 1). *Rheumatology* 2003; 42: 97–101.
 45. Dijkerman HC, Letswaart M, Johnston M, MacWalter RS. Does motor imagery training improve hand function in chronic stroke patients? A pilot study. *Clin Rehab* 2004; 18: 538–549.
 46. Moseley GL. Graded motor imagery is effective for long-standing complex regional pain syndrome. *Pain* 2004; 108: 192–198.
 47. Moseley GL. Is successful rehabilitation of complex regional pain syndrome due to sustained attention to the affected limb? A randomised clinical trial. *Pain* 2005; 114: 54–61.
 48. Taub E, Uswatte G, Pidikiti R. Constraint-Induced Movement Therapy: A new family of techniques with broad application to physical rehabilitation – a clinical review. *J Rehabil Res Dev* 1999; 36: 237–251.
 49. Sathian K, Greenspan AI, Wolf SL. Doing it with mirrors: a case study of a novel approach to neurorehabilitation. *Neurorehabil Neural Repair* 2000; 14: 73–76.
 50. Garry MI, Loftus A, Summers JJ. Mirror, mirror on the wall: view ing a mirror reflection of unilateral hand movements facilitates ipsilateral M1 excitability. *Exp Brain Res* 2005; 163: 118–122.
 51. Acerra NE, Souvlis T, Moseley GL. Does mirror-box therapy improve sensory and motor changes in the early post-stroke population? A randomised-controlled trial. In: *Joint Conference of the National Neurology and Gerontology Groups of the Australian Physiotherapy Association*; 2005 November; Melbourne, Australia: Australian Physiotherapy Association; 2005, p. 23.
 52. Dickstein R, Dunsky A, Marcovitz E. Motor imagery for gait rehabilitation in post-stroke hemiparesis. *Phys Ther* 2004; 84: 1167–1178.
 53. Jackson PL, Lafleur MF, Malouin F, Richards CL, Doyon J. Functional cerebral reorganization following motor sequence learning through mental practice with motor imagery. *Neuroimage* 2003; 20: 1171–1180.
 54. Yue G, Cole KJ. Strength increases from the motor program: comparison of training with maximal voluntary and imagined muscle contractions. *J Neurophysiol* 1992; 67: 1114–1123.
 55. Rizzolatti G, Fadiga L, Gallese V, Fogassi L. Premotor cortex and the recognition of motor actions. *Brain Res Cogn Brain Res* 1996; 3: 131–141.
 56. Sterr A, Freivogel S. Intensive training in chronic upper limb hemiparesis does not increase spasticity or synergies. *Neurology* 2004; 63: 2176–2177.
 57. Liepert J, Bauder H, Miltner WHR, Taub E, Weiller C. Treatment-induced cortical reorganization after stroke in humans. *Stroke* 2000; 31: 1210–1216.
 58. Sterr A, Muller M, Elbert T, Rockstroh B, Taub E. Development of cortical reorganization in the somatosensory cortex of adult

- Braille students. Functional neuroscience: evoked potentials and magnetic fields. *Electroencephalogr Clin Neurophysiol Suppl* 1999; 49: 292–298.
59. Pantev C, Engelien A, Candia V, Elbert T. Representational cortex in musicians: plastic alterations in response to musical practice. *Ann N Y Acad Sci*. 2001; 930: 300–14.
60. Sterr A, Freivogel S. Motor-improvement following intensive training in low-functioning chronic hemiparesis. *Neurology* 2003; 61: 842–844.
61. Stanton-Hicks M, Baron R, Boas R, Gordh T, Harden N, Hendler N, et al. Complex regional pain syndromes: guidelines for therapy. *Clin J Pain* 1998; 14: 155–166.
62. Butler SH. Disuse and CRPS. In: Harden RN, Baron R, Janig W, eds. *Complex regional pain syndrome: progress in pain research and management*. Seattle: IASP Press; 2001, p. 141–150.
63. Sluka KA, Walsh D. Transcutaneous electrical nerve stimulation: basic science mechanisms and clinical effectiveness. *J Pain* 2003; 4: 109–121.
64. Waylett-Rendall J. Desensitization of the traumatized hand. In: Hunter JM, Schneider LH, Mackin EJ, Callahan AD, eds. *Rehabilitation of the hand: surgery and therapy*. Third edn. St Louis: Mosby; 1990, p. 1258.