

Trends in potential exposure to Australian bat lyssavirus in South East Queensland, 1996 to 2003

Megan K Young,¹ Bradley J McCall²

Abstract

This study examined trends in notifications of potential exposure to Australian bat lyssavirus reported to the Brisbane Southside Public Health Unit, Australia between 1 November 1996 and 31 January 2003. Notification rates declined among all population groups and potential exposures were notified more promptly. Concern exists regarding possible under-reporting of potential exposure to Australian bat lyssavirus especially among volunteer bat carers. *Commun Dis Intell* 2004;28:258–260.

Keywords: lyssavirus; epidemiology; Australia

Introduction

Australian bat lyssavirus (ABL) is a member of the Rhabdoviridae family, possessing marked similarity to classic rabies virus on both serotyping and molecular sequencing.¹ To date, two cases of fatal ABL infection have been reported in Australia, one in 1996 and the second in 1998.² The epidemiology of potential exposure to ABL has been previously described.³ The key feature, in a predominantly urban population in South East Queensland, was that potential exposures were likely to be the result of human-initiated contact by people with some professional or volunteer interest in caring for bats and/or flying foxes (53% of potential exposures). A lower proportion of potential exposures (35%) were reported by members of the general community.

Potential exposures to ABL continue to occur despite consistent information and reminders to the community about the dangers of handling flying foxes and insectivorous bats.⁴ This paper examines population trends in potential exposure to ABL reported to the Brisbane Southside Public Health Unit (BSPHU) between 1 November 1996 and 31 January 2003.

Methods

Since 1 November 1996, all persons reporting a potential exposure to ABL have been asked to complete a standard questionnaire. The details of the questionnaire, methods of study and results until 31 January 1999, have been described.³ However,

the geographic boundaries of the area served by the BSPHU have changed since 1999. They now include South Brisbane (part of the Brisbane City Council Area), Logan, Redlands, Ipswich, Laidley, Boonah and Esk Local Government Areas (Figure) with an estimated resident population of 920,680 as at 30 June 2000.⁵

The time frame of this study was divided into period 1 (the initial study period from 1 November 1996 to 31 January 1999, which included the two human cases of ABL infection) and period 2 (1 February 1999 to 31 January 2003). To allow comparison of data across these time periods, the original study data was restricted to include only that related to people who resided within the current Brisbane Southside Public Health Unit boundaries. SPSS version 11.5 was used for analysis.⁶

Results


Two hundred and forty-six notifications were reported between 1 November 1996 and 31 January 2003. One hundred and thirty-six notifications of potential ABL exposure from the re-defined study area were reported to the BSPHU in period 1 (duration = 27 months), an average annual notification rate of 6.56 per 100,000. One hundred and ten notifications were reported in period 2 (duration = 48 months), an average annual notification rate of 2.98 per 100,000. The proportion of notifications from females fell from 60 per cent in period 1 to 46 per cent in period 2.

1. Public Health Registrar, Brisbane Southside Public Health Unit, Brisbane, Queensland

2. Public Health Physician, Brisbane Southside Public Health Unit, Brisbane, Queensland

Corresponding author: Dr Megan K Young, Public Health Registrar, Brisbane Southside Public Health Unit, PO Box 333, Archerfield, QLD 4108. Telephone: +61 7 3000 9148. Facsimile: +61 7 3000 9130. Email: megan_young@qld.health.gov.au

Figure. The geographical area covered by the Brisbane Southside Public Health Unit



There was no significant difference between the age distributions ($\chi^2 = 8.13$, 6 df, $p = 0.23$), with the peak age group between 20 and 49 years in both time periods.

The median time interval between potential exposure and notification to the BSPHU fell from 14.5 days (range 0 to 3,636; 25th, 75th centiles: 2, 79.8) in period 1 to one day (range 0 to 1,860; 25th, 75th centiles: 0,3) in period 2.

The Table describes the circumstance of potential exposure to ABL, the treatment received and the history of previous vaccination in the two time periods. There was a decline in the average annual number of reported potential exposures for all population groups (professionals, volunteer bat carers and their family members, community members). However, the proportion of potential exposures reported by community members increased (from 40% to 71%). The proportion of potential exposures reported by professional handlers and volunteer bat carers fell substantially, with the greatest fall among volunteer bat carers (from 36% to 11%). Professional handlers and volunteer bat carers reporting potential exposures in period 2 were more likely to have been previously vaccinated. Only 16 per cent of professional and volunteer handlers reported no previous vaccination in period 2 compared to 86 per cent in period 1.

Table. Circumstance, treatment and history of previous vaccination of potential exposures to Australian bat lyssavirus for each time period, Brisbane Southside Public Health Unit area, 1 November 1996 to 31 January 2003*

Circumstance	Period 1 (1/11/96 – 31/1/99) n=136		Period 2 (1/2/99 – 31/1/03) n=110	
	n	%	n	%
Community member bat initiated contact	3	2.2	17	15.5
Community member intentionally handled bat	51	37.5	61	55.5
Professional handlers	17	12.5	10	9.0
Volunteer bat carers	49	36.0	12	10.9
Family member of volunteer bat carers	13	9.6	3	2.7
Treatment				
Nil (bat tested negative)	5	3.7	40	36.4
Course ceased (bat tested negative)	18	13.2	2	1.8
2 doses of vaccine	12	8.8	12	10.9
5 doses of vaccine	69	50.7	8	7.3
Rabies immunoglobulin and 5 doses of vaccine	31	22.8	47	42.7
Recommended treatment but declined	0	0.0	1	0.9
Previous vaccination				
Nil	121	89.0	85	77.3
Pre-exposure prophylaxis	3	2.2	10	9.1
Pre-exposure prophylaxis and booster/s	0	0.0	4	3.6
Previous post-exposure prophylaxis	10	7.4	6	5.5

* Percentages may not total 100 because of missing values.

Discussion

The notification rate of potential exposure to ABL fell markedly during the study. This rate may more accurately estimate baseline potential exposure rates since earlier rates were inflated by the large number of retrospective reports received after the initial recognition of this disease in humans in 1996. However, it is possible that a proportion of recent potential exposures are not being notified. This may arise from waning concern about the risks of bat bites or scratches among the community or medical profession, especially as no cases of human ABL infection have been reported since 1998.

Professional handlers and volunteer bat carers may be unwilling to notify potential exposures within their groups, a reluctance that has been communicated to both authors. This unwillingness may be an undesired consequence of the public health requirement to euthanase and test all bats involved in human potential exposures. Unwillingness to notify potential exposure may also reflect preformed opinions about the risk posed by scratches, the protection afforded by pre-exposure vaccination, or the level of risk associated with the clinical appearance of the bat. These suggestions are further supported by the finding that volunteer bat carers have the largest decrease in proportion of notifications from time period 1 to period 2 [36% (n=49) to 11% (n=12)], with a corresponding increase in community notifications [40% (n=54) to 71% (n=78)], despite no recognisable change to volunteer bat carer numbers in the Brisbane Southside area over recent years (personal communication, Allan McKinnon, Manager, Moggill Koala Hospital, Queensland Parks and Wildlife Services).

The reduction observed in notifications among females may also be explained by the fall in the proportion of volunteer bat carer notifications. The high proportion of females in period 1 was influenced by the high proportion of female volunteer bat carers in the study population.

The time between potential exposure and notification decreased substantially over the study, with fewer long-term retrospective reports of potential exposure in period 2. A corresponding decrease was observed in the proportion of people requiring treatment after their potential exposure. Post exposure treatment may be delayed for 48 hours pending the results of tests on the bat involved.⁷ In period 2, potential exposures were notified more promptly. This allowed a greater proportion of the bats involved to be tested and the negative results to be obtained within the required 48 hours. This represented an important cost saving through the reduced use of rabies immune globulin (RIG), rabies vaccine and fewer doctors' visits.

Future public health messages should continue to emphasise the need for the community to maintain a safe distance from all bats or flying foxes, even if they are orphaned or distressed. Messages should reinforce that it is usually impossible for an untrained person to handle a bat without sustaining a bite or scratch, even if protective measures are used. Members of the public can be of most help to orphaned or injured bats by contacting a trained, vaccinated bat handler. Volunteer bat carers must also be aware of the potential risks associated with bites or scratches from apparently healthy looking bats, and seek medical advice regardless of their pre-exposure prophylaxis or the nature of the wound.

Acknowledgements

Staff of Public Health Virology, Queensland Health Scientific Services, Archerfield, Queensland, Australia

Staff of Brisbane Southside Public Health Unit, Archerfield, Queensland

Allan McKinnon and staff at Wildlife Rangers Conservation Services, Queensland Parks and Wildlife Services, Moggill, Queensland

References

1. Gould AR, Hyatt AD, Lunt R, Kattenbelt JA, Hengstberger S, Blacksell SD. Characterisation of a novel lyssavirus isolated from Pteropid bats in Australia. *Virus Res* 1998;54:165–187.
2. Hanna JN, Carney IK, Smith GA, Tannenberg AE, Deverill JE, Botha JA, *et al.* Australian bat lyssavirus infection: a second human case, with a long incubation period. *Med J Aust* 2000;172:597–599.
3. McCall BJ, Epstein JH, Neill AS, Heel K, Field H, Barrett J, *et al.* Potential exposure to Australian bat lyssavirus, Queensland, 1996–1999. *Emerg Infect Dis* 2000;6:259–264.
4. Scott JG. Australian bat lyssavirus: the public health response to an emerging infection. *Med J Aust* 2000;172:573–574.
5. Australian Bureau of Statistics. 2000 estimated resident population by Statistical Local Area. Australian Bureau of Statistics Catalogue No. 3235.3. Canberra, Australia: The Organization, 2000.
6. SPSS Inc. SPSS® for Windows™ version 11.5: A Statistical Package for Social Sciences (computer program). 2002. Chicago, Illinois.
7. Australian Technical Advisory Group on Immunisation. *Australian Immunisation Handbook* 8th edition. Australian Government Printing Service, 2003. pp. 65–72.