

Geophysical Research Abstracts
Vol. 12, EGU2010-1283, 2010
EGU General Assembly 2010
© Author(s) 2009



High resolution record of the Last Glacial Maximum in eastern Australia

Lynda Petherick, Patrick Moss, and Hamish McGowan

School of Geography, Planning & Environmental Management, The University of Queensland, St Lucia, Australia
(l.petherick@uq.edu.au)

A continuous, high resolution (average ca. 22 year) record encompassing the Last Glacial Maximum (LGM) has been developed using multiple proxies (aeolian sediment flux, grain size, pollen and charcoal) in lake sediment from Tortoise Lagoon (TOR), North Stradbroke Island, Queensland, Australia. The presence of Asteraceae tubilifloreae and spineless Asteraceae (common indicators of glacial conditions in Australia) at TOR indicates significantly cooler temperatures (mean annual temperature up to 6°C lower than today).

In addition to the palaeoclimatic reconstruction, a record of palaeodust transport pathways for eastern Australia was developed using ICP-MS trace element analysis and geochemical “fingerprinting” of TOR aeolian sediment to continental dust source areas. Vectors between dominant dust source areas and North Stradbroke Island allowed the reconstruction of the position and intensity of LGM dust transport pathways. Furthermore, changes in likely synoptic scale conditions can be postulated based on the position of the dust transport corridors.

Similarities between the vegetation at TOR during the LGM and that at temperate sites e.g. Caledonia Fen, Victoria (Kershaw et al. 2007), Redhead Lagoon, New South Wales (Williams et al. 2006) and Barrington Tops, New South Wales (Sweller and Martin 2001) suggests that this record reflects regional conditions across southeastern Australia. The TOR record also correlates well with that from nearby Native Companion Lagoon which suggests that the LGM was actually an extended period of ca. 8 – 10 kyr, characterised by 2 periods of increased aridity (ca. 30 – 26.5 kyr and 21 – 19.5 kyr) (Petherick et al. 2008). A growing number of records from across the Southern Hemisphere e.g. New Zealand (Suggate and Almond 2003; Alloway et al. 2007; Newnham et al. 2007), Chile (Denton et al. 1999), Antarctica (Röthlisberger et al. 2002; EPICA 2006) and Australia (Smith 2009) also show evidence that the LGM encompassed a longer period of time than traditionally accepted, and was not uniformly cool and dry.

Alloway, B. V., D. J. Lowe, D. J. A. Barrell, R. M. Newnham, P. C. Almond, P. C. Augustinus, N. A. N. Bertler, L. Carter, N. J. Litchfield, M. S. McGlone, J. Shulmeister, M. J. Vandergoes, P. W. Williams and N.-I. members (2007). Towards a climate event stratigraphy for New Zealand over the past 30 000 years (NZ-INTIMATE project). *Journal of Quaternary Science* 22(1): 9-35.

Denton, G. H., T. V. Lowell, C. J. Heusser, C. Schluchter, B. G. Andersen, L. E. Heusser, P. I. Moreno and D. R. Marchant (1999). Geomorphology, stratigraphy, and radiocarbon chronology of Llanquihe Drift in the area of the Southern Lake District, Seno Reloncavi, and Isal Grande de Chiloe, Chile. *Geografiska Annaler* 81A: 167-229.

EPICA (2006). One-to-one coupling of glacial climate variability in Greenland and Antarctica. *Nature* 444: 195-198.

Kershaw, A. P., G. M. McKenzie, N. Porch, R. G. Roberts, J. Brown, H. Heijnis, M. L. Orr, G. Jacobsen and P. R. Newall (2007). A high-resolution record of vegetation and climate through the last glacial cycles from Caledonia Fen, southeastern highlands of Australia. *Journal of Quaternary Science* 22(5): 481-500.

Newnham, R. M., D. J. Lowe, T. Giles and B. V. Alloway (2007). Vegetation and climate of Auckland, New Zealand, since ca. 32 000 cal. yr ago: support for an extended LGM *Journal of Quaternary Science* 22(5):

517-534.

Petherick, L. M., H. A. McGowan and P. T. Moss (2008). Climate variability during the Last Glacial Maximum in eastern Australia: Evidence of two stadials? *Journal of Quaternary Science* 23(8): 787-802.

Röthlisberger, R., R. Mulvaney, E. W. Wolff, M. A. Hutterli, M. Bigler, S. Sommer and J. Jouzel (2002). Dust and sea salt variability in central East Antarctica (Dome C) over the last 45 kyr and its implications for southern high latitude climate. *Geophysical Research Letters* 29(20): Art # 1963.

Smith, M. A. (2009). Late Quaternary landscapes in Central Australia: sedimentary history and palaeoecology of Punitjarra rock shelter. *Journal of Quaternary Science* 24(7): 747-760.

Suggate, R. P. and P. C. Almond (2003). The LGM in western South Island, New Zealand. Australasian Quaternary Association Biennial Conference. Westport, New Zealand.

Sweller, S. and H. A. Martin (2001). A 40,000 year vegetation history and climatic interpretations of Burraga Swamp, Barrington Tops, New South Wales. *Quaternary International* 83-85: 233-244.

Williams, N. J., K. J. Harle, S. J. Gale and H. Heijnis (2006). The vegetation history of the last glacial-interglacial cycle in eastern New South Wales, Australia. *Journal of Quaternary Science* 21(7): 735-750.