

# Solvent effect and fluorescence response of the 7-*tert*-butylpyrene-dipicolyl amine linkage for the selective and sensitive response toward Zn(II) and Cd(II) ions

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

Zannatul Kowser,<sup>a</sup> Hirotsugu Tomiyasu,<sup>a</sup> Xuekai Jiang,<sup>a</sup> Ummei Rayhan, Carl Redshaw<sup>b</sup> and Takehiko Yamato<sup>\*a</sup>

The different binding behaviour of 7-*tert*-butylpyrene based chemosensors bearing dipicolylamine (Dpa) linkages at the 1,3-positions was investigated in various solvents for the sensing of Zn(II) and Cd(II). The potential mono-chelating ligand **L1** follows the same binding pattern in both THF and methanol-water solvent systems, exhibiting high selectivity and sensitivity for Cd(II) than Zn(II) mainly in THF solvent system. The potential bis-chelate ligand **L2** can selectively bind both Zn(II) and Cd(II) in a 1:1 ratio in THF, whereas in methanol-water (7:3) at pH = 7.0; a 1:2 binding ratio was observed. In THF, two sites of ligand **L2** can only selectively and sensitively bind one Zn(II) or Cd(II). The different complexation behaviours of **L1** and **L2** in different solvents were studied by means of fluorescence spectra and <sup>1</sup>H-NMR titration experiments in the presence of Zn(II) and Cd(II).

## Introduction

The design and synthesis of molecular receptors for the detection of environmentally and biologically important species has attracted growing interest in recent years.<sup>1</sup> Amongst them, chemosensors whose fluorescence emission is sensitive to the environment and solvent media are especially important.<sup>2-5</sup> In this regard, many fluorescence mechanisms have also been reported by probing sensing properties based on Photoinduced Electron Transfer (PET), Intermolecular Charge Transfer (ICT), Chelation Enhanced Fluorescence (CHEF). Indeed, their application in the field of supramolecular chemistry has been elegantly illustrated.<sup>6</sup> In case of PET,<sup>7</sup> there is little or no change of the spectral shifts with changes of emission intensities, whereas both spectral shifts and intensity changes are observed for ICT<sup>8</sup>, whilst CHEF<sup>9</sup> also exhibited fluorescence enrichment with or without accompanying spectral changes.

The detection of Zn<sup>2+</sup> is important both *in vitro* and *in vivo* due to its biological relevance.<sup>10,11</sup> It is an indispensable element for the human body and in many physiological and pathological processes, it performs an essential role.<sup>12</sup> It has been reported that its deficiency give rise to acrodermatitis enteropathica,<sup>13</sup> but it is detrimental when present in excess, causing severe health problems such as superficial skin diseases, prostate cancer, diabetes and brain diseases. Unfortunately spectroscopically silent Zn<sup>2+</sup> is difficult to detect directly.<sup>14</sup> By contrast, a trace amount of Cd<sup>2+</sup> is highly toxic

towards the human body. Its intake causes serious diseases such as renal dysfunction, calcium metabolism disorders and prostate cancer.<sup>15</sup>

It is known that fluorescence quenching sometimes creates an unfavourable condition for a high signal output upon recognition of ions and also interferes with temporal separation of spectrally similar complexes with time-resolved fluorometry.<sup>16</sup> Thus, our main focus is to design a chemosensor that does not quench the fluorescence upon binding with a metal ion. In this regard, the PET which is responsible for fluorescence quenching is minimized in the signaling moiety upon binding and results in the enhancement of the fluorescence.

Recently, pyrene has been utilized widely as a fluorophore to detect ion pairs, cations, anions<sup>17</sup> and neutral species,<sup>18</sup> because of the photoluminescence properties and chemical stabilities associated with pyrene. Given this, we have developed chemosensors that contain a 7-*tert*-butylpyrene as a fluorophore moiety and dipicolylamine as a receptor moiety connected through a C-N bond. Such an efficient and simple ligand system was also proposed by Ojida *et al.*<sup>19</sup> They synthesized the binuclear anthracene complex Zn(II)-Dpa, and used it as an anion sensor for phosphorylated peptides. In our present work, we have established the ligands as efficient cation sensors which reveal different behaviour in different solvent systems.



The purpose of this work is to shed light on the mechanism of the different fluorescence response of receptor **L2** with Zn<sup>2+</sup>

and  $\text{Cd}^{2+}$  in various solvent systems. Interestingly, a 1:1 ligand to metal binding ratio was observed in case of THF for both  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ions, whereas when using a methanol-water solvent system, it can selectively interact with  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  ions in a 1:2 (ligand/metal) stoichiometry. In case of methanol-water, **L2** exhibits a significant fluorescence enhancement for  $\text{Zn}^{2+}$ , which is twice that observed for the THF solvent system. However, the potentially mononuclear receptor **L1** is highly selective in coordinating with  $\text{Zn}(\text{II})$  and  $\text{Cd}(\text{II})$ .


## Results and discussions

We have designed and successfully synthesized **L1** and **L2** using the reaction pathway shown in scheme 1. The fluorogenic molecule **L2** is synthesized from 7-*tert*-butylpyrene-1,3-dicarbaldehyde by treatment with 2,2'-dipicolylamine, following which, the Schiff base is reduced by the gradual addition of  $\text{NaBH}_4(\text{OAc})_3$  to obtain **L2** in 82 % yield. Following the same reaction pathway, the potentially mono-chelate **L1** has also been prepared from 7-*tert*-butylpyrene-1-carbaldehyde in order to compare the binding affinities for  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  in different solvent systems. The characterization of these compounds was confirmed by  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy and by High-Mass spectrometry. In the absence of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ion, both **L1** and **L2** only afford weak fluorescence because of PET; the lone pair electrons from the amino group are transferred to the excited pyrenyl moiety and are presumed to quench the emission intensity of the pyrenyl fluorophore. After addition of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  at small concentrations, preferential binding with dipicolylamine occurs to terminate the PET. In this way, the 7-*tert*-butylpyrene binuclear-Dpa complex exhibits a significant fluorescent enhancement for  $\text{Zn}^{2+}$  and can detect both  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ions upon changing the solvent system. Addition of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ions using THF as solvent reveals a fluorescence at 402 nm. On the other hand, ligand **L2** can only

$\text{Zn}^{2+}$ . **L2** itself exhibits very weak fluorescence. It was then found that a large fluorescence enhancement (8-fold) was observed upon addition of



Scheme 1: Synthesis of receptors **L1**, **L2** and **L3**.




detect  $\text{Zn}^{2+}$  ion with almost twice the fluorescence enhancement with on changing the solvent media, ie methanol-water instead of THF.

Firstly, the fluorescence properties of the receptor **L2** were investigated in different solvents (Fig. 1a) following addition of

(a)

(b)

**Fig. 1** (a) Fluorescence response of ligand **L2** (7  $\mu\text{M}$ ) upon addition of  $\text{Zn}^{2+}$  in different solvent systems with excitation at 353 nm. (b) Fluorescence spectra of **L1**, **L2** and **L3** in  $\text{CH}_3\text{OH}/\text{H}_2\text{O}$  (10mM HEPES/ $\text{CH}_3\text{OH}$  = 3:7, pH = 7.0) with excitation at 347 and 353 nm, respectively.




It was also used in combination with [(2,2'-dipicolylamino)butyl]pyrene as the  $\text{Zn}^{2+}$  receptor, and pyrophosphate works as the bridging substrate for the excimer formation.<sup>20</sup>

To investigate the sensitivity of **L2** toward  $\text{Zn}^{2+}$  ion, 9,10-bis[(2,2'-dipicolylamino)methyl]anthracene, **L3** was also synthesized. As


**Fig. 2** Fluorescence intensity changes of receptor **L1**, **L2** and **L3** (7  $\mu\text{M}$ ) in  $\text{CH}_3\text{OH}/\text{H}_2\text{O}$  (10 mM HEPES/ $\text{MeOH}$  = 3:7, pH = 7.0) at 298K.  $I$  is the fluorescence intensity after addition of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  (100  $\mu\text{M}$ ) and  $I_0$  is fluorescence intensity for free receptor.

indicated in Fig. 1b, like **L2**, neither **L3** nor **L1** exhibit a distinct fluorescence emission after addition of  $\text{Zn}^{2+}$  (10 equiv) in methanol-water (10 mM HEPES/ $\text{MeOH}$  = 7:3, pH = 7.0). These observations suggest that in methanol-water, the ligand **L2** was highly sensitive toward the  $\text{Zn}^{2+}$  ion. Fig. 2 shows the selective fluorescence enhancement after addition of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ion. As shown in fig. 2, receptor **L1** was more selective for  $\text{Cd}^{2+}$  ion than  $\text{Zn}^{2+}$  unlike receptors **L2** and **L3**. Fig. 3a reveals that the fluorescence emission intensity of **L2** become approximately 7 times greater than that of **L1** upon addition of 10 equiv. of  $\text{Zn}^{2+}$  and that ligand **L1** exhibits

greater fluorescence enhancement than does **L2** in the presence of  $\text{Cd}^{2+}$  ligand pres



To verify the fluorescence intensity changes in different solvents, fluorescence titration experiments and job's plot were carried out. Figure 4 illustrates a gradual enhancement of fluorescence upon the addition of  $\text{Zn}^{2+}$  in **L2** (7 $\mu\text{M}$ ) was observed at 406 nm when excited at 353 nm. The change was almost terminated after addition of 2 equiv. of  $\text{Zn}^{2+}$ , which suggested a 1:2 stoichiometry for the metal-ligand complex. This was again confirmed by the Job's plot analysis. The fluorescence intensity exhibited a maximum at the mole fraction 0.65, suggestive of 1:2 complexation. The association constant for the complexation of **L2** with  $\text{Zn}^{2+}$  was determined to be  $3.3 \times 10^4 \text{ M}^{-1}$  (Fig SI 31). Figure 5a shows the fluorescence titrations of  $\text{Zn}^{2+}$  with **L1** in THF. Stepwise addition of  $\text{Zn}^{2+}$  led to an increase of the fluorescence intensity until the complete addition of 1 equiv. of  $\text{Zn}^{2+}$ . To confirm the binding sites of the sensor, the stoichiometries of **L1** with  $\text{Zn}^{2+}$  were calculated using the Job's plot, for which there was a maximum at 0.5 mole fraction, indicative of a 1:1 stoichiometry. Figure 5b presents the change of the fluores




(a)

(b)

**Fig. 3** Fluorescence response of ligand **L1** and **L2** (7.0  $\mu\text{M}$ ) in (a)  $\text{CH}_3\text{OH}/\text{H}_2\text{O}$  (10mM HEPES/MeOH = 3:7, pH = 7.0) (b) THF solvent at 298 K after addition of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ion with excitation at 347 nm and 353 nm, respectively.

These results indicated that ligands **L2** and **L1** exhibit similar behaviour and binding toward  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ions in THF. The US Environmental Protection Agency (EPA) set the maximum contaminant levels of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  in drinking water at 7.6 and 4.5




<sup>a</sup>Measured at 27 °C by fluorescence titration experiments (Figure SI. 31–34); host concentration was 7  $\mu\text{M}$ .

to be highly selective for the detection of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  (Table 1). Figure 6a shows the selectivity among various metal ions. Probe **L2** exhibited high selectivity toward  $\text{Zn}^{2+}$  over ( $\text{Cu}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$ ,  $\text{Hg}^+$ ,

(a)

$\text{K}^+$ ,  $\text{Li}^+$  (as their perchlorate salts) and  $\text{Co}^{3+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Ni}^{2+}$  including  $\text{Cd}^{2+}$  (as nitrate salts). Therefore, the affinity of **L1** was observed with each of the respective metal cations and the results implied that **L1** can selectively detect both  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  ions, but with a slightly higher affinity for  $\text{Cd}^{2+}$  versus  $\text{Zn}^{2+}$ . Figure 6b reveals that **L1** and **L2** were more sensitive toward  $\text{Cd}^{2+}$  than  $\text{Zn}^{2+}$  when using THF as solvent. By contrast, the addition of other cations ( $\text{Cu}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$ ,  $\text{Hg}^+$ ,  $\text{K}^+$ ,  $\text{Li}^+$ ,  $\text{Co}^{3+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Na}^+$ ,  $\text{Li}^+$ ) showed almost no fluorescence enhancement. These results indicated that **L1** and **L2** exhibit selective emission enhancement toward  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  both in THF and methanol-water solvents. On the other hand, observations for the fluorescence emissions for the **L2** (7  $\mu\text{M}$ ) and  $\text{Zn}^{2+}$  (100  $\mu\text{M}$ ) system, indicated that most of the competitive cations such as  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$ ,  $\text{Hg}^+$ ,  $\text{K}^+$ ,  $\text{Li}^+$ ,  $\text{Co}^{3+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cd}^{2+}$  caused no obvious change at higher concentration (100 $\mu\text{M}$ ) (figure SI 22.). However,  $\text{Cu}^{2+}$ ,  $\text{Ag}^+$ ,  $\text{Hg}^+$  all strongly quenched the fluorescence in the **L2**+ $\text{Zn}^{2+}$  system. These results suggested that the co-ordination of  $\text{Zn}^{2+}$  with **L2** is more selective than other metal ions, with the exception of  $\text{Cu}^{2+}$ ,  $\text{Ag}^+$  and  $\text{Hg}^{2+}$ .




in  $\text{MeOH}/\text{H}_2\text{O}$  (10 mM HEPES/MeOH = 3:7, pH = 7.0) at 298 K with excitation at 353 nm.


The  $^1\text{H}$  NMR spectroscopic analysis of **L2** provided further evidence of the 1:1 and 1:2 binding mode in methanol-water and THF. In methanol-water, receptor **L2** is not fully soluble in the 3:7 mixture of  $\text{D}_2\text{O}/\text{CD}_3\text{OD}$ . Therefore, a 1:9 ratio of  $\text{D}_2\text{O}/\text{CD}_3\text{OD}$

(b)



Compd.	Solvent	Binding model, $\text{L}:\text{M}^{2+}$	$K_a$ ( $\text{M}^{-1}$ )
<b>L2</b> → $\text{Zn}^{2+}$	$\text{MeOH}-\text{H}_2\text{O}$	1:2	$3.3 \times 10^4$
<b>L2</b> → $\text{Zn}^{2+}$	THF	1:1	$6.6 \times 10^4$
<b>L1</b> → $\text{Zn}^{2+}$	THF	1:1	$5.0 \times 10^5$
<b>L2</b> → $\text{Cd}^{2+}$	THF	1:1	$5.0 \times 10^5$




(7  $\mu\text{M}$ ) (b) ligand **L2**(7  $\mu\text{M}$ ) in addition with  $\text{Zn}^{2+}$  at 298 K. The excitation was performed at 347 nm for **L1** and 353 nm for **L2**.



**Fig. 6** Fluorescence intensity changes of ligand **L1** and **L2** (7  $\mu\text{M}$ ) in (a)  $\text{CH}_3\text{OH}/\text{H}_2\text{O}$  (10 mM HEPES/MeOH = 3:7, pH = 7.0) and (b) THF solvent at 298 K after addition of various metal ions (100  $\mu\text{M}$ ).  $I$  is the fluorescence intensity after addition of metal ions and  $I_0$  is fluorescence intensity for free receptor.



**Fig. 7** Partial  $^1\text{H-NMR}$  titration of **L2**/guest (H/G = 1:2); (a) Free ligand **L2** ( $1.5 \times 10^{-2}$  M); (b) **L2** $\rightarrow$ **Cd** $^{2+}$  (1 equiv.); (c) **L2** $\rightarrow$ **Cd** $^{2+}$  (2 equiv.). Solvent: **L2** ( $0.5 \times 10^{-3}$  M); (b) **L2** $\rightarrow$ **Cd** $^{2+}$  (0.5 equiv.); (c) **L2** $\rightarrow$ **Cd** $^{2+}$  (1 equiv.); (d)  $\text{CD}_3\text{OD-D}_2\text{O}$  (9:1, v/v, pD = 7.0). 300 MHz at 298 K.



**Table 2.** Chemical shift of dipicolylamine and methylene protons of free **L2** and **L2** with  $\text{Zn}^{2+}$  or  $\text{Cd}^{2+}$ .

$\Delta\delta$  values are the difference of the chemical shift between **L2** and  $\text{Zn}^{2+}$  or  $\text{Cd}^{2+}$  at 27 °C. Here, minus sign (-) denotes a shift to higher magnetic field.

(pD = 7.0) was applied for these analysis. The  $^1\text{H}$  NMR signals reveal the aromatic and methylene regions of **L2** (Fig. 7 and Figure SI 28). After addition of 2 equiv. of  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$ , the proton signals of **L2**, when in the presence of  $\text{Zn}^{2+}$  ion, undergo a larger downfield shift than when the  $\text{Cd}^{2+}$  ion was present. Moreover, two sets of four methylene  $\text{H}_a$  protons were split into two peaks and broadened following binding with  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$ . The other proton

	Chemical Shift, $\delta_{\text{ppm}}$ in MeOH-H <sub>2</sub> O (H/G = 1:2)					Chemical Shift, $\delta_{\text{ppm}}$ in THF (H/G = 1:1)		
	Free <b>L2</b>	<b>L2</b> $\rightarrow$ <b>Cd</b> $^{2+}$	$\Delta\delta$	<b>L2</b> $\rightarrow$ <b>Zn</b> $^{2+}$	$\Delta\delta$	Free <b>L2</b>	<b>L2</b> $\rightarrow$ <b>Cd</b> $^{2+}$	$\Delta\delta$
$\text{H}_a$	3.85	3.80, 4.21	-0.05, 0.36	3.88, 4.36	0.03, 0.61	3.87	3.81, 4.16	-0.06, 0.29
$\text{H}_b$	4.28	4.68	0.40	4.64	0.36	4.39	4.25, 4.56	-0.14, 0.17
$\text{H}_c$	7.42	7.38	-0.06	7.49	0.07	7.48	7.23	-0.25
$\text{H}_d$	8.35	8.75	0.40	8.83	0.48	8.46	8.68	0.22

signals overlapped with each other among the four pyridine rings of the two sets of Dpas as does of pyrene ring protons and leads to a downfield shift which is due to the decrease of electron density by the metal-nitrogen co-ordination.<sup>21</sup> The  $\text{H}_d$  protons of adjacent pyrene rings underwent a significant downfield shift ( $\delta$  8.35 to 8.75 and 8.83 ppm) for  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  ions respectively. Furthermore, two sets of two methylene  $\text{H}_b$  protons also broadened and underwent a large downfield shift. These results suggested that two sets of dpas were equally assigned for making a co-ordination bond with two metal ions and confirmed a 1:2 metal-ligand stoichiometry.<sup>19</sup> The  $^1\text{H}$  NMR analysis also revealed larger chemical shift differences for **L2** for the complexation with  $\text{Zn}^{2+}$

versus the  $\text{Cd}^{2+}$  ion.

In contrast, when using THF as solvent, there is no such change after addition of 1 equiv. of  $\text{Cd}^{2+}$  ion which confirmed the 1:1 binding mode for the complexation of **L2** with  $\text{Cd}^{2+}$  (Fig. 8). Here, the same  $\text{H}_d$  protons of the adjacent four pyridine rings undergo a smaller downfield shift (from  $\delta$  8.46 to 8.68 ppm) than in methanol-water solvent after addition of 1 equiv. of  $\text{Cd}^{2+}$ . Another three protons ( $\text{H}_c$ ,  $\text{H}_e$  and  $\text{H}_f$ ) also experience a downfield shift. Moreover, two sets of four methylene  $\text{H}_a$  protons are split into two broad peaks from  $\delta$  3.87 to 3.81 and 4.16 ppm following binding with  $\text{Cd}^{2+}$  akin to the methanol-water system. On the other hand, the  $\text{H}_g$  proton of the pyrene ring exhibits a large upfield shift from  $\delta$  8.35 to 7.64 ppm, and unlike the methanol-water system, the  $\text{H}_b$  protons split into two peaks from  $\delta$  4.39 to 4.25 and 4.56 ppm, which suggested that the methylene  $\text{H}_b$  and pyrene  $\text{H}_g$  protons directly contribute to the binding with the metal ion. This phenomenon is only possible when the  $\text{Cd}^{2+}$  is positioned at the centre between the two binding sites. However, in THF, addition of  $\text{Zn}^{2+}$  induces vigorous precipitation which does not allow for analysis using  $^1\text{H}$  NMR spectra for elucidation of the binding mode. Moreover, the fluorescence spectra and Job's plot confirmed the 1:1 binding mode of a **L2** $\rightarrow$ **Zn** $^{2+}$  complex.

The above NMR and fluorescence spectra together with the Job's plot suggested that in methanol-water solvent system, two binding sites equally co-ordinate with two metal ions. On the other hand,  $\text{Zn}^{2+}$  or  $\text{Cd}^{2+}$  is positioned between two binding sites in THF. Given the shape of THF (a five membered ring), it can lead to a pronounced pseudorotational effect which is responsible for the stable twisted conformation. It is assumed that this structural property plays an important role in the 1:1 ligand to metal binding system.

## Conclusion

In conclusion, the novel fluorogenic molecules **L1** and **L2** based on 7-*tert*-butylpyrene have been synthesized. The binding of  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  ions at the pyrene linked dipicolylamine moieties was investigated by using fluorescence and  $^1\text{H}$  NMR titration experiments. It was found that receptor **L1** exhibits a similar binding toward  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  in both solvent systems. Herein, **L1** displayed higher fluorescence sensitivity for  $\text{Cd}^{2+}$  versus  $\text{Zn}^{2+}$ . On the other hand, receptor **L2** exhibited different binding behaviour in different solvent systems. When the molecule was dissolved in methanol-water solvent system, it selectively detected  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  with a 1:2 (ligand:metal) binding ratio. It was noticeable that **L2** had the strongest affinity for binding with  $\text{Zn}^{2+}$  ion versus  $\text{Cd}^{2+}$  and all the other competitive metal ions. In contrast, using THF as solvent,  $\text{Zn}^{2+}$  or  $\text{Cd}^{2+}$  is positioned between two binding sites and followed a 1:1

binding mode. It was concluded that ligands **L1** and **L2** exhibited similar binding behaviour in THF.

## Experimental Section

### General

**General :** Unless otherwise stated, all other reagents used were purchased from commercial sources and were used without further purification. Compounds **1**,<sup>25</sup> **3**,<sup>22</sup> **4**<sup>22</sup> and receptor **L3**<sup>19a</sup> were prepared following the reported procedures. All the solvents used were dried and distilled by the usual procedures before use. All melting points were determined using a Yanagimoto MP-S1. JEOL FT-300 NMR spectrometer and Varian-400MR-vmrs400 with SiMe<sub>4</sub> as an internal reference: *J*-values are given in Hz. UV-vis spectra were recorded using a Shimadzu UV-3150UV-vis-NIR spectrophotometer. Fluorescence spectroscopic studies of compounds in solution were performed in a semimicro fluorescence cell (Hellma®, 104F-QS, 10 × 4 mm, 1400 μL) with a Varian Cary Eclipse spectrophotometer. Mass spectra were obtained with a Nippon Denshi JMS-HX110A Ultrahigh Performance mass spectrometer at 75 eV by using a direct-inlet system.

### Synthesis of Compound 2

To 7-*tert*-butylpyrene (500 mg, 1.93 mmol), 1,1-dichloromethyl methyl ether (333 mg, 2.90 mmol) was added in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) at 0 °C with stirring for 15 min. A solution of TiCl<sub>4</sub> (0.53 ml, 4.8 mmol) was added drop wise to the stirred solution over 10 min. After this addition, the reaction mixture was continuously stirred for 3 h at room temperature. Then, the reaction mixture was quenched with ice water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 50 mL). The organic layer was washed with water (2 × 200 mL), brine (2 × 200 mL), dried over MgSO<sub>4</sub> and then evaporated. The crude product was recrystallized from hexane to obtain 7-*tert*-butylpyrene-1-carbaldehyde **2** as a yellow powder (400 mg, 72 %). The <sup>1</sup>H NMR spectrum agreed with the reported values.<sup>23</sup> <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 1.60 (9H, s, *t*Bu), 8.06 (1H, d, *J* = 7.83 Hz, pyrene-*H*<sub>5</sub>), 8.20 (1H, d, *J* = 4.83 Hz, pyrene-*H*<sub>4</sub>), 8.23 (1H, d, *J* = 3.8 Hz, pyrene-*H*<sub>9</sub>), 8.29 (1H, d, *J* = 9.2 Hz, pyrene-*H*<sub>10</sub>), 8.34 (2H, d, *J* = 3.2 Hz, pyrene-*H*<sub>6,8</sub>), 8.39 (1H, d, *J* = 7.9 Hz, pyrene-*H*<sub>3</sub>), 9.38 (1H, d, *J* = 9.0 Hz, pyrene-*H*<sub>2</sub>) and 10.78 (1H, s, *CHO*) ppm.

### Synthesis of Receptor L2

To a solution of 7-*tert*-butylpyrene-1,3-dicarbaldehyde (336 mg, 1.07 mmol) in 1,2-dichloroethane (18 mL), 2,2'-dipicolylamine (436 mg, 2.18 mmol) was added drop wise. Then the mixture was stirred for 18 h at 45 °C. After that, sodium triacetoxyborohydride (1.35 g, 6.42 mmol) was added, and the mixture was further stirred for 24 h at 50°C. Then, the reaction mixture was quenched with ice water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 × 100 mL). The organic layer was washed with water (2 × 200 mL), brine (2 × 200 mL), dried over MgSO<sub>4</sub> and then evaporated. The crude product was purified by column chromatography eluting with acetone-methanol (1:1) to afford an orange gummy substance **L2** (600 mg, 82 %). Mp. 65–66 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 1.55 (9H, s, *t*Bu), 3.82 (8H, s, *CH*<sub>2</sub>),

4.36 (4H, s, *CH*<sub>2</sub>), 7.12–7.07 (4H, m, pyridine-*H*), 7.45 (4H, d, *J* = 7.8 Hz, pyridine-*H*), 7.56 (4H, ddd, *J* = 1.8, 7.8, 15.2 Hz, pyridine-*H*), 7.99 (2H, d, *J* = 9.3 Hz, pyrene-*H*<sub>4,10</sub>), 8.18 (2H, s, pyrene-*H*<sub>6,8</sub>), 8.21 (1H, s, pyrene-*H*<sub>2</sub>), 8.29 (2H, d, *J* = 9.2 Hz, pyrene-*H*<sub>5,9</sub>) and 8.52–8.50 (4H, m, pyridine-*H*) ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ = 31.9, 35.1, 57.1, 60.5, 121.9, 122.2, 123.1, 123.3, 123.9, 125.4, 126.9, 129.2, 130.2, 130.8, 131.7, 136.3, 148.8, 148.9 and 159.6 ppm; HRMS: *m/z* calcd. for C<sub>46</sub>H<sub>44</sub>N<sub>6</sub> 681.3706; found 681.3707 [M<sup>+</sup>].

### Synthesis of receptor L1

To a solution of 7-*tert*-butylpyrene-1-carbaldehyde (225 mg, 0.79 mmol) in 1,2-dichloroethane (18 mL), 2,2'-dipicolylamine (156 mg, 0.79 mmol) was added drop wise. Then the mixture was stirred for 18 h at 45°C. After that, sodium triacetoxyborohydride (500 mg, 2.36 mmol) was added, and the mixture was further stirred for 24 h at 50 °C. The reaction mixture was quenched with ice water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 × 100 mL). The organic layer was washed with water (2 × 200 mL), brine (2 × 200 mL), dried over MgSO<sub>4</sub> and then evaporated. The crude product was purified by column chromatography eluting with ethyl acetate-hexane (3:1) to afford a yellow solid (230 mg, 62 %). Mp: 134–135 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 1.58 (9H, s, *t*Bu), 3.92 (4H, s, *CH*<sub>2</sub>), 4.39 (2H, s, *CH*<sub>2</sub>), 7.14–7.09 (2H, m, pyridine-*H*), 7.47 (2H, d, *J* = 7.8 Hz, pyridine-*H*), 7.60 (2H, ddd, *J* = 1.74, 7.7, 15.2 Hz, pyridine-*H*), 7.96 (2H, s, pyrene-*H*<sub>9,10</sub>), 8.04 (1H, d, *J* = 9.33 Hz, pyrene-*H*<sub>3</sub>), 8.07 (2H, s, pyrene-*H*<sub>4,5</sub>), 8.19 (2H, dd, *J* = 1.7, 6.3 Hz, pyrene-*H*<sub>7,8</sub>), 8.33 (1H, d, *J* = 9.2 Hz, pyrene-*H*<sub>2</sub>) and 8.53 (2H, d, *J* = 4.9 Hz, pyridine-*H*) ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ = 31.9, 35.2, 57.1, 60.4, 122.0, 122.1, 122.3, 122.9, 123.3, 123.9, 124.3, 124.9, 127.2, 127.3, 127.9, 129.6, 130.5, 130.6, 131.1, 132.3, 136.4, 148.8, 148.9 and 159.6 ppm. HRMS: *m/z* calcd. for C<sub>33</sub>H<sub>31</sub>N<sub>3</sub> 470.2596; found 470.2596 [M<sup>+</sup>].

### Determination of the Association Constants

The association constants were determined by using the fluorescent titration experiment of **L1** and **L2** in a constant concentration of host receptor (7 × 10<sup>-6</sup> M) and varying the guest concentration (0–20 × 10<sup>-6</sup> M). The association constant (*K*<sub>a</sub>) for the complexes of receptor **L1** and **L2** were calculated by observing the integral intensities of the complex and of free host molecules using nonlinear curve-fitting analysis according to the literature procedure.<sup>24</sup>

### <sup>1</sup>H NMR Titration Experiments

A solution of Zn(ClO<sub>4</sub>)<sub>6</sub>·6H<sub>2</sub>O or Cd(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O in D<sub>2</sub>O (1.5 × 10<sup>-2</sup> M) was added to a CD<sub>3</sub>OD–D<sub>2</sub>O (11:1, v/v) solution of receptor **L2** in the absence or presence of Zn<sup>2+</sup> and Cd<sup>2+</sup> ion in an NMR tube (300 MHz NMR). Similarly, a solution of Zn(ClO<sub>4</sub>)<sub>6</sub>·6H<sub>2</sub>O or Cd(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O in THF-*d*<sub>8</sub> (0.5 × 10<sup>-3</sup> M) was added to a THF-*d*<sub>8</sub> solution of **L2** (400 MHz NMR). <sup>1</sup>H NMR spectra were recorded after addition of the reactants and the temperature of the NMR probe was kept constant at 27 °C.

**Supporting information:** Detailed fluorescence and  $^1\text{H}$  NMR titration data.

#### Acknowledgements

This work was performed under the Cooperative Research Program of “Network Joint Research Center for Materials and Devices (Institute for Materials Chemistry and Engineering, Kyushu University)”. We would like to thank the OTEC at Saga University and the International Cooperation Projects of Guizhou Province (No. 20137005) for financial support. The EPSRC is thanked for an overseas travel grant (to CR).

#### Notes and references

<sup>a</sup> Department of Applied Chemistry, Faculty of Science and Engineering, Saga University, Honjo-machi 1, Saga 840-8502 Japan, E-mail: [yamatot@cc.saga-u.ac.jp](mailto:yamatot@cc.saga-u.ac.jp)

<sup>b</sup> Department of Chemistry, The University of Hull, HU6 7RX, UK.

† Electronic Supplementary Information (ESI) available: Details of the  $^1\text{H}/^{13}\text{C}$  NMR spectra,  $^1\text{H}$  NMR spectroscopic and UV-vis titration experimental data, the Bensei-Hilderbrand plot and Job's plot, See DOI: 10.1039/b000000x/

- (a) V. Amendola, L. Fabbrizzi, F. Forti and M. Liccheli, *Coord. Chem. Rev.*, 2006, **250**, 273–299; (b) A. Ojida, H. Nanoka, Y. Miyahara, S. Tamaura, K. Sada and I. Hamachi, *Angew. Chem. Int. Ed.*, 2006, **45**, 5518–5521.
- J. R. Lakowicz, *Principles of Fluorescence Spectroscopy*, 2<sup>nd</sup> ed.; Kulwer Academic: New York, 1999.
- E. M. Goldys, Ed. *Fluorescence and applications in biotechnology and life sciences*; Wiley Blackwell: New Jersey, 2009.
- B. N. G. Giepmans, S. R. Adams, M. H. Ellisman and R. Y. Tsien, *Science*, 2006, **312**, 217–224.
- A. Baker, *Environ. Sci. Technol.*, 2001, **35**, 948–953.
- B. Valeur, *Molecular Fluorescence Principles and Applications*, Wiley-VCH, New York, 2002.
- M. Sauer, *Angew. Chem., Int. Ed.*, 2003, **42**, 1790–1793.
- (a) K. Hanaoka, Y. Muramatsu, Y. Urano, T. Terai and T. Nagano, *Chem.-Eur. J.*, 2010, **16**, 568–572; (b) Y. Chen, C. Zhu, Z. Yang, J. Li, Y. Chiao, W. He, J. Chen and Z. Guo, *Chem. Commun.*, 2012, **48**, 5094–5096; (c) J. Wu, W. Liu, J. Ge, H. Zhang and P. Wang, *Chem. Soc. Rev.*, 2011, **40**, 3483–3495.
- (a) D. A. Pearce, N. Jotterand, I. S. Carrico and B. Imperiali, *J. Am. Chem. Soc.*, 2001, **123**, 5160–5161; (b) M. Mameli, M. C. Aragoni, M. Arca, C. Caltagirone, F. Demartin, G. Farruggia, G. D. Filippio, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis, S. Murgia, L. Prodi, A. Pintus and N. Zaccheroni, *Chem.-Eur. J.*, 2010, **16**, 919–930; (c) Y. Bao, B. Liu, H. Wang, F. Du and R. Bai, *Anal. Methods*, 2011, **3**, 1274–1276; (d) Y. Bao, B. Liu, F. Du, J. Tian, H. Wang and R. Bai, *J. Mater. Chem.*, 2012, **22**, 5291–5294.
- R. J. Wandell, A. H. Younes and L. Zhu, *New J. Chem.*, 2010, **34**, 2176–2182.
- L. Zhang, C. S. Murphy, G. -C. Kuang, K. L. Hazelwood, M. H. Constantino, M. W. Davidson and L. Zhu, *Chem. Commun.*, 2009, **47**, 7408–7410.
- B. L. Vallee and H. K. Falchuk, *Physiol. Rev.*, 1993, **73**, 79–118.
- S. Küry, B. Dréno, S. Bézieau, S. Giraudet, M. Kharfi, R. Kamoun and J. -P. Moisan, *Nat. Genet.* 2002, **31**, 239–240.
- (a) M. J. Berg, Y. Shi, *Science*, 1996, **271**, 1081–1085; (b) E. Manandhar, H. Broome, J. Myrick, W. Lagrone, P. J. Cragg and K. J. Wallace, *Chem. Commun.*, 2011, **47**, 8796–8798.
- S. Satarug, R. J. Baker, S. Urbenzapol, M. Haswell-Elkins, B. E. P. Reilly, J. D. Williams and R. M. Moore, *Toxicol. Lett.*, 2003, **137**, 65–83.
- K. Rurack, U. Resch-Genger and W. Rettig, *J. Photochem. Photobiol. A. Chem.*, 1998, **118**, 143–149.
- (a) Z. Xu, N. J. Singh, J. Lim, J. Pan, H. N. Kim, S. Park, K. S. Kim and J. Yoon, *J. Am. Chem. Soc.*, 2009, **131**, 15528–15533; (b) Y. Yanga, X. Gou, J. Blecha and H. Cao, *Tetrahedron Lett.*, 2010, **51**, 3422–3425.
- K. Fujimoto, S. Yamada and M. Inouye, *Chem. Commun.*, 2009, **46**, 7164–7166.
- (a) A. Ojida, Y. Mito-Oka, M. Inoue and I. Hamachi, *J. Am. Chem. Soc.*, 2002, **124**, 6256–6258; (b) A. Ojida, Y. Mito-Oka, M. Inoue and I. Hamachi, *J. Am. Chem. Soc.*, 2004, **126**, 2454–2463.
- K. Baek, M. S. Eom, S. Kim and M. S. Han, *Tetrahedron Lett.*, 2013, **54**, 1654–1657.
- S. Sumiya, Y. Shirashi and T. Hiarai, *J. Phys. Chem.*, 2013, **117**, 1474–1482.
- J. Inoue, K. Fukuzi, T. Kubo, S. Nakazawa, K. Sato, D. Shiomi, Y. Morita, K. Yamamoto, T. Takui and K. Nakasuji, *J. Am. Chem. Soc.*, 2001, **123**, 12702–12703.
- G. Liu, Z. Li, D. Wu, W. Xue, T. Li, S. H. Liu and J. Yin, *J. Org. Chem.*, 2014, **79**, 643–652.
- H. A. Benesi and J. H. Hildebrand, *J. Am. Chem. Soc.*, 1949, **71**, 2703–2707.
- T. M. Figueira-Duarte, S. C. Simon, M. Wagner, S. I. Druzhinin, K. A. Zachariasse and K. Mullen, *Angew. Chem. Int. Ed.*, 2008, **47**, 10175–10178.



