

1 **Managed realignment for habitat compensation: use of a new intertidal**
2 **habitat by fishes**

3

4 A. D. NUNN^{1,*}, D. CLIFTON-DEY² and I. G. COWX¹

5

6 ¹*Hull International Fisheries Institute, School of Biological, Biomedical &*
7 *Environmental Sciences, University of Hull, Hull, HU6 7RX, UK.*

8 ²*Environment Agency, Kings Meadow House, Kings Meadow Road, Reading, RG1*
9 *8DQ, UK.*

10

11 Running headline: Use of a new intertidal habitat by fishes

12

13 Key words: climate change; coastal squeeze; flood defence; habitat creation; habitat
14 loss; mudflat development.

15

16 **ABSTRACT**

17 Managed realignment has become an increasingly common mechanism to increase the
18 efficiency and sustainability of flood defences, reduce defence costs or compensate for
19 habitat losses. This study investigated the use by fishes of a new intertidal habitat,
20 created by managed realignment, intended to compensate for the loss of mudflat
21 associated with a major port development. Although broadly similar, statistically
22 significant differences in fish species composition, abundance, biomass, size structure,
23 diversity and diet composition indicate that the managed realignment is not yet

*Author to whom correspondence should be addressed. Tel.: +44 (0)1482 466429; fax: +44 (0)1482 465458; email: a.d.nunn@hull.ac.uk

24 functioning in an identical manner to the mudflat in the adjacent estuary, most likely
25 due to differences in habitat between sites. Notwithstanding, similarity in the species
26 composition of fyke catches in the managed realignment and estuary increased annually
27 during the 5-year study period, suggesting that the mudflat in the realignment is still
28 developing. Indeed, the site will inevitably change over time with accretion,
29 establishment of vegetation and possibly development of creeks. This will not
30 necessarily prevent the aim of the realignment scheme being achieved, as long as
31 sufficient suitable mudflat remains.

32

33 INTRODUCTION

34 Intertidal habitats support high biological productivity (McLusky *et al.*, 1992;
35 Ysebaert *et al.*, 2003), contribute to flood defence (Dixon *et al.*, 1998) and provide
36 important habitats for fishes (Elliott *et al.*, 2007; Ramos *et al.*, 2012) and birds
37 (Atkinson *et al.*, 2004; Mander *et al.*, 2007). Many intertidal areas, however, are
38 subjected to a range of anthropogenic pressures. Of particular importance is land
39 claim for industrial development (McLusky *et al.*, 1992; Esteves, 2014). Land claim
40 can have direct negative impacts on intertidal biota, and profound implications for
41 ecosystem functioning through the role of the biological communities in sediment
42 dynamics, biogeochemical cycling, benthic metabolism and trophic interactions
43 (Herringshaw & Solan, 2008). Loss of intertidal areas can also increase the risk of
44 flooding, which is likely to be exacerbated by the effects of climate change, especially
45 in areas already experiencing coastal squeeze (Mazik *et al.*, 2007; Pontee, 2013;
46 Esteves, 2014). It is therefore desirable, sometimes necessary, to compensate for
47 habitat losses due to land claim, especially those predicted to compromise the
48 integrity of designated conservation areas (Morris, 2013; Esteves, 2014).

49

50 Managed realignment – the deliberate process of realigning river, estuary or coastal
51 flood defences – has become an increasingly common mechanism to increase the
52 efficiency and sustainability of flood defences, reduce defence costs or compensate
53 for habitat losses (e.g. Ledoux *et al.*, 2005; Garbutt *et al.*, 2006; Mazik *et al.*, 2007;
54 Rupp-Armstrong & Nicholls, 2007; Shih & Nicholls, 2007; Esteves, 2013; Morris,
55 2013; Pétillon *et al.*, 2014). Managed realignment also has the potential to enhance
56 fish diversity, recruitment and production by increasing the availability and diversity
57 of intertidal habitats, such as mudflats and salt marshes (Dixon *et al.*, 1998; Colclough

58 *et al.*, 2005; French, 2006). It is essential, however, that the physical characteristics
59 and biological communities of managed realignments replicate those being lost if
60 habitat compensation is to be truly successful (Mazik *et al.*, 2010).

61

62 A port and logistics centre is being developed on the north bank of the Thames
63 Estuary, England. The development includes a container terminal to accommodate the
64 largest deep-sea container ships, and was considered likely to have an adverse impact
65 on the integrity of the Thames Estuary and Marshes Special Protection Area (SPA)
66 and Ramsar Site. Predicted direct impacts of the development on physical habitats
67 included: (1) conversion of 5 ha of designated intertidal habitat to shallow subtidal
68 habitat; (2) destruction of 25 ha of undesignated intertidal habitat; (3) changes in
69 accretion over 60 ha of intertidal habitat, potentially converting 10 ha of mudflat to
70 saltmarsh; (4) long-term impacts on 90 ha of subtidal habitat affected by capital
71 dredging; and (5) temporary damage to >1700 ha of subtidal habitat outside the SPA
72 and Ramsar Site (Morris & Gibson, 2007). To compensate for part of the impacts on
73 the Thames Estuary and Marshes SPA and Ramsar Site and ensure the overall
74 coherence of the Natura 2000 network is maintained, a minimum of 74 ha of new
75 intertidal mudflat is being created through managed realignment (Morris & Gibson,
76 2007). Habitat creation and improvement of flood defences are common objectives of
77 managed realignment schemes (French, 2006; Esteves, 2013), but few studies have
78 assessed their use by fishes (e.g. Colclough *et al.*, 2005). The aim of this study was to
79 advance the understanding of the use by fishes of intertidal habitats created through
80 managed realignment by investigating changes over a 5-year period. The hypothesis
81 was that the species composition, size structure, abundance, biomass and diet
82 composition of fishes in the realignment and adjacent estuary would increase in

83 similarity as the mudflat in the realignment developed. High similarities in these
84 parameters in the two sites should suggest that the realignment is functioning in a
85 similar manner to the mudflat in the adjacent estuary, and that the aim of the
86 realignment scheme, namely to compensate for losses of mudflat associated with port
87 development, is being achieved (*cf.* Mazik *et al.*, 2007, 2010; Mossman *et al.*, 2012).

88

89 **METHODOLOGY**

90

91 **Sampling strategy, methods and techniques**

92 London Gateway Site A managed realignment (51.50232 °N, 0.44799 °E; also
93 known as Stanford Wharf Nature Reserve) is located to the east of Mucking Creek,
94 near Stanford-le-Hope, on the north bank of the Thames Estuary, England. The site
95 was created in 2010 by reducing the level of 27 ha of former agricultural land and
96 creating a 300-m-wide breach in the sea defences to the south. Fish surveys were
97 conducted during spring tides in October and November 2010 and April, June and
98 August 2011-2014. These timeframes coincide with the larval and juvenile periods of
99 many fishes, thus enabling assessment of the function of the habitat (e.g. nursery) for
100 specific species (*cf.* Nunn *et al.*, 2007). The sampling frequency therefore accounts for
101 temporal variations in fish community structure associated with the phenology of fish
102 hatching and ontogenetic and seasonal shifts in habitat use. A combination of active
103 (seine, epibenthic trawl) and passive (fyke) gear types with replicated sampling
104 stations was included in the design, to provide as accurate an assessment as possible
105 of the species composition, size structure, density and biomass of fishes in the
106 realignment and adjacent estuary (immediately to the east of the realignment); using a
107 range of methods at fixed stations in a seasonal format is recommended to obtain a

108 robust assessment of intertidal fish communities (Colclough *et al.*, 2005). Gear types
109 were selected based on the potential operational constraints imposed by realignment
110 sites (e.g. deep mud, benthic obstructions, semi-permanent flooding regimes, deep
111 creeks) and the usual development of newly created intertidal areas (e.g. accretion,
112 establishment of vegetation). Fine-meshed gears were employed due to the expected
113 dominance of small-sized species or individuals in the fish assemblages using newly
114 created intertidal areas. Multi-method approaches, recognised as European best
115 practice (Hemingway & Elliott, 2002), have been successfully employed elsewhere to
116 examine the use of intertidal areas by fishes, including in managed realignments, and
117 as a tool for assessing the ecological status of estuaries (e.g. Laffaille *et al.*, 2000;
118 Colclough *et al.*, 2002, 2005; Coates *et al.*, 2007). Up to 50 individuals of each fish
119 species were measured (total length, L_T , mm) and weighed (0.01 g) for each sample,
120 with the remainder identified and counted. There were no significant differences in
121 water temperature (paired t -test, d.f. = 13, $t = 0.929$, $P = 0.370$) or salinity (paired t -
122 test, d.f. = 11, $t = 0.150$, $P = 0.884$), recorded at 15-minute intervals using an Aqua
123 TROLL 200 data logger, in the realignment and adjacent estuary.

124

125 ***Fyke netting***

126 Fykes were deployed at four stations in the realignment and two in the estuary, and
127 left for one tidal cycle. The nets were emptied as they became exposed by the
128 receding tide and then left for another tidal cycle, thereby allowing separate analysis
129 of diurnal and nocturnal catches (total $n = 180$). Each gear consisted of two fykes (53-
130 cm entrance, 10-m central panel, 14-mm mesh) joined entrance-to-entrance by their
131 leader panels; data from each gear were expressed as the abundance and biomass of
132 fishes per 'fyke-hour' (i.e. the number of hours that the gear was inundated). Fykes

133 were set at the same shore height in the realignment and estuary to ensure they
134 sampled comparable water depths, allowing an assessment of the larger fishes using
135 the area (Colclough *et al.*, 2005).

136

137 ***Seine netting***

138 A micromesh beach seine (25-m long, 3-m deep, 3-mm hexagonal mesh) was set at
139 eight stations in the realignment and two in the estuary; data from each sample (total n
140 = 150) were expressed as the abundance and biomass of fishes per m^2 . The area
141 sampled by the seine was calculated from direct *in situ* measurements (i.e. length \times
142 width of the area enclosed by the net). This method allowed an assessment of the
143 smaller fishes using the area (Cowx *et al.*, 2001; Colclough *et al.*, 2002, 2005; Coates
144 *et al.*, 2007).

145

146 ***Trawling***

147 Trawling was conducted using an epibenthic sledge fitted with a tickle chain and a
148 0.5-mm-meshed cod-end (Nitex cloth), to target benthic species and individuals for
149 which the fyke mesh was too large (Colclough *et al.*, 2002, 2005; Coates *et al.*, 2007).

150 The trawl was pulled by hand at $\sim 1 \text{ m s}^{-1}$; data from each sample (total $n = 135$) were
151 expressed as the abundance and biomass of fishes per m^2 . The area sampled by the
152 trawl was calculated by multiplying the width of the trawl entrance (1 m) by the
153 length of each transect (20 m). Three replicates were collected at each of three stations
154 in the realignment (nine trawls in total); trawling was not conducted in the estuary due
155 to safety issues.

156

157 **Data analysis**

158 The relative abundance of each fish species in the managed realignment and the
159 estuary was calculated for the entire study period and each gear type. Bray-Curtis
160 similarity matrices (Bray & Curtis, 1957) were calculated using the relative
161 abundance of each fish species and ordinated using non-metric multidimensional
162 scaling (MDS) to investigate similarities in the species composition of fyke and seine
163 catches in the realignment and estuary. The matrices were then submitted to
164 permutational multivariate analysis of variance (PERMANOVA) (9999 random
165 permutations) to assess the statistical significance of any differences in the species
166 composition of fyke and seine catches in the realignment and estuary (Anderson,
167 2001; Anderson *et al.*, 2008). In addition, similarity percentages (SIMPER) analysis
168 was used to calculate the percentage contributions of key fish species to dissimilarities
169 in fyke and seine catches in the realignment and estuary (Clarke & Warwick, 2001).
170 Mean Shannon-Wiener diversity (H') and Pielou's evenness (J) were compared for
171 fyke and seine catches in the realignment and estuary using independent samples t -
172 tests (Washington, 1984).

173

174 Mean lengths of the most abundant fish species were compared for fyke and seine
175 catches in the realignment and estuary, and diurnal and nocturnal fyke catches, using
176 independent samples t -tests (Dytham, 2003). Length distributions of the most
177 abundant species were compared for fyke and seine catches in the realignment and
178 estuary, and diurnal and nocturnal fyke catches, using two-sample Kolmogorov-
179 Smirnov tests (Dytham, 2003). For seine and trawl catches, the density (fish m^{-2}) and
180 biomass (g m^{-2}) of fishes in each sample were calculated by dividing their abundance
181 and biomass, respectively, by the area sampled. For fyke catches, abundance and
182 biomass were expressed, respectively, as catch-per-unit-effort (CPUE; fish h^{-1}) and

183 biomass-per-unit-effort (BPUE; g h^{-1}). Mean densities, biomasses, CPUE and BPUE
184 were compared between the realignment and estuary, and diurnal and nocturnal fyke
185 catches, using independent samples *t*-tests (Dytham, 2003).

186

187 For each sampling occasion, the stomach contents were removed from a sample of
188 juvenile bass (*Dicentrarchus labrax* (L.)) ($n = 139$, realignment L_T range 14-103 mm,
189 estuary L_T range 17-110 mm) and common goby (*Pomatoschistus microps* (Krøyer))
190 ($n = 167$, realignment L_T range 11-51 mm, estuary L_T range 11-46 mm) captured in
191 the realignment and estuary. Catches of other species were insufficient for a
192 comparison of diet composition in the realignment and estuary in all 5 years. Prey
193 items were identified to the lowest practicable taxonomic level and recorded as
194 percent volume. The diet composition of the most abundant fish species in the
195 realignment and estuary was then compared using PERMANOVA and SIMPER
196 analysis, as described for fish composition.

197

198 **RESULTS**

199 A total of 39 376 specimens of 16 fish species was captured during the study.
200 Common goby was the most abundant species, accounting for 62% of the total catch,
201 followed by herring (*Clupea harengus* L.) (24%). Other species captured were bass,
202 eel (*Anguilla anguilla* (L.)), flounder (*Platichthys flesus* (L.)), plaice (*Pleuronectes*
203 *platessa* L.), sand goby (*Pomatoschistus minutus* (Pallas)), sand smelt (*Atherina*
204 *presbyter* Cuvier), smelt (*Osmerus eperlanus* (L.)), sole (*Solea solea* (L.)), sprat
205 (*Sprattus sprattus* (L.)), ten-spined stickleback (*Pungitius pungitius* (L.)), thick-lipped
206 grey mullet (*Chelon labrosus* (Risso)), thin-lipped grey mullet (*Liza ramada* (Risso)),

207 three-spined stickleback (*Gasterosteus aculeatus* L.) and whiting (*Merlangius*
208 *merlangus* (L.)).

209

210 **Species composition**

211 There was a significant difference in the species composition of fyke catches in the
212 realignment and estuary (Fig. 1; PERMANOVA, d.f. = 1, $F = 5.277$, $P < 0.001$).

213 Catches in both sites were dominated by bass and flounder (76% in the realignment,
214 66% in the estuary), but the relative abundances of bass and eel were higher in the
215 realignment, whereas those of flounder, smelt and sole were higher in the estuary

216 (Table 1). Notwithstanding, similarity between the realignment and estuary increased
217 annually during the study period, from 29% in 2010 to 43% in 2014 (2011 = 33%,

218 2012 = 34%, 2013 = 41%). There was no significant difference in species
219 composition between years (PERMANOVA, d.f. = 4, $F = 1.801$, $P = 0.120$), and there

220 was no significant interaction between site and year (PERMANOVA, d.f. = 4, $F =$
221 0.854 , $P = 0.604$). Although the relative abundances of bass and flounder were


222 highest during daylight and those of eel and sole were highest at night (Table 2), there
223 were no statistically significant differences in the species composition of diurnal and

224 nocturnal fyke catches in the realignment (PERMANOVA, d.f. = 1, $F = 0.623$, $P =$
225 0.718) or estuary (PERMANOVA, d.f. = 1, $F = 1.646$, $P = 0.188$). Over the 5-year

226 study period, the mean diversity of fyke catches was significantly higher in the
227 estuary than the realignment (independent samples t -test, d.f. = 86, $t = 3.252$, $P =$

228 0.002), but there was no significant difference in evenness (independent samples t -
229 test, d.f. = 86, $t = 1.756$, $P = 0.083$).

230



231

232 **Fig. 1.** Non-metric multidimensional scaling (MDS) ordination plot comparing the
 233 fish species composition of fyke catches (2010-2014 centroids with trajectories) in the
 234 managed realignment and estuary.

235

236 **Table 1.** Similarity percentages (SIMPER) analysis of the mean relative abundances
 237 of key fish species and their percentage contributions to dissimilarities in fyke and
 238 seine catches in the managed realignment (R) and estuary (E).

Species	Fyke			Species	Seine		
	R	E	%		R	E	%
Bass	47.9	18.2	32.6	Bass	24.3	37.9	27.0
Flounder	27.7	47.6	27.7	Common goby	32.0	30.3	26.7
Eel	16.4	9.5	14.8	Herring	17.2	8.4	15.3
Smelt	4.2	15.2	13.2	Three-spined stickleback	8.0	8.1	9.8
Sole	0.5	6.6	6.0	Thin-lipped grey mullet	7.2	7.8	8.8
				Flounder	2.4	4.9	4.7

Mean **56.3** **Mean** **73.3**
dissimilarity **dissimilarity**

239


240 **Table 2.** Similarity percentages (SIMPER) analysis of the mean relative abundances
 241 of key fish species and their percentage contributions to dissimilarities in diurnal (D)
 242 and nocturnal (N) fyke catches in the managed realignment and estuary.

Species	Realignment			Species	Estuary		
	D	N	%		D	N	%
Bass	52.4	35.2	34.9	Flounder	52.2	49.5	32.2
Flounder	28.5	32.2	28.2	Bass	20.0	14.1	21.8
Eel	11.9	25.6	24.7	Smelt	12.2	13.1	16.6
Smelt	4.1	3.6	6.3	Eel	8.8	10.1	13.1
				Sole	4.8	11.5	13.0
Mean			53.9	Mean			53.4
dissimilarity				dissimilarity			

243

244 In seine catches, the relative abundance of herring was highest in the realignment
 245 whereas those of bass and flounder were highest in the estuary (Table 1), but there
 246 were no statistically significant differences in species composition between sites
 247 (PERMANOVA, d.f. = 1, $F = 1.341$, $P = 0.240$) or years (PERMANOVA, d.f. = 4, F
 248 = 0.820, $P = 0.660$) (Fig. 2); there was also no significant difference in the
 249 composition of trawl catches between years (PERMANOVA, d.f. = 4, $F = 1.237$, $P =$
 250 0.353). Over the 5-year study period, there were no significant differences in the mean
 251 diversity (independent samples t -test, d.f. = 130, $t = 1.318$, $P = 0.190$) or evenness

252 (independent samples *t*-test, d.f. = 130, *t* = 1.271, *P* = 0.206) of seine catches in the
253 realignment and estuary.
254



255
256 **Fig. 2.** Non-metric multidimensional scaling (MDS) ordination plot comparing the
257 fish species composition of seine catches (2010-2014 centroids with trajectories) in
258 the managed realignment and estuary.

259

260 **Size structure**


261 Overall, the mean lengths of bass (independent samples *t*-test, d.f. = 860, *t* = 4.875,
262 *P*<0.001) and flounder (independent samples *t*-test, d.f. = 281, *t* = 7.202, *P*<0.001) in
263 fyke catches and common goby (independent samples *t*-test, d.f. = 1102, *t* = 14.016,
264 *P*<0.001) in seine catches were significantly larger in the realignment than the
265 estuary, whereas bass in seine catches were larger in the estuary (independent samples
266 *t*-test, d.f. = 1183, *t* = 9.015, *P*<0.001). In addition, bass (independent samples *t*-test,
267 d.f. = 706, *t* = 2.056, *P* = 0.040) and eel (independent samples *t*-test, d.f. = 118, *t* =
268 2.030, *P* = 0.045) in fyke catches in the realignment were significantly larger during

269 daylight than at night, but there were no other diel differences in the mean lengths of
270 bass, eel and flounder in the realignment or estuary (independent samples *t*-tests, all
271 $P > 0.05$).

272

273 Modes representing the 0+ age class were present in the length distributions of bass,
274 common goby, flounder and herring in all years, with juveniles of most other species
275 also caught in some years. Overall, there were significant differences in the length
276 distributions of bass (two-sample Kolmogorov-Smirnov test, $Z = 3.388$, $P < 0.001$) and
277 flounder (two-sample Kolmogorov-Smirnov test, $Z = 4.350$, $P < 0.001$) in fyke catches
278 and bass (two-sample Kolmogorov-Smirnov test, $Z = 6.509$, $P < 0.001$) and common
279 goby (two-sample Kolmogorov-Smirnov test, $Z = 5.653$, $P < 0.001$) in seine catches in
280 the realignment and the estuary (Fig. 3), and also of bass (two-sample Kolmogorov-
281 Smirnov test, $Z = 1.849$, $P = 0.002$) and flounder (two-sample Kolmogorov-Smirnov
282 test, $Z = 1.390$, $P = 0.042$) in fyke catches in the realignment during the day and at
283 night. Data were insufficient for between-site and diel comparisons of length
284 distributions for other species.

285



286

287 **Fig. 3.** Length distributions of (a) bass and flounder in fyke catches and (b) bass and
 288 common goby in seine catches in the managed realignment and estuary.


289

290 **Abundance and biomass**

291 With the exceptions of BPUE in 2010 and density and CPUE in 2014, mean annual
 292 catches were always highest in the realignment (Fig. 4), and overall mean densities

293 (independent samples *t*-test, d.f. = 126, *t* = 2.327, *P* = 0.022), biomasses (independent
 294 samples *t*-test, d.f. = 117, *t* = 2.437, *P* = 0.016), CPUE (independent samples *t*-test,
 295 d.f. = 77, *t* = 3.171, *P* = 0.002) and BPUE (independent samples *t*-test, d.f. = 84, *t* =
 296 4.142, *P*<0.001) were significantly higher in the realignment than the estuary. In
 297 addition, mean CPUE (independent samples *t*-test, d.f. = 85, *t* = 2.947, *P* = 0.004) and
 298 BPUE (independent samples *t*-test, d.f. = 63, *t* = 5.299, *P*<0.001) in the realignment
 299 and CPUE in the estuary (independent samples *t*-test, d.f. = 40, *t* = 2.126, *P* = 0.040)
 300 were significantly higher during daylight than at night, but there was no significant
 301 diel difference in BPUE in the estuary (independent samples *t*-test, d.f. = 50, *t* =
 302 1.719, *P* = 0.092).

303



304



305 **Fig. 4.** Mean (\pm S.D.) fish (a) density and (b) biomass in seine catches and (c) catch-
306 per-unit-effort (CPUE) and (d) biomass-per-unit-effort (BPUE) in fyke catches in the
307 managed realignment (white bars) and estuary (black bars), 2010-2014.

308

309 **Diet composition**

310 In 2010, the diets of bass in the realignment were dominated by harpacticoid
311 copepods, with palaemonids and gammarids also consumed; insufficient fish were
312 captured from the estuary for analysis of diet composition (Fig. 5a). In 2011, bass in
313 the realignment preyed mainly upon oligochaetes, mysids and corophiids, while
314 mysids and corophiids dominated diets in the estuary (Fig. 5a). Corophiids dominated
315 the diets of bass in both the realignment and estuary in 2012, with small amounts of
316 mysids also consumed in both habitats (Fig. 5a). In 2013, bass in the realignment
317 preyed mainly upon corophiids and polychaetes, although mysids, oligochaetes and
318 harpacticoid copepods were also consumed; insufficient fish were captured from the
319 estuary for analysis (Fig. 5a). Corophiids were the main prey of bass in the estuary in
320 2014, whereas corophiids, polychaetes and mysids were consumed in the realignment
321 (Fig. 5a). There were no consistent differences in the diets of bass in the realignment
322 and estuary (PERMANOVA, d.f. = 1, $F = 1.741$, $P = 0.184$), although the mean
323 relative abundances of polychaetes, harpacticoid copepods and oligochaetes were
324 higher in the realignment than the estuary, whereas corophiids were more abundant in
325 the estuary (Table 3).

326



327

328 **Fig. 5.** Diet composition of juvenile (a) bass and (b) common goby in the managed
 329 realignment (R) and estuary (E), 2010-2014.

330

331

332 **Table 3.** Similarity percentages (SIMPER) analysis of the mean relative abundances
 333 of key prey taxa and their percentage contributions to dissimilarities in the diets of
 334 juvenile bass and common goby in the managed realignment (R) and estuary (E).

Taxa	Bass			Taxa	C. goby		
	R	E	%		R	E	%
Corophiidae	45.5	70.4	38.3	Corophiidae	30.3	55.9	38.9
Mysidacea	11.1	17.7	18.8	Harpacticoida	50.8	30.2	36.6
Polychaeta	15.5	4.2	14.6	Oligochaeta	7.3	0.8	6.2
Harpacticoida	10.4	2.0	9.9				
Oligochaeta	5.3	0.0	4.4				
Mean			60.6	Mean			63.2
dissimilarity				dissimilarity			

335
 336 In 2010 and 2013, the diets of common goby in the realignment were dominated by
 337 harpacticoid copepods, whereas corophiids were dominant in the estuary (Fig. 5b). In
 338 2011, corophiids, harpacticoid copepods and oligochaetes characterised the diets in
 339 both the realignment and estuary, with gastropods also important at the latter site (Fig.
 340 5b). Corophiids dominated the diets of common goby in both the realignment and
 341 estuary in 2012, although harpacticoid copepods were also consumed, especially in
 342 the realignment (Fig. 5b). The diets in 2014 were similar to those in 2011, with
 343 harpacticoid copepods the most abundant prey in both the realignment and estuary
 344 (Fig. 5b). There was a significant difference in the diets of common goby in the
 345 realignment and estuary (PERMANOVA, d.f. = 1, $F = 7.730$, $P = 0.004$), with the
 346 mean relative abundances of harpacticoids and oligochaetes higher in the realignment
 347 than the estuary, whereas corophiids were more abundant in the estuary (Table 3).

348

349 **DISCUSSION**

350 French (2006) concluded, from a review of the literature, that fish use of suitable
351 managed realignments and reference sites is virtually identical. In this study, however,
352 there was a significant difference in the species composition of fyke catches in the
353 realignment and estuary. In addition, mean densities, CPUE, biomasses and BPUE
354 were higher in the realignment than the estuary, whereas the mean diversity of fyke
355 catches was higher in the estuary. Catches in the realignment are necessarily
356 dependent upon the fishes present in the adjacent estuary, as the site drains at low
357 water, so the causes of the differences are not immediately obvious. It is possible that
358 the manner in which the site floods, or where the gears were deployed in relation to
359 the routes that certain fish species use to enter and leave the site, may have had an
360 influence on the catches. For example, it is possible that fishes enter the drainage
361 ditches with the flooding tide and then disperse across the realignment when the
362 ditches over-top, as observed elsewhere (Colclough *et al.*, 2005; Fonseca *et al.*, 2011).
363 Indeed, densities in seine catches in November 2010 were substantially higher than at
364 any other time during the study because large numbers of fishes were aggregated, and
365 efficiently captured, in a drainage ditch that did not over-top. It is also possible that
366 the deployment of the fykes close to ditches and the breach effectively increased their
367 efficiency relative to those in the estuary, because fishes using the realignment must
368 pass the gears when entering and leaving the site, whereas fishes in the estuary may
369 only pass the gears once. However, the species composition of fyke catches in the
370 realignment and estuary increased in similarity annually during the study period,
371 suggesting that there are differences in habitat between sites but, moreover, that the
372 mudflat in the realignment is still developing. By contrast, there was no significant

373 difference in the species composition of seine catches in the realignment and estuary,
374 possibly because small fishes (targeted by the seine) moved passively into the
375 sampling areas, whereas larger individuals (targeted by the fykes) exhibited active
376 habitat selection (Colclough *et al.*, 2002; Gibson, 2003).

377

378 The majority of catches were dominated by juvenile individuals, demonstrating the
379 importance of the realignment as a nursery area; a similar observation was made by
380 Colclough *et al.* (2005). Larger fishes, especially bass and flounder, also used the
381 realignment, presumably to forage on the abundant juvenile fishes and crustaceans in
382 the site. Overall, the mean lengths of bass and flounder in fyke catches and common
383 goby in seine catches were significantly larger in the realignment than the estuary,
384 whereas bass in seine catches were larger in the estuary. These were unlikely to have
385 been caused by spatial differences in growth rate linked to temperature regime or food
386 availability because the site drains at low tide, so any fishes using the site will
387 necessarily mix with others in the estuary. More likely is that it was caused by size-
388 related differences in habitat use (Gibson, 2003; Colclough *et al.*, 2005; Elliott *et al.*,
389 2007) linked to differences in habitat characteristics in the realignment and estuary.

390

391 Although the mean relative abundance of bass was highest during daylight and that of
392 eel was highest at night, there was no statistically significant difference in the species
393 composition of diurnal and nocturnal fyke catches in the realignment (or the estuary).
394 Contrary to expectations, however, mean CPUE and BPUE in the realignment and
395 CPUE in the estuary were significantly higher during daylight than at night, and the
396 mean lengths of bass and eel in the realignment were significantly larger during
397 daylight than at night (due to an absence of the largest individuals at night). These

398 results suggest that fewer fishes entered the sampling area at night than during
399 daylight, and that there were size-specific, but not species-specific, differences in diel
400 use of the realignment. By contrast, Colclough *et al.* (2005) observed that large bass
401 entered Abbots Hall managed realignment (Blackwater Estuary, England) at night,
402 possibly because the water was too shallow for larger fish to risk entering during
403 daylight. Nocturnal surveys should therefore be considered when assessing the use of
404 managed realignment sites by fishes, as resource use may be substantially greater over
405 the diel cycle than during daylight or darkness alone (Copp, 2008).

406

407 Bass and common goby had relatively narrow diet spectra, with small numbers of
408 taxa, mainly corophiids, copepods, gastropods, mysids or polychaetes, accounting for
409 the majority of the diet; similar results have been obtained elsewhere (Hampel &
410 Cattrijsse, 2004; Laffaille *et al.*, 2001; Fonseca *et al.*, 2011; Nunn *et al.*, 2012; Leclerc
411 *et al.*, 2014). There were no consistent differences in the diets of bass in the
412 realignment and estuary, but there was a significant difference in the diets of common
413 goby, with the mean relative abundances of harpacticoid copepods and oligochaetes
414 higher in the realignment than the estuary, whereas corophiids were more abundant in
415 the estuary. Such differences could be caused by spatial variations in prey abundance,
416 prey size, fish size, foraging behaviour and/or microhabitat characteristics. Regarding
417 the latter possibility, the sediment in parts of the realignment appears to have changed
418 little since the site was breached (A. D. Nunn, *pers. obs.*), and may not yet support
419 high densities (or large sizes) of certain benthic species; macroinvertebrate abundance
420 in Paull Holme Strays managed realignment (Humber Estuary, England) was still an
421 order-of-magnitude lower than in the adjacent mudflat 5 years after the site was first
422 flooded (Mazik *et al.*, 2010). Similarly, Fonseca *et al.* (2011) observed that 30-59 mm

423 bass consumed benthic prey in natural saltmarshes, but mainly copepods in artificial
424 saltmarshes (managed realignments), which was assumed to have been due to
425 differences in microhabitat characteristics and prey availability.

426

427 Although broadly similar, statistically significant differences in fish species
428 composition, abundance, biomass, size structure, diversity and diet composition
429 indicate that the managed realignment is not yet functioning in an identical manner to
430 the mudflat in the adjacent estuary, most likely due to differences in habitat between
431 sites. Notwithstanding, similarity in the species composition of fyke catches in the
432 managed realignment and estuary increased annually during the 5-year study period,
433 suggesting that the mudflat in the realignment is still developing. Indeed, the site will
434 inevitably change over time with accretion, establishment of vegetation and possibly
435 development of creeks (Dixon *et al.*, 1998; French, 2006; Garbutt *et al.*, 2006; Mazik
436 *et al.*, 2010; Kadiri *et al.*, 2011; Mossman *et al.*, 2012; Spencer *et al.*, 2012; Morris,
437 2013; Pétillon *et al.*, 2014). The eastern and northern edges of the site have already
438 accumulated relatively deep mud, similar in depth but of a different consistency to in
439 the estuary, whereas other areas appear largely unchanged since the site was breached
440 (A. D. Nunn, *pers. obs.*). Large numbers of fishes were captured in isolated pools and
441 a drainage channel in 2010, demonstrating the importance of such habitats to fishes in
442 intertidal areas, and similar results have been reported elsewhere (e.g. Colclough *et*
443 *al.*, 2005). However, the depth of water in the pools and drainage channels at low
444 water is now very shallow (due to accretion), and is likely to provide shelter only for
445 small numbers of gobies and juvenile flatfishes; creeks could therefore provide refuge
446 for small fishes at low water and areas of deeper water for larger fishes at high water
447 (Kelley, 1988; Desmond *et al.*, 2000; Laffaille *et al.*, 2001; Colclough *et al.*, 2005;

448 Fonseca *et al.*, 2011). Little vegetation has established to date, although it is likely
449 that coverage will increase in the future, especially along the eastern edge of the site,
450 which is more sheltered from wave action than the western edge and area around the
451 breach; the rate and extent of colonisation will be partly determined by propagule
452 pressure, the elevation of the site, the rate of accretion and the redox potential of the
453 sediment (Mossman *et al.*, 2012). Establishment of (some) vegetation will increase
454 habitat complexity, and not necessarily prevent the aim of the realignment scheme
455 being achieved, as long as sufficient suitable mudflat remains.

456

457 **Acknowledgements**

458 The authors would like to thank the Environment Agency and DP World for their
459 support in the project. Many thanks also to Natalie Angelopoulos, Mike Bailey, Chris
460 Baulcomb, Jon Bolland, Jon Butterfield, Liam Byrne, Terry Coddington, Jamie Dodd,
461 Dave Mitchell, Taya Parker-Mian, Rafa Pérez-Domínguez, Tim Smith, Marie Taylor,
462 Ryan Taylor, Karen Twine, Dan Upsher, Sam Walton and Connor Whiteley for
463 assistance in the field. The work was funded by DP World and administered by the
464 Environment Agency. The views expressed in the paper are those of the authors and
465 not necessarily those of DP World or the Environment Agency.

466

467 **REFERENCES**

- 468 Anderson, M. J. (2001). Permutation tests for univariate or multivariate analysis of
469 variance and regression. *Canadian Journal of Fisheries and Aquatic Sciences*
470 **58**, 626-639.
- 471 Anderson, M. J., Gorley, R. N. & Clarke, K. R. (2008). PERMANOVA+ for
472 PRIMER: Guide to Software and Statistical Methods. PRIMER-E, Plymouth.

473 Atkinson, P. W., Crooks, S., Drewitt, A., Grant, A., Rehfisch, M. M., Sharpe, J. &
474 Tyas, C. J. (2004). Managed realignment in the UK – the first 5 years of
475 colonization by birds. *Ibis* **146**, 101-110.

476 Bray, J. R. & Curtis, J. T. (1957). An ordination of the upland forest communities of
477 Southern Wisconsin. *Ecological Monographs* **27**, 325-349.

478 Clarke, K. R. & Warwick, R. M. (2001). *Change in Marine Communities: an*
479 *Approach to Statistical Analysis and Interpretation*. Second Edition.
480 PRIMER-E, Plymouth.

481 Coates, S., Waugh, A., Anwar, A. & Robson, M. (2007). Efficacy of a multi-metric
482 fish index as an analysis tool for the transitional fish component of the Water
483 Framework Directive. *Marine Pollution Bulletin* **55**, 225-240.

484 Colclough, S. R., Gray, G., Bark, A. & Knights, B. (2002). Fish and fisheries of the
485 tidal Thames: management of the modern resource, research aims and future
486 pressures. *Journal of Fish Biology* **61** (Suppl. A), 64-73.

487 Colclough, S., Fonseca, L., Astley, T., Thomas, K. & Watts, W. (2005). Fish
488 utilisation of managed realignments. *Fisheries Management and Ecology* **12**,
489 351-360.

490 Copp, G. H. (2008). Putting multi-dimensionality back into niche: diel vs. day-only
491 niche breadth separation in stream fishes. *Fundamental and Applied*
492 *Limnology* **170**, 273-280.

493 Cowx, I. G., Nunn, A. D. & Harvey, J. P. (2001). Quantitative sampling of 0-group
494 fish populations in large lowland rivers: point abundance sampling by electric
495 fishing versus micromesh seine netting. *Archiv für Hydrobiologie* **151**, 369-
496 382.

497 Desmond, J. S., Zedler, J. B. & Williams, G. D. (2000). Fish use of tidal creek
498 habitats in two southern California saltmarshes. *Ecological Engineering* **14**,
499 233-252.

500 Dixon, A. M., Leggett, D. J & Weight, R. C. (1998). Habitat creation opportunities for
501 landward coastal re-alignment: Essex case studies. *Journal of the Chartered*
502 *Institution of Water and Environmental Management* **12**, 107-112.

503 Dytham, C. (2003). *Choosing and Using Statistics: a Biologist's Guide*. Second
504 Edition. Blackwell Science, Oxford, 248 pp.

505 Elliott, M., Whitfield, A. K., Potter, I. C., Blaber, S. J. M., Cyrus, D. P., Nordlie, F. G.
506 & Harrison, T. D. (2007). The guild approach to categorizing estuarine fish
507 assemblages: a global review. *Fish and Fisheries* **8**, 241-268.

508 Esteves, L. S. (2013). Is managed realignment a sustainable long-term coastal
509 management approach? *Journal of Coastal Research*, Special Issue No. 65,
510 933-938.

511 Esteves, L. S. (2014). *Managed Realignment: a Viable Long-term Coastal*
512 *Management Strategy?* Springer, New York.

513 Fonseca, L., Colclough, S. & Hughes, R. G. (2011). Variations in the feeding of 0-
514 group bass *Dicentrarchus labrax* (L.) in managed realignment areas and
515 saltmarshes in SE England. *Hydrobiologia* **672**,15-31.

516 French, P. W. (2006). Managed realignment – the developing story of a comparatively
517 new approach to soft engineering. *Estuarine, Coastal and Shelf Science* **67**,
518 409-423.

519 Garbutt, R. A., Reading, C. J., Wolters, M., Gray, A. J. & Rothery, P. (2006).
520 Monitoring the development of intertidal habitats on former agricultural land

521 after the managed realignment of coastal defences at Tollesbury, Essex, UK.
522 *Marine Pollution Bulletin* **53**, 155-164.

523 Gibson, R. N. (2003). Go with the flow: tidal migration in marine animals.
524 *Hydrobiologia* **503**, 153-161.

525 Hampel, H. & Cattrijsse, A. (2004). Temporal variation in feeding rhythms in a tidal
526 marsh population of the common goby *Pomatoschistus microps* (Krøyer,
527 1838). *Aquatic Sciences* **66**, 315-326.

528 Hemingway, K. L. & Elliott, M. (2002). Field methods. In: *Fishes in Estuaries* (eds
529 M. Elliott & K. L. Hemingway). Blackwell Science, Oxford, pp. 410-509.

530 Herringshaw, L. G. & Solan, M. (2008). Benthic bioturbation in the past, present and
531 future. *Aquatic Biology* **2**, 201-205.

532 Kadiri, M., Spencer, K. L., Heppell, C. M. & Fletcher, P. (2011). Sediment
533 characteristics of a restored saltmarsh and mudflat in a managed realignment
534 scheme in Southeast England. *Hydrobiologia* **672**, 79-89.

535 Kelley, D. F. (1988). The importance of estuaries for sea-bass *Dicentrarchus labrax*
536 (L.). *Journal of Fish Biology* **33** (Supplement A), 25-33.

537 Laffaille, P., Feunteun, E. & Lefeuvre, J. C. (2000). Composition of fish communities
538 in a European macrotidal saltmarsh (the Mont Saint-Michel Bay, France).
539 *Estuarine, Coastal and Shelf Science* **51**, 429-438.

540 Laffaille, P., Lefeuvre, J. C., Schricke, M. T. & Feunteun, E. (2001). Feeding ecology
541 of 0-group sea bass, *Dicentrarchus labrax*, in saltmarshes of Mont Saint
542 Michel Bay (France). *Estuaries* **24**, 116-125.

543 Leclerc, J., Riera, P., Noël, L. M. J., Leroux, C. & Andersen, A. C. (2014). Trophic
544 ecology of *Pomatoschistus microps* within an intertidal bay (Roscoff, France),

545 investigated through gut content and stable isotope analyses. *Marine Ecology*
546 **35**, 261-270.

547 Ledoux, L., Cornell, S., O’Riordan, T., Harvey, R. & Banyard, L. (2005). Towards
548 sustainable flood and coastal management: identifying drivers of, and
549 obstacles to, managed realignment. *Land Use Policy* **22**, 129-144.

550 Mander, L., Cutts, N. D., Allen, J. & Mazik, K. (2007). Assessing the development of
551 newly created habitat for wintering estuarine birds. *Estuarine, Coastal and*
552 *Shelf Science* **75**, 163-174.

553 Mazik, K., Smith, J. E., Leighton, A. & Elliott, M. (2007). Physical and biological
554 development of a newly breached managed realignment site, Humber estuary,
555 UK. *Marine Pollution Bulletin* **55**, 564-578.

556 Mazik, K., Musk, W., Dawes, O., Solyanko, K., Brown, S., Mander, L. & Elliott, M.
557 (2010). Managed realignment as compensation for the loss of intertidal
558 mudflat: a short term solution to a long term problem? *Estuarine, Coastal and*
559 *Shelf Science* **90**, 11-20.

560 McLusky, D. S., Bryant, D. M. & Elliott, M. (1992). The impact of land-claim on
561 macrobenthos, fish and shorebirds on the Forth Estuary, eastern Scotland.
562 *Aquatic Conservation: Marine and Freshwater Ecosystems* **2**, 211-222.

563 Morris, R. K. A. (2013). Managed realignment as a tool for compensatory habitat
564 creation: a re-appraisal. *Ocean and Coastal Management* **73**, 82-91.

565 Morris, R. K. A. & Gibson, C. (2007). Port development and nature conservation:
566 experiences in England between 1994 and 2005. *Ocean and Coastal*
567 *Management* **50**, 443-462.

- 568 Mossman, H. L., Davy, A. J. & Grant, A. (2012). Does managed coastal realignment
569 create saltmarshes with 'equivalent biological characteristics' to natural
570 reference sites? *Journal of Applied Ecology* **49**, 1446-1456.
- 571 Nunn, A. D., Harvey, J. P. & Cowx, I. G. (2007). Benefits to 0+ fishes of connecting
572 man-made waterbodies to the lower River Trent, England. *River Research and
573 Applications* **23**, 361-376.
- 574 Nunn, A. D., Tewson, L. H. & Cowx, I. G. (2012). The foraging ecology of larval and
575 juvenile fishes. *Reviews in Fish Biology and Fisheries* **22**, 377-408.
- 576 Pétilion, J., Potier, S., Carpentier, A. & Garbutt, A. (2014). Evaluating the success of
577 managed realignment for the restoration of salt marshes: lessons from
578 invertebrate communities. *Ecological Engineering* **69**, 70-75.
- 579 Pontee, N. (2013). Defining coastal squeeze: a discussion. *Ocean and Coastal
580 Management* **84**, 204-207.
- 581 Ramos, S., Amorim, E., Elliott, M., Cabral, H. & Bordalo, A. A. (2012). Early life
582 stages of fishes as indicators of estuarine ecosystem health. *Ecological
583 Indicators* **19**, 172-183.
- 584 Rupp-Armstrong, S. & Nicholls, R. J. (2007). Coastal and estuarine retreat: a
585 comparison of the application of managed realignment in England and
586 Germany. *Journal of Coastal Research* **23**, 1418-1430.
- 587 Shih, S. C. W. & Nicholls, R. J. (2007). Urban managed realignment: application to
588 the Thames Estuary, London. *Journal of Coastal Research* **23**, 1525-1534.
- 589 Spencer, T., Friess, D. A., Möller, I., Brown, S. L., Garbutt, R. A. & French, J. R.
590 (2012). Surface elevation change in natural and re-created intertidal habitats,
591 eastern England, UK, with particular reference to Freiston Shore. *Wetlands
592 Ecology and Management* **20**, 9-33.

593 Washington, H. G. (1984). Diversity, biotic and similarity indices: a review with
594 special relevance to aquatic ecosystems. *Water Research* **18**, 653-694.

595 Ysebaert, T., Herman, P. M. J., Meire, P., Craeymeersch, J., Verbeek, H. & Heip, C.
596 H. R. (2003). Large-scale spatial patterns in estuaries: estuarine macrobenthic
597 communities in the Schelde estuary, NW Europe. *Estuarine, Coastal and Shelf*
598 *Science* **57**, 335-355.