

# Liquid Chromatography Mass Spectrometry Analysis and Cytotoxicity of *Asparagus adscendens* Roots against Human Cancer Cell Lines

Kashif Maqbool Khan<sup>1,2,3</sup>, Lutfun Nahar<sup>1</sup>, Abdul Mannan<sup>2</sup>, Muhammad Arfan<sup>4</sup>, Ghazanfar Ali Khan<sup>5</sup>, Afaf Al-Groshi<sup>1</sup>, Andrew Evans<sup>1</sup>, Nicola M. Dempster<sup>1</sup>, Fyaz M. D. Ismail<sup>1</sup>, Satyajit D. Sarker<sup>1</sup>

<sup>1</sup>School of Pharmacy and Biomolecular Sciences, Medicinal Chemistry and Natural Products Research Group, Liverpool John Moores University, England, UK, <sup>2</sup>Department of Pharmacy, COMSATS Institute of Information Technology, Abbottabad, <sup>3</sup>Institute of Pharmaceutical Sciences, University of Veterinary and Animal Sciences, Lahore, <sup>4</sup>Department of Chemistry, School of Natural Sciences, National University of Science and Technology, Islamabad, <sup>5</sup>Department of Research and Development, Drug Regulatory Authority, Islamabad, Pakistan

Submitted: 10-04-2017

Revised: 03-06-2017

Published: 31-01-2018


## ABSTRACT

**Background:** *Asparagus adscendens* Roxb. (Asparagaceae), is native to the Himalayas. This plant has been used in the prevention and effective treatment of various forms of cancers. **Objective:** This paper reports, for the first time, on the cytotoxicity of the methanol (MeOH) extract of the roots of *A. adscendens* and its solid-phase extraction (SPE) fractions against four human carcinoma cell lines and LC-ESI-QTOF-MS analysis of the SPE fractions. **Materials and Methods:** Finely powdered roots of *A. adscendens* were macerated in methanol and extracted through SPE using gradient solvent system (water: methanol) proceeded for analysis on LC-ESI-QTOF-MS and cytotoxicity against four human carcinoma cell lines: breast (MCF7), liver (HEPG2), lung (A549), and urinary bladder (EJ138), using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide assay. **Results:** The MeOH extract and four SPE fractions exhibited cytotoxicity against all cell lines with the IC<sub>50</sub> values ranging from 6 to 79 µg/mL. As observed in other *Asparagus* species, the presence of saponins and saponinins in the SPE fractions was evident in the liquid chromatography-mass spectrometry data. **Conclusion:** It is reasonable to assume that the cytotoxicity of the MeOH extract of the roots of *A. adscendens* and its SPE fractions, at least partly, due to the presence of saponins and their aglycones. This suggests that *A. adscendens* could be exploited as a potential source of cytotoxic compounds with putative anticancer potential.

**Key words:** 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide, asparagaceae, *Asparagus adscendens*, cancer, cytotoxicity, liquid chromatography-mass spectrometry, saponinins, saponin

## SUMMARY

- The MeOH extract and all solid-phase extraction (SPE) fractions exhibited various levels of cytotoxicity against all cell lines with the IC<sub>50</sub> values ranging from 6 to 79 µg/mL
- The presence of saponins and saponinins in the SPE fractions was evident in the Liquid chromatography-mass spectrometry data
- Due to the presence of saponins and their aglycones, suggest that *A. adscendens* could be exploited as a potential source of cytotoxic compounds with putative anticancer potential.



**Abbreviation used:** SPE: Solid-phase extraction, MCF7: Breast cancer cell line, HEPG2: Liver cancer cell line, A549: Lung liver cancer cell line, EJ138: Urinary bladder cancer cell line, MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide, LC-MS: Liquid chromatography-mass spectrometry.

## Correspondence:

Dr. Kashif Maqbool Khan,  
Department of Pharmacy, COMSATS Institute of Information  
Technology, Abbottabad 22060, Pakistan.  
E-mail: [kashifmkhan007@gmail.com](mailto:kashifmkhan007@gmail.com)

Dr. Satyajit D. Sarker,  
Medicinal Chemistry and Natural Products Research  
Group, School of Pharmacy and Biomolecular  
Sciences, Liverpool John Moores University, James  
Parsons Building, Byrom Street, Liverpool L3 3AF,  
England, UK.

E-mail: [S.Sarker@ljmu.ac.uk](mailto:S.Sarker@ljmu.ac.uk)  
**DOI:** 10.4103/jpm.pm\_136\_17

Access this article online

Website: [www.phcog.com](http://www.phcog.com)

Quick Response Code:



## INTRODUCTION

*Asparagus adscendens* Roxb. (Asparagaceae), commonly known as “safed mushi” in Pakistan and “Shatawari,” “Shatavar,” “Shatamuli,” “Sahasrapal,” or “Sainsarbuti” in India, is native to the Himalayas.<sup>[1]</sup> This plant grows 1–2 m tall and prefers to take root in gravelly, rocky soils high up in piedmont plains, at 1300–1500 m above the sea level.<sup>[2]</sup> This plant that has been used for a long time as a component of various complementary and alternative medicinal preparations in India and Pakistan.<sup>[3]</sup> Conventionally, *A. adscendens* is considered to be a general health promoting tonic and has been used to treat various sexual disorders in men.<sup>[4]</sup>

The genus *Asparagus* comprises of about 300 species, and most of the European species are used as vegetables.<sup>[5]</sup> Of the species that grow in the

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

**Cite this article as:** Khan KM, Nahar L, Mannan A, Arfan M, Khan GA, Al-Groshi A, *et al.* Liquid chromatography mass spectrometry analysis and cytotoxicity of *Asparagus adscendens* roots against human cancer cell lines. *Phcog Mag* 2017;13:S890-4.

Himalayan region of Pakistan, *A. adscendens* and *A. racemose*, are the most commonly used species in traditional medicines. Aliphatic, nitrogenous and phenolic compounds, saponins, steroids, and triterpenoids have been reported from *A. adscendens* of Indian origin;<sup>[1]</sup>  $\beta$ -sitosterol glucoside, spirostanol glycosides (asparanin C and asparanin D), and furostanol glycosides (asparoside C and asparoside D) were isolated;<sup>[6]</sup> steroidal saponins and glycosides, and various lipophilic compounds were found in the tuberous roots and leaves;<sup>[2]</sup> sarsasapogenin, diosgenin,  $\beta$ -sitosterol glucoside, spirostanol glycosides (asparanin A and B), and furostanol glycosides (asparoside A and B) [Figure 1], were isolated from this plant.<sup>[7-9]</sup>

Due to its aphrodisiac and immunomodulatory properties, *A. adscendens* has become one of the most commercially exploited species in India and Pakistan.<sup>[10]</sup> Nowadays, *A. adscendens* has been used in the prevention and effective treatment of various forms of cancers.<sup>[11]</sup> In addition, several pharmacological properties of the plants of the genus *Asparagus* have been well-documented, including antimicrobial,<sup>[12]</sup> anti-inflammatory, antitussive,<sup>[13]</sup> hepatoprotective,<sup>[14]</sup> immunomodulatory activity,<sup>[10]</sup> antistress,<sup>[3]</sup> anti-secretory, and antiulcer activity.<sup>[5,15]</sup> A study of ancient classical Ayurvedic literature claimed several therapeutic attributes for the root of *Asparagus* and has been especially recommended in cases of threatened abortion and as a galactagogue.<sup>[16]</sup> However, with the only exception of the report on the isolation and identification of conyopodioid from *A. adscendens*,<sup>[17]</sup> and significant acetylcholinesterase- and butyrylcholinesterase-inhibitory activity (and inactivity against monkey kidney epithelial cell (LCMK-2) and mice hepatocytes),<sup>[1]</sup> to the best of our knowledge, there has been no systematic pharmacological and phytochemical work performed on *A. adscendens* native to Pakistan. Therefore, this study was undertaken to explore potential cytotoxicity of the MeOH extract of the roots of *A. adscendens* and its solid-phase extraction (SPE) fractions against four human carcinoma cell lines: breast (MCF7), liver (HepG2), lung (A549), and urinary bladder (EJ138) using the *in vitro* 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide (MTT) cytotoxicity/viability assay and to carry out LC-ESI-QTOF-MS analysis of the SPE fractions.

## MATERIALS AND METHODS

### Reagents and chemicals

All chemicals were purchased from Sigma-Aldrich (Dorset, UK). Solvents were obtained from Fischer Scientific (Loughborough, UK). All cell culture reagents were purchased from Biosera (Nauaille, France).

### Plant materials


Plant sample was collected from Nathia Gali, region of Khyber Pakhtunkhwa, Pakistan and identified as *A. adscendens* Roxb. by Dr. Muhammad Zafar, Herbarium Botanist, Department of Plant Sciences, Quaid-I-Azam University, Islamabad, Pakistan. A herbarium specimen (voucher number: Acc no. PAC1001) has been deposited and retained in the above herbarium.

### Extraction and preparation of plant samples

Shade-dried and finely powdered roots (2.5 kg) of *A. adscendens* were macerated in MeOH (5 L) for 10 days at room temperature, filtered, and the solvent was evaporated under vacuum using a rotatory evaporator (<45°C) to get a concentrated gummy crude extract.

### Solid-phase extraction and sample purification

The procedure was similar to that described.<sup>[18]</sup> In brief, a portion of the dried MeOH extract (2 g) was suspended in 20 mL of HPLC grade water and loaded onto a Strata C-18 cartridge (20 g), initially washed



**Figure 1:** Major saponins and sapogenins previously isolated from *Asparagus adscendens*

with MeOH (50 mL) followed by equilibration with water (100 mL). The cartridge was eluted with MeOH-water mixture of decreasing polarity to generate four fractions: 20, 50, 80, and 100% MeOH in water (250 mL each), coded respectively, as AAMF1, AAMF2, AAMF3, and AAMF4, evaporated to dryness using a rotary evaporator and a freeze-dryer, re-dissolved in MeOH (10 mg/mL), centrifuged at 12,000 rpm for 3 min, filtered through 0.20  $\mu$ m sterile syringe filter for injection (10  $\mu$ L) into the Liquid chromatography-mass spectrometry (LC-MS) system.

### Liquid chromatography-mass spectrometry

An Alliance HPLC System 2695 (Waters) was used. Reversed-phase chromatography was performed on a Phenomenex Gemini-NX 5  $\mu$ m C<sub>18</sub> column (250 mm  $\times$  4.6 mm). The column temperature was set at 25°C. A variable wavelength UV-Vis detector was set at 220 nm. An elution gradient was applied with solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid in MeOH). The initial mobile phase composition was 70% of A and 30% B at 0 min, then linear gradient to 100% of B over 30 min and held at that composition for 5 min (flow rate of 1 mL/min).

The LC system was connected to a quadrupole time-of-flight (TOF) mass spectrometer (Waters Micromass LCT) having an electrospray ion source. The response was recorded in real time by the mass spectrometer data system (Waters MassLynx version 4.1). The tuning parameters were set as follows: electrospray interface 3000 V, rangefinder lens 250 V, extraction cone 3 V, desolvation temperature 20°C, source temperature 100°C, nebulizer gas flow 20 L/h, desolvation gas flow 760 L/h, and TOF tube 4687 V. Data acquisition method was set as follows: cycle time 1 s, scan duration 0.9 s, inter-scan delay 0.1 s, mass range 100–1600, and centroid mode. Positive ion mode was operated with many cone voltage settings of 40V.

### Cell lines, cell cultures, and the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide assay

The potential cytotoxicity of the MeOH extract of the roots of *A. adscendens* and its SPE fractions was studied against four human

carcinoma cell lines: breast (MCF7), liver (HepG2), lung (A549), and urinary bladder (EJ138) using the MTT assay.<sup>[18-20]</sup> The cells were washed by phosphate buffer saline and harvested by trypsinization. All cell lines were cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum. All cells were cultured at 37°C in 95% air and 5% CO<sub>2</sub>. For the MTT assay, cells were seeded into 24 well plates at density 1.2 × 10<sup>4</sup> cells/well in a working volume of 1 mL/well and allowed to grow for 24 h before the commencement of each experiment.

The cells were treated for 24 h with different concentrations of test samples (the MeOH extract and SPE fractions; 0, 0.8, 4, 20, 100, and 500 µg/mL). Dilution of stock solutions was made in culture medium yielding final sample concentrations with a final dimethyl sulfoxide concentration of 0.1%, including the control. Each sample was used to treat four wells of cells in each 24-well plate. After 24-h treatment period, the cytotoxicity of the samples on each carcinoma cell line was quantified using MS Excel. To achieve this, the medium in each well was replaced by MTT solution (500 µg/mL) and incubated for 2 h. Toxicity was assessed by the ability of the cells to reduce the yellow dye MTT to a blue formazan product.<sup>[21]</sup> MTT reagent was removed, and the formazan crystals produced by viable cells were dissolved in isopropanol and absorbance (560 nm) was determined with the microplate reader (CLARIO Star Microplate reader, BMG Labtech, UK). The average absorbance obtained from all the control wells (without test sample) on each plate was arbitrarily set at 100% and the absorbance value for the average of wells of cells treated with each test samples was expressed as a percentage of this control. Each assay was performed on a minimum of three separate occasions, and the IC<sub>50</sub> values were for each sample on each cell line were calculated using Microsoft Excel version 2013 (Redmond, WA, USA).

## Statistical analysis

The data were expressed as mean values ± standard error of the mean (SEM) of three parallel replicates. The graph was plotted using nonlinear regression with the use of Microsoft Excel version 2013. The means were separated at confidence level  $P \leq 0.05$  using analysis of variance with Tukey's range test.

## RESULTS AND DISCUSSION

The MeOH extract of the roots of *A. adscendens* and its SPE fractions (AAMF1, AAMF2, AAMF3 and AAMF4) displayed different levels of cytotoxicity against four human carcinoma cell lines, breast (MCF7), liver (HepG2), lung (A549) and urinary bladder (EJ138) in the *in vitro* MTT cytotoxicity/viability assay [Table 1]. The MeOH extract exhibited the highest level of cytotoxicity against the breast cancer cell line (MCF7; IC<sub>50</sub> = 6 µg/mL), but it was also active against three other cell lines, HepG2, EJ138, and A549 (IC<sub>50</sub> = 13, 30 and 63 µg/mL, respectively) [Table 1]. The SPE fraction AAMF1, which had the most polar components of the parent MeOH extract, showed

the most significant cytotoxicity against MCF7 (IC<sub>50</sub> = 8 µg/mL), and considerable cytotoxicity against HepG2, EJ138, and A549 with the IC<sub>50</sub> values of 17, 34, and 65 µg/mL, respectively. This SPE fraction was almost as potent as its parent MeOH extract in terms of cytotoxicity against the urinary bladder cell line EJ138. The SPE fraction AAMF2 showed most prominent cytotoxicity against the breast cancer cell line MCF7 (IC<sub>50</sub> = 10 µg/mL), and also activity against other three cell lines, HepG2, EJ138, and A549 (IC<sub>50</sub> = 18, 36 and 67 µg/mL, respectively). The cytotoxicity pattern of the SPE fraction AAMF3 was quite similar to that of the AAMF1 and the MeOH extract; it showed most significant cytotoxicity against the HepG2 cell line (IC<sub>50</sub> = 15 µg/mL). This fraction was also cytotoxic to MCF7, EJ138, and A549 cell lines with the IC<sub>50</sub> values of 17, 39, and 73 µg/mL, respectively. The SPE fraction AAMF4, which contained the least polar components of the parent MeOH extract, exhibited notable cytotoxicity against the HepG2 cell line (IC<sub>50</sub> = 19 µg/mL), and was also active against the MCF7, EJ138, and A549 cell lines (IC<sub>50</sub> = 27, 43 and 79 µg/mL, respectively). With the exception of AAMF3 and AAMF4, two other SPE fractions showed the highest level of cytotoxicity against the MCF7 cell line, as was observed with their parent MeOH extract.

This is the first report on cytotoxicity of the MeOH extract of the roots of *A. adscendens* and its SPE fractions against any carcinoma cell lines. The current finding is in line with the findings of a few other previous studies on cytotoxicity of some other species of the genus *Asparagus*.<sup>[22-26]</sup> The bioactive components of the genus *Asparagus* belong predominantly to the chemical classes of sapogenins and saponins [Figure 1], which are well-known to exhibit cytotoxicity. Most of the compounds isolated previously from the genus *Asparagus* as described in the literature are steroidal sapogenins and saponins. Sarsasapogenin, diosgenin, β-sitosterol and its glucoside, spirostanol glycoside, and furostanol glycoside were reported from *A. adscendens*.<sup>[2,7,9]</sup> In another study, the steroidal saponins from the roots of *A. filicinus* showed significant cytotoxic activities against human lung carcinoma (A549) and breast adenocarcinoma (MCF7) tumor cell lines,<sup>[22]</sup> compounds isolated from this species were cytotoxic to human breast adenocarcinoma cell line MDA-MB-231 (IC<sub>50</sub> 3.4–6.6 µM).<sup>[23]</sup> *A. officinalis* displayed significant activity against HeLa and BEL-7404 cells in a dose-dependent manner *in vitro* at 10 mg/mL.<sup>[25]</sup> The alkaloid, aspastilaine, isolated from *A. stipularis* was shown to be active against MCF7 (IC<sub>50</sub> = 47.7 µM).<sup>[26]</sup> A series of sarsasapogenin and diosgenin derived steroidal constituents, isolated from *A. racemosus*, were screened for their ability to induce cell death and apoptosis against HCT-116 human colon carcinoma cell line and the carbohydrates moieties linked to the steroid backbone were found to strongly influence cytotoxic activity and cell death mode.<sup>[24]</sup>

LC-MS studies on the SPE fractions of the MeOH extract of the roots of *A. adscendens* were carried out to get an insight into the possible chemical composition of the fractions, particularly, to have an indication whether they contain saponins and sapogenins as possible contributors to the significant cytotoxicity of the extract and its fractions. The chromatographic conditions were optimized by method development. A linear gradient elution with water and MeOH containing 0.1% formic acid as the mobile phase offered the best resolution. Typical chromatograms of fractions with mass spectrometric detection in positive ion mode exhibited quite complex patterns of peaks [Table 2], and only the possible presence of saponins (e.g., Paris saponins VII at  $t_R$  18.64 [Figure 2], [M + 1] + m/z 1032) and spirostanol (e.g., Type VI daucosterol at  $t_R$  34.82, [M + 1] + m/z 576) in all SPE fractions could be suggested from the retention times and the MS spectral data of the separated peaks.<sup>[27-30]</sup> The presence of saponins and their aglycones was in agreement with that of other *Asparagus* species.

**Table 1:** The IC<sub>50</sub> values of the MeOH extract of the roots of *Asparagus adscendens* and its solid-phase extraction fractions against four carcinoma cell lines

Cell lines	IC <sub>50</sub> value (µg/mL)				
	MeOH extract	SPE fractions			
		AAMF1	AAMF2	AAMF3	AAMF4
A549	63	65	67	73	79
EJ138	30	34	36	39	43
HepG2	13	17	18	15	19
MCF-7	6	8	10	17	27

IC<sub>50</sub> of etoposide (reference drug) against lung cancer cell line (A549) was 100 µM. SPE: Solid-phase extraction