



# Open Research Online

---

The Open University's repository of research publications and other research outputs

## Mauritius on fire: Tracking historical human impacts on biodiversity loss

### Journal Item

How to cite:

Gosling, William D.; de Kruif, Jona; Norder, Sietze J.; de Boer, Erik J.; Hooghiemstra, Henry; Rijdsijk, Kenneth F. and McMichael, Crystal N. H. (2017). Mauritius on fire: Tracking historical human impacts on biodiversity loss. *Biotropica*, 49(6) pp. 778–783.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: Accepted Manuscript

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.1111/btp.12490>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

1 For submission to: **Biotropica as an Insight**

2

3 Left running header: **Gosling *et al.***

4 Right running header: **Mauritius on fire**

5

6 Title: **Mauritius on fire: Tracking historical human impacts on biodiversity**

7 **loss**

8

9 **Authors:** William D. Gosling<sup>1, 2\*</sup>, Jona de Kruif<sup>1</sup>, Sietze J. Norder<sup>1, 3</sup>, Erik J. de Boer<sup>1</sup>,

10 Henry Hooghiemstra<sup>1</sup>, Kenneth F. Rijdsdijk<sup>1</sup> & Crystal N.H. McMichael<sup>1</sup>

11

12 **Institutions:**

13 1 = Institute for Biodiversity & Ecosystem Dynamics, University of Amsterdam, P.O.

14 Box 94248, 1090 GE, The Netherlands.

15 2 = School of Environment, Earth & Ecosystem Sciences, The Open University,

16 Walton Hall, Milton Keynes, MK7 6AA, UK.

17 3 = Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity

18 Group, Faculdade de Ciências, Universidade de Lisboa, 1749-016, Lisboa, Portugal.

19

20 \* corresponding author: [W.D.Gosling@uva.nl](mailto:W.D.Gosling@uva.nl)

21

22

23

24

25 Received \_\_\_\_\_ ; revision accepted \_\_\_\_\_.

26 **Abstract**

27 Fire was rare on Mauritius prior to human arrival (AD 1598); subsequently three  
28 phases of elevated fire activity occurred: c. 1630-1747, 1787-1833, and 1950-modern.  
29 Elevated fire frequency coincided with periods of high human impact evidenced from  
30 the historical record, and is linked to the extinction of island endemics.

31

32 **Tweetable abstract:** A 1000 year charcoal record shows humans brought frequent  
33 fire to Mauritius, fire is linked to extinction events #Dodo

34

35 **Key words:** Anthropocene; charcoal; Dodo; ecosystem degradation; extinction;  
36 introduced species; islands; habitat degradation; Mare Tatos; sub-tropical

37

38 **Text**

39 THE ISLAND OF MAURITIUS BORE WITNESS TO PROBABLY THE MOST ICONIC  
40 SINGLE ACT OF HUMAN ECOSYSTEM INTERFERENCE OF ALL TIME WHEN, IN AD 1690,  
41 THE DODO WAS DRIVEN TO EXTINCTION (Roberts & Solow 2003, Hume *et al.* 2004).  
42 Yet the relative importance of the multiple anthropogenic factors responsible for the  
43 loss of the Dodo, and other extinctions on Mauritius, remain ambiguous. Prior to the  
44 first landing of the Dutch on Mauritius in AD 1598 (all dates hereafter in years AD)  
45 the island naturally supported a palm or semi-dry woodland vegetation and was  
46 virtually untouched by humans, with the only possible earlier visitors being Arab or  
47 Portuguese traders (Cheke & Hume 2008, de Boer *et al.* 2014). Mauritius is therefore  
48 an ideal location to explore the role of humans in extinction events because it is a rare  
49 case of an island where human impact on ecosystems and written historical records  
50 commence simultaneously (Vaughan & Wiehe 1937, Brouard 1963, Grove 1996,  
51 Moree 1998, Grihault 2005, Cheke & Hume 2008, Floore & Jayasena 2010, Norder *et*  
52 *al.* 2017).

53

54 The Dutch first settled Mauritius 40 years after their initial landings. During the  
55 subsequent four centuries the island was occupied and abandoned by the Dutch (1638-  
56 1710), French (1721-1810), and British (1810-1968), until it achieved independence  
57 in 1968. Since the arrival of humans, Mauritian ecosystems have become degraded  
58 through the overexploitation of species for food, the introduction of exotic species  
59 (such as rats and cats), and the transformation of the landscape through clearance and  
60 agriculture (Brouard 1963, Cheke & Hume 2008, Florens *et al.* 2012, Hume 2013,  
61 Florens 2013, Norder *et al.* 2017).

62

63           Throughout the centuries, Mauritius has been deforested to harvest timber for  
64 construction purposes, to create space for agriculture, and to supply energy for  
65 cooking, railways, and the sugar industry. Fire was an integral part of the  
66 deforestation process on Mauritius, and consequently charcoal deposited into  
67 sedimentary sequences can be used to assess the scale of human impact through time.  
68 Here we reconstruct a c. 1000 year fire history from Mauritius, and relate it to  
69 historical records of deforestation and demography, species introductions, and  
70 extinction events. The integration of empirical palaeoecological data with historically  
71 documented events provides a unique insight into the relative impact of the different  
72 phases of human activity on Mauritian ecosystems. Although charcoal has been used  
73 to track human impacts on tropical island ecosystems elsewhere in the world (e.g.  
74 Burney 1987, Premathilake 2006, Rull *et al.* 2015) the unique combination of the  
75 Mauritius data with historical records highlights the accuracy with which  
76 palaeoecological reconstructions can capture past human impacts.

77

78           In 2010, a sediment core was recovered from the Mare Tatos wetland on  
79 Mauritius (20° 12' 44" S, 57° 46' 22" E, 21 m above sea level) using a Russian corer  
80 (de Boer *et al.* 2014). Mare Tatos is located c. 40 km from the site of the first Dutch  
81 landings at Port de Warwick (present day Vieux Grand Port). Five radiocarbon dates  
82 indicate the upper 100 cm of the Mare Tatos core represents sediments that were  
83 deposited during the last c. 1000 years (Table 1). The most probable chronology for  
84 the core was established using the ShCal13 and post bomb radiocarbon calibration  
85 curves (Hogg *et al.* 2013, Hua *et al.* 2013, Reimer *et al.* 2013) with Bacon source  
86 code (Blaauw & Christen 2011) in R (Version 3.2.2; R Core Team 2015) (Fig. 1a).

87

88           We extracted 46 samples from the Mare Tatos core for macro-charcoal analysis.  
89   For each sample, charcoal particles were identified using standard laboratory  
90   procedures (Whitlock & Larsen 2001). Two size fractions of charcoal fragments were  
91   quantified (75-160  $\mu\text{m}$  and  $>160 \mu\text{m}$ ); particle counts for both size fractions, and  
92   additionally surface area for particles  $>160 \mu\text{m}$ . The smaller size fraction represents  
93   fires in the landscape (10's of km), while the larger size fraction reflects fires at the  
94   study site (Clark & Patterson 1997). Area calculations were made for the larger  
95   fraction to standardise charcoal particle size relative to abundance. Periods of high,  
96   landscape scale, fire frequency were classified as “fire zones” when the number of  
97   small particles (75-160  $\mu\text{m}$ ) per cubic centimetre exceeded 20% of the maximum  
98   value (1093 particles/ $\text{cm}^3$ ). Fire was assumed to be virtually absent from the  
99   ecosystems when the number of particles dropped below 5% of the maximum values  
100   at the landscape (75-160  $\mu\text{m}$ ), and local ( $>160 \mu\text{m}$ ) scales (Kelly *et al.* 2011). Data  
101   were plotted using C2 version 1.7.2 (Juggins 2005).

102

103           The charcoal data from Mare Tatos indicate three fire zones: (1) c. 1630-1747;  
104   (2) c. 1787-1833; and (3) c. 1950 to modern (Fig. 1a). The three fire zones coincide  
105   with periods of increasing human impact on Mauritius inferred from historical maps  
106   and documented population growth (Fig. 1). Fire frequency and abundance sharply  
107   increased after the first colonization of Mauritius by the Dutch, and peaks around the  
108   transition between Dutch and French governances (fire zone 1). After an initial peak  
109   in fire activity, during most of the French governance, fire activity remains low. The  
110   second fire zone, between c. 1787 and 1833, commences during the latter stages of  
111   the French governance with the expansion of sugar cane agriculture, and includes the  
112   arrival of the British and the industrialisation of agriculture (Norder *et al.* 2017).

113 During much of the British governance fire activity remained low. The third fire zone,  
114 c. 1950 to modern, is broadly coincident with the destruction of forest remnants, and  
115 diversification of the economy following independence.

116

117 Our data suggest that for c. 500 years prior to the arrival of the Dutch on  
118 Mauritius, very little fire activity occurred at Mare Tatos, or within the surrounding  
119 landscape (Fig. 1a). The rarity of pre-human fire is concordant with a c. 8000-year  
120 reconstruction of micro-charcoal (c. 200 year between samples) from Mare Tatos (de  
121 Boer *et al.* 2014). Fire became a major element of the landscape only after the Dutch  
122 settled the island, which supports previous assertions that there were no significant  
123 landings of people on Mauritius prior to the arrival of the Dutch (Cheke & Hume  
124 2008, Floore & Jayasena 2010).

125

126 The first period of elevated fire activity on Mauritius (fire zone 1; c. 1630-1747)  
127 is coincident with the first Dutch settlement on the island. Despite a population of just  
128 a few hundred people, the charcoal signal evidences fire occurring at both regional  
129 (<160  $\mu\text{m}$ ) and local (>160  $\mu\text{m}$ ) scales (Fig. 1a). The charcoal record detects fires  
130 reflecting early clearance and cultivation in the bay area just c. 5 km north-west and  
131 south-east of Mare Tatos (Floore & Jayasena 2010; Fig 1b, 1685 map). During fire  
132 zone 1 forests were increasingly cut and burnt to provide space to grow crops, such as  
133 sugar cane, rice, tobacco, indigo, vegetables and citrus trees (Brouard 1963, Grihault  
134 2005). The peaks and troughs in the local charcoal signal within fire zone 1 could  
135 reflect sporadic visits by people to Mare Tatos, possibly related to foraging  
136 expeditions and/or selective logging for ebony trees.

137

138           During fire zone 1 all large flightless birds on Mauritius became extinct,  
139 including Dodo (*Raphus cucullatus*), along with the Raven Parrot (*Lophopsittacus*  
140 *mauritianus*), and two species of endemic giant tortoise (*Cylindraspis inepta* and *C.*  
141 *triserrata*) (Fig. 1a). It is interesting to note that the extinction events lag the first  
142 introduction of rats to the island by Arab traders by many hundred years, the  
143 subsequent wave of introduced species that occurred with the first Dutch landings by  
144 c. 82, 92 and 102 years respectively, and the occupation and introduction of  
145 widespread fire by c. 50, 60 and 70 years. The differential duration that species  
146 persisted post human arrival attests to the cumulative nature of human impacts, and  
147 the species-specific nature of extinction debt (Gaston & Blackburn 1995, Triantis *et*  
148 *al.* 2010). Although it is unlikely that the fire events directly caused extinctions,  
149 the data indicates that even localised settlement of humans, their cultivation and  
150 burning activities, affected ecosystems in a wider region than the settlements  
151 themselves. The destruction of ecosystems (habitat) likely put additional  
152 pressure on already stressed wild animal populations.

153

154           Once established on Mauritius, the Dutch colonists faced a series of natural  
155 (plagues, heavy storm damage) and political (lack of support from the mainland,  
156 workers rebellion) difficulties, and abandoned the island in 1710 (Moree 1998). After  
157 a decade of practically no human habitation the French settled Mauritius (Cheke &  
158 Hume 2008). Fire remained a significant feature of the Mauritian landscape after the  
159 governance transition (fire zone 1). The first major lull in fire activity on Mauritius  
160 after European colonization occurred between c. 1747-1787, preceding the French  
161 revolution (Fig. 1a), when land-based agricultural activities were of minor economic



162 importance compared to the provisioning of goods and services to passing ships  
163 (Allen 1989, Addison & Hazareesingh 1999).

164

165         The second major increase in fire (fire zone 2; c. 1787-1833) is coincident with  
166 an increase of the human population. In 1787 the population was c. 40,000 and grew  
167 to c. 60,000 in the 1790's to support the rapidly growing sugar cane industry (Lutz &  
168 Wils 1994). Historical archives document the cutting down of vegetation and  
169 subsequent burning (Brouard 1963) focused in the northern part of the island near  
170 Mare Tatos (Fig. 1b).

171

172         A species of endemic fruit bat (*Pteropus subniger*), which roosted in tree  
173 hollows and cliffs, went extinct in the wild c. 1795, eight years after the onset of fire  
174 zone 2 (Cheke & Hume 2008). The extant fruit bat species on Mauritius (*P. niger*) is  
175 currently listed as vulnerable due to habitat loss and hunting (Hutson & Racey 2013,  
176 Florens 2015). *P. subniger* was also heavily hunted for food and making torches  
177 (Cheke & Hume 2008), but the coincidence of its extinction with the rapid increases  
178 in fire (Fig. 1a) suggests that, in addition to predation, habitat destruction may have  
179 played a role in its demise.

180

181         The major drop in fire activity c. 30 years after the onset of the major expansion  
182 in sugar cane area in 1825 is coincident with the projected loss of c. 50% of the  
183 natural vegetation on Mauritius (Fig. 1a). By c. 1850 most of the land suitable for  
184 sugar cane agriculture on the northern part of Mauritius had been cleared, leaving no  
185 natural vegetation left to burn (Fig. 1b, 1872 map). Management of the land for sugar  
186 cane from 1833-1950 has left no significant charcoal record despite the practice of a

187 seasonal burning of the fields (Lalljee & Facknath 2008). The absence of a charcoal  
188 (fire) signal likely reflects a shift from burning woody vegetation to burning crop  
189 stubble (grass), because non-woody vegetation tends to produce less charcoal  
190 (Whitlock & Larsen 2001).

191

192         It is not until the time of independence that fire activity again increased (c.  
193 1950). Fire zone 3 reflects the transition to the current situation of c. 98 % loss of  
194 natural vegetation on Mauritius. The elevated charcoal at Mare Tatos in recent times  
195 likely reflects the burning of woody material brought into the area by humans,  
196 possibly harvested elsewhere on the island, as by this point little local natural  
197 vegetation remained (Fig. 1b). In the last decades, island wide degradation of  
198 ecosystems is mainly linked to growing human populations, and an increase in  
199 tourism and textile industries (Ramessur 2002).

200

201         The fire history from the Mare Tatos wetland was able to precisely detect the  
202 documented phases of human arrival, occupation, abandonment, governance and  
203 demographic change. Our finding adds weight to inferences made in the absence of  
204 corroborating historical evidence that elevated charcoal (fire) on other tropical islands  
205 indicates the arrival of human populations (e.g. Burney 1987, Premathilake 2006, Rull  
206 *et al.* 2015). The Mare Tatos fire record supports the assertion that, prior to 1598,  
207 there was no human occupation of Mauritius. If a conservation goal on the island is to  
208 restore ecosystem processes to a pre-human state then fire management is  
209 consequently key.

210

211           During c. 500 years of human occupation the ecosystems of Mauritius were  
212 degraded through forest fragmentation, fires, introduction of exotic species, and  
213 overexploitation of natural resources (Cheke & Hume 2008, Rijdsdijk *et al.* 2011,  
214 Florens *et al.* 2012, Rijdsdijk *et al.* 2015). Our research reveals that many extinction  
215 events (including Raven parrot, Dodo, giant tortoise and fruit bat) over the last 500  
216 years on Mauritius occurred during the periods of elevated human activity indicated  
217 by frequent fire events recorded in the Mare Tatos sediments (Fig. 1a). All of these  
218 species had suffered significant environmental pressures in the past, such as major  
219 drought events, and survived (de Boer *et al.* 2015). Yet within c. 50 years of human  
220 occupation, and the introduction of frequent fire events, the extinctions had begun  
221 (Fig. 1a). The timings of the individual extinction events highlights the high  
222 sensitivity of taxa to human activity, and the species-specific response to fire and  
223 habitat loss. The coincidence of the greatest species loss with the first period of  
224 elevated fire likely reflects the wide-ranging and high-level human interference on  
225 Mauritius during this period. The reconstructed fire history of Mauritius suggests  
226 that charcoal records are a helpful proxy for understanding human impacts on  
227 island ecosystems and environments.  
228  
229

230 **Acknowledgements**

231 Sediments were collected as part of the Netherlands Science Foundation (NWO)  
232 project (ALW 2300155042) granted to HH. SN received support from the Portuguese  
233 National Funds, through FCT - Fundação para a Ciência e a Tecnologia, within the  
234 project UID/BIA/00329/2013 and the research Fellowship PD/BD/114380/2016. We  
235 thank the Mauritius Sugar Industry Research Institute for logistic support in the field.  
236 We thank Claudia Baider (The Mauritius Herbarium, Réduit) for her continuous  
237 support, Julian Hume and another anonymous referee for their detailed and insightful  
238 comments which improved the quality of the manuscript.

239

240 **Data availability statement**

241 The data used in this study are archived in Data Dryad (DOI:  
242 10.5061/dryad.f1c31).

243

244 **Literature cited**

- 245 ADDISON, J., HAZAREESINGH, K. 1999. A new history of Mauritius, 3rd edn. Rose-  
246 Hill, Editions de l'océan Indien, Mauritius.
- 247 ALLEN, R.B. 1989. Economic marginality and the rise of the free population of colour  
248 in Mauritius, 1767–1830. *Slavery & Abolition* 10: 126-150. DOI:  
249 10.1080/01440398908574981
- 250 BLAAUW, M., CHRISTEN, J.A. 2011. Flexible paleoclimate age-depth models using an  
251 autoregressive gamma process. *Bayesian Analysis* 6: 457-474. DOI:  
252 10.1214/11-BA618
- 253 BROUARD, N.R. 1963. A history of woods and forests in Mauritius. J.E. Feilix, I.S.O.,  
254 Government Printer, Port Louis, Mauritius.

255 BURNEY, D.A. 1987. Late Quaternary Stratigraphic Charcoal Records from  
256 Madagascar. *Quaternary Research* 28: 274-280. DOI : 10.1016/0033-  
257 5894(87)90065-2

258 CHEKE, A., HUME, J.P. 2008. *Lost land of the Dodo: An ecological history of the*  
259 *Mascarene Islands*, 1st edn. Poyser, London.

260 CLARK, J.S., PATTERSON III, W.A. 1997. Background and local charcoal in sediments:  
261 Scales of fire evidence in the paleorecord. *In* J.S. Clark, H. Cachier, J.G.  
262 Goldammer & B. Stocks (Eds.) *Sediment Records of Biomass Burning and*  
263 *Global Change*. NATO ASI Series. Series 1: Global Environmental Change vol.  
264 51, pp. 23-48. Springer, Berlin.

265 DE BOER, E.J., VÉLEZ, M.I., RIJSDIJK, K.F., DE LOUW, P.G., VERNIMMEN, T.J., VISSER,  
266 P.M., TJALLINGII, R., HOOGHIEMSTRA, H. 2015. A deadly cocktail: How a  
267 drought around 4200 cal. yr BP caused mass mortality events at the infamous  
268 ‘Dodo swamp’ in Mauritius. *The Holocene* 25: 758-771. DOI:  
269 10.1177/0959683614567886

270 DE BOER, E.J., TJALLINGII, R., VÉLEZ, M.I., RIJSDIJK, K.F., VLUG, A., REICHAERT, G.,  
271 PRENDERGAST, A.L., DE LOUW, P.G.B., FLORENS, F.B.V., BAIDER, C.,  
272 HOOGHIEMSTRA, H. 2014. Climate variability in the SW Indian Ocean from an  
273 8000-yr long multi-proxy record in the Mauritian lowlands shows a middle to  
274 late Holocene shift from negative IOD-state to ENSO-state. *Quaternary Science*  
275 *Reviews* 86: 175-189. DOI: 10.1016/j.quascirev.2013.12.026

276 FLOORE, P.M., JAYASENA, R.M. 2010. In want of everything? Archaeological  
277 perceptions of a Dutch outstation on Mauritius (1638-1710). *Post-Medieval*  
278 *Archaeology* 44: 320-340. DOI: 10.1179/174581310X12810074246708

279 FLORENS, F. 2013. Conservation in Mauritius and Rodrigues: Challenges and  
280 achievements from two ecologically devastated oceanic islands. *In* N.S. Sodhi,  
281 L. Gibson, P.H. Raven (Eds.) Conservation Biology: Voices from the Tropics,  
282 pp. 40-50. John Wiley & Sons, Ltd, Oxford.

283 FLORENS, F.B.V., BAIDER, C., MARTIN, G.M.N., STRASBERG, D. 2012. Surviving 370  
284 years of human impact: What remains of tree diversity and structure of the  
285 lowland wet forests of oceanic island Mauritius? *Biodiversity and Conservation*  
286 21: 2139-2167. DOI: 10.1007/s10531-012-0304-4

287 FLORENS, F.B.V. 2015. Flying foxes face cull despite evidence. *Science* 350: 1325-  
288 1325. DOI: 10.1126/science.350.6266.1325-a

289 GASTON, K.J., BLACKBURN, T.M. 1995. Birds, body size and the threat of extinction.  
290 *Philosophical Transactions of the Royal Society B: Biological Sciences* 347:  
291 205. DOI: 10.1098/rstb.1995.0022

292 GRIHAULT, A. 2005. Dodo: The bird behind the legend. Imprimerie & Papeterie  
293 Commerciale, Mauritius.

294 GROVE, R.H. 1996. Green Imperialism: Colonial expansion, tropical island Edens and  
295 the origins of environmentalism. Cambridge University Press, Cambridge, UK.

296 HOGG, A.G., HUA, Q., BLACKWELL, P.G., BUCK, C.E., GUILDERSON, T.P., HEATON,  
297 T.J., NIU, M., PALMER, J., REIMER, P.J., REIMER, R., TURNEY, C.S.M.,  
298 ZIMMERMAN, S.R.H. 2013. ShCal13 Southern Hemisphere calibration, 0-50,000  
299 cal yr BP. *Radiocarbon* 55: 1889-1903. DOI: 10.2458/azu\_js\_rc.55.16783

300 HUA, Q., BARBETTI, M., RAKOWSKI, A.Z. 2013. Atmospheric radiocarbon for the  
301 period 1950-2010. *Radiocarbon* 55: 2059-2072. DOI:  
302 10.2458/azu\_js\_rc.v55i2.16177

303 HUME, J.P. 2013. A synopsis of the pre-human avifauna of the Mascarene Islands. *In*  
304 U.B. Göhlich & A. Kroh (Eds.) *Paleornithological Research 2013: Proceedings*  
305 *of the 8th International Meeting of the Society of Avian Paleontology and*  
306 *Evolution, Vienna, 2012, pp. 195-238. Natural History Museum Vienna,*  
307 *Vienna, Austria.*

308 HUME, J.P., MARTILL, D.M., DEWDNEY, C. 2004. *Palaeobiology: Dutch diaries and the*  
309 *demise of the Dodo. Nature 429: 245. DOI: 10.1038/nature02688*

310 HUTSON, A.M., RACEY, P.A. 2013. *Pteropus niger*. The IUCN Red List of Threatened  
311 *Species 2013 e.T18743A22084054.*

312 JUGGINS, S. 2005. C2 Program version 1.5. Department of Geography, University of  
313 *Newcastle, Newcastle upon Tyne, UK.*

314 KELLY, R.F., HIGUERA, P.E., BARRETT, C.M., HU, F.S. 2011. A signal-to-noise index  
315 *to quantify the potential for peak detection in sediment-charcoal records.*  
316 *Quaternary Research 75: 11-17. DOI: 10.1016/j.yqres.2010.07.011*

317 LALLJEE, B., FACKNATH, S. 2008. A study of the historical and present day changes in  
318 *land use profile, and their driving forces in Mauritius. In M.T. Mrityunjay &*  
319 *R.B. Singh (Eds.) Land use: reflection on spatial informatics agriculture and*  
320 *development, pp. 231-256. Concept Publishing Company, New Delhi, India.*

321 LUTZ, W., WILS, A.B. 1994. *People on Mauritius: 1638-1991. In W. Lutz (Ed.)*  
322 *Population-environment-development: Understanding their interactions in*  
323 *Mauritius, pp. 75-97. Springer, Berlin/Heidelberg, Germany.*

324 MOREE, P.J. 1998. *A concise history of Dutch Mauritius, 1598-1710. Kegan and Paul*  
325 *International, London, UK.*

326 NORDER, S.J., SEIJMONSBERGEN, A.C., RUGHOOPUTH, S.D.D.V., VAN LOON, E.E.,  
327 *TATAYAH, V., KAMMINGA, A.T., RIJSDIJK, K.F. 2017. Assessing temporal*

328 couplings in social–ecological island systems: Historical deforestation and soil  
329 loss on Mauritius (Indian Ocean). *Ecology and Society* 22: 29. DOI:  
330 10.5751/ES-09073-220129

331 PREMATHILAKE, R. 2006. Relationship of environmental changes in central Sri Lanka  
332 to possible prehistoric land-use and climate changes. *Palaeogeography,*  
333 *Palaeoclimatology, Palaeoecology* 240: 468-496. DOI:  
334 10.1016/j.palaeo.2006.03.001

335 RAMESSUR, R. 2002. Anthropogenic-driven changes with focus on the coastal zone of  
336 Mauritius, south-western Indian Ocean. *Regional Environmental Change* 3: 99-  
337 106. DOI: 10.1007/s10113-002-0045-0

338 R CORE TEAM. 2015. A language and environment for statistical computing, version  
339 3.2.2. R Core Team, R Foundation for Statistical Computing, Vienna, Austria.  
340 <https://www.R-project.org>.

341 REIMER, P.J., BARD, E., BAYLISS, A., BECK, J.W., BLACKWELL, P.G., RAMSEY, C.B.,  
342 BUCK, C.E., CHENG, H., EDWARDS, R.L., FRIEDRICH, M., GROOTES, P.M.,  
343 GUILDERTSON, T.P., HAFLIDASON, H., HAJDAS, I., HATTE, C., HEATON, T.J.,  
344 HOFFMANN, D.L., HOGG, A.G., HUGHEN, K.A., KAISER, K.F., KROMER, B.,  
345 MANNING, S.W., NIU, M., REIMER, R.W., RICHARDS, D.A., SCOTT, E.M.,  
346 SOUTON, J.R., STAFF, R.A., TURNEY, C.S.M., VAN DER PLICHT, J. 2013.  
347 Intcal13 and Marine13 Radiocarbon Age Calibration Curves 0-50,000 years cal  
348 BP. *Radiocarbon* 55: 1869-1887. DOI: 10.2458/azu\_js\_rc.55.16947

349 RIJSDIJK, K.F., HUME, J.P., LOUW, P.G.B.D., MEIJER, H.J.M., JANOO, A., DE BOER,  
350 E.J., STEEL, L., DE VOS, J., VAN, D.S., HOOGHIEMSTRA, H., FLORENS, F.B.V.,  
351 BAIDER, C., VERNIMMEN, T.J.J., BAAS, P., VAN HETEREN, A.H., RUPEAR, V.,  
352 BEEBEEJAUN, G., GRIHAULT, A., VAN, D.P., BESSELINK, M., LUBEEK, J.K.,



353 JANSSEN, M., KLUIVING, S.J., HOLLUND, H., SHAPIRO, B., COLLINS, M., BUCKLEY,  
354 M., JAYASENA, R.M., PORCH, N., FLOORE, R., BUNNIK, F., BIEDLINGMAIER, A.,  
355 LEAVITT, J., MONFETTE, G., KIMELBLATT, A., RANDALL, A., FLOORE, P.,  
356 CLAESSENS, L.P.A.M. 2015. A review of the Dodo and its ecosystem: Insights  
357 from a vertebrate concentration Lagerstätte in Mauritius. *Journal of Vertebrate*  
358 *Paleontology* 35: 3-20. DOI: 10.1080/02724634.2015.1113803

359 RIJSDIJK, K.F., ZINKE, J., DE LOUW, P.G.B., HUME, J.P., VAN, D.P., HOOGHIEMSTRA,  
360 H., MEIJER, H.J.M., VONHOF, H.B., PORCH, N., FLORENS, F.B.V., BAIDER, C.,  
361 VAN GEEL, B., BRINKKEMPER, J., VERNIMMEN, T., JANOO, A. 2011. Mid-  
362 Holocene (4200 kyr BP) mass mortalities in Mauritius (Mascarenes): Insular  
363 vertebrates resilient to climatic extremes but vulnerable to human impact. *The*  
364 *Holocene* 21: 1179-1194. DOI: 10.1177/0959683611405236

365 ROBERTS, D.L., SOLOW, A.R. 2003. Flightless birds: When did the Dodo become  
366 extinct? *Nature* 426: 245. DOI: 10.1038/426245a

367 RULL, V., CAÑELLAS-BOLTÀ, N., MARGALEF, O., SÁEZ, A., PLA-RABES, S., GIRALT, S.  
368 2015. Late Holocene vegetation dynamics and deforestation in Rano Aroi:  
369 Implications for Easter Island's ecological and cultural history. *Quaternary*  
370 *Science Reviews* 126: 219-226. DOI: 10.1016/j.quascirev.2015.09.008

371 TRIANTIS, K.A., BORGES, P.A.V., LADLE, R.J., HORTAL, J., CARDOSO, P., GASPAR, C.,  
372 DINIS, F., MENDONÇA, E., SILVEIRA, L.M.A., GABRIEL, R., MELO, C., SANTOS,  
373 A.M.C., AMORIM, I.R., RIBEIRO, S.P., SERRANO, A.R.M., QUARTAU, J.A.,  
374 WHITTAKER, R.J. 2010. Extinction debt on oceanic islands. *Ecography* 33: 285-  
375 294. DOI: 10.1111/j.1600-0587.2010.06203.x

376 VAUGHAN, R.E., WIEHE, P.O. 1937. Studies on the vegetation of Mauritius: I. A  
377 preliminary survey of the plant communities. *Journal of Ecology* 25: 289-343.

378 WHITLOCK, C., LARSEN, C. 2001. Charcoal as a fire proxy. *In* J.P. Smol, H.J.B. Birks,  
379 W.M. Last (Eds.) *Tracking Environmental Change Using Lake Sediments*  
380 *Volume 3: Terrestrial, Algal and Siliceous Indicators*, pp. 75-98. Kluwer  
381 Academic Publishers, Dordrecht/Boston/London.  
382

383 **Tables**

384 **TABLE 1:** Radiocarbon ( $^{14}\text{C}$ ) dates of bulk sediment samples from Mare Tatos

385 (Mauritius) used to establish the age vs. depth relationship (Fig. 1). Radiocarbon  
 386 calibrations were done using ShCal13 and post bomb (Hogg *et al.* 2013, Hua *et*  
 387 *al.* 2013, Reimer *et al.* 2013). pMC = percentage modern carbon, where modern  
 388 is AD 1950. For further information on the calibration methods see online  
 389 Supporting Information. \* = first published de Boer *et al.* (2014).

390

Laboratory code	Depth in cm	pMC ( $1\sigma$ )	$^{14}\text{C}$ years before present ( $1\sigma$ )	$1\sigma$ range in calendar years AD/BC (probability)
D-AMS 017458	14.0	114.06 ( $\pm 0.26$ )	modern	AD 1994-1992 (0.628) AD 1992-1992 (0.035) AD 1991-1991 (0.054) AD 1990-1990 (0.017) AD 1960-1959 (0.266)
D-AMS 017459	36.0	97.41 ( $\pm 0.41$ )	211 ( $\pm 34$ )	AD 1955-1950 (0.036) AD 1805-1728 (0.747) AD 1689-1664 (0.217)
D-AMS 017460	56.0	101.41 ( $\pm 0.34$ )	modern	AD 1957-1957 (0.714) AD 1956-1956 (0.286)
GrA-51609*	66.5	-	260 ( $\pm 30$ )	AD 1796-1780 (0.270) AD 1770-1763 (0.063) AD 1775-1745 (0.112) AD 1672-1643 (0.562)
GrA-51610*	180.5	-	2450 ( $\pm 30$ )	401-510 BC (1.000)

391

392 **List of Figures**

393

394 **FIGURE 1:** (A) Fire history of Mare Tatos wetland (Mauritius) since c. AD 1000  
395 compared with key historical and ecological events. Historical data: dashed  
396 black line = percentage of modern human population (1,219,265 in 2014), and  
397 solid black line = percentage of natural vegetation. Open circle = first Dutch  
398 landing in Mauritius. Open square = French revolution. Grey horizontal bars =  
399 fire zones, defined by charcoal >20% of maximum of counts in size fraction 75-  
400 160  $\mu\text{m}$  (landscape scale). Dark grey vertical dashed lines = presence/absence of  
401 fire threshold set at 5% of maximum charcoal values. Red icons and dotted  
402 horizontal lines = extinction events in the wild: Raven parrot (1680), Dodo  
403 (1690), giant tortoise (1700), and fruit bat (1795). Blue icons and dotted  
404 horizontal lines = introduction events: rats (14<sup>th</sup> Century), cats (1598), locust  
405 (1720), and Java sparrow (1740). Information on extinction and introduction  
406 dates was obtained from: Cheke & Hume (2008) and Hume (2013). \* = depth of  
407 radiocarbon date. (B) Mauritian land-cover based on historical documents. Open  
408 circle = Mare Tatos. Open square = Vieux Grand Port. Note: Historical data  
409 and maps are modified from Norder *et al.* (2017) and Floore & Jayasena (2010).  
410