

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the peer-reviewed, manuscript version of an article published in *Animal Reproduction Science*. The version of record is available from the journal site:

<http://dx.doi.org/10.21451/1984-3143-AR859>.

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Supplementing α -Linolenic acid in the in vitro maturation media improves nuclear maturation rate of oocytes and early embryonic development in the Nili Ravi buffalo

AUTHORS: Azam, A; Shahzad, Q; Asma-UI-Husna; Qadee S; Ejaz, R; Fouladi-Nashta, A A; Khalid, M; Ullah, N; Akhtar, T; Akhter, S

JOURNAL: Animal Reproduction Science

PUBLISHER: Elsevier Masson

PUBLICATION DATE: Dec. 2017

DOI: 10.21451/1984-3143-AR859

Supplementing α -Linolenic acid in the *in vitro* maturation media improves nuclear maturation rate of oocytes and early embryonic development in the

Nili Ravi buffalo

A. Azam¹, Q. Shahzad², A. Husna¹, S. Qadeer¹, R. Ijaz¹, A. A. Fouladi-Nashta³, M. Khalid³, N. Ullah⁴, T. Akhtar² and S. Akhter^{1,*}

¹Animal Physiology Laboratory, Department of Zoology, Pir Mehr Ali Shah-Arid Agriculture University, Rawalpindi-46300, Pakistan

² Buffalo Research Institute Pattoki, Kasur, Lahore-54000, Pakistan

³Reproduction, Genes and Development Research Group, The Royal Veterinary College, Hawkshead Lane, Hatfield, Herts AL9 7TA, UK.

⁴ Department of Animal Reproduction, Riphah International University, Lahore-54000 Pakistan

*** Corresponding Author:**

Shamim Akhter (Chairperson)

Associate Professor, Department of Zoology, Pir Mehr Ali Shah Arid Agriculture University, Rawalpindi-46300, Pakistan

Email: shamim@uaar.edu.pk

Cell: + 923335206056

Article Type: Applied Research

Running Title: Evaluation of α -Linolenic acid for *in vitro* maturation of buffalo oocytes

Abstract

The present study was conducted to investigate the effect of omega-3 poly unsaturated fatty acid (PUFA), α -Linolenic acid (ALA; 18:3 *n*-3) on the *in vitro* maturation (IVM) of buffalo oocytes and subsequent embryonic development. Buffalo cumulus-oocyte complexes (COCs; n=2282) were *in vitro* matured in TCM-199 (0.6 % fatty acid free bovine serum albumin, 0.02 Units/mL FSH, 1 μ g/mL 17- β -estradiol, 50 μ g/mL gentamicin) supplemented with 0 (control), 25, 50, 100, 150 or 300 μ M ALA under an atmosphere of 5 % CO₂ in air at 38.5 °C for 22-24 h. The matured oocytes were then fertilized in Tyrode's Albumin Lactate Pyruvate (TALP) medium and cultured in synthetic oviductal fluid (SOF) medium. Concentrations up to 100 μ M ALA improves ($P \leq 0.05$) the cumulus expansion compared to control. Higher percentage of oocytes reaching MII stage was observed at 50 μ M and 100 μ M of ALA compared to control ($P \leq 0.05$). Concentrations of 150 and 300 μ M ALA were detrimental both for cumulus expansion and nuclear maturation rate of buffalo oocytes. Moreover, supplementation with 100 μ M ALA improves ($P \leq 0.05$) cleavage rate compared to control and treatment with 50 μ M and 100 μ M ALA yielded significantly higher morulae compared to control. The results of present study indicate that the supplementation with 100 μ M to the IVM medium improves nuclear maturation rate of buffalo oocytes and subsequent early embryonic development.

Key words: Buffalo, α -Linolenic acid, omega-3 PUFAs, nuclear maturation, embryonic development

Introduction

The Nili-Ravi buffalo is considered to be one of the best milk producers among other breeds of buffaloes in the world (Warriach *et al.*, 2008). Although, this breed has the potential of producing more than 5000 liters of milk /lactation (Bilal *et al.*, 2006), the average milk yield is quite low, which raises the opportunities of genetic improvement through assisted

reproductive technologies like artificial insemination, and multiple ovulation and embryo transfer (MOET). However, the implementation of these technologies is hampered due to some inherent reproductive problems in buffalo such as fewer primordial follicles, which results in smaller number of recruitable follicles, high level of atresia, poor estrus behavior detection and a poor response to superovulation protocols (Madan, 1990). Recently, the emphasis has now been shifted towards the *in vitro* production (IVP) of buffalo embryos (Hansen, 2006) utilizing germ plasm of both male and female animals simultaneously (Barakat *et al.*, 2012).

In vitro maturation (IVM) is a key step during which oocytes undergo all the necessary changes required for successful fertilization and further embryonic development (Wang *et al.*, 1997). Therefore, the maturation medium is crucial, not only for the maturation of oocyte itself but also for its further competence after IVF (Bavister *et al.*, 1992; Kharche *et al.*, 2006). Previously, several studies have evaluated different types of media (Totey *et al.*, 1993; Abdoon *et al.*, 2001; Zicarelli *et al.*, 2003), and its supplements including different sources of protein (Chauhan *et al.*, 1998), hormones (Totey *et al.*, 1993), thiol compounds (Gasparrini *et al.*, 2006; Zicarelli *et al.*, 2005), growth factors (Purohit *et al.*, 2005) and antioxidants (Ullah *et al.*, 2006) for *in vitro* maturation of buffalo oocytes with variable success rates and blastocyst production rate of no more than 10-20% (Kumar and Anand, 2012) which is relatively lower when compared to 30-40 % blastocyst production rate in cattle (Rizos *et al.*, 2008).

Fatty acids are essential for numerous physiological functions (Wathes *et al.*, 2007), such as membrane biosynthesis (Sturmey *et al.*, 2009), signal transduction and gene expression (Sampath and Ntambi, 2005). They also provide a potent energy source and prevent the lipotoxic effects result in from increased cellular contents of saturated fatty acids through lipid storage and β -oxidation (Dunning *et al.*, 2014). Fatty acids are also precursors for prostaglandins and progesterone synthesis and therefore play an important role in the regulation of normal reproductive function (Abayasekara and Wathes, 1999; Mattos *et al.*, 2000).

Polyunsaturated fatty acids (PUFAs) play a significant role in increasing the number (Lucy *et al.*, 1991) and size of ovarian follicles (Zeron *et al.*, 2002), level of LH (Lucy *et al.*, 1991) and progesterone in follicular fluid (Ryan *et al.*, 1992), oocyte quality (Zeron *et al.*, 2002), regulation of ovulation, CL function (Abayasekara and Wathes, 1999; Mattos *et al.*, 2000) and pregnancy rate (Bellows *et al.*, 1999).

PUFAs must be provided by diet, since their *in vivo* synthesis is not possible due to absence of proper enzymes (Gurr *et al.*, 2002). Change in the composition of dietary fatty acids not only modifies fatty acid composition in the blood plasma but also of the reproductive tissues including, follicular fluid, cumulus cells and the oocytes (Zeron *et al.*, 2002; Ferguson and Leese, 1999; Bilby *et al.*, 2006; Child *et al.*, 2008; Fouladi-Nashta *et al.*, 2009; Wonnacott *et al.*, 2010), which can directly influence the competence of oocytes for further development and/or fertility (Wonnacott *et al.*, 2010; Armstrong *et al.*, 1990; Burke *et al.*, 1997; Petit *et al.*, 2001). For instance, dietary fish oil supplementation has been shown to alter the specific n-3 and n-6 fatty acids in follicular fluid, cumulus cells and oocytes in both beef heifers (Childs *et al.*, 2008) and ewes (Zeron *et al.*, 2002; Wonnacott *et al.*, 2010). Such changes in the fatty acid profile may improve oocyte maturation, which is essential for successful fertilization and further embryo development (Marei *et al.*, 2009) Fatty acids supplemented into IVM medium yielded positive effects on oocyte maturation, fertilization, and embryonic development in the rat (Khandoker and Tsujii, 1999), goat, sheep (Veshkini *et al.*, 2015) and cattle (Kim *et al.*, 2001).

Alpha-Linolenic acid (ALA; C18:3) is the dietary precursor for the long-chain omega-3 PUFAs (Brenna *et al.*, 2009). It plays an important role in folliculogenesis, developmental competence of oocyte (Moallem *et al.*, 2013), fertilization rate and embryo quality (Thangavelu *et al.*, 2007). As ALA is produced by the ovarian follicles and the amount of ALA increases as the ovarian follicles enlarge (Veshkini *et al.*, 2015); a role of specific unsaturated fatty acids

like ALA may be speculated in the oocyte maturation and/or follicular growth. In fact, ALA has been shown to improve the fertility rate in both cattle and sheep by improving folliculogenesis and fertilization rate *in vivo* (Moallem *et al.*, 2013). When supplemented in the *in vitro* maturation medium it was reported to regulate the molecular mechanism leading to increased number of MII stage oocytes and improved their subsequent development into early embryos in both cattle (Fouladi-Nashta *et al.*, 2009; Marei *et al.*, 2009) and sheep (Ghaffarilaleh *et al.*, 2014). However, the role of ALA supplementation of IVM media for Nili Ravi buffalo oocytes has not been studied yet. In this study we hypothesized that the supplementation of ALA in IVM media improves the quality of *in vitro* matured buffalo oocytes and their subsequent development post-fertilization. The main objective of the study was to evaluate the effects of ALA supplementation of the IVM media on the *in vitro* oocyte maturation and subsequent embryonic development in buffalo.

Materials and Methods

Collection of Oocytes from Buffalo Ovaries

Buffalo ovaries (n = 2500) were collected from the local abattoir and within two hours of collection transferred to the laboratory in a thermos containing sterilized phosphate buffered saline (PBS) kept at 33-35 °C. Fresh PBS was used to wash the ovaries immediately after arrival. Sterile disposable plastic syringe (10 mL) fitted with 18 gauge needle was used to aspire cumulus-oocyte complexes (COCs) from antral follicles (2-8 mm). Searching for COCs was done under stereomicroscope in PBS and was classified as grade A, B, C and D, on the basis of their cumulus investment and ooplasm homogeneity (Sabasthin *et al.*, 2013). Number of

COCs (n = 2282) with homogenous ooplasm and more than 4 compact layers of cumulus cells were selected for the experiments.

In vitro Maturation (IVM) of Oocytes

COCs were in vitro matured in 100 µL drops covered with sterile mineral oil for 24 h at 38.5 °C in 5 % CO₂ in air with 95 % humidity. All the media and culture dishes were equilibrated at 38 °C in CO₂ incubator for at least 1-2 h before experiment.

Assessment of cumulus cell expansion

After 24 h of maturation, cumulus cell expansion was assessed by visual assessment using stereomicroscope as 1) not expanded (No expansion observed in cumulus cells) 2) Partially expanded (Expansion of outer layers of cumulus cells only) or 3) Fully expanded (Expansion of all layers of cumulus cells) (Azam *et al.*, 2017).

Oocyte staining and determination of stage of nuclear maturation

For determination of nuclear stage in meiosis, COCs were completely denuded. Denuded oocytes were washed twice in PBS and fixed overnight in aceto-ethanol. Oocytes were placed on a grease free glass slide and covered with a cover slip. Oocytes were slightly compressed onto the glass slide with a needle until drop containing oocytes touched the cover slip. The oocytes were then stained with 1% aceto-orcein and de-stained with aceto-glycerol as described by Azam *et al.* (2017). Oocyte's nuclear maturation was evaluated with a phase contrast microscope at 200X to 400X magnification.

Based on morphology, nuclear maturation was categorized as described by Azam *et al.* (2017). Oocyte nucleus with a nucleolus and filamentous chromatin is at germinal vesicle (GV) stage. Oocyte without nucleolus, nuclear membrane and shortening of chromosomes is at germinal vesicle breakdown (GVBD) stage. At MI stage chromosomes look as thick dots arranged at metaphase plate. The segregation of chromosomes start at anaphase-I and at Telophase-I, complete segregation of chromosomal sets occur. First polar body is released at M-II.

In vitro Fertilization (IVF) of Oocytes

Frozen semen was used for *in vitro* fertilization (IVF). Three 0.5 mL straws of cryopreserved buffalo semen were thawed in water bath at 37 °C for 30 seconds. Thawed semen was placed in a 15 mL conical tube. Spermatozoa with maximum motility were collected by swim up technique (Parrish *et al.*, 1986). About 250 µL of thawed semen was deposited at the bottom of four 15 mL tubes containing 3 mL of pre warmed sperm wash medium (TALP: modified calcium-free Tyrode's Albumin Lactate Pyruvate with 6 mg/ml BSA fraction-V). Tubes were incubated at 45° angle for 30 minutes. Supernatant from each tube was removed and transferred into another 15 mL conical tube and centrifuged at 1600 rpm for 10 minutes. The pellet obtained after centrifugation of supernatant was assessed for sperm motility, and concentration was determined in an improved Neubauer counting chamber. Subsequently, sperm suspension was resuspended in pre-warmed fertilization TALP (supplemented with 0.1mM hypotaurine, 0.2mM penicillamine, 0.01mM epinephrine, 10 µg/mL heparin) to a final concentration of 2×10^6 cells mL⁻¹.

After 24 h of IVM, buffalo oocytes were washed in fertilization media and were placed in fertilization droplets (5 COCs / 50 µL droplet) of pre warmed fertilization media under mineral oil. Oocytes and spermatozoa were co-incubated for 20 h at 38.5 °C under 5 % CO₂

with maximum humidity (Gasparrini *et al.*, 2008). The day of fertilization was defined as Day 0.

In vitro Culture (IVC) of embryos

Following fertilization, presumptive zygotes were denuded by vigorous pipetting in PBS and transferred to 25 μ L drops containing IVC media (SOF) supplemented with BME, MEM and 5% fetal calf serum (FCS). Embryo culture was performed at 38.5 °C in a humidified incubator with 5 % CO₂ in air. On day 2 the cleavage rate (number of oocytes cleaved/total COCs incubated \times 100) was evaluated. Further developmental stages (4-8 cell stage, > 8 cell stage and morula) were evaluated and recorded every other day.

Experimental Design

Linolenic acid (ALA; Stock solution in DMSO) was added at different levels (0 μ M (control), 25 μ M, 50 μ M, 100 μ M, 150 μ M, 300 μ M) to the oocytes in maturation medium (TCM-199) supplemented with 0.6 % fatty acid free-bovine serum albumin (FAF-BSA), 0.02 IU/mL FSH, 1 μ g/mL estradiol-17 β (E2) and 50 μ g/mL gentamicin.

Experiment 1

Oocytes (1200) were randomly allocated to the experimental groups as described: 1) 0 μ M ALA (control); 2) 25 μ M ALA (25); 3) 50 μ M ALA (50); 4) 100 μ M ALA (100); 5) 150 μ M ALA (150); or 6) 300 μ M ALA (300). The experiment was replicated for seven times.

Experiment 2

Oocytes (1082) were matured as in the Experiment 1. Matured oocytes were fertilized and cultured. The experiment was replicated seven times. Embryonic development was assessed as cleaved, 4-8 cell stage, > 8 cell stage and morula formation.

Statistical Analysis

To compare the effect of different doses of ALA data on COCs expansion, MII stage, cleavage rate and developmental stages (2 cell stage, 4-8 cell stage, > 8 cell stage and morula) were recorded and analyzed by one-way analysis of variance (ANOVA). Results were considered significant at ($P < 0.05$). Duncan's multiple range test (DMRT) was used to compare the treatment means.

Results

Experiment 1

The data on the effect of ALA supplementation of maturation media on cumulus expansion are shown in Fig. 1 and Fig. 2. The addition of 25 μM ALA in maturation media did not show improvement ($P > 0.05$) in cumulus expansion compared to control. However, supplementation with 50 and 100 μM ALA resulted in increased rate of cumulus expansion ($P \leq 0.05$) compared to control. The higher levels of ALA, *i.e.*, 150 and 300 μM resulted in significant decrease ($P < 0.05$) in expansion of cumulus cells compared to other groups (Fig. 1 and 2).

Data of the effects of ALA supplementation in maturation media on percentage of oocytes at different stages of nuclear maturation are shown in Fig. 3. The higher percentage of oocytes reaching MII stage ($P \leq 0.05$) was observed with 50 μM ALA ($67.0\% \pm 2.2$) compared to control ($57.4\% \pm 2.5$) and 25 μM ALA ($60.3\% \pm 1.9$) in IVM medium. Further improvement ($P \leq 0.05$) was observed with 100 μM ALA ($76.3\% \pm 2.5$). The higher concentrations of ALA i.e., The ALA concentrations of 150 μM ($51.3\% \pm 1.8$) and 300 μM ($29.8\% \pm 2.3$) resulted in a dose dependent decrease in percentage of oocytes reaching MII stage compared to 100 μM ALA. The percentage of oocytes remaining at GVBD and MI stage were recorded higher in 300 μM ALA ($26.2\% \pm 2.2$ and $39.1\% \pm 1.4$) compared to control ($17.7\% \pm 3.1$ and $20.7\% \pm 2.4$), respectively. Percentage of degenerative oocytes was higher ($p < 0.05$) with 300 μM ALA ($4.9\% \pm 2.5$) compared to control, 25 μM ALA and 150 μM ALA ($1.9\% \pm 1.2$, $1.8\% \pm 1.2$, $1.0\% \pm 1.0$) respectively. No degenerative oocytes were observed with 50 μM ALA and 100 μM ALA.

Experiment 2

Data on the effect of ALA supplementation on percentage of cleavage, 4-8 cells, > 8 cells and morula are presented in Table 1. Supplementation with 25, 50 and 150 μM ALA (49.7 to 56.9%) did not improved ($P > 0.05$) the cleavage rate compared to the controls ($50.4\% \pm 2.0$). The addition of 100 μM ALA (49.7% to 57.8%) raised ($P < 0.05$) the cleavage rate compared to the controls and 150 μM ALA, but did not do so when compared to 25 and 50 μM ALA. A significant decrease ($P < 0.05$) in cleavage rate was recorded when maturation media was supplemented with 300 μM ALA ($21.0\% \pm 2.5$).

Embryos reaching 4-8 cell stage were higher ($P \leq 0.05$) when the oocytes were matured in the presence of ALA at 100 μM (47.0 ± 2.9) compared to control (34.0 ± 1.1) and to 25 μM

ALA (36.6 ± 3.8). At 50 μM ALA (42.3 ± 1.5) embryos developed up to 4-8 cell stage differ significantly ($P < 0.05$) with control and to 150 μM ALA ($31.2\% \pm 0.8$), while did not differ ($P > 0.05$) from 25 μM and 100 μM . The higher concentration used in this study, 300 μM (14.3 ± 1.8) resulted in the worst decrease ($P \leq 0.05$) on number of 4-8 cell embryos.

Recorded number of embryos that reached more than 8 cell stage was higher ($P \leq 0.05$) when the oocytes matured in media supplemented with 100 μM ALA ($38.0\% \pm 2.2$) followed by 50 μM ALA (31.1 ± 1.6) and by control (22.2 ± 2.9) and 25 μM ALA (23.1 ± 2.4). The increase in ALA supplementation up to 150 μM (14.4 ± 1.8) and 300 μM (1.2 ± 1.2) compromised ($P \leq 0.05$) the embryo development to >8 cell embryos compared to other groups.

The percentage of morulae was increased ($P < 0.05$) with the supplementation with 50 μM ALA (17.8 ± 2.3) and 100 μM (21.0 ± 2.6) compared to control ($P \leq 0.05$) percentage of morulae compared to the control (11.2 ± 2.6) and 25 μM ALA (12.1 ± 2.1), while 150 μM ALA (7.8 ± 0.2) in maturation media resulted in decreased ($P \leq 0.05$) number of morulae. None of the embryos developed to morula stage ($P > 0.0$) when the oocytes were matured with 300 μM ALA.

Discussion

The fatty acid content of developing oocytes changes depending on the environment in which they develop (Wakefield *et al.*, 2008). Changing the micro-environment of oocyte also changes intra-cytoplasmic concentration of fats. Therefore, PUFAs in the micro-environment may replace saturated FAs in the oocyte cytoplasm and make the oocyte membrane comparatively more permeable for sperm entry and increase fertilization rate and developmental competence of the oocyte. Present study shows that supplementation of maturation media with 100 μM ALA increased the number of fully expanded oocytes.

Supporting our current findings, a previous study (Marei *et al.*, 2009) demonstrated that supplementation of maturation medium with ALA at a concentration of lower than 100 μM had no adverse effect on cumulus cell expansion in cattle. The deleterious effect of ALA at a concentration of 150 and 300 μM in the present study is consistent with the previous studies (Coyral- Castel *et al.*, 2010; Ghaffarilaleh *et al.*, 2014; Veshkini *et al.*, 2015) that reported decreased viability of granulosa cells with higher concentration of ALA after 24 h of maturation. In the present study, high levels of ALA in maturation medium might have been outside the tolerance threshold of the cells, exerted toxic effect leading to a reduction in cell viability (Andrade *et al.*, 2005).

Cumulus cells are important for keeping the oocytes under meiotic arrest, inducing meiotic resumption and supporting cytoplasmic maturation. These functions have been attributed to their gap junctions and their specific metabolizing capabilities (Tanghe *et al.*, 2002). It is anticipated that cumulus cells mediate signals and regulate the synthesis of important cytoplasmic factors which support nuclear maturation (Albertini *et al.*, 2001). However, it is not fully understood that expansion of cumulus cells is prerequisite for nuclear maturation or both processes are not inter-related. It has been reported in the ovine that cumulus cell expansion is prerequisite for nuclear maturation (Amini *et al.*, 2016), at the same time no such connection has also been reported in the bovine (Khalil *et al.*, 2013; Marei *et al.*, 2010).

Meiotic competence of oocytes determines the developmental capacity for subsequent embryo development. In the present study, supplementation of maturation medium with 100 μM ALA increased the MII rate of buffalo oocytes. High doses of ALA in IVM medium inhibited maturation progression, manifested by high percentage of oocytes arrested at GVBD or MI stage and a concomitant decrease in percentage of oocytes that completed second meiotic division metaphase. Our results are in agreement with the results of Veshkini *et al.* (2015) who reported improved nuclear maturation in sheep oocytes at 100 μM ALA in maturation medium.

The positive effect of ALA on oocyte's meiotic competence may be mediated directly by improvement of cytoplasmic maturation via the mitogen-activated protein kinase pathway and indirectly through PGE2 synthesis (Marei *et al.*, 2009). The PGE2 is a key paracrine and/or autocrine regulator of cumulus cell functions and has been proposed to play a major role in oocyte nuclear maturation (Elvin *et al.*, 2000). An elevated concentration of PGE2 positively stimulated the extent of cumulus expansion and nuclear maturation by enhancing the phosphorylation of MAPK1 and MAPK2 in both oocytes and cumulus cells through the elevation of cAMP levels (Marei *et al.*, 2009).

In addition to the positive effects observed on cumulus expansion and oocyte nuclear maturation, improvement in all developmental stages of buffalo embryos was observed when oocytes were matured in the presence of 50-100 μ M ALA in the maturation media. Similar to the findings of our study, an increased rate of embryonic development has been observed with supplementation of ALA in maturation media in the bovine (Marei *et al.*, 2009), sheep (Ghaffarilaleh *et al.*, 2014) and goat oocytes (Veshkini *et al.*, 2015). Provision of ALA in maturation media might have improved the cellular processes during different phases of maturation, which in turn influenced the developmental competence of subsequent embryos (Krisher *et al.*, 1998). In another study on the bovine, addition of ALA to the culture medium did not improve the cleavage and blastocyst rates (Al Darwich *et al.*, 2010), however, the viability of the embryos was influenced in a positive manner by the presence of ALA in culture medium (Al Darwich *et al.*, 2010). It is relevant to mention that the improvement in embryo development rate in present study can only be attributed to supplementation of ALA during oocyte maturation as maturation medium was serum free and there was no supplementation during *in vitro* embryo culture. These findings suggest that ALA might have been more important to the oocytes if added to the maturation medium rather than to the culture medium after fertilization.

In the present study, ALA at higher concentrations (300 μ M ALA) resulted in reduced number of cleaved embryos and no morula stage embryos. It can be anticipated that supplementation of PUFAs (such as ALA) in the maturation medium resulted in their accumulation in oocytes cell membranes. Since PUFAs are more susceptible to peroxidation than monounsaturated and saturated fatty acids, supplementation at higher levels might have increased the oxidative stress (Song *et al.*, 2000; Gladine *et al.*, 2007). Higher levels of ALA might have disturbed the balance of unsaturated and saturated fatty acids, so it might be the ratio of saturated vs unsaturated FAs that is more critical in the oocyte's micro-environment rather than a certain concentration of the unsaturated FAs per se. Therefore, one may speculate that the higher concentrations of ALA in our study might have generated oxidative stress after oxidative phosphorylation of ALA (Wakefield *et al.*, 2008) which might have compromised the developmental competence of these oocytes and be responsible for the observed reduction of embryonic growth. Further studies may be suggested to investigate whether antioxidant supplementation of ALA-supplemented IVM medium could abrogate negative effects as observed for higher concentrations of ALA. In conclusion, the results of the present study have shown that supplementation of *in vitro* maturation media with ALA at 100 μ M improves the *in vitro* maturation of buffalo oocytes that in turn improves early embryonic development.

Acknowledgements

The authors thank Higher Education Commission, Pakistan, for financial support under PhD 5000 indigenous Fellowship Program.

References

- Abayasekara DR, Wathes DC.** 1999. Effects of altering dietary fatty acid composition on prostaglandin synthesis and fertility. *Prostaglandins Leukot. Essent. Fatty Acids*, 61:275-87.
- Abdoon ASS, Kandil OM, Otoi T, Suzuki T.** 2001. Influence of oocyte quality, culture media and gonadotropins on cleavage rate and development of *in vitro* fertilized buffalo embryos. *Anim Reprod Sci*, 65:215-223.
- Albertini DF, Combelles CM, Benecchi E, Carabatsos MJ.** 2001. Cellular basis for paracrine regulation of ovarian follicle development. *Reproduction*, 121: 647-653.
- Al Darwich A, Perreau C, Petit MH, Papillier P, Dupont J, Guillaume D, Mermillod P, Guignot F.** 2010. Effect of PUFA on embryo cryoresistance, gene expression and AMPK alpha phosphorylation in IVF-derived bovine embryos. *Prostag Oth Lipid M*, 93:30-6.
- Amini E, Asadpour R, Roshangar L, Jafari-Joozani R.** 2016. Effect of linoleic acid supplementation on in vitro maturation, embryo development and apoptotic related gene expression in ovine. *Int J Reprod BioMed*, 14:255-262.
- Armstrong JD, Goodall EA, Gordon FJ, Rice DA, McCaughey WJ.** 1990. The effects of levels of concentrate offered and inclusion of maize gluten or fish meal in the concentrate on reproductive performance and blood parameter of dairy cows. *Anim Prod*, 50:1-10.
- Azam A, Husna AU, Shahzad Q, Akhtar T, Haq E, Ullah N, Akhter S.** 2017. Efficiency of Fatty Acid-Free Defined System for In Vitro Maturation of Buffalo Oocytes. *J. Anim. Plant Sci*, 27(1): 112-118.
- Barakat IAH, El-Ashmaoui HM, Barkawi A, Kandeal SA, EL-Nahass E.** 2012. Ultra-structural study of Egyptian Buffalo oocytes before and after *in vitro* maturation. *Afr J Biotech*, 11:7592-7602.

Bavister BD, Rose-Hellekant TA, Pinnyomminter T. 1992. Development of *in vitro* and matured *in vitro* fertilized bovine embryos into morula and blastocyst in defined culture media. *Theriogenology*, 37:127-146.

Bellows RA, Simms DD, Grings EE, Phelps DA, Bellows SE, Bellows NR, Short RE, Funston RN, Geary TW. 1999 Effects of feeding supplemental fat during gestation on reproduction in primiparous beef heifers. *J Anim Sci*, 77:236.

Bilal MQ, Suleman M, Raziq A. 2006. Buffalo: Black gold of Pakistan. *Livest Res Rural Dev*, 18:140-151.

Bilby TR, Block J, Amaral BCD, Filho OS, Silvestre FT, Hansen PJ, Staples CR, Thatcher WW. 2006. Effects of dietary unsaturated fatty acids on oocyte quality and follicular development in lactating dairy cows in summer. *J Dairy Sci*, 89:3891-3903.

Brenna JT, Salem NJ, Sinclair AJ, Cunnane SC. 2009. alpha-Linolenic acid supplementation and conversion to n-3 long-chain polyunsaturated fatty acids in humans. *Prostaglandins Leukot Essent Fatty Acids*, 80:85-91.

Burke JM, Staples CR, Risco CA, Sota RLDL, Thatcher WW. 1997. Effect of ruminant grade menhaden fish meal on reproductive performance of lactating dairy cows. *J Dairy Sci*, 80:3386-3398.

Chauhan MS, Singla SK, Palta P, Manik RS, Madan ML. 1998. *In vitro* maturation and fertilization, and subsequent development of buffalo (*Bubalus bubalis*) embryos: effects of oocyte quality and type of serum. *Reprod Fertil Dev*, 10:173-177.

Childs S, Hennessy AA, Sreenan JM, Wathes DC, Cheng Z, Stanton C, Diskin MG, Kenny DA. 2008. Effect of level of dietary n-3 polyunsaturated fatty acid supplementation on systemic and tissue fatty acid concentrations and on selected reproductive variables in cattle. *Theriogenology*, 70:595-611.

Dunning KR, Russell DL, Robker RL. 2014. Lipids and oocyte developmental competence: the role of fatty acids and β -oxidation. *Reproduction*, 148:15-27.

Ferguson EM, Leese HJB. 1999. Triglyceride content of bovine oocytes and early embryos. *J Reprod Fertil*, 116:373-378.

Fouladi-Nashta AA, Wonnacott KE, Gutierrez CG, Gong JG, Sinclair KD, Garnsworthy PC, Webb R. 2009. Oocyte quality in lactating dairy cows fed on high levels of n-3 and n-6 fatty acids. *Reproduction*, 138:771-781.

Gasparrini B, Boccia L, Marchandise J, Palo RD, George F, Donnay I, Zicarelli L. 2006. Enrichment of *in vitro* maturation medium for buffalo (*Bubalus bubalis*) oocytes with Thiol compounds: effects of cystine on glutathione synthesis and embryo development. *Theriogenology*, 65:275-287.

Gasparrini B, Rosa AD, Attanasio L, Boccia L, Palo RD, Campanile G, Zicarelli L. 2008. Influence of the duration of *in vitro* maturation and gamete co-incubation on the efficiency of *in vitro* embryo development in Italian Mediterranean Buffalo (*Bubalus bubalis*). *Anim Reprod Sci*, 105:354-364.

Ghaffarilaleh V, Fouladi-Nashta A, Paramio M. 2014. Effect of α -linolenic acid on oocyte maturation and embryo development of pre-pubertal sheep oocytes. *Theriogenology*, 82:686-696.

Gurr MI, Harwood JL, Frayn KN. 2002. Lipid biochemistry: an introduction, 5th Ed, Blackwell Science, Oxford, United Kingdom.

Hansen PJ. 2006. Realizing the promise of IVF in cattle, an overview. *Theriogenology*, 65:119-125.

Khalil WA, Marei WFA, Khalid M. 2013. Protective effects of antioxidants on linoleic acid-treated bovine oocytes during maturation and subsequent embryo development. *Theriogenology*, 80:161-823.

- Khandoker MAM, Tsujii H.** 1999. Effect of exogenous fatty acids on *in vitro* development of rat embryos. *Asian Australas J Anim Sci*, 12:169-173.
- Kharche SD, Goel AK, Jindal SK, Sinha NK.** 2006. *In vitro* maturation of caprine oocytes in different concentrations of estrous goat serum. *Small Rumin Res*, 64:186-189.
- Kim JY, Kinoshita M, Ohnishi M, Fukui Y.** 2001. Lipid and fatty acid analysis of fresh and frozen thawed immature and *in vitro* matured oocytes. *Reproduction*, 122:131-138.
- Krisher RL, Bavister BD.** 1998. Responses of oocytes and embryos to the culture environment. *Theriogenology*, 49:103-114.
- Kumar D, Anand T.** 2012. *In vitro* Embryo Production in Buffalo: Basic Concepts. *Journal of Buffalo Science*, 1:50-54.
- Lopes CN, Scarpa AB, Cappellozza BI, Cooke RF, Vasconcelos JLM.** 2009. Effects of rumen-protected polyunsaturated fatty acid supplementation on reproductive performance of *Bos indicus* beef cows. *J Anim Sci*, 87:3935-3943.
- Lucy MC, Staples CR, Michel FM, Thatcher WW.** 1999. Energy balance and size and number of ovarian follicles detected by ultrasonography in early postpartum dairy cows. *J Dairy Sci*, 74:473-482.
- Madan ML.** 1990. Conservation of germplasm through embryo transfer. In: Proceedings of the 13th Inter Dairy Congress, Montreal. p. 302-314.
- Marei WF, Wathes DC, Fouladi-Nashta AA.** 2009. The effect of linolenic acid on bovine oocyte maturation and development. *Biol Reprod*, 81:1064-1072.
- Marei WF, Wathes DC, Fouladi-Nashta AA.** 2010. Impact of linoleic acid on bovine oocyte maturation and embryo development. *Reproduction*, 139:97-88.
- Marei WF, Wathes DC, Fouladi-Nashta AA.** 2012. Differential effects of linoleic and alpha-linolenic fatty acids on spatial and temporal mitochondrial distribution and activity in bovine oocytes. *Reprod Fertil Dev*, 24:679-690.

Mattos R, Staples CR, Thatcher WW. 2000. Effects of dietary fatty acids on reproduction in ruminants. *Rev Reprod*, 5:38-45.

Moallem U, Shafran A, Zachut M, Dekel I, Portnick Y, Arieli A. 2013. Dietary α -linolenic acid from flaxseed oil improved folliculogenesis and IVF performance in dairy cows, similar to eicosapentaenoic and docosahexaenoic acids from fish oil. *Reproduction*, 146:603-614.

Parrish JJ, Susko-Parrish JL, Leibfreid-Rutledge ML, Crister ES, Eystone WH, First NL. 1986. Bovine *in vitro* fertilization with frozen-thawed semen. *Theriogenology*, 25:591-600.

Petit HV, Dewhurst RJ, Proulx JG, Khalid M, Haresign W, Twagiramungu H. 2001. Milk production, milk composition, and reproductive function of dairy cows fed different fats. *Can J Anim Sci*, 81:263-271.

Purohit GN, Brady MS, Sharma SS. 2005. Influence of EGF and IGF-1 on maturation and fertilization of buffalo cumulus oocyte complex in serum free media and their subsequent development *in vitro*. *Anim Reprod Sci*, 87:229-239.

Rizos D, Clemente M, Bermejo-Alvarez P, Fuente J de La, Lonergan P, Gutierrez-Adan A. 2008. Consequences of *In vitro* Culture Conditions on Embryo Development and Quality. *Reprod Dom Anim*, 43:44-50.

Ryan DP, Spoon RA, Williams GL. 1992. Ovarian follicular characteristics, embryo recovery, and embryo viability in heifers fed high fat diets and treated with follicle-stimulating hormone. *Journal of Animal Science*, 70:3505-3513.

Sabasthin A, Nandi S, Kumar VG, Gowda PSUS, Murthy VC. 2013. Effect of sera of normal cycling, pregnant and repeat breeding buffaloes (*Bubalus bubalis*) on *in vitro* maturation of buffalo, sheep and goat oocytes. *Asian Pac J Reprod*, 2:110-113.

Sampath H, Ntambi JM. 2005. Polyunsaturated fatty acid regulation of genes of lipid metabolism. *Annu Rev Nutr*, 25:317-340.

- Sturmey R, Reis A, Leese H, McEvoy T.** 2009. Role of fatty acids in energy provision during oocyte maturation and early embryo development. *Reprod Domest Anim*, 44:50-8.
- Tanghe S, Soom AV, Nauwynck H, Coryn M, DeKruif A.** 2002. Minireview: Functions of the cumulus ophorus during oocyte maturation, ovulation and fertilization. *Mol Reprod Dev*, 61:414-424.
- Thangavelu G, Colazo MG, Ambrose DJ, Oba M, Okine EK, Dyck MK.** 2007. Diets enriched in unsaturated fatty acids enhance early embryonic development in lactating Holstein cows. *Theriogenology*, 68:949-957.
- Totey SM, Pawshe CH, Singh GP.** 1993. *In vitro* maturation and fertilization of buffalo oocytes (*Bubalus bubalis*): Effect of media, hormone and sera. *Theriogenology*, 39:1153-1171.
- Ullah I, Jalali S, Shami SA, farooq K, Khan MI.** 2006. Effect of 2-Mercaptoethanol on Nili Ravi buffalo oocytes on *in vitro* maturation. *J Anim Vet Adv*, 5: 380-385.
- Veshkini A, Asadi H, Khadem AA, Mohammadi-Sangcheshmeh A, Khazabi S, Aminafshar M, Deldar H, Soleimani M, Cinar MU.** 2015. Effect of Linolenic acid during *in vitro* maturation of ovine oocytes: embryonic developmental potential and mRNA abundances of genes involved in apoptosis. *J Assist Reprod Genet*, 32:653-9.
- Wakefield SL, Lane M, Schulz SJ, Hebart ML, Thompson JG, Mitchell M.** 2008. Maternal supply of omega-3 polyunsaturated fatty acids alters mechanisms involved in oocyte and early embryo development in the mouse. *Am J Physiol Endocrinol Metab*, 294:425-434.
- Wakle SS, Bakshi SA, Deopurkar VL.** 2002. *In vitro* maturation of buffalo oocyte in TCM-199 and DPBS media supplemented with BSA. *Indian Vet J*, 79:28-29.
- Wang WH, Hosoe M, Shioya Y.** 1997. Induction of cortical granule exocytosis of pig oocytes by spermatozoa during meiotic maturation. *J Reprod Fertil*, 109:247-255.

Warriach HM, Channa AA, Ahmad N. 2008. Effect of oestrus synchronization methods on oestrus behaviour, timing of ovulation and pregnancy rate during the breeding and low breeding season in Nili-Ravi buffaloes. *Anim Reprod Sci*, 107:62-67.

Wathes DC, Abayasekara DR, Aitken RJ. 2007. Polyunsaturated fatty acids in male and female reproduction. *Biol Reprod*, 77:190-201.

Wonnacott KE, Kwong WY, Hughes J, Salter AM, Lea RG, Garnsworthy PC, Sinclair KD. 2010. Dietary omega-3 and omega-6 polyunsaturated fatty acids affect the composition and development of sheep granulosa cells, oocytes and embryos. *Reproduction*, 139:57-69.

Zeron Y, Sklan D, Arav A. 2002. Effect of polyunsaturated fatty acid supplementation on biophysical parameters and chilling sensitivity of ewe oocytes. *Mol Reprod Dev*, 61:271-278.

Zicarelli L, Donnay I, De Rosa A, Boccia L, Monaco E, Attanasio L, Gasparrini B. 2005. Use of thiol compounds during *in vitro* maturation of buffalo oocytes: effects on embryo development. *Ital J Anim Sci*, 4:304-306.

Zicarelli L, Neglia G, di Brienza VC, Papaccio G, Esposito L, Gasparrini B. 2003. Buffalo (*Bubalus bubalis*) *in vitro* embryo production in two different defined culture media. *Italian J Anim Sci*, 1:136-138.

List of abbreviations used

ALA	α -Linolenic acid
BSA	Bovine Serum Albumin
COCs	Cumulus Oocyte Complexes
CO ₂	Carbon dioxide
EGF	Epidermal Growth Factor
FAF-BSA	Fatty acid free-Bovine serum albumin

FCS	Fetal Calf Serum
FA	Fatty Acid
GVBD	Germinal Vesicle Breakdown
IVC	<i>In vitro</i> culture
IVEP	<i>In vitro</i> embryo production
IVF	<i>In vitro</i> fertilization
IVM	<i>In vitro</i> maturation
M199	Medium 199
PBS	Phosphate buffered saline
SOF	Synthetic oviductal fluid
TALP	Tyrode's Albumin Lactate Pyruvate medium
μg	Micro gram
μM	Micro mole
μL	Micro litre
μg/μL	Micro gram per micro litre
mM	Milli Mole
mL ⁻¹	Per milli litre
α	Alpha
β	Beta

List of figures

Fig. 1: Representative pictures of COCs after 24 h of maturation.

Fig. 2: Cumulus expansion rate of buffalo COCs after 24 h of maturation in IVM medium supplemented with different concentrations of ALA. Data are shown as Mean percentage \pm SEM. Bars with different superscripts differ significantly ($P<0.05$). 0 (Control), 25 (supplementation with 25 μ M ALA in the maturation media), 50 (supplementation with 50 μ M ALA in the maturation media), 100 (supplementation with 100 μ M ALA in the maturation media), 150 (supplementation with 150 μ M ALA in the maturation media), 300 (supplementation with 300 μ M ALA in the maturation media)

Fig. 3: Nuclear maturation of buffalo oocytes after 24 h of maturation in IVM medium supplemented with different concentrations of ALA. Data are shown as Mean percentage \pm SEM. Bars with different superscripts differ significantly ($P<0.05$). GV: Germinal Vesicle, GVBD: Germinal Vesicle Breakdown, MI: Metaphase I, MII: Metaphase II, Deg: Degenerated.

Table 1: Effect of supplementation of IVM medium with different concentrations of α -Linolenic acid on the cleavage rate and subsequent embryo development of buffalo oocytes after *in vitro* fertilization.

α -Linolenic acid	Developmental stages (Mean percentage \pm SEM)				
	No. of	Cleavage	4-8 cell embryos	>8 cell embryos	Morulae
	COCs	N (%)	N (%)	N (%)	N (%)
0 μ M	182	92 (50.4 \pm 2.0) ^b	62 (34.0 \pm 1.1) ^c	40 (22.2 \pm 2.9) ^c	20 (11.2 \pm 2.6) ^b
25 μ M	180	92 (51.1 \pm 2.5) ^{ab}	66 (36.6 \pm 3.8) ^{bc}	42 (23.1 \pm 2.4) ^c	22 (12.1 \pm 2.1) ^b
50 μ M	180	102 (56.9 \pm 2.0) ^{ab}	76 (42.3 \pm 1.5) ^{ab}	56 (31.1 \pm 1.6) ^b	32 (17.8 \pm 2.3) ^a
100 μ M	180	104 (57.8 \pm 1.0) ^a	84 (47.0 \pm 2.9) ^a	68 (38.0 \pm 2.2) ^a	38 (21.0 \pm 2.6) ^a
150 μ M	180	90 (49.7 \pm 2.8) ^b	56 (31.2 \pm 0.8) ^c	26 (14.4 \pm 1.8) ^d	14 (7.8 \pm 0.2) ^b
300 μ M	180	38 (21.0 \pm 2.5) ^c	26(14.3 \pm 1.8) ^d	2 (1.2 \pm 1.2) ^e	0 (0.0 \pm 0.0) ^c

Data were collected in seven independent repeats.

^{a,b,c,d,e}The values with different superscripts in the same column differ significantly ($P < 0.05$).




Figure 1: Representative pictures of COCs (A: Fully expanded, B: Partially expanded, C: Not Expanded) after 24 h of maturation.





Figure 2: Cumulus expansion rate of buffalo COCs after 24 h of maturation in IVM medium supplemented with different concentrations of ALA. Data are shown as Mean percentage \pm SEM. Bars with different superscripts differ significantly ($P < 0.05$). 0 (Control), 25 (supplementation with 25 μ M ALA in the maturation media), 50 (supplementation with 50 μ M ALA in the maturation media), 100 (supplementation with 100 μ M ALA in the maturation media), 150 (supplementation with 150 μ M ALA in the maturation media), 300 (supplementation with 300 μ M ALA in the maturation media)



1
 2 **Figure 3:** Nuclear maturation of buffalo oocytes after 24 h of maturation in IVM medium supplemented with different concentrations of
 3 ALA. Data are shown as Mean percentage ± SEM. Bars with different superscripts differ significantly (P < 0.05). GV: Germinal
 4 Vesicle, GVBD: Germinal Vesicle Breakdown, M1: Metaphase I, M11: Metaphase II, Deg: Degenerated.
 5