

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the peer-reviewed, manuscript version of an article published in *Acta Tropica*. The version of record is available from the journal site:

<http://dx.doi.org/10.1016/j.actatropica.2017.12.026>.

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Brucellosis in West and Central Africa: A review of the current situation in a changing landscape of dairy cattle systems

AUTHORS: Craighead, L; Meyer, A; Chengat, B; Musallam, I; Akakpo, J; Kone, P; Guitian, J; Haesler, B

JOURNAL: Acta Tropica

PUBLISHER: Elsevier

PUBLICATION DATE: March 2018

DOI: 10.1016/j.actatropica.2017.12.026

1 Brucellosis in West and Central Africa: A
2 review of the current situation in a
3 changing landscape of dairy cattle
4 systems

5
6 L. Craighead^A, A. Meyer^A, B. Chengat^A, I. Musallam^A, J. Akakpo^B, P. Kone^B, J. Guitian^A, B. Häsler^A

7 ^ARoyal Veterinary College, Hawkshead Lane, Hatfield Herts, AL9 7TA

8 ^BEcole Inter-Etats des Sciences et Médecine Vétérinaires de Dakar, Dakar Université Cheikh Anta Diop

9

10 Corresponding author : Laura Craighead, lcraighead2@rvc.ac.uk

11 Abstract

12 Brucellosis is a neglected endemic zoonosis in West and Central Africa. In this narrative review,
13 evidence of livestock and human infection is presented along with details of past and current
14 control strategies in 14 selected countries. Data from available literature is combined with expert
15 opinion elicited during a regional workshop on brucellosis diagnostics. Demographic changes that
16 affect both the epidemiology of brucellosis and the success of control or surveillance are also
17 considered. The evidence suggests that brucellosis prevalence in emerging peri-urban dairy cattle
18 systems may be higher than that found in traditional transhumant extensive systems. Accurate
19 microbiological and epidemiological evidence across the region is lacking but it appears there is
20 inherent interest in controlling the disease. There are many data gaps which require collaborative
21 future research to evaluate fully the social and economic impact of the disease in an evolving
22 livestock sector heavily influenced by high rates of urbanisation and regional population growth.

23 1. Introduction

24 The dairy sector in West and Central Africa is evolving. Demand for dairy products is growing and
25 consumer preferences and behaviours are changing; in order to capitalise on the opportunities, the
26 sector must adapt and develop. The growth of the dairy industry in the region is undoubtedly
27 positive in terms of increased nutrition and diversity in the diets of consumers, as a driving force for
28 poverty alleviation amongst farmers and as an avenue for national and regional economic growth
29 (Mathias and Mundy, 2010). However, this burgeoning demand is not without inherent risks in
30 terms of the spread of zoonotic diseases and exposure to food borne pathogens. Of these risks,
31 brucellosis is perhaps one of the most significant threats due to the potential impact on both
32 human health and animal health and productivity (Moreno, 2014).

33 Brucellosis is a zoonotic disease caused by gram negative coccobacilli bacteria belonging to the
34 *Brucella* genus. Four of the eleven *Brucella* species are recognised human zoonoses, these being *B.*
35 *abortus*, *B. melitensis*, *B. suis* and *B. canis* (Corbel, 2006). As one of the world's most widespread
36 zoonosis, brucellosis is responsible for a vast global burden imposed on poor people through

37 disease, disability and impaired livestock productivity (J. McDermott et al., 2013; WHO, 2005).
38 However, brucellosis is rarely prioritized by decision makers within health systems and the World
39 Health Organisation (WHO) includes it as one of its seven neglected endemic zoonoses (WHO,
40 2005). In humans, brucellosis causes flu-like symptoms and chronic debilitating illness. It often
41 manifests as recurrent bouts of fever, which can be misdiagnosed as drug-resistant malaria and
42 lead to underestimation of its incidence (Chabasse et al., 1983). The main routes for human
43 infection are consumption of contaminated dairy products and contact with infected ruminants
44 (Charters, 1980). In livestock, brucellosis decreases productivity by causing abortions, reducing
45 fertility and decreasing milk yield (Corbel, 1988). In many western European countries, elimination
46 of *B. abortus* in cattle has been achieved by stringent test and slaughter regimes and continued
47 surveillance such as those practiced in the UK leading to 'brucellosis free' status being granted in
48 the 1980's (Young and Corbel, 1989). In other areas, reduced human incidence has come through
49 pasteurization of dairy products alongside mass livestock vaccination programmes.

50 Globally, WHO reported foodborne disability-adjusted life years (DALYs) due to brucellosis to be
51 124,884 (95% C.I. 43,153 - 2,910,416) in 2010, acknowledging that in resource-scarce settings at
52 least as many DALYs could be due to the non-foodborne route (WHO, 2015). However, the
53 estimation of the DALYs for brucellosis is criticised due to under reporting of the non-specific
54 symptoms often associated with *Brucella* infection in humans. As a neglected endemic zoonosis,
55 accurate epidemiological data are often scarce; consequently meaningful estimates of the
56 economic and/or social burden are mostly unavailable. Across the African continent, while there
57 are examples of research studies investigating the prevalence of brucellosis (Akakpo and Bornarel,
58 1987; Ducrotoy et al., 2015; J. J. McDermott et al., 2013), there is very limited if any consistent
59 surveillance data in humans, domestic or wild animals. The limited available literature mostly
60 considers seroprevalence and often lacks evidence of epidemiological rigour in the study design
61 and presentation of results. Spread widely over time and geographical landscapes, the literature
62 does not offer a full interpretation of the specific epidemiology of brucellosis across the continent,

63 which it seems, may differ considerably to other continents. Of the four zoonotic *Brucella* species,
64 *B. melitensis* and *B. abortus* are globally considered the predominant species of importance due to
65 the impact on human health and animal productivity. Generally, *B. abortus* is associated with cattle
66 while *B. melitensis* is associated with small ruminants and a higher pathogenicity in humans (Young
67 and Corbel, 1989). Despite the host predilection of the *Brucella* species, there is evidence of cross
68 infectivity as shown by infection of sheep with *B. abortus* in the absence of *B. melitensis* (Allsup,
69 1969; Luchsinger and Anderson, 1979; Shaw, 1976). This may be relevant in the case of sub Saharan
70 Africa where little evidence of *B. melitensis* has been found and mixed farming systems are
71 common. More work is needed however, to establish the epidemiological situation before using
72 these assumptions to inform control policy decisions. To date, the limited number of studies
73 isolating bacterial species have given preference to sampling cattle rather than small ruminants or
74 humans (Akakpo and Bornarel, 1987; Domenech et al., 1982; Pilo-Moron et al., 1979; Sacquet,
75 1955).

76 The West African region spans an area of approximately 5 million square kilometres with diverse
77 geographical and cultural characteristics. It has many disparities in food requirements and
78 production (Dixon et al., 2001). Migration of people, livestock and wildlife across the region has
79 long been commonplace. Many initiatives have been established at various times to combine forces
80 politically and economically in order to take advantage of economies of scale and the opportunities
81 the diversity of states brings. Initiatives such as the Economic Community of West African States
82 (ECOWAS) Regional Agricultural Policy (ECOWAP) and the Comprehensive Africa Agriculture
83 Development Programme (CAADP) over the past decade have shown the renewed interest and
84 emphasis on the role and importance agriculture has on the economic development and growth of
85 the region. While there have undoubtedly been many benefits and positive outcomes of such
86 initiatives there are unsurprisingly varying levels of success in different member states. In many
87 instances priority setting is affected by food crisis such as those seen after the financial crash in
88 2008 or following numerous civil wars and unrest in many countries.

89 Economies of scale are important when the region's dairy industry faces fierce international
90 competition from imports of dried milk powder from Europe. Such regional cooperation can
91 facilitate procurement of vaccines, medicines and diagnostic capacity for diseases that reduce
92 productivity and pose risks to human health. In conjunction with animal health policy and
93 interventions, market strengthening in terms of improved logistics, processing and distribution are
94 key to putting regional production on a more even playing field with cheap imports of milk powder
95 into the region (Ndamb et al., 2007).

96 The predominant driver for an increase in demand for dairy products is population growth. Levels
97 of population growth are amongst the highest in Africa and specifically in the West African region
98 where the population has more than doubled over the last thirty years, growing by 2.7% annually
99 with a projection of 736 million people by 2050 (UNDESA, 2015). In conjunction with population
100 growth, urbanisation and changes in consumer preference have a big influence on overall demand
101 for dairy products and types of products purchased. Urbanisation is also transforming where and
102 how products are bought. Urbanisation rates in West Africa are rapid even when compared to
103 other African regions.

104 In many countries, the burgeoning dairy potential is recognised and there have been various
105 attempts to capitalise on it. Many governments in countries such as Senegal and Ivory Coast have
106 implemented breed enhancement programmes often utilising imported European breeds with
107 higher milk yields since 1990-2000. These programmes often see limited success due to disease
108 susceptibility amongst such breeds. Concurrently, there has been an increase in commercial closed
109 dairy farms utilising modern dairy milking machines and housing onsite processing plants. Such
110 processing plants have also had limited success due to the inability to produce adequate quantities
111 of milk from single herds and the difficulties in forming milk co-operatives due to poor cold chain
112 and transport logistics.

113 There is serological evidence of widespread brucellosis infection of cattle in the West and Central
114 African region (Akakpo and Ndour, 2013). There is consistent correlation between abortion levels
115 and seropositive herds as well as carpal hygroma being a quasi-pathognomonic sign of brucellosis
116 in cattle (Domenech et al., 1980). Hygromas are often considered the main symptom of brucellosis
117 in cattle besides abortion in sub-Saharan Africa (SSA) as cattle are often kept within a herd for the
118 entirety of their natural lives allowing for persistence of chronically infected animals. In central
119 Africa a linear correlation between abortion rate and carpal hygroma prevalence has been shown;
120 it has been suggested this can be used as a proxy to establish whether a brucellosis control
121 programme is necessary when other diagnostic methods are not available (Domenech et al., 1982,
122 1980) .

123 To date there is no published review of West Africa collating brucellosis data in animals and people
124 for the region and the magnitude of the disease burden is unclear. Without such baseline data it is
125 impossible to conduct epidemiological and economic assessments on mitigation strategies
126 (Rushton et al., 1999) for brucellosis and/or for policy makers to make informed decisions on
127 disease management (McInerney, 1996). Therefore, in this review, we aimed to present currently
128 available data and information on the brucellosis situation and a description of key demographic
129 drivers in 14 francophone West and Central African countries, namely Benin, Burkina Faso,
130 Cameroon, Central African Republic, Côte d'Ivoire, Gabon, Ghana, Mali, Mauritania, Niger, Rwanda,
131 Senegal, Chad & Togo.

132 The objectives were to 1) collate data on prevalence of brucellosis in cattle, humans and small
133 ruminants; 2) provide an overview of existing control strategies and policies; and 3) describe the
134 demographic evolution of human and cattle populations and the impact this has on the dairy sector
135 structure now and in the future.

136 2. Methods

137 2.1 Overview

138 A narrative literature review, a method suitable to illustrate current knowledge and gaps on a topic
139 where scarce evidence exists (Baumeister and Leary, 1997), was conducted by integrating data from
140 open source population statistics databases and scientific literature. Moreover, expert opinion was
141 collated to complement the secondary data available.

142 2.2 Literature search and review

143 2.2.1 Literature search

144 Three scientific databases (Google scholar, Pubmed, Science direct) were searched for relevant
145 articles using the following search terms: (brucellosis OR brucella OR dairy zoonosis) AND (Africa OR
146 West Africa OR Central Africa OR *individual country names*), with a publication limitation of 1950 to
147 the 30 June 2016. The searches were conducted in both French and English. All databases were
148 searched using title and abstract search. Duplicate entries were identified and removed before
149 selection of articles was begun.

150 2.2.2 Selection process

151 The articles were sorted by two reviewers with a combined fluency in English and French.

152 All reports were classified into one of two categories, based on their abstracts: Category 1:

153 Relevant – articles related to human/cattle or small ruminant brucellosis in relation to brucellosis
154 infection in populations (i.e. disease frequency) or articles related to diagnostic capacity or disease
155 control methods; Category 2: Irrelevant - articles related to non-brucellosis dairy zoonosis; articles
156 addressing topics not related to the current review, such as genetics or experimental laboratory
157 animal studies.

158 The abstracts of papers belonging to Category 1 and meeting the following criteria were retained:
159 published between 1950 and 30 June 2016, prevalence or incidence data included, and containing
160 information relating to diagnostic tests, *Brucella* species isolation results or evaluation of social or
161 economic impact of disease in livestock or human populations.

162 The selected papers were reviewed in full by both reviewers and information on study area, study
163 population, seroprevalence estimates and bacterial isolation as well as diagnostic tests utilised and
164 discussion of control methods or surveillance were tabulated.

165 2.3 Collection of expert opinion

166 Expert opinion was elicited in order to supplement the scientific literature. A regional workshop by
167 a joint research project established between the Royal Veterinary College (RVC) the Ecole Inter-
168 Etats des Sciences et Médecine Vétérinaires de Dakar (EISMV), London School of Hygiene and
169 Tropical Medicine (LSHTM), Animal and Plant Health Agency (APHA) and GALVMED, under the
170 Zoonosis in Emerging Livestock Systems (ZELS) initiative was run in Dakar in June 2015. In
171 attendance were 28 participants from 14 countries with the majority of those invited working in
172 teaching, government or diagnostic institutions (see appendix 1 for list of delegates' countries and
173 institutions). A snowballing sampling of experts was conducted to identify participants through the
174 alumni and contact network of EISMV. During the workshop, delegates were divided into three
175 groups of 9-10 people. Semi-structured group interviews were carried out with each group covering
176 three main topics; namely 1) Current knowledge of brucellosis situation and its impact; 2)
177 Brucellosis control programmes; and 3) Brucellosis diagnostic capacity (see
178 Appendix/supplementary materials for the question guide).

179 Free discussion in French was encouraged by a facilitator who took notes. At the end of the
180 discussion, participants were given questionnaires to fill in to summarise their opinion on the topic
181 that had been discussed (see appendix 2/3/4 for questions). Data were analysed by means of
182 collating, translating and tabulating questionnaire responses and coding of discussion notes to
183 highlight common themes and opinions on the topics discussed.

184 2.4 Extraction of information from databases

185 Data on livestock parameters were obtained from the FAOSTAT database (FAO, n.d.). Human
186 demographic data were obtained through the open source Health Nutrition and Population
187 Statistics database of the World Bank for each of the countries. Data sets were downloaded in excel

188 format, relevant data was extrapolated to formulate graphical representations concerning trends in
189 populations and dairy production.

190 3. Results

191 3.1 Literature review

192 A total of 68 publications, chapters or theses were included in the review. Of these 48 were written
193 in French and 20 in English. They spanned from the 1950s to 2015, with only three studies being
194 carried out within the last five years. As data was relatively limited older pieces were not excluded
195 and as such included two publications from the 1950's, one from the 1960's, seven from the
196 1970's, 21 from the 1980's, five from the 1990's and 17 from the 2000's.

197 3.2 Brucellosis Impact

198 3.2.1 General overview


199 Tables 1 and 2 show the literature findings in cattle and human populations. A more detailed
200 summary of individual countries can be found in the supplementary material (appendix 5). The
201 highest number of publications was from Ivory Coast, Mali, Niger and Senegal with a large
202 proportion dating back over two decades ago. Across all 14 countries, there were numerous
203 national surveys in the 1970's and 80's, in contrast more recent studies tend to concentrate on
204 particular geographic areas. On occasion, it appears there are conflicting findings in the same areas
205 such as in Benin and Burkina Faso, however in the case of Burkina Faso there were long time gaps
206 between studies and they may not be directly comparable (Table 1). Only few papers were found
207 that outline in full the farming systems within the studies. Seroprevalence estimates ranged from
208 0% to 55.2% \pm 11.9%. Numerous high seroprevalence estimates were reported in peri-urban areas
209 of Burkina Faso, extensive systems in Cameroon, Chad, Niger and Togo in the late 1970's and early
210 80's. The recent highest seroprevalence estimate was reported for peri-urban farms around Dakar
211 in Senegal 2012 (Table 1).

212 Ten of the 14 countries had literature on studies in humans (Table 2). Most of the studies were
213 carried out in at risk groups such as farmers or abattoir workers or those presenting at health

214 facilities with fever or abortions. There was no evidence of seropositivity in a none-exposed group
215 in Benin or in any groups of a small sample of 77 people in a study in Ghana.


216 A lack of agreement in results was identified between the Rose Bengal and ELISA techniques in
217 Benin, Cameroon and Chad (Table 1).

218 In many cases, a situation of low individual seroprevalence and moderate to high herd
219 seroprevalence is reported (Akakpo, 1987; McDermott and Arimi, 2002) suggesting a state of
220 endemicity with relatively low transmission rates. Factors such as low reproductive rates as well as
221 environmental conditions of transhumant herds moving in dry and hot conditions have been
222 hypothesised as explanations for



223

224 *Figure 1 Literature estimates of cattle herd seroprevalence for Brucella spp. in West and Central Africa*



225

226 *Figure 2 Brucella isolates in West and Central Africa listed in literature from 1955 to 2011.*

227 the low transmission rates (Ducrotoy et al., 2015). These findings however are in sharp contrast to
 228 recent studies in peri-urban farming systems (Mugizi et al., 2015; Tialla et al., 2014), where reports
 229 of both high individual and herd seroprevalence estimates suggest a different epidemiological
 230 picture to that outlined above. These farming systems are often more intensive with little
 231 transhumance which may account for the more favourable transmission conditions and therefore a
 232 greater burden of brucellosis in these emerging farming systems. In personal communication with
 233 an author of the study in Senegal it was noted that the affected farms are ‘renewed by
 234 transhumant animals especially the males but the females remained within the herd for extended
 235 periods, sometimes up to 10 years. In some farms visited, there are females with hygromas. After,
 236 our visit, most of the farmers slaughtered positive animals.’ In a report in 2012 by the International
 237 Livestock Research Institute (ILRI, 2012), they highlighted a higher risk of brucellosis in intensive
 238 systems and list Togo, Mali, Ivory Coast, Niger, Cameroon and Burundi amongst the top hot spots
 239 for the disease globally.

240 These findings contradict those found in a review in East Africa (Njeru et al., 2016), where a lower
 241 seroprevalence in large urban centres such as Nairobi was noted in comparison to extensive
 242 pastoralist systems. There are numerous things to consider when hypothesising the reasons behind

243 these differing findings, perhaps different climate, production systems or the mixing of extensively
244 managed herds with a plethora of wildlife in East Africa which is far less common in West Africa.

Year	Sampling	Diagnostic Test	Outcome <i>Brucella spp</i> prevalence/species isolation (plus confidence intervals or standard errors where given)	Area / system /subpopulation sampled and additional findings	Reference
BENIN: 114.8 thousand Sq.km, neighbouring Nigeria on the southern coast of the region. 10.88 million people; urban growth rate 3.6%, 2.2 m cattle, of which 12% dairy cattle					
1980-81	Non-probabilistic	TFC, TRB	10.4% ± 1.97%	National survey. Traditional farming	(Akakpo A.J., Bornarel P., 1984)
2001	Risk based, probabilistic	TRB	6.2% ± 1.8%	Bétécoucou, Kpinnou, Okpara and Samiondji (southern half of the country)	(Koutinhouin et al., 2003)
		ELISA	15.2% ± 2.6%		
2000-03	Specific population, all tested	TFC, iELISA, TRB	1.91% ± 0.25%	State farms of Okpara, Bétécoucou and Samiondji	(Adehan et al., 2003)
BURKINA FASO: A medium sized land locked country in the middle of the region, having five direct land borders with other countries it is well positioned for regional trade. 18.11 million people; urban growth rate 5.7%, 9 m cattle, of which 15% dairy cattle					
1970-73	Non-probabilistic	SAW, TFC	8.1% ± 3.3%	Gaoua and Markoye	(Gidel et al., 1974)
		RT	11.5% ± 1.7%	Banfora, Dori	
1981-82	Unknown	TRB, TFC	12.3% ± 1.8%	7 régions out of 11, traditional farming	(Bessin, 1982)
		TRB, TFC	55.2% ± 11.9%	Ouagadougou, urban area	
2001-02	Non-probabilistic	TRB	13.1% ± 3.9%	Hamdallaye borough of Ouagadougou. Potential risk factors: concentration of animals and manure accumulation near household	(Traore et al., 2004)
2004-05	Unknown	TRB, iELISA	8.2% ± 2.8%.	Rural extensive	Dairy farm survey in and around Bobo-Dioulasso
			8.1% ± 2.9%.	Intra-urban extensive	
			0.31% ± 0.35%	(Semi-) intensive	

2012	Probabilistic	TRB and iELISA	7.3% (95% C.I. 3.5%-14.7%)	Transhumant herds from Burkina Faso in the Northern Savannah region of Togo	(Dean et al., 2013)
Cameroon: Cameroon has a relatively large land mass bordering Nigeria with a small coastal border within Central Africa. 23.34 million people; urban growth rate 3.5%, 6 m cattle, of which 5% dairy cattle					
1976-80	Unknown	TRB	30.8% ± 1.0 %	Breeding females in extensive farming systems (sedentary and transhumant)	(Domenech et al., 1982)
1980	Unknown	TFC, TRB	22.2% ± 6.3%	Region of Diamaré	(Akakpo, 1987)
			19.7% ± 4.5%	Region of Bénoué	
			4.8% ± 1.9%	Region of Adamaoua	
			12.5% ± 2.1%	National survey	
2002-03	Non-probabilistic	iELISA	9.64%	Abattoir of Dschang (West Cameroon)	(Shey-Njila et al., 2005)
		TRB	4.88%		
2009	Unknown	iELISA	8.4% <i>No CI given</i>	21 villages in Cameroon, Holstein cattle	(Bayemi et al., 2009)
2015	Unknown	iELISA	5.2% <i>No CI given</i>	North-West region of Cameroon	(Bayemi et al., 2015)
Central African Republic: Central African Republic (CAR) is a sizeable landlocked country in Central Africa bordering Cameroon on its westerly side. Despite its size, the total population in 2015 was only 4.9 million. Civil war and displacement in the country have hindered development and consequently CAR ranks as one of the world's top 10 most poor countries. urban growth rate 2.7%, 4.35 m cattle, of which 5% dairy cattle					
2004	Unknown	TRB	3.3%		(Nakoun, 2004)
Chad: Chad is one of the large inland countries of central Africa, neighbouring Niger, CAR and Cameroon. 14 million people; urban growth rate 3.8%, 8 m cattle, of which 10% dairy cattle					
1955	Unknown		<i>B. abortus</i> isolated from infected cattle		(Sacquet, 1955)
		SAW	12%.	National Survey Regional variations: from 7.4% in Fianga to 23.8% in Massakori	
1976-80	Unknown	TRB	31.9% ± 1.1 %	Breeding females in extensive farming (sedentary and transhumant) Prevalence of clinical signs in females 13% (abortion, hygroma or arthritis)	(Domenech et al., 1982)
			<i>B. abortus</i> biotypes 3, 6 and intermediate 3/6 isolated from cattle samples.		
1999-2000	Non-probabilistic	TRB	7.3 % [0.9-17.0]	Chari-Baguirmi and Kanem regions	(Schelling et al., 2004)

2000	Probabilistic	TRB, TFC	2.6% ± 1.2%	Sedentary herds in Abéché peri-urban area, producing milk for local market.	(Delafosse et al., 2002)
2008-09	Unknown		7%	National level reported in OIE questionnaire	(Akakpo et al., 2009)
2014	Unknown	TRB	5.7% (95% C.I.: 3.8-7.6%)	South-eastern shore of Chad (only the islands, not the mainland)	(Abakar et al., 2014)
		iELISA	11.9% (95% C.I.: 9.3%-14.6%)		
Gabon: Gabon is one of the smaller countries included in the study, the majority of this population are urban dwellers (almost 90%). Gabon is the most southerly of the countries and has a substantial coastal border. 1.73 million people; urban growth rate 2.5%, 38 thousand cattle, of which 30% dairy cattle.					
Ghana: Ghana is a medium size land mass along the south coast of the region, despite its moderate size it has a large population with over half of the population reported as urban dwellers. In contrast, Ghana has amongst the lower population of cattle in the region, reporting just under 1.7 million head of cattle in 2014 27 million people; urban growth rate 3.5%, 1.7 m cattle, of which 20% dairy cattle					
1992	Unknown	RBT	9.3%	Coastal savannah area	(Turkson and Boadu, 1992)
2000	Unknown	RBT	6.6%	Akwapin-South district	(Kubuafor et al., 2000)
2012	Probabilistic	RBT	2.93%	Dormaa and Kintampo districts, Brong Ahafo region.	(Folitse et al., 2014)
Ivory Coast: The Ivory Coast neighbours Ghana on the southern coast of the region. Slightly larger in land area it has a smaller total population than Ghana but a similar proportion (55%) of citizens residing in urban areas. 22 million people; urban growth rate 3.7%, 1.6 m cattle, of which 20% dairy cattle					
1970-73	Non-probabilistic	RT	42.9% ± 2.7%	Bouaké, Korogho, Man and Odiene	(Gidel et al., 1974)
		SAW, TFC	15.5% ± 2.6%		
1975-77	Non-probabilistic	TRB, SAW, TFC	10.0% ± 0.5%.	National survey Herd prevalence 75 %. 38% of cows with abortions are seropositive. Important regional variations, from 1.1 to 40%	(Pilo-Moron et al., 1979)
			<i>B. abortus</i> biovar 1 and 6 isolated from infected cattle		
1976-78	Unknown	TRB	28.3%	North region Herd prevalence: 41% (RT), 75% (TRB). In positive herds: higher abortion rate, reduced fertility, higher mortality in calves.	(Camus, 1980)
1984	Unknown	TRB, SAW	5.4% (centre) to 19.5% (north)	North and Centre regions Overestimates true prevalence because some vaccination is practised	(Angba et al., 1987)

2004	Unknown	Bayesian analysis of the results of TRB, iELISA, SAW and TFC	3.6%	Dairy farms (training and private farms)	Peri-urban forest area around Abidjan. Farms providing milk products for Abidjan	(Thys E., Yahaya M.A., Walravens K., Baudoux C., Bagayoko I., Berkvens D., 2005)
			4.3%	Traditional farms		
2005	Probabilistic	Bayesian analysis of the results of TRB, iELISA and SAW.	8.8 % [5.0-16.4]	Nzi, Comoé and Lake regions		(Sanogo et al., 2008)
2012	Non-probabilistic, high risk groups	TRB and iELISA	10.3% (95% C.I.: 8.4%-12.4%).	Savannah-forest region of Ivory Coast		(Sanogo et al., 2012)
			<i>B. abortus</i> biovar 3	UNPUBLISHED*		
Mali: Mali is one of the largest inland countries in the region, landlocked it shares borders with many other countries. Mali reports the second largest national herd of cattle and the largest dairy herd of the study countries. 17.6 million people; urban growth rate 4.9%, 10 m cattle, of which 20% dairy cattle						
1991	Unknown	ELISA	23.3% ± 2.5%	National survey. Reported prevalence from 11 to 20%, in previous unpublished surveys. West Mali more affected (subhumid area): herd prevalence up to 87% in Koulikouro, 74% in Kayes. Tombouctou area is less affected (arid climate)		(Toukara et al., 1994)
				<i>B. abortus.</i>		
1998-01	Risk based	TRB	Prevalence among suspect cases rose from 8.7 to 19.5% during study period	Central Veterinary Laboratory, suspect cases (mostly abortions and pre-sale check-up). Herd prevalence rose from 31.4 to 65%.		(Bonfoh et al., 2003)
2008-09	Unknown		22%	National level reported in OIE questionnaire		(Akakpo et al., 2009)
2011	Unknown	TRB	1%	Cinzana region		(Sow et al., n.d.)
		ELISA	0%	2 of 204 cattle tested positive to TRB; neither were positive to iELISA.		
Mauritania: Mauritania is another large country in the North of the region neighbouring Mali but also having an expanse of coastal border in the West. In contrast to Mali, it has a much smaller human and cattle population. 4 million people; urban growth rate 3.5%, 1.9 m cattle, of which 20% dairy cattle						

Niger: Niger, like Mali is a large landlocked country in the northern edge of the region. With a population of nearly 20 million in 2015 it has a very high urban population growth rate of 5.4% annually but currently less than 20% of its population reported to be urban dwellers. Niger has the largest national herd of cattle. 3.5%, 11.4 m cattle, of which 13% dairy cattle					
1970-73	Non-probabilistic	RT	21.2% ± 5.1%	Niamey area	(Gidel et al., 1974)
		SAW, TFC	2.4% ± 4.6%		
1980-81	Unknown	TRB, FC	35.3% ± 3.6%	Niamey (more humid than Zinder) Prevalence in other species: 6.6% ovine, 2.0% caprine, 8.6% camels.	(Saley, 1983)
		TRB, FC	12.1% ± 5.1%.		
1980-81	Unknown	TFC, TRB	30.9% ± 3.2%	National survey. Traditional farming, Most affected area in Niger is the department of Niamey	(Akakpo, 1987)
			<i>B. abortus</i> 3 or 3/6		
1989-90	Probabilistic	TRB	1.4% ± 0.4%	National level survey. Endemicity in all cattle rearing areas	(Bloch and Diallo, 1991)
2013	Probabilistic	iELISA	1.3% (95% C.I.: 0.9% - 1.8%)	Niamey	(Boukary et al., 2013)
Rwanda: Rwanda is a small landlocked country in the East of Africa but is included in this work due to the links and interest of ministerial vets there and their participation in the regional workshop. Despite its small geographical size compared to other countries, it has a total population of over.11 million people; urban growth rate 5.8%, 1.1 m cattle, of which 20% dairy cattle					
1982-83	Unknown	TFC, TRB	34.9% ± 3.7%	National survey, traditional farming	(Akakpo, 1987)
			<i>B. abortus</i> 3 or 3/6		
2008-2009	Unknown		1.7%	National level, reported in OIE questionnaire	(Akakpo et al., 2009)
Senegal: Senegal is a medium sized country with a large proportion of its border being coastal on the region's western coast. 15 million people; urban growth rate 3.8%, 3.5 m cattle, of which 20% dairy cattle					
1960-62	Unknown	SAW	13% ± 2.1%.	National survey. Casamance and Thies are infected. Diourbel seems free. Prevalence of clinical signs 8.5% (abortion, hygroma)	(Chambron, 1965)
			<i>B. abortus</i> and <i>B. intermedia</i> isolated from infected cattle		
1974-75	Probabilistic	SAW	12.0% ± 1.9 %	Dakar's abattoir (cattle from northern half of Senegal) 17.2% of sera are positive at least once in SAW, TFC, Coombs or TRB	(Chantal and Thomas, 1976)
1976	Unknown	TRB	14.4% ± 3.5%		(Fensterbank et al., 1977)
		Skin test	8.5% ± 2.8%		

			<i>B. abortus</i> isolated from infected cattle	1 village in Basse-Casamance. All animals with hygromas (except 1) are seropositive.	
1979	Non-probabilistic		180 <i>B. abortus</i> biovar 3/6 and 1 <i>B. abortus</i> biovar 1	181 strains isolated from cattle arthritis samples	(Verger et al., 1979)
1984	Non-probabilistic		<i>B. abortus</i> 1 and 3/6		(Verger and Grayon, 1984)
1987	Non-probabilistic		<i>B. abortus</i> 3 or 3/6	Casamance region	(Akakpo, 1987)
2001-03	Probabilistic	RBT and TFC	0.6% (95% C.I. 0% -1.3%)	Bassin Arachidier region	(Unger et al., 2003)
2007-08	Unknown	Competition ELISA, TRB	1.5% ± 2.1%	Dairy farms in Thiès region	(Kouamo et al., 2010)
2008-09	Unknown		20 %	National level. Reported in OIE questionnaire	(Akakpo et al., 2009)
2012	Probabilistic	TFC, TRB	25.0% ± 4.9%	Peri-urbane dairy farms around Dakar. Supply Dakar with unpasteurized milk and milk products. Correlation of seropositivity with age, breed, abortion and hygroma. Herd prevalence >95%.	(Tialla et al., 2014)
Togo: Togo is a relatively small country sitting between Ghana and Benin on the southern coast of the region, although Togo only has a small coastal border. 7.3 million people; urban growth rate 3.9%, 520 thousand cattle, of which 15% dairy cattle					
1977	Unknown	TFC, TRB	40.9% ± 3.0%	National survey, traditional farming. Lomé and Avetonou are the regions most affected by brucellosis	(Akakpo, 1987)
			<i>B. abortus</i> 3 or 3/6		
1977-79	Non-probabilistic	TRB, SAW	41.2% ± 2.9%	National survey. Regional variations: increase from North to South. Some seropositive among livestock keepers	(Sonhayé, 1980)
1984	Non-probabilistic		<i>B. abortus</i> 3/6		(Verger and Grayon, 1984)
1990	Non-probabilistic	TRB, TFC	16.6% ± 3.3%	National survey (but small sample) Higher prevalence in Maritime and Plateaux regions (but vaccination in the latter)	(Kponmassi, 1991)
2011	Probabilistic	TRB and iELISA	9.2%, (95% CI: 4.3-18.6%)	Northern Savannah region of Togo	(Dean et al., 2013)
			<i>B. abortus</i>	isolated from 3 bovine hygroma samples	

				Northern Savannah region of Togo – sample also includes transhumant herds from Burkina Faso (approx. 1/3 herds)	
--	--	--	--	---	--

247 *Supplementary work referred to within the paper but unpublished itself

248 **TFC Complement fixation, TRB Rose Bengal test, ELISA Enzyme linked immunosorbent assay, SAW Sero agglutination of Wright, RT Ring test, iELISA Indirect enzyme linked immunosorbent
249 assay, RBT Rose Bengal test, IFI Indirect immunofluorescence, C.I confidence interval

250 Table 2 Literature findings in humans for brucella spp. in West and Central Africa

Country	Population	Test	Outcome	Comment	Reference
Benin	1. Livestock keepers	TRB, IFI	17.7% ± 6.4 %	<i>Gouka and Cotonou area. No positive among non-exposed persons</i>	(Audurier A., Fayomi B., Laudat P., 1987)
	2. Slaughterhouse employees		0		
	3. Non-exposed group				
Burkina Faso	Village surveys	SAW, TFC	2.5%	<i>Banfara, Dori, Gaoua and Markoye Correlation with livestock keeping, esp. in pastoralists</i>	(Gidel et al., 1974)
Chad	Livestock keepers	TRB	1.8 %	<i>Chari-Baguirmi and Kanem regions Risky behaviours reported (assistance with calving, contact with abortion material, consumption of non-pasteurized milk)</i>	(Schelling et al., 2004)
Ghana	High risk groups	RBT	0	<i>No evidence of human brucellosis was detected by antibody screening in selected risk groups (6 veterinarians, 6 butchers, 21 herdsman and 12 environmental health inspectors) nor in the control group of 30 people</i>	(Kubuafor et al., 2000)
Ivory Coast	Village surveys	SAW, TFC	0.45%	<i>Bouaké, Korogho, Man and Odiene No positive in Tabou (no cattle in that area). Correlation with cattle keeping, esp. in pastoralists</i>	(Gidel et al., 1974)
		Not specified		<i>Known human focal outbreaks in these areas Odiene, Korogho and Bouaké</i>	(Armand, 2001)
Mali		TRB, IFI	0.5%	<i>Fourou, Mostly agriculture, few contacts with livestock</i>	(Tasei et al., 1982)

			24.4%	<i>Gourma, Pastoralists, raw milk consumption</i>	
	Unknown	RBT	24.4%	<i>Rural areas</i>	(Bonfoh et al., 2003)
			4.6%	<i>Bamako</i>	
	Mopti town. Survey among febrile patients	SAW	67%	<i>Significant risk factors: proximity of ruminants and consumption of non-pasteurized milk.</i>	(Dao et al., 2009)
	Village surveys	TRB	0.9% (2/213)	Cinzana region	(Sow et al., n.d.)
		iELISA	0.04% (1/213)		
Niger	Village surveys	SAW, TFC	1.4 %	<i>Niamey area, Correlation with cattle keeping, esp. in pastoralists</i>	(Gidel et al., 1974)
			8.6%.	<i>90% of households consume raw milk, 77% manipulate abortion materials.</i>	(Adamou, 2014)
Rwanda	Prevalence of positive serology in women presenting with abortion and/or stillbirth	TRB	25% 15/60 women were positive		(Rujeni and Mbanzamihiho, 2014)
Senegal	Abattoir workers in Dakar	TRB	8.6%		(Diop, 1975)
Togo	Village surveys- Fulani ethnicity	TRB and iELISA	2.4% (95% C.I.: 1.0-5.6%)	<i>Northern savannah region of Togo</i>	(Dean et al., 2013)
	Village surveys- Non-Fulani ethnicity		0.2% (95% C.I.: 0-1.6%).		

251
252

**TFC Complement fixation, TRB Rose Bengal test, ELISA Enzyme linked immunosorbent assay, SAW Sero agglutination of Wright, RT Ring test, iELISA Indirect enzyme linked immunosorbent assay, RBT Rose Bengal test, IFI Indirect immunofluorescence, C.I confidence interval

253 3.3 Brucellosis control

254 3.3.1 Vaccination & surveillance –Workshop findings

255 During workshop discussions, the consensus was that there are minimal surveillance and control
256 activities for brucellosis taking place across the region. No participant reported extensive
257 countrywide surveillance. Four delegates reported that vaccines were available in their country
258 (Table 3), but none of these countries had a structured vaccination programme in place, all
259 reporting voluntary vaccination with the onus falling on the farmer.

260 *Table 3 Vaccines currently available in study countries according to workshop delegates*

Country	Vaccines available
Gabon	S19, RB51
Ghana	S19, RB51
Ivory Coast	S19, Rev1
Rwanda	RB51

261

262 There was also consensus that cattle vaccines were needed, but opinions were mixed on whether
263 vaccination in small ruminants was necessary.

264 Five countries reported control activities, namely Chad, Gabon, Rwanda, Mali and Niger.

265 Chad reported there to be a written policy for brucellosis in the country but no actual
266 implementation or activity. The delegate believed that there was interest in control programmes in
267 some government managed farms and in the peri-urban farms emerging which are often owned by
268 wealthy people from the cities.

269 Gabon has only recently (2014) begun any animal health service activities as animal production is
270 very minimal in the country. There is reportedly test and slaughter carried out by a large cattle
271 producer in the south of the country. There is no national programme and it is down to farmers to
272 utilize private veterinarians for advice and buying vaccines.

273 While there is no government run control in Mali, serum samples have been taken through a
274 private breed improvement programme but have not been tested as there was a lack of reagents
275 available. The delegate believed that vaccination should be implemented nationwide as
276 government services do not have money to compensate farmers for test and slaughter methods.

277 In Niger, there is routine testing of workers in the central veterinary laboratory and in cattle
278 breeding centers, but no reported testing in the general population of animals.

279 Rwanda has the most extensive control activities. Large dairy farms are routinely tested once per
280 year; smaller farms only being tested if there are cases of abortions. Samples are also collected
281 twice yearly at vaccination drives where vaccines for various diseases are disseminated and
282 samples are collected simultaneously. There is a government initiative of giving either a cow or
283 small ruminant to poor families, all animals used in this programme are tested and slaughtered if
284 found to be positive. Animals found to be positive on farm are sent to abattoirs so the farmer
285 receives only the normal going rate and no additional compensation to cover the difference in
286 market value of a dairy animal and its slaughter price. The delegate from Rwanda hypothesized that
287 although not routinely tested, small farms could have high levels of brucellosis. He described a case
288 where a bull from a large farm was being used to serve cows on small local farms who then
289 introduced the disease onto the large farm. It seems likely given that there is no compensation
290 paid, farmers on the smaller farms would be less likely to report abortions and risk losing valuable
291 dairy animals. Approximately 5000 vaccines per year are administered mainly in the Kigali and
292 Gishwati dairy areas. The vaccine costs the farmer 3500Frw per dose (approx. 4.25 USD). The
293 vaccines are produced by MSD in South Africa and imported.

294 Senegal and Ghana reported historical control programmes that are no longer in existence.

295 3.3.2 Vaccination & surveillance – Literature findings

296 In literature only two papers were found that detailed any structured control activities in the past,
297 these being from Burkina Faso and Ivory Coast. In Burkina Faso, Bessin (1982) highlights in his

298 thesis the detection and slaughter of positive animals as well as vaccination in large peri-urban
299 herds. There is no further evidence of this after the early 1980's.

300 In Ivory Coast, Angba et al (1987), Camus (1995) , Chambron (1965), Armand (2001) and Thys et al
301 (2005) all described vaccine control implemented in the north of the country from 1978 and
302 described a decrease in abortion rate of 37% after the first year of the campaign. By the early
303 1990's however there is no further evidence in the literature of the continuation of any control
304 activity.

305 3.3.3 Diagnostic capacity

306 During workshop discussions on the diagnostic methods for detecting brucellosis all countries
307 reported having the capacity to carry out Rose Bengal tests and ELISA, however many countries did
308 not have the access to reagents required to utilise these methods. While those working in the
309 animal health sector were generally reasonably familiar with the diagnostic techniques available for
310 *Brucella* testing, those working in the human health sector were not and none had heard of any
311 testing for brucellosis within routine healthcare screening of patients. Many alluded to fever
312 profiles utilised for febrile patients and expressed interest in being able to incorporate *Brucella*
313 testing within this profile. However, considering recent findings in East Africa (de Glanville et al.,
314 2017) in regards to misdiagnosis by the commonly used febrile antigen *Brucella* agglutination test
315 (FBAT), careful consideration would be required before implementing routine testing in the West
316 African setting.

317 While discussing constraints in *Brucella* diagnostics, the most frequent reason noted in animal
318 diagnostics was the lack of minor equipment such as reagents, while in human diagnostics major
319 equipment lacking was considered the most frequently cited constraint.

320 Most countries were said to have the capacity to carry out PCR if they were provided with
321 protocols, primers and controls. However, many participants expressed apprehensions about
322 having the appropriate expertise to do so.

323 There was no relevant literature that detailed diagnostic capacity within the countries that were
 324 reviewed.

325 3.4 Evolution of human and animal demographics

326 The combined human population of the study countries in 2015 according to World Bank statistics
 327 was 198.7 million people. This ranged from Ghana with the largest population (27.4 million) to
 328 Gabon with a population of 1.7 million (Fig 3).




Figure 3 Country wide human population statistics extrapolated from World Bank Database

Urban population growth was 4.5% annually, whereas rural population growth was 1.8%. This trend is predicted to continue into 2050 with urban population growth projected to be 3.7% per annum compared to only 0.5% in rural areas (UNDESA, 2015). The majority of countries now have over 40% of their population as

339 urban dwellers with Gabon now nearing 90% of its population living in urban areas.

340 Countries across the region
 341 vary strongly in their sizes and
 342 agricultural potentials. Figure 4
 343 shows the larger inland
 344 countries having the largest
 345 cattle populations. While the
 346 smaller coastal countries have
 347 the lowest cattle populations





Figure 4 Cattle population statistics extrapolated from FAOSTAT


348 but generally the highest human urban populations in the region. In all countries, steady increases
 349 in overall cattle populations have been occurring over the last decades. While there is ambiguity
 350 over reporting of specifically dairy cattle, Figure 5 shows the percentage increases over the 2003-
 351 2013 period and the preceding 10-years. Burkina Faso, Gabon, Niger, Ghana, Togo and Benin all
 352 had much higher percentage increase in



353
 354 *Figure 5 Percentage increases in dairy cattle numbers by country extrapolated from FAOSTAT*
 355 dairy cattle numbers during the 1993-2003 period than the period from 2003-2013. Conversely,
 356 Senegal has seen dramatic growth of 57% in the last 10 years compared to only 8% in the 10-year
 357 period preceding. Similarly, Mali and Cameroon have both also seen relatively higher growths more
 358 recently than during the previous period. Both Mauritania and Central African Republic have seen
 359 smaller and more consistent growth over both the time periods.

360 Despite the relative increases in dairy cattle and milk production across the countries, consumer
 361 demand is not met by local production as shown by the import figures for dry milk powder in Figure
 362 6. All countries have seen increasing value of dry milk imports since the turn of the century with
 363 marked fluctuation during the financial crisis of 2008; this highlights the vulnerability of countries
 364 heavily reliant on imports.

365



366

367 *Figure 6 extrapolated from FAOSTAT*

368 4. Discussion

369 It is evident that although variable and scarce, available literature suggests a reasonably high
 370 burden of brucellosis in the cattle populations of the 14 study countries. There is insufficient data
 371 to conclude any meaningful estimates of the burden however in small ruminant or human
 372 populations.

373 Although the literature is scarce for the region, the high seroprevalence estimates seen in
 374 developing systems such as those reported in Dakar, Senegal (Tialla et al., 2014) are concerning.
 375 This evolving brucellosis epidemiological picture may be true in many of the study countries that
 376 are witnessing the same changes in production styles as a result of urbanisation. Dakar in particular
 377 represents one of West Africa's most developed cities and is most likely a good indicator of future
 378 directions of other major cities in the region.

379 With the highest rates of urbanisation, the West African cattle sector is far more likely to have
 380 higher proportions of emerging farming systems around urban centres that, as suggested

381 previously, may have a much higher burden of brucellosis and different epidemiological picture to
382 the extensive systems predominantly seen elsewhere.

383 This review, in line with others (Ducrotoy et al., 2015) highlights the degree of the lack of reliable
384 and up to date data on the epidemiological picture of brucellosis within West and Central Africa.
385 With no up to date bacterial isolation in any species and very limited human and small ruminant
386 studies it is paramount that microbiological research establishes the detail in a region where mixed
387 farming systems, informal food markets and migration of people and animals are commonplace.

388 The brucellosis prevalence estimates obtained from the literature in this review must be viewed
389 with caution. Most data reviewed was at least five years old in countries where the farming
390 systems and brucellosis prevalence show changing patterns within a setting experiencing rapid
391 rates of urbanization (Unger et al., 2003). Many of the studies were carried out in particular
392 geographic or industry specific settings and are not generalizable to the entire population as
393 suggested. Others have not been sampled in a probabilistic manner and are not representative
394 therefore of an accurate estimate. Likewise, the human and cattle population data utilised in this
395 review gives some good preliminary information and indicators for the region's trends over the last
396 decades, but the accuracy and consistency of the data portrayed is uncertain. Both FAOSTAT and
397 the World Bank databases rely on many different sources to compile datasets, some being
398 household surveys while others coming from countries own reporting and estimations by different
399 organisations.

400 With an obvious scarcity of literature from the human health field there is a need for further
401 research to assess the burden of the disease on humans within this setting. Although human health
402 workshop delegates felt they lacked knowledge of brucellosis presentation and diagnostics, they
403 were very encouraged by spending time with colleagues from the animal health sector discussing
404 the issues surrounding the disease.

405 Although diagnostic techniques for *Brucella* are not routinely used in most countries, the capacity is
406 largely there and small stumbling blocks could be overcome with regional co-operation. As was
407 demonstrated at diagnostic training sessions during the workshop, delegates were comfortable
408 with the techniques and quickly adapted to new concepts brought forward for alterations in field
409 testing protocols.

410 It is clear that the study countries while all experiencing urbanisation, population growth and
411 changing consumer demands, vary a lot in their size, capacity and progression through structural
412 adjustments on the path to development. While the inland countries are generally much larger
413 geographically they all so have more favourable livestock rearing conditions. The smaller coastal
414 countries however are often much more densely populated with very large urban populations and
415 high demand for dairy and value added products in general. We have only examined the
416 francophone countries of the region and must not forget the likely influence that Nigeria has over
417 regional affairs given its sheer size and dominance of both population and economy in the region.

418 In all countries, urbanisation is associated with changes in employment patterns and an increase in
419 the 'middle classes' with higher disposable incomes. Increasingly both sexes are working outside of
420 the home often with long commutes across large cities. There is therefore less time available to
421 people for shopping, processing and preparing food. In turn, this is driving trends for convenient
422 processed food that can be purchased and often consumed away from the home. As predicted by
423 Bennett's Law (BENNETT, 1954) there is a strong relationship between increasing incomes and a
424 disproportionate rise in the share of non-staple foods in the overall food budget of a household.
425 This is most notably seen in increased demand for high protein products such as meat and dairy.
426 Historically raising GDP of a country was linked with increased consumption of animal fats and
427 proteins. However, more recent food balance sheets from the FAO seem to show that this link is
428 less pronounced, with increased consumption of animal products even in those countries with low
429 GDP and high levels of urbanisation. This suggests that drivers besides those purely economic

430 influence consumption patterns within modern urban settings. Explanation could come in the form
431 of modern retailing within most African countries and the high penetration of internet technologies
432 in this setting. Most large urban areas in West Africa have a myriad of international and national
433 supermarket chains, which utilise mass media advertising and offer a more luxurious shopping
434 experience for the growing middle classes. As well as targeting the higher income population, many
435 supermarkets recognise the large proportions of low-income residents in urban dwellings and offer
436 small package low cost items. As is evident in many developed countries, the supermarket's
437 influence over agricultural policy can be very powerful. As they take over a larger market share,
438 their ability to dictate market prices and place demands in terms of food safety and value addition
439 shapes the direction of many sectors. The influence of such players in West Africa over agricultural
440 policy and knock on effects for animal health implications in the future is not be overlooked.

441 Despite the rise in number and popularity of supermarkets within West Africa little data is available
442 regionally outlining the market coverage of the sector at this time and the relative amounts of
443 locally produced versus imported goods sold by supermarkets in the region.

444 Companies such as Arla and Nestle have established processing plants in West Africa and import
445 large quantities of European milk powder for reconstitution in the region before widespread
446 distribution. Locally produced milk cannot compete with the cheap imports due the inconsistent
447 supply over different seasons and the logistical difficulties with the transport and storage of short
448 shelf life fresh produce. This reliance on imported goods reduces the potential for development of
449 the regional dairy industry. When the increasing consumer demand is being supplied by imports,
450 there is less impetus for policy change and development to strengthen the sector.

451 None the less there is clear indication of a vast interest, effort and need for further development of
452 the dairy sector in the region. There are many issues that the countries face in developing the dairy
453 sector in order to compete on a more level playing field with cheap European imports of dry milk
454 powder. However, there are great drivers to do so not only for regional economic growth but also

455 to alleviate poverty and to enhance nutrition in line with the sustainable development goals. One
456 of the most important areas in the development of the dairy sector is animal health, due not only
457 to the productivity limiting nature of endemic diseases but also the human health risks associated
458 with many.

459 From review of the literature and workshop discussions, it would appear that the largest and most
460 difficult gap to overcome in terms of animal health and particularly brucellosis control is the lack of
461 policy or legislation. This ultimately requires reliable research in to the current epidemiological
462 picture in the emerging peri-urban systems and assessment of the economic impacts of disease and
463 benefits of control.

464 5. Conclusion

465 At present, the true social and economic burden of brucellosis in West and Central Africa is
466 unknown. Scarce and aged data would suggest that the epidemiological picture is evolving as
467 demographic changes occur at varying levels across the region. In order to assess the impact and
468 evaluate control methods, systematic research is required to investigate seroprevalence in
469 emerging livestock systems and in human populations. Only then can meaningful impact
470 evaluations inform policy decisions as to appropriate and sustainable control and surveillance
471 strategies for the future.

472 6. Acknowledgements

473 The authors would like to thank all the participants of the workshop who gave their time and
474 insight. The staff at the EISMV who graciously helped with all the organisation of the workshop
475 logistics are also greatly appreciated.

476 Funding: This work was supported by the Zoonosis in Emerging Livestock Systems programme
477 (ZELS)

478

479 References

- 480 Abakar, M.F., Naré, N.B., Schelling, E., Hattendorf, J., Alfaroukh, I.O., Zinsstag, J., 2014.
481 Seroprevalence of Rift Valley fever, Q fever, and brucellosis in ruminants on the southeastern
482 shore of Lake Chad. *Vector Borne Zoonotic Dis.* 14, 757–62. doi:10.1089/vbz.2014.1585
- 483 Adamou, H.H., 2014. Evaluation de trois tests de dépistage de la brucellose bovine pour une aide
484 décisionnelle de contrôle de la maladie dans le bassin laitier de Niamey (Niger). University of
485 Dakar. doi:10.1017/CBO9781107415324.004
- 486 Adehan, R., KOUTINHOIN, B., BABA-MOUSSA, L.S., AIGBE, L., AGBADJE, P.M., YOUSAO, I.A.K.,
487 2003. Prévalence de la brucellose bovine dans les Fermes d'Etat du Bénin de 2000 à 2003. *Rev.*
488 *Africaine Santé Prod. Anim.* 3, 200–204.
- 489 Akakpo, A., Bornarel, P., 1987. Epidémiologie des brucelloses animales en Afrique tropicale:
490 Enquêtes clinique, sérologique et bactériologique. *Rev. Sci. Techn. Off. Int. Epiz* 981–1027.
- 491 Akakpo, A., Teko-Agbo, A., Kone, P., 2009. The impact of brucellosis on the economy and public
492 health in Africa. pp. 85–98.
- 493 Akakpo, A.J., 1987. Brucelloses animales en Afrique tropicale. Particularités épidémiologique,
494 clinique et bactériologique. *Rev. Elev. Med. Vet. Pays Trop.* 40, 307–20.
- 495 Akakpo, J.A., Ndour, A.P.N., 2013. La brucellose bovine en Afrique de l'ouest et du centre : *Rev.*
496 *Africaine Santé Prod. Anim.* 11, 23–28.
- 497 Allsup, T., 1969. Abortion in sheep associated with *Brucella abortus* infection. *Vet. Rec.* 84, 104–108.
- 498 Angba, A., Traore, A., Fritz, P., 1987. Situation de la brucellose animale en Côte d'Ivoire'. *Rev. Elev.*
499 *Med. Vet. Pays Trop.* 325–29.
- 500 Armand, C., 2001. Etude bibliographique des zoonoses en Côte-d'Ivoire. University Paul-Sabatier,
501 Toulouse.
- 502 Audurier A., Fayomi B., Laudat P., Z., 1987. Diagnostic sérologique de la brucellose humaine au
503 Bénin. *Rev. Elev. Med. Vet. Pays Trop.* 40, 347–348.
- 504 Baumeister, R.F., Leary, M.R., 1997. Writing narrative literature reviews. *Rev. Gen. Psychol.* 1, 311–
505 320. doi:10.1037/1089-2680.1.3.311
- 506 Bayemi, P.H., Mah, G.D., Ndamukong, K., Nsongka, V.M., Leinyuy, I., Unger, H., Ndoumbe, N.M.,
507 Webb, E.C., Achukwi, M.D., Hakoue, F., Luogbou, N.D., 2015. Bovine Brucellosis in Cattle
508 Production Systems in the Western Highlands of Cameroon. *Int. J. Anim. Biol.* 1, 38–44.
- 509 Bayemi, P.H., Webb, E.C., Nsongka, M. V, 2009. Prevalence of *Brucella abortus* antibodies in serum
510 of Holstein cattle in Cameroon. *Trop Anim Heal. Prod* 41, 141–144. doi:10.1007/s11250-008-
511 9184-8
- 512 BENNETT, M.K., 1954. The world's food. A study of the interrelations of world populations, national
513 diets, and food potentials. *world's food. A study Interrelat. world Popul. Natl. diets, food*
514 *potentials.*
- 515 Bessin, R., 1982. Contribution à l'Etude de la brucellose bovine en Haute Volta. These Docteur
516 Veterinaire. University of Dakar.
- 517 Bloch, N., Diallo, I., 1991. Enquête sérologique et allergologique sur les bovins au Niger. *Rev. Elev.*
518 *Med. Vet. Pays Trop.* 44, 117–22.
- 519 Bonfoh, B., Fane, A., Konate, A., Sidibe, S., Niang, M., Simbe, C., Oumar Alfaroukh, I., Akakpo, J.A.,

- 520 Farah, Z., Nicolet, J., Zinsstag, J., 2003. La brucellose bovine: une zoonose endémique au Mali.
521 Séminaire sous-régional "Lait Sain pour le Sahel.
- 522 Boukary, A.R., Saegerman, C., Abatih, E., Fretin, D., Deken, R. De, Harouna, H.A., Yenikoye, A., Thys,
523 E., 2013. Seroprevalence and Potential Risk Factors for *Brucella* Spp . Infection in Traditional
524 Cattle , Sheep and Goats Reared in Urban , Periurban and Rural Areas of Niger 8, 1–12.
525 doi:10.1371/journal.pone.0083175
- 526 Boussini, H., Traoré, A., Tamboura, H.H., Bessin, R., 2012. Prévalence de la tuberculose et de la
527 brucellose dans les élevages bovins laitiers intra-urbains et périurbains de la ville d '
528 Ouagadougou au Burkina Faso. Rev. Sci. Tech. Int. Off. Epizoot. 31, 943–951.
- 529 Camus, E., 1995. Evaluation of trypanosomiasis and brucellosis control in cattle herds of Ivory coast.
530 Agric. Human Values 12, 90–94. doi:10.1007/BF02217299
- 531 Camus, E., 1980. Incidence clinique de la brucellose bovine dans le Nord de la Côte d'Ivoire. Rev.
532 d'élevage médecine vétérinaire des pays Trop.
- 533 Chabasse, D., Roure, C., Rhaly, A., Maiga, D., Traore, M., Tounkara, A., Dumon, H., Ranque, P., 1983.
534 Evaluation de l'état sanitaire des populations nomades et semi-nomades du Gourma-Mali:
535 approche epidemiologique II. Resultats globaux et conclusion. Med Trop 43, 127–135.
- 536 Chambron, J., 1965. La brucellose bovine au Sénégal. Rev. Elev. Med. Vet. Pays Trop. 19–38.
- 537 Chantal, J., Thomas, J., 1976. Etude sérologique sur la brucellose bovine aux abattoirs de Dakar. Rev.
538 d'élevage Med. Vet. des pays Trop. 101–108.
- 539 Charters, A., 1980. Brucellosis. Australian Family Physician.
- 540 Corbel, 2006. Brucellosis in humans and animals Brucellosis in humans and animals. WHO 89.
- 541 Corbel, M., 1988. Fertility and Infertility in Veterinary Practice, 4th edition, J.A. Laing. Br. Vet. J. 146,
542 190–221. doi:10.1016/S0007-1935(11)80015-1
- 543 Dao, S., Traore, M., Sangho, A., Dantoume, K., Oumar, A.A., Maiga, M., Bougoudogo, F., 2009.
544 Seroprevalence of Human Brucellosis in Mopti , Mali. Rev. Tunisienne d'Infectiologie 2, 23–26.
- 545 de Glanville, W.A., Conde-Álvarez, R., Moriyón, I., Njeru, J., Díaz, R., Cook, E.A.J., Morin, M.,
546 Bronsvoort, B.M. de C., Thomas, L.F., Kariuki, S., Fèvre, E.M., 2017. Poor performance of the
547 rapid test for human brucellosis in health facilities in Kenya. PLoS Negl. Trop. Dis. 11, e0005508.
548 doi:10.1371/journal.pntd.0005508
- 549 Dean, A.S., Bonfoh, B., Kulo, A.E., Boukaya, G.A., Amidou, M., Hattendorf, J., Pilo, P., Schelling, E.,
550 2013. Epidemiology of Brucellosis and Q Fever in Linked Human and Animal Populations in
551 Northern Togo. PLoS One 8, 1–8. doi:10.1371/journal.pone.0071501
- 552 Delafosse, a., Goutard, F., Thébaud, E., 2002. Epidémiologie de la tuberculose et de la brucellose des
553 bovins en zone périurbaine d ' Abéché , Tchad. Rev. Elev. Méd. Vét. Pays Trop 55, 5 – 13.
- 554 Diop, P.E.H., 1975. Ecole Inter-Etats De Sciences Et De Medecine Veterinaires De La Brucellose
555 Bovine Au Sénégal. Univ. Dakar.
- 556 Dixon, J., Gulliver, A., Gibbon, D., Hall, M., 2001. Farming Systems and Poverty.
- 557 Domenech, J., Lucet, P., Vallat, B., Stewart, C., Bonnet, J., Hentic, A., 1982. La brucellose bovine en
558 Afrique centrale. III. Résultats statistiques des enquêtes menées au Tchad et au Cameroun.
559 Rev. Elev. Med. Vet. Pays Trop. 35, 15–22.
- 560 Domenech, J., Lucet, P., Vallat, B., Stewart, C., Bonnet, J.B., Bertaudiere, L., 1980. Bovine brucellosis

561 in central Africa. II. Clinical and epidemiological survey: regional characteristics and problems of
562 semi-intensive husbandry. *Rev. Elev. Med. Vet. Pays Trop.* 33, 277–284.

563 Ducrotoy, M., Bertu, W.J., Matope, G., Cadmus, S., Conde-Álvarez, R., Gusi, A.M., Welburn, S.,
564 Ocholi, R., Blasco, J.M., Moriyón, I., 2015. Brucellosis in Sub-Saharan Africa: Current challenges
565 for management, diagnosis and control. *Acta Trop.* doi:10.1016/j.actatropica.2015.10.023

566 FAO, n.d. FAOSTAT [WWW Document]. 2017. URL <http://www.fao.org/faostat/en/#home> (accessed
567 7.5.17).

568 Fensterbank, R., Doutre, M.P., Sagna, F., 1977. Etude de la brucellose bovine dans un village de
569 Basse-Casamance (Sénégal). II. Diagnostic allergique. *Rev. d'élevage médecine vétérinaire des
570 pays Trop.* 30, 353–358. doi:10.19182/remvt.8072

571 Folitse, R.D., Boi-Kikimoto, B.B., Emikpe, B.O., Atawalna, J., 2014. The prevalence of Bovine
572 tuberculosis and Brucellosis in cattle from selected herds in Dormaa and Kintampo Districts,
573 Brong Ahafo region, Ghana. *Arch. Clin. Microbiol.* 5, 1–6. doi:10.3823/280

574 Gidel, R., Albert, J., Le Mao, G., Retif, M., 1974. La brucellose en Afrique occidentale et son incidence
575 sur la santé publique. Résultats de dix enquêtes épidémiologiques effectuées en Côte-d'Ivoire,
576 Haute-Volta et Niger de 1970 à 1973. *Rev. Elev. Med. Vet. Pays Trop.*

577 ILRI, 2012. Mapping of poverty and likely zoonoses hotspots. Zoonoses Project 4. Report to
578 Department for International Development, UK. 119 pp.

579 Kouamo, J., Habimana, S., Bada, R.A., GJ Sawadogo, Ouedraogo, G., 2010. Séroprévalences de la
580 brucellose, de la BVD et de l'IBR et impact sur la reproduction des femelles zébus Gobra et
581 croisements inséminées en milieu traditionnel dans la région de Thiès au Sénégal. *Rev. Med.
582 Vet. (Toulouse).* 314–321.

583 Koutinhoun, B., Youssao, a K.I., Agbadje, a E.H.P.M., 2003. Prévalence de la brucellose bovine dans
584 les élevages traditionnels encadrés par le Projet pour le Développement de l' Elevage (PDE)
585 au Bénin. *Rev. Med Vet* 154, 271–276.

586 Kponmassi, T., 1991. Epidémiologie des affections abortives des bovins au togo. enquête serologique
587 sur la brucellose, la chlamydie et la fièvre Q. University of Dakar.

588 Kubuafor, D.K., Awumbila, B., Akanmori, B.D., 2000. Seroprevalence of brucellosis in cattle and
589 humans in the Akwapim-South district of Ghana: public health implications. *Acta Trop.* 76, 45–
590 48. doi:10.1016/S0001-706X(00)00088-7

591 Luchsinger, D.W., Anderson, R., 1979. Longitudinal studies of naturally acquired *Brucella abortus*
592 infection in sheep. *Am. J. Vet. Res.* 40, 1307–1312.

593 Mathias, E., Mundy, P., 2010. Adding value to livestock diversity. *FAO Anim. Prod. Heal. Pap.*

594 McDermott, J., Grace, D., Zinsstag, J., 2013. Economics of brucellosis impact and control in low-
595 income countries. *Rev. sci. tech* 249–262.

596 McDermott, J.J., Arimi, S., 2002. Brucellosis in sub-Saharan Africa: epidemiology, control and
597 impact. *Vet. Microbiol.* 90, 111–134. doi:10.1016/S0378-1135(02)00249-3

598 McDermott, J.J., Grace, D., Zinsstag, J., 2013. Economics of brucellosis impact and control in low-
599 income countries. *Rev. Sci. Tech.* 32, 249–61.

600 McInerney, J., 1996. OLD ECONOMICS FOR NEW PROBLEMS -LIVESTOCK DISEASE: PRESIDENTIAL
601 ADDRESS. *J. Agric. Econ.* 47, 295–314. doi:10.1111/j.1477-9552.1996.tb00695.x

- 602 Moreno, E., 2014. Retrospective and prospective perspectives on zoonotic brucellosis. *Front.*
603 *Microbiol.* 5, 213. doi:10.3389/fmicb.2014.00213
- 604 Mugizi, D.R., BOQVIST, S., NASINYAMA, G.W., WAISWA, C., IKWAP, K., ROCK, K., LINDAHL, E.,
605 MAGNUSSON, U., ERUME, J., 2015. Prevalence of and factors associated with *Brucella* sero-
606 positivity in cattle in urban and peri-urban Gulu and Soroti towns of Uganda. *J. Vet. Med. Sci.*
607 77, 557–564. doi:10.1292/jvms.14-0452
- 608 Nakoun, E., 2004. Serological surveillance of brucellosis and Q fever in cattle in the Central African
609 Republic 92, 147–151. doi:10.1016/j.actatropica.2004.06.007
- 610 Ndamb, O., Hemme, T., Latacz-Lohmann, 2007. Dairying in Africa - Status and recent developments.
611 *Livest. Res. Rural Dev.* 19.
- 612 Njeru, J., Wareth, G., Melzer, F., Henning, K., Pletz, M.W., Heller, R., Neubauer, H., 2016. Systematic
613 review of brucellosis in Kenya: disease frequency in humans and animals and risk factors for
614 human infection. *BMC Public Health* 16, 853. doi:10.1186/s12889-016-3532-9
- 615 Pilo-Moron, E., Pierre, F., Kouame, J., 1979. La Brucellose bovine en Côte-d'Ivoire: Epidémiologie'.
616 *Rev. Elev. Med. Vet. Pays Trop.* 325–333.
- 617 Rujeni, N., Mbanzamihiho, L., 2014. Prevalence of Brucellosis among Women Presenting with
618 Abortion / Stillbirth in Huye , Rwanda 2014.
- 619 Rushton, J., Thornton, P.K., Otte, M.J., 1999. Methods of economic impact assessment. *Rev. Sci.*
620 *Tech.* 18, 315–42.
- 621 Sacquet, E., 1955. La brucellose bovine au Tchad - note préliminaire. *Rev. Elev. Med. Vet. Pays Trop.*
622 8, 5–7.
- 623 Saley, H., 1983. Contribution à l'étude des brucelloses au Niger. University of Dakar.
- 624 Sanogo, M., Abatih, E., Thys, E., Fretin, D., Berkvens, D., Saegerman, C., 2012. Risk factors associated
625 with brucellosis seropositivity among cattle in the central savannah-forest area of Ivory Coast.
626 *Prev. Vet. Med.* 107, 51–56. doi:10.1016/j.prevetmed.2012.05.010
- 627 Sanogo, M., Cissé, B., Ouattara, M., Walravens, K., Praet, N., Berkvens, D., Thys, E., 2008. Prévalence
628 réelle de la brucellose bovine dans le centre de la Côte d'Ivoire. *Name Rev. d'Elevage Médecine*
629 *Vétérinaire des Pays Trop.* 61, 147–151.
- 630 Schelling, E. [b1] (analytic), Diguimbaye, C. [b2] (analytic), DAOUD, S. [b3] (analytic), NICOLET, J.
631 (analytic), ZINSSTAG, J. [b1] (analytic), Institut tropical Suisse. Bâle, C.H.E., 2004.
632 Séroprévalences des maladies zoonotiques chez les pasteurs nomades et leurs animaux dans le
633 Chari-Baguirmi du Tchad. *Médecine Trop.* 64, 474–477.
- 634 Shaw, W., 1976. *Brucella abortus* infection in sheep. I. Field case. *Br. Vet. J.* 132, 18–27.
- 635 Shey-Njila, O., D., Nye, E., Zoli, P.A., Walravens, K., Godfroid, J., Geerts, S., 2005. Serological Survey
636 of Bovine Brucellosis in Cameroon. *Rev. Elev. Med. Vet.* 20, 139–143.
- 637 Sonhaye, A.S., 1980. Contribution a l'étude de la brucellose bovine au Togo. *Ec. Inter-Etats Des Sci.*
638 *Med. Vet. Dakar.*
- 639 Sow, I., Niang, M., Makita, K., Costard, S., Grace, D., Bonfoh, B., n.d. Investigation on the risk of
640 brucellosis linked to the production and consumption of milk in rural Cinzana , Mali . Present.
641 First Int. Congr. Pathog. Human-Animal Interface (ICOPHAI), Addis Ababa, Ethiop.
- 642 Tasei, J.-P., Ranque, P., Balique, H., 1982. La brucellose humaine au Mali : résultats d'une enquête

643 séro-épidémiologique [WWW Document]. *Acta Trop.* URL
644 <http://retro.seals.ch/cntmng?pid=act-001:1982:39::429> (accessed 11.24.15).

645 Thys E., Yahaya M.A., Walravens K., Baudoux C., Bagayoko I., Berkvens D., G.S., 2005. Etude de la
646 prévalence de la brucellose bovine en zone forestière de la Côte d'Ivoire = Study of the
647 prevalence of bovine brucellosis in the forest zone of Cote d'Ivoire= Estudio de la prevalencia
648 de la brucelosis bovina en zonas forestales de la Costa de. *Rev. Elev. Med. Vet. Pays Trop.* 58.

649 Tialla, D., Koné, P., Kadja, M.C., Kamga-Waladjo, A., Dieng, C.B., Ndoye, N., Kouame, K.G.G., Bakou,
650 S., Akakpo, A.J., 2014. Prevalence of bovine brucellosis and related risk behavior in the
651 suburban area of Dakar, Senegal = Prévalence de la brucellose bovine et comportements à
652 risque associés à cette zoonose dans la zone périurbaine de Dakar au Sénégal = Prevalencia de
653 la bruc. *Rev. Elev. Med. Vet. Pays Trop.* 67, 62–72.

654 Tounkara, K., Maiga, S., Traoré, A., Seck, B.M., Akakpo, A.J., 1994. [Epidemiology of bovine
655 brucellosis in Mali: serologic investigation and initial isolation of strains of *Brucella abortus*].
656 *Rev. Sci. Tech.* 13, 777–86.

657 Traore, A., Tamboura, H.H., Bayala, B., Rouamba, D.W., Yameogo, N., Sanou, M., 2004. Prevalence
658 globale des pathologies majeures liées à la production laitière bovine en système d'élevage
659 intraurbain à Hamdallaye (Ouagadougou). *Biotechnol. Agron. Soc. Environ.* 8, 3–8.

660 Turkson, P.K., Boadu, D.Q., 1992. Epidemiology of bovine brucellosis in the coastal savanna zone of
661 Ghana. *Acta Trop.* 52, 39–43.

662 UNDESA, 2015. World Population Prospects: The 2015 Revision, Volume I: Comprehensive Tables.
663 doi:(ST/ESA/SER.A/379).

664 Unger, F., Münstermann, A., Goumou, C.N., Apia, M., Konte, M., A, H., 2003. Risk associated with
665 bovine brucellosis in selected study herds and market places in four countries of West Africa.

666 Verger, J.M., Grayon, M., 1984. [Characteristics of 273 strains of *Brucella abortus* of African origin].
667 *Dev. Biol. Stand.* 56, 63–71.

668 Verger, J.M., Grayon, M., Doutre, M.P., Sagna, F., Grayon, M., Doutre, M.P., Sagna, F., 1979. *Brucella*
669 *abortus* d'origine bovine au Sénégal identification et typage. *Rev. Elev. Méd. Vét. Pays Trop*
670 32, 25–32.

671 WHO, 2015. WHO estimates of the global burden of foodborne diseases Foodborne diseases burden
672 epidemiology reference group 2007-2015.

673 WHO, 2005. The Control of Neglected Zoonotic Diseases.

674 Young, E.J., Corbel, M.J., 1989. *Brucellosis : clinical and laboratory aspects.* CRC Press.

675