

# The British Journal of Radiology

## Enhanced pre-operative axillary staging using intradermal microbubbles and contrast-enhanced ultrasound to detect and biopsy sentinel lymph nodes in breast cancer: A potential replacement for axillary surgery

--Manuscript Draft--

<b>Manuscript Number:</b>	BJR-D-17-00626R1
<b>Full Title:</b>	Enhanced pre-operative axillary staging using intradermal microbubbles and contrast-enhanced ultrasound to detect and biopsy sentinel lymph nodes in breast cancer: A potential replacement for axillary surgery
<b>Short Title:</b>	Microbubble and CEUS core biopsy of sentinel nodes in breast cancer
<b>Article Type:</b>	Full Paper
<b>Corresponding Author:</b>	Karina Cox, FRCS MD Maidstone Hospital Kent, ME16 9QQ UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Maidstone Hospital
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Karina Cox, FRCS MD
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Karina Cox, FRCS MD Sian Taylor-Phillips, MPhys PhD Nisha Sharma, BSc FRCR Jennifer Weeks, MBBS Philippa Mills, FRCR Ali Sever, PhD Adrian Lim, FRCR MD Isobel Haigh, FRCR Mohamed Hashem, MRCS Tania De Silva, MRCS Keshthra Satchithananda, FRCS FRCR Mengxing Tang, PhD Matthew Wallis, FRCR
<b>Order of Authors Secondary Information:</b>	
<b>Abstract:</b>	<p><b>Objectives</b> To compare the experience of 4 UK Centres in the use of intradermal microbubbles and contrast enhanced ultrasound (CEUS) to pre-operatively identify and biopsy sentinel lymph nodes (SLN) in patients with breast cancer.</p> <p><b>Methods</b> In all centres, breast cancer patients had a microbubble/ CEUS SLN core biopsy prior to axillary surgery and patients in Centres 1 and 2 had a normal grey-scale axillary ultrasound. Data was collected between 2010 and 2016; 1361 from Centre 1 (prospective, sequential), 376 from Centre 2 (retrospective, sequential), 122 from Centre 3 (retrospective, selected) and 48 from Centre 4 (prospective, selected).</p> <p><b>Results</b></p>

SLN were successfully core biopsied in 80% (Centre 1), 79.6% (Centre 2), 77.5% (Centre 3) and 88% (Centre 4). The sensitivities to identify all SLN metastases were 46.9% (95% CI 39.4-55.1), 52.5% (95% CI 39.1-65.7), 46.4% (95% CI 27.5-66.1) and 45.5% (95% CI 16.7-76.6) respectively. Specificities; 99.7% (95% CI 98.9-100), 98.1% (95% CI 94.5-99.6), 100% (95% CI 93.2-100%) and 96.3% (95% CI 81-99.9) respectively. Negative predictive values; 87.0% (95% CI 84.3-89.3), 84.5% (95% CI 78.4-89.5), 86.9% (95% CI 82.4-90.3) and 86.2% (95% CI 78.4-91.5) respectively. At Centres 1 and 2, 12/730 (1.6%) and 7/181 (4%) respectively of patients with a benign microbubble/ CEUS SLN core biopsy had 2 or more lymph node (LN) macrometastases found at the end of primary surgical treatment.

#### Conclusions

The identification and biopsy of SLN using CEUS is a reproducible technique.

#### Advances in knowledge

In the era of axillary conservation, microbubble/ CEUS SLN core biopsy has the potential to succeed surgical staging of the axilla.

Dear Reviewers,

Many thanks for your constructive review, which was very welcome. I have now addressed your comments.

Reviewer no. 1

1. As suggested, an additional paragraph has been added to the discussion and conclusion to emphasise the added value of CEUS and biopsy to a normal grey-scale axillary ultrasound with an additional reference.
2. A recent publication has been referenced with a sentence (in the discussion) explaining that a vacuum assisted biopsy does not increase the sensitivity of CEUS as a test to identify sentinel node metastases.
3. The primary data regarding evidence of a core biopsy in surgically excised lymph nodes has been obtained and added to the results sections and the discussion.
4. In the discussion section, an explanation has been given on how to avoid overtreatment of the axilla if CEUS sentinel node biopsy is adopted into routine practice. In the conclusion section more evidence and context has been added (including another reference) to justify the proposition that microbubble/ CEUS SLN core biopsy has the potential to replace surgical staging of the axilla particularly if it is combined with a grey-scale ultrasound.
5. The missing value in figure 3 has been added.

Reviewer no. 2

1. A sentence in the discussion has been added about training. In addition, a cost analysis directly comparing surgical sentinel node excision with microbubble/ CEUS SLN core biopsy has been added to the conclusion. A full economic assessment has not been carried out to look at the cost of surgical morbidity following SLNE such as ITU admission if the patient develops anaphylaxis to blue dye.

In terms of operating time, a SLNE takes approximately 30 minutes in addition to the breast procedure but the patient remains under general anaesthesia. Although modern anaesthesia is safe most anaesthetists prefer short procedures especially if the patient is obese or has other co-morbidities so reducing the length by 30 minutes may prove important. Also, if 30 minutes can be saved per appropriate patient over the course of a full operating day, this may gain enough time for another case to be added and improve theatre utilisation. A sentence to this effect has been added to the conclusion.



**Enhanced pre-operative axillary staging using intradermal microbubbles and contrast-enhanced ultrasound to detect and biopsy sentinel lymph nodes in breast cancer: A potential replacement for axillary surgery**

**Short title:** Microbubble and CEUS core biopsy of sentinel nodes in breast cancer

**Type of manuscript:** Full paper

**Authors;** Karina Cox FRCS MD<sup>1</sup>, Sian Taylor-Phillips MPhys PhD<sup>2</sup>, Nisha Sharma Bsc FRCR<sup>3</sup>, Jennifer Weeks MBBS<sup>1</sup>, Philippa Mills FRCR<sup>1</sup>, Ali Sever PhD<sup>4</sup>, Adrian Lim FRCR MD<sup>5</sup>, Isobel Haigh FRCR<sup>3</sup>, Mohamed Hashem MRCS<sup>1</sup>, Tania D'Silva MRCS<sup>6</sup>, Keshthra Satchithananda FRCS FRCR<sup>4</sup>, Mengxing Tang PhD<sup>7</sup> and Matthew Wallis FRCR<sup>8</sup>,

**Corresponding author:** Karina Cox, Peggy Wood Breast Unit, Maidstone Hospital, Hermitage Lane, Maidstone, Kent ME16 9QQ, UK.

Telephone: 00 44 1622 22 4111

Email: [karina.cox@nhs.net](mailto:karina.cox@nhs.net)

**Institutions**

1. Peggy Wood Breast Unit, Maidstone Hospital, Hermitage Lane, Maidstone, Kent ME16 9QQ, UK.
2. WMS - Population Evidence and Technologies, University of Warwick, Coventry, CV4 7AL, UK.

3. Leeds Breast Unit, St James's University Hospital, Beckett Street, Leeds, LS9 7TF, UK.
4. Breast Care Unit, Kings College Hospital, Ground floor, Cheyne Wing, Denmark Hill, London, SE5 9RS, UK.
5. Breast Unit, Charing Cross Hospital, Fulham Palace Road, W6 8RF, UK.
6. School of Surgery, Health Education Kent, Surrey and Sussex, Stewart House, 32 Russell Square, London WC1B 5DN, UK.
7. Department of Bioengineering, Imperial College London, London SW7 2AZ, UK.
8. Breast Unit, Addenbrooke's Treatment Centre, Keith Day Rd, Cambridge CB2 0SL, UK.

**Disclosure:**

Dr Adrian Lim receives some funding from Toshiba.

A proportion of this data was presented at the 102<sup>nd</sup> Scientific Assembly and Annual Meeting of the Radiological Society of North America 2016 (abstract SSA02-06).

**Acknowledgements:**

We would like to acknowledge the support of patients and staff at all 4 Centres.

1 **Enhanced pre-operative axillary staging using intradermal microbubbles**  
2 **and contrast-enhanced ultrasound to detect and biopsy sentinel lymph**  
3 **nodes in breast cancer: A potential replacement for axillary surgery**  
4  
5  
6  
7

8 **Short title:** Microbubble and CEUS core biopsy of sentinel nodes in breast  
9 cancer  
10  
11  
12  
13  
14  
15

16 **Abstract**

17 **Objectives**

18 To compare the experience of 4 UK Centres in the use of intradermal  
19 microbubbles and contrast enhanced ultrasound (CEUS) to pre-operatively  
20 identify and biopsy sentinel lymph nodes (SLN) in patients with breast cancer.  
21  
22  
23  
24  
25  
26  
27

28 **Methods**

29 In all centres, breast cancer patients had a microbubble/ CEUS SLN core biopsy  
30 prior to axillary surgery and patients in Centres 1 and 2 had a normal grey-scale  
31 axillary ultrasound. Data was collected between 2010 and 2016; 1361 from  
32 Centre 1 (prospective, sequential), 376 from Centre 2 (retrospective, sequential),  
33 122 from Centre 3 (retrospective, selected) and 48 from Centre 4 (prospective,  
34 selected).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 **Results**

46 SLN were successfully core biopsied in 80% (Centre 1), 79.6% (Centre 2), 77.5%  
47 (Centre 3) and 88% (Centre 4). The sensitivities to identify all SLN metastases  
48 were 46.9% (95% CI 39.4-55.1), 52.5% (95% CI 39.1-65.7), 46.4% (95% CI 27.5-  
49 66.1) and 45.5% (95% CI 16.7-76.6) respectively. Specificities; 99.7% (95% CI  
50 98.9-100), 98.1% (95% CI 94.5-99.6), 100% (95% CI 93.2-100%) and 96.3%  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

(95% CI 81-99.9) respectively. Negative predictive values; 87.0% (95% CI 84.3-89.3), 84.5% (95% CI 78.4-89.5), 86.9% (95% CI 82.4-90.3) and 86.2% (95% CI 78.4-91.5) respectively. At Centres 1 and 2, 12/730 (1.6%) and 7/181 (4%) respectively of patients with a benign microbubble/ CEUS SLN core biopsy had 2 or more lymph node (LN) macrometastases found at the end of primary surgical treatment.

### **Conclusions**

The identification and biopsy of SLN using CEUS is a reproducible technique.

### **Advances in knowledge**

In the era of axillary conservation, microbubble/ CEUS SLN core biopsy has the potential to succeed surgical staging of the axilla.



## Introduction

1 Sentinel lymph node excision (SLNE) is the axillary staging method of choice for  
2 breast cancer patients with normal axillary lymph nodes (LN) on grey-scale  
3 ultrasound or a benign biopsy of morphologically abnormal LN (1,2). Although  
4 SLNE has less reported morbidity than axillary lymph node dissection (ALND) it  
5 remains a surgical procedure performed under general anaesthesia with  
6 recognised immediate complications such as infection (11%) and long term  
7 problems with sensory loss (11%) and arm lymphoedema (5%) at 12 months (3)  
8 The operation is also reliant upon 2 tracers (radioactive isotope and blue dye) to  
9 locate SLN and maintain a false negative rate of 6% (4). The blue dye carries a  
10 hazard of anaphylaxis (0.9%) (5) and there are logistical challenges in obtaining  
11 medical grade radioisotopes.

12 Recently, the practice of removing all malignant axillary LN to achieve local  
13 control has been challenged by the results of a trial where patients with SLN  
14 metastases were randomized to a completion ALND or no further axillary  
15 surgery (6). The local recurrence rate in the axilla was low with no difference  
16 between the groups despite the fact that 27.3% of patients in the ALND arm had  
17 further lymph node metastases retrieved (6). These results emphasize the role  
18 of modern adjuvant treatment in preventing loco-regional disease recurrence  
19 and have led to the introduction of conservative surgical management of the  
20 axilla for patients with up to 2 SLN macrometastases (7).

21 In patients with breast cancer, SLN can be identified and percutaneously  
22 biopsied in the clinic using intradermally injected microbubbles and contrast-  
23 enhanced ultrasound (CEUS) (8, 9, 10). The technique was originally described in  
24 a swine melanoma model (11) and following a trial period, was incorporated  
25

1 into routine practice at Centre 1 for all invasive breast cancer patients with a  
2 normal grey-scale axillary ultrasound (12,13) to aid treatment planning such as  
3 the selection of appropriate axillary surgery (14), the initiation of neo-adjuvant  
4 systemic therapy (15) and reconstructive decisions for patients who may  
5 consequently benefit from post-mastectomy radiotherapy (16). The  
6 identification of SLN using microbubbles and CEUS was previously validated  
7 against the surgical detection of SLN using blue dye and radioisotope and  
8 concordance was found in 93% of cases that had a core biopsy (13). Other  
9 Centres in the UK also adopted the microbubble/ CEUS SLN biopsy technique for  
10 use in their own practice.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 Previous work from Centre 1 has shown that patients with invasive breast  
24 cancer and a normal grey-scale ultrasound and benign microbubble/ CEUS SLN  
25 core biopsy are unlikely to have extensive metastatic axillary disease that is both  
26 grey-scale ultrasound occult and missed on SLN core biopsy (14). These results  
27 suggest that complete radiological staging of the axilla might be feasible.  
28  
29  
30  
31  
32  
33  
34

35 We therefore aimed to assess the reproducibility of the microbubble/ CEUS SLN  
36 core biopsy procedure by comparing the test accuracy of Centre 1 with a smaller  
37 population of consecutive patients from Centre 2 as well as 2 other UK centres  
38 using patients, which were selected on the basis of tumour clinico-pathological  
39 features. We also examined the technical performance of individual radiologists  
40 at Centre 1. Lastly, we measured the volume of axillary metastases in patients  
41 from Centres 1 and 2 to determine the proportion of patients with a false  
42 negative microbubble/ CEUS SLN core biopsy who had 2 or more axillary LN  
43 macrometastases at the end of primary surgical treatment.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1           **Methods**

### 2           *Study Design*

3  
4  
5  
6           Data was collected from 4 Breast Units across the UK (figure 1). In Centres 1 and  
7  
8  
9           2, patients were included with newly diagnosed breast cancer and an initial grey-  
10  
11           scale axillary ultrasound +/- biopsy of indeterminate/ abnormal LN (17) and  
12  
13           only those with normal axillary imaging/ benign pathology proceeded to have a  
14  
15           microbubble/ CEUS SLN core biopsy. At Centre 3, patients with newly diagnosed  
16  
17           grade 2 or 3 invasive breast cancer and a normal grey-scale axillary ultrasound  
18  
19           or indeterminate LN that were inadequate (not enough cells to make a diagnosis)  
20  
21           or benign on cytology were included. At Centre 4, patients with newly diagnosed  
22  
23           T2 or larger tumours, grade 2 or 3 invasive breast cancer with grey-scale  
24  
25           ultrasound indeterminate axillary LN were included. Centres 3 and 4 selected  
26  
27           patients at the discretion of the multi-disciplinary team. In all centres, pregnant  
28  
29           patients and those with locally advanced/ metastatic disease were excluded. At  
30  
31           Centre 1, 1361 consecutive patients were identified from a prospective database  
32  
33           of patients having microbubble/ CEUS guided SLN core biopsy between  
34  
35           December 2010 and November 2016. A proportion of this data (570 patients)  
36  
37           has been previously analysed and published (14). Retrospective data was  
38  
39           collected on 376 consecutive patients from Centre 2 between July 2010 and July  
40  
41           2014. Retrospective data was collected on 122 selected patients from Centre 3  
42  
43           between July 2013 and December 2015. Prospective data was collected on 48  
44  
45           selected patients from Centre 4 between June 2013 and February 2015.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*Identification and biopsy of SLN using intradermal microbubbles and CEUS*

1 Following a diagnosis of breast cancer, patients had an injection of ultrasound  
2  
3 contrast agent (Sonovue, BRACCO Imaging S.p.A, Italy). 0.2ml of ultrasound  
4  
5 contrast agent was injected intra-dermally in the periareolar, upper outer  
6  
7 quadrant position using a 26G needle with a 1 ml tuberculin syringe at Centre 1,  
8  
9 Centre 2 and Centre 3. Centre 4 used a 24G needle with a 1 ml tuberculin syringe.  
10  
11 Lymphatic channels were visualised on contrast pulse sequencing and tracked  
12  
13 into the axilla. Areas of contrast accumulation were also imaged with grey scale  
14  
15 or live dual images (figure 2). Further injections of contrast up to 1ml and 3  
16  
17 consecutive injections together with injection site massage for 10-30 seconds  
18  
19 were performed if progress of contrast through the lymphatic vessels was slow/  
20  
21 not immediately evident. Once identified, the SLN was biopsied using a core  
22  
23 biopsy technique, Centres 1 and 2 (Achieve automatic biopsy device, Carefusion,  
24  
25 San Diego, California, USA. 2-3x 14-16G), Centre 3 (Trucut biopsy device, San  
26  
27 Diego, California, USA. 3-4x 14G) and Centre 4 (Trucut biopsy device. 3-4x 18G).  
28  
29 Microbubble/ CEUS guided SLN core biopsies were performed by consultant  
30  
31 breast radiologists; 7 at Centre 1, 4 at Centre 2, 2 at Centre 3 and a single  
32  
33 consultant breast radiologist at Centre 4. Three of the 7 radiologists at Centre 1  
34  
35 also worked at Centre 2. Very rarely 2 SLN were visualised and both were  
36  
37 biopsied at Centre 1, Centre 2 and Centre 3. At Centre 4, only the first enhancing  
38  
39 SLN was biopsied. Centre 1 ultrasound examinations were performed with a  
40  
41 Sequoia 512 Acuson (Siemens Medical Systems, Issaquah, Wash, USA) or LOGIQ  
42  
43 9 (GE Healthcare, Fairfield, CT, USA) using a linear transducer operating at 4.5 to  
44  
45 15MHz. Centre 2 ultrasound examinations were performed with a LOGIQ 9 using  
46  
47 a linear transducer operating at 4.5 to 15MHz, Centre 3 ultrasound examinations  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 were performed with a Siemens Antaris or Acuson S1000 (Siemens Medical  
2 Systems, Issaquah, Wash, USA) and Centre 4 ultrasound examinations were  
3 performed with a Toshiba Aplio 500 (Toshiba Medical Systems Europe B.V.,  
4 Zilverstraat 1, 2718RP, Zoetermeer, The Netherlands) using a mid-range linear  
5 probe (PLT-704SBT Linear). All ultrasound machines provided: conventional  
6 grey-scale, pulse-inversion harmonic grey scale, contrast specific sonographic  
7 imaging with live dual images of tissue only and contrast agent images. In order  
8 to reduce microbubble destruction, low mechanical index values were applied  
9 (MI: 0.1-0.4). Each centre performed histological analysis on biopsy samples.  
10 Centres 1 and 2 also used pancytokeratin (MNF116) immunohistochemical  
11 staining for patients with a lobular phenotype.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

### 28 *Surgical Management of the axilla*

29 Initially, at Centre 1 and Centre 2 if the microbubble/ CEUS SLN core biopsy  
30 contained malignant cells, patients were advised to have an ipsilateral ALND.  
31  
32 Later in the study period (Centre 1 and Centre 2) and throughout the study  
33 period at Centre 3 and Centre 4, to avoid surgical overtreatment, patients were  
34 recommended to have ALND only if a macrometastasis (>2mm) was seen in the  
35 core biopsy specimen and those with core biopsy isolated tumour cells (ITC) or  
36 micrometastases (<2mm) had SLNE. Patients also had SLNE if the core biopsy  
37 identified indeterminate cellular changes, normal lymphoid tissue, inadequate  
38 tissue sampling or the patient declined primary ALND. Before surgery, patients  
39 had an injection of radioactive isotope (Nanocoll, G.E Healthcare, Chicago, USA),  
40 between 20 and 40mbeq in the peri-areolar, upper outer quadrant position.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 anaesthetic induction, patients received a 2ml injection of blue dye (Bleu Patente  
2 V 2.5%, Guerbet, France) sub-dermally in the periareolar, upper outer quadrant  
3 position. A gamma probe (Navigator GPS, RMD Instruments, Watertown, USA)  
4 was used to identify SLN. All SLN within the axilla were excised and sent for local  
5 histological analysis including immunohistochemical staining for patients with a  
6 lobular phenotype at Centres 1 and 2. The total volume of axillary metastases at  
7 the end of surgical treatment was determined for each patient using the  
8 following scoring system: 1 (LN containing a macrometastasis), 0.5 (LN  
9 containing a micrometastasis) and 0.2 (LN containing isolated tumour cells)(12).  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

### 23 *Statistics*

24  
25 The proportion of patients who had SLN identified was calculated. Using  
26 contingency tables, the sensitivity and specificity of a successful (LN tissue  
27 retrieved) CEUS guided SLN core biopsy, as the index test to identify SLN  
28 metastases in breast cancer patients with invasive disease was determined with  
29 axillary surgery (SLNE/ ALND) as the reference standard. The estimated values  
30 were calculated along with corresponding 95% confidence intervals (CI)  
31 illustrating the uncertainty in the results. All CI were calculated using the exact  
32 binomial method. The prevalence of SLN metastases in each population was  
33 derived from the reference standard. For centres 1 and 2, positive and negative  
34 predictive values were calculated directly, as they both employed consecutive  
35 enrolment so the study prevalence will be representative of the population of  
36 interest. For centres 3 and 4, which only included a non-randomly selected  
37 subset of eligible patients, Bayesian methods were used to calculate positive and  
38 negative predictive values, using the prevalence from centre 1 (the largest centre  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

with consecutive enrolment). The Mann-Whitney test was used to compare the volume of axillary metastases between patients with a false negative and true positive microbubble/ CEUS SLN biopsy. All analysis was completed using Stata software (Stata version 13.1; Stata Corp LP, College Station, Tx, USA).

### *Ethics*

Kent Research Ethics Committee, UK approved the original trials at Centre 1 (reference numbers: 04/Q1801/25 and 11/H1101/1) and the Medicines and Healthcare Products Regulatory Agency (MHRA) sanctioned the use of microbubbles by intraparenchymal injection (Eudract Number: 2004-002423-41). All participating Centres were NHS Trusts and the New Procedures Committees at Centre 1, Centre 2, and Centre 3 and the Breast Services Committee at Centre 4 each approved the use of intradermal microbubbles and CEUS to identify and biopsy SLN in patients with breast cancer at their institution.

## Results

### Visualisation and targeted core biopsy of SLN using intradermal microbubbles and CEUS

A total of 1361 consecutive patients were identified from a prospectively maintained database at Centre 1. Complete data was not available for 12 patients and 1 patient had a fine needle aspiration biopsy (FNAB) rather than a core biopsy. Sentinel lymph nodes were clearly visualised by CEUS in 1216 of the 1348 patients (90%). In the 132 cases where the intradermally injected microbubbles failed to traffic through lymphatics to identify SLN, 25 (19%) had previous surgery on the breast/ axilla, a post breast biopsy haematoma or malignant involvement of the nipple areolar complex. A successful SLN core biopsy was achieved in 1083 of the 1348 patients (80%). The performance statistics of the 7 Centre 1 radiologists are presented in Table 1.

A total of 376 patients were identified retrospectively at Centre 2. Complete data was not available for 15 patients, 3 patients had a FNAB and 15 patients did not have the microbubbles/ CEUS procedure. Sentinel lymph nodes were clearly visualised by CEUS in 290 of 343 patients (84.5%). A successful core biopsy of SLN was achieved in 273 of 343 patients (79.6%).

At Centre 3, 121 patients were identified retrospectively. Complete data was not available for one patient. Sentinel lymph nodes were clearly visualised by CEUS in 109 of the 120 patients (90.8%). A successful core biopsy of SLN was achieved in 93 of 120 patients (77.5%). Prospective data was collected on 48 patients at Centre 4. There were technical problems with the initial 5 patients, as the lymphatics could not be visualised. However, once the settings on the ultrasound machine were optimised, SLN were clearly visualised and successfully core



1 biopsied in 38 of 43 patients (88%). In all centres, there were no allergic  
2 reactions following the administration of contrast agent and only one significant  
3 bleeding complication (Centre 1) of a large haematoma after SLN core biopsy  
4 which was evacuated 2 weeks later at the time of primary surgical treatment.  
5  
6  
7  
8  
9

10 **Identification of SLN metastases in patients with invasive breast cancer and**  
11 **a successful microbubble/ CEUS SLN core biopsy that underwent primary**  
12 **surgical treatment**  
13  
14  
15  
16

17 In patients with a successful microbubble/ CEUS SLN core biopsy, the following  
18 were excluded from final analysis; those with pre-invasive disease (DCIS), those  
19 who had primary systemic or endocrine therapy and patients who did not  
20 proceed with axillary surgery because of choice or inability to tolerate a general  
21 anaesthetic. At Centres 1 and 2, patients with un-biopsied grey-scale abnormal  
22 LN were excluded and 6 patients at Centre 1 were also excluded because of  
23 incomplete surgical data.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 At Centre 1, 816 of the 1083 patients with a successful microbubble/ CEUS SLN  
37 core biopsy went on to have primary surgical treatment. Evidence of a core  
38 biopsy tract was seen in the excised SLN of 656 (80%) patients, 127 (16%)  
39 patients did not have a core biopsy tract visualized in excised SLN, 2 (0.2%)  
40 patients had a core biopsy tract seen in non-SLN and data regarding the presence  
41 of a core biopsy tract was not recorded for 31 (3.8%) patients. For the other  
42 centres; 215 of 273 patients (Centre 2), 80 of 93 patients (Centre 3) and 38 of 38  
43 patients (Centre 4) with a successful microbubble/ CEUS SLN core biopsy went  
44 on to have primary surgical treatment. Comprehensive data documenting  
45 evidence of previous biopsy in excised SLN was not available for Centres 2, 3 and  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

4. For each centre, the accuracy of a successful microbubble/ CEUS SLN core biopsy to identify SLN metastases in patients with invasive breast cancer is presented in Table 2.

**Volume of axillary metastases at the end of primary surgical treatment in patients with invasive breast cancer, radiologically normal axillary LN and a successful microbubble/ CEUS SLN core biopsy in Centres 1 and 2**

At Centre 1, 95 patients had a false negative microbubble/ CEUS SLN core biopsy with metastases found in the excised SLN. Of these, evidence of a core biopsy tract was seen in the excised SLN of 69 (72.6%) patients, 23 (24.2%) patients did not have a core biopsy tract visualised in excised SLN and data regarding the presence of a core biopsy tract was not recorded for 3 (3.2%) patients. Sixty-one patients (64%) with a false negative microbubble/ CEUS SLN core biopsy result went on to have a completion ALND and 34 (36%) had no further axillary surgery (axillary conservation). At the end of primary surgical treatment, 12 patients (12.6%) were found to have 2 or more LN macrometastases and 82 patients (87.4%) had less than 2 LN macrometastases. Of the 730 patients with an initial benign microbubble/ CEUS SLN core biopsy, only 12 (1.6%) had 2 or more LN macrometastases found at the end of primary surgical treatment (Table 3). Eighty-four patients at Centre 1 had a true positive microbubble/ CEUS SLN core biopsy and 81 (96%) had an ALND whereas 3 (4%) had a SLNE. Of these 84 patients, 42 (50%) had 2 or more axillary LN macrometastases at the end of primary surgical treatment.

At Centre 2, 28 patients had a false negative microbubble/ CEUS SLN core biopsy with metastases found in the excised SLN. Nineteen patients (68%) went on to

1 have a completion ALND and 9 (32%) had axillary conservation. At the end of  
2 primary surgical treatment, 7 patients (25%) were found to have 2 or more  
3 axillary macrometastases and 22 patients (75%) had a malignant axillary LN  
4 score less than 2 or had ITC in multiple LN (one patient with ITC in 13 LN). Of the  
5 181 patients with an initial benign microbubble/ CEUS SLN core biopsy, 7 (4%)  
6 had 2 or more axillary macrometastases found at the end of primary surgical  
7 treatment (Table 4). Thirty-one patients at Centre 2 had a true positive  
8 microbubble/ CEUS SLN core biopsy and 30 (97%) had an ALND whereas 1 (3%)  
9 had a SLNE. Of these 31 patients, 14 (45%) had 2 or more axillary LN  
10 macrometastases at the end of primary surgical treatment.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 For both centres, at the end of surgical treatment the difference between the  
24 final malignant LN score of false negative versus true positive microbubble/  
25 CEUS SLN core biopsies was statistically significant (figure 3).  
26  
27  
28  
29  
30  
31  
32

### 33 **Discussion**

34 Despite the use of different ultrasound machines and variations in the methods  
35 and patient selection, the identification and core biopsy of SLN using intradermal  
36 microbubbles and CEUS in patients with breast cancer is a reproducible  
37 technique across multiple centres. Overall, the visualisation of SLN across the 4  
38 centres ranged from 84.5% to 90.8% and a successful core biopsy from 77.5% to  
39 88%. Factors that appeared to adversely affect the visualisation of SLN at Centre  
40 1 included previous surgery and disease involvement of the nipple areolar  
41 complex.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 There is undoubtedly a learning curve associated with the procedure and  
56 familiarity with the equipment is important as demonstrated by the data from  
57  
58  
59  
60  
61  
62  
63  
64  
65

Centre 4 where they experienced technical problems with the first 5 patients.

1 The performance of Centre 1 radiologists also highlight the distinct  
2 competencies of the 2 components of the procedure, namely SLN identification  
3 and SLN core biopsy. Even though identical ultrasound equipment and methods  
4 were used, the percentage of procedures with successful visualisation of SLN  
5 varied from 97% to 73%, which suggests either that not all radiologists at Centre  
6 1 received adequate procedural training before performing the test or some  
7 found it difficult to visualise microbubbles trafficking through lymphatic  
8 channels. Six of the 7 radiologists were fairly consistent in their ability to  
9 successfully core biopsy visualised SLN, but 1 obviously struggled and only  
10 successfully retrieved lymphoid tissue in 70% of cases. Anecdotally, most  
11 radiologists accustomed to the procedure recommend that novices observe 3  
12 cases, then perform 10 cases supervised before undertaking 30 independent  
13 procedures with an audit of their results. Once proficient, the whole procedural  
14 time is 15 to 30 minutes.

15 There is scope to improve the technology of CEUS. In swine models, LN  
16 metastases can be identified as areas devoid of contrast agent (11) and in a  
17 recent study of breast cancer patients, the sensitivity of CEUS as a test to identify  
18 SLN metastases using only enhancement patterns (no biopsy) was 81.8% (18).  
19 Innovations such as ultrafast ultrasound (19), super resolution imaging (20) and  
20 improved lymphatic microbubble transit (21) may improve the ability of  
21 clinicians to visualise SLN and achieve a reliable standard.

22 The sensitivity of a microbubble/ CEUS core biopsy as a test to identify SLN  
23 metastases in patients with invasive breast cancer and a normal grey-scale  
24 axillary ultrasound/ benign axillary LN biopsy is consistently around 50% with

Centres 2, 3 and 4 within the 95% confidence intervals of Centre 1. As grey-scale axillary ultrasound can usefully identify approximately 50% of LN metastases (22), the addition of a microbubble/ CEUS SLN core biopsy substantially increases the overall detection rate for metastatic axillary LN. Consequently, the negative predictive value of the test is high and <5% of patients (Centres 1 and 2) with a normal grey-scale ultrasound and benign microbubble/ CEUS SLN core biopsy had 2 or more LN macrometastases detected by axillary surgery.

We have previously speculated that the technique has a high false negative rate because the core biopsy fails to pick up small metastatic deposits in SLN (13) and this may be the reason why very few patients with a false negative microbubble/ CEUS core biopsy at Centres 1 and 2 had two or more axillary LN macrometastases found at the end of primary surgical treatment. Yet, retrieving more LN tissue with a vacuum-assisted biopsy technique does not appear to appreciably increase the sensitivity of microbubbles and CEUS (23).

Alternatively, it is usual for only one SLN to be visualised and biopsied with CEUS but the median number of SLN retrieved with a surgical excision is 2 (3). In this series, 80% of surgically excised LN at Centre 1 showed evidence of a previous core biopsy and this proportion dropped to 72.6% in patients with a false negative benign microbubble/ CEUS SLN core biopsy. This raises the possibility that in the false negative cases, the second or subsequent SLN contained the metastases and perhaps more than one SLN should be actively sought with the microbubbles/ CEUS procedure.

When compared to Centre 1, a higher proportion of Centre 2 patients with a benign microbubble/ CEUS SLN core biopsy had two or more axillary macrometastases found at the end of surgical treatment (1.8 vs 4%). This may

1 be related to the smaller patient sample size in Centre 2 or the higher prevalence  
2 of LN metastases in patients with a successful microbubble/ CEUS SLN core  
3 biopsy in Centre 2 (27% vs 22% at Centre 1). As the patient and  
4 clinicopathologic features of the tumours were similar in both centres, the  
5 greater prevalence of LN metastases at Centre 2 may be a consequence of a  
6 lower initial metastatic LN detection rate with grey-scale axillary ultrasound.  
7

8  
9  
10  
11  
12 The difference in the volume of axillary disease between patients with a false  
13 negative and true positive microbubble/ CEUS SLN core biopsy at the end of  
14 primary surgical treatment was statistically significant in Centres 1 and 2. When  
15 compared to the false negative groups in Centres 1 and 2, more patients with a  
16 true positive microbubble/ CEUS SLN core biopsy had complete axillary surgery  
17 (ALND) rather than axillary conservation (SLNE). The retrieval of a greater  
18 number of LN in the true positive groups from Centres 1 and 2 may therefore  
19 have increased the total number of axillary LN metastases found at the end of  
20 surgical treatment and influenced the final metastatic score. However, in a  
21 previous publication from Centre 1 (14) comparing only patients with complete  
22 axillary surgery (ALND), the difference in the volume of axillary disease between  
23 those with a false negative and true positive microbubble/ CEUS SLN core biopsy  
24 remained statistically significant.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 An argument against using CEUS to biopsy SLN in routine practice is that  
46 patients with a biopsy containing malignant cells are committed to a primary  
47 ALND for what may be a low burden of axillary disease. This can be mitigated  
48 against by offering SLNE to patients with micrometastases in the core biopsy  
49 specimen. It should also be noted that 50% of patients at Centre 1 and 45% of  
50 patients at Centre 2 with a true positive microbubble/ CEUS SLN core biopsy had  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

2 or more axillary macrometastases found at the end of surgical treatment and therefore using the test for patients who are not eligible for axillary conservation (7) is beneficial.

## **Conclusion**

The results of the American College of Surgeons Oncology Group Z0011 trial (6) have changed practice by showing that loco-regional control of axillary metastases is not solely dependant upon surgical excision and residual disease can be treated with adjuvant therapy. In addition, anatomic staging of breast cancer is likely to become less relevant to treatment decisions as tumour genomic and molecular assays are better understood (24). Based on recent information obtained from Centre 1, a SLNE costs 3.6x the cost of a microbubble/ CEUS SLN core biopsy (£671.63 vs £189 respectively). Omitting axillary surgery will improve theatre utilisation (potentially allowing more cases to be added to a list) as well as reducing the anaesthetic time for each patient. In the era of axillary conservation, molecular medicine and dwindling resources the combination of grey-scale axillary ultrasound and microbubble/ CEUS SLN core biopsy has the potential to succeed surgical staging of the axilla. Further work now needs to be undertaken to refine the procedure with protocols, standard setting and training.

1           **References**  
2  
3

- 4           **1.** Veronesi U, Paganelli G, Viale G, et al. A randomised comparison of  
5           sentinel-node biopsy with routine axillary dissection in breast cancer. N  
6           Engl J Med 2003;349:546-553.  
7  
8  
9  
10  
11           **2.** CG80 Early and locally advanced breast cancer: full guideline. United  
12           Kingdom, National Institute for Health and Clinical Excellence, February  
13           2009.           (Accessed           July           2016           at  
14           [https://www.nice.org.uk/guidance/cg80/resources/early-and-locally-](https://www.nice.org.uk/guidance/cg80/resources/early-and-locally-advanced-breast-cancer-diagnosis-and-treatment-975682170565)  
15           [advanced-breast-cancer-diagnosis-and-treatment-975682170565](https://www.nice.org.uk/guidance/cg80/resources/early-and-locally-advanced-breast-cancer-diagnosis-and-treatment-975682170565)).  
16  
17  
18  
19  
20  
21  
22  
23           **3.** Mansel RE, Fallowfield L, Kissin M, Goyal A, Newcombe RG, Dixon J M et al.  
24           Randomized Multicenter Trial of Sentinel Node Biopsy Versus Standard  
25           Axillary Treatment in Operable Breast Cancer: The Almanac Trial.  
26           J Nat Can Inst 2006; 98: 599-609.  
27  
28  
29  
30  
31  
32  
33           **4.** Pesek S, Ashikaga T, Krag L E, Krag D. The false-negative rate of sentinel  
34           node biopsy in patients with breast cancer: a meta-analysis. World J Surg  
35           2012; 36: 2239-2251.  
36  
37  
38  
39  
40  
41           **5.** Barthelmes L, Goyal A, Newcombe RG, McNeill F, Mansel RE. Adverse  
42           reactions to patent blue V dye – The NEW START and ALMANAC  
43           experience. Eur J Surg Oncol 2010; 36: 399-403.  
44  
45  
46  
47  
48           **6.** Giuliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PW,  
49           Blumencranz PW, et al. Axillary Dissection vs No Axillary Dissection in  
50           Women With Invasive Breast Cancer and Sentinel Node Metastasis. JAMA  
51           2011; 305: 569-575.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
7. Association of Breast Surgery Consensus Statement Management of the Malignant Axilla in Early Breast Cancer, March 2015. (Accessed June 2017 at [http://www.associationofbreastsurgery.org.uk/media/50934/axilla\\_abs\\_consensus\\_statement\\_16\\_3\\_15.pdf](http://www.associationofbreastsurgery.org.uk/media/50934/axilla_abs_consensus_statement_16_3_15.pdf)).
  8. Omoto K, Matsunaga H, Take N, Hozumi Y, Takehara M, Omoto Y et al. Sentinel Node detection method using contrast-enhanced ultrasonography with sonazoid in breast cancer: preliminary clinical study. *Ultrasound Med Biol* 2009; 35: 1249-1256.
  9. Sever A, Jones S, Cox K, Weeks J, Mills P, Jones P. Preoperative localization of sentinel lymph nodes using intradermal microbubbles and contrast-enhanced ultrasonography in patients with breast cancer. *Br J Surg* 2009; 96:1295-1299.
  10. Rautiainen S, Sudah M, Joukainen S, Sironen R, Vanninen R and Sutela A. Contrast-enhanced ultrasound-guided axillary lymph node core biopsy: Diagnostic accuracy in preoperative staging of invasive breast cancer. *Eur J Radiol* 2015; 84: 2130-2136.
  11. Goldberg BB, Merton DA, Liu JB et al. Sentinel lymph nodes in a swine model with melanoma: contrast-enhanced lymphatic US. *Radiology* 2004;230:324-330.
  12. Sever A, Broillet A, Schneider M, Cox K, Jones S et al. Dynamic visualisation of lymphatic channels and sentinel lymph nodes using intradermal microbubbles and contrast enhanced ultrasound in a swine model and patients with breast cancer. *J Ultrasound Med* 2010; 29: 1699-704.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
13. Cox K, Sever A, Jones S, Weeks J, Mills P, Devalia H et al. Validation of a technique using microbubbles and contrast enhanced ultrasound (CEUS) to biopsy sentinel lymph nodes (SLN) in pre-operative breast cancer patients with a normal grey-scale axillary ultrasound. *Eur J Surg Oncol* 2013; 39: 760-765.
  14. Cox K, Weeks J, Mills P, Chalmers R, Devalia H, Fish D, Sever A. Contrast-Enhanced Ultrasound Biopsy of Sentinel Lymph Nodes in Patients with Breast Cancer: Implications for Axillary Metastases and Conservation. *Ann Surg Oncol* 2016; 23: 58-64.
  15. Kilbride KE, Lee MC, Nees AV, Cimmino VM, Diehl KM, Sabel MS, Hayes DF, Schott AF, Kleer CG, Chang AE, Newman LA. Axillary staging prior to neoadjuvant chemotherapy for breast cancer: predictors of recurrence. *Ann Surg Oncol* 2008; 15: 3252-3258.
  16. Su YL, Li SH, Chen YY, Chen HC, Tay Y, Huang CH, Chou FF, Wu SC, Rau KM. Post-mastectomy radiotherapy benefits subgroups of breast cancer patients with T1-2 tumor and 1-3 axillary lymph node(s) metastasis. *Radiol Oncol* 2014; 48 (3): 314-322.
  17. Yang WT, Metreweli C, Lam PK, Chang J. Benign and malignant breast masses and axillary nodes: evaluation with echo enhanced color power Doppler Ultrasound. *Radiology* 2001; 220:795-802.
  18. Xie F, Zhang D, Cheng L, Yu L, Yang L, Tong F et al. Intradermal microbubbles and contrast-enhanced ultrasound (CEUS) is a feasible approach for sentinel lymph identification in early-stage breast cancer. *World J Surg Oncol* 2015; 13: 319.
  19. Tanter M, Fink M. Ultrafast imaging in biomedical ultrasound. *IEEE Trans*

Ultrason Ferroelectr Freq Control 2014; 61: 102-119.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
20. Christensen-Jeffries K, Browning RJ, Tang M-X, Dunsby CW; Eckersley RJ, In Vivo Acoustic Super-Resolution and Super-Resolved Velocity Mapping Using Microbubbles. IEEE TRANSACTIONS ON MEDICAL IMAGING 2015; 34: 433-440
21. Gorce J-M, Arditi M, Schneider M. Influence of bubble size distribution on the echogenicity of ultrasound contrast agents. Investigative Radiology 2000; 35: 661-671.
22. Diepstraten SC, Sever AR, Buckens CF, Veldhuis WB, van Dalen T, van den Bosch MA, Mali WP and Verkooijen HM. Value of preoperative ultrasound-guided axillary lymph node dissection in breast cancer: a systematic review and meta-analysis. Ann Surg Oncol 2014; 21: 51-59.
23. Britton P, Willsher P, Taylor K, Kilburn-Toppin F, Provenzano E, Forouhi P, Benson J, Agrawal A, Forman JR and Wallis MG. Microbubble detection and ultrasound-guided vacuum-assisted biopsy of axillary lymph nodes in patients with breast cancer. Clin Radiol 2017; 72: 772-779.
24. Donovan CA, Giuliano AE. Evolution of the Staging System in Breast Cancer. Ann Surg Oncol 2017; 24: 3469-3470.

1  
2  
3 **Figure Captions**  
4  
5

6 **Figure 1**

7  
8 Diagram showing the selection and flow of participants from each centre  
9 through the study. Lymph node (LN), Sentinel Lymph Node (SLN),  
10 Contrast enhanced ultrasound (CEUS), Ductal Carcinoma in situ (DCIS),  
11 isolated tumour cells (ITC), surgical sentinel lymph node biopsy (SLNB),  
12 axillary lymph node dissection (ALND) and POSNOC – (POSitive Sentinel  
13 NOde: adjuvant therapy alone versus adjuvant therapy plus Clearance or  
14 axillary radiotherapy. A randomised controlled trial of axillary treatment  
15 in women with early stage breast cancer who have metastases in one or  
16 two sentinel nodes).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31  
32  
33 **Figure 2**

34  
35 A. Ultrasound contrast pulse sequencing image of a SLN (white arrow)  
36 after injection of intradermal microbubbles (between 0.2 and 1 ml  
37 injected using a 26G needle with 1ml tuberculin syringe) into the UOQ  
38 periareolar area of the breast. B. Grey scale ultrasound image of the same  
39 SLN as visualised in A. Images provided by Centre 3.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 **Figure 3**

52  
53 A. Volume of axillary disease at the end of primary surgical treatment for  
54 individual patients at Centres 1 and 2 who had a false negative (FN) or  
55 true positive (TP) microbubble/ CEUS core biopsy of SLN. Axillary lymph  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 node dissection (ALND) and sentinel lymph node excision (SLNE). B. The  
2 total volume of axillary metastases at the end of surgical treatment was  
3 determined using a scoring system (isolated tumour cells = 0.2, each LN  
4 micrometastasis = 0.5 and each LN macrometastasis = 1). The Mann-  
5 Whitney test was used to compare the volume of axillary metastases  
6 between patients with a false negative and true positive microbubble/  
7 CEUS SLN biopsy in both centres.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **Table 1**

19 Performance statistics of 7 consultant breast radiologists at Centre 1  
20 (Total 1349 of 1361 procedures). In 12 cases the data was incomplete and  
21 the name of the radiologist was not recorded. For radiologist no.1, one  
22 successful procedure was a fine needle aspiration biopsy rather than a  
23 core biopsy.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

### 36 **Table 2**

37 Test accuracy of CEUS guided SLN biopsy using intradermally injected  
38 microbubbles in women with normal (centres 1 and 2) or indeterminate  
39 results from previous grey scale ultrasound (with or without previous  
40 biopsy). Reference standard is Sentinel Lymph node excision (SLNE) or  
41 axillary lymph node dissection (ALND). Positive predictive value (PPV)  
42 and negative predictive value (NPV) were calculated directly for centres 1  
43 and 2 who employed consecutive recruitment, and using Bayesian  
44 methods with 22% prevalence at centres 3 and 4 as these were not a  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

consecutively or randomly selected group. True positive (TP), false positive (FP), true negative (TN) and false negative (FN).

**Table 3**

Age and clinicopathological characteristics of all patients at Centre 1 (first column) with a successful microbubble/CEUS SLN core biopsy before primary surgical treatment and Centre 1 patients with false negative microbubble/ CEUS SLN core biopsies sub-divided into micrometastases (<2mm)/ isolated tumour cells (ITC), low volume metastases and high volume metastases identified at the end of primary surgical treatment. Data are expressed as *n* (%). Estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (Her2), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC).

**Table 4**

Age and clinicopathological characteristics of all patients at Centre 2 (first column) with a successful microbubble/CEUS SLN core biopsy before primary surgical treatment and Centre 2 patients with false negative microbubble/ CEUS SLN core biopsies sub-divided into micrometastases (<2mm)/ isolated tumour cells (ITC), low volume metastases and high volume metastases identified at the end of primary surgical treatment. Data are expressed as *n* (%). Estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (Her2), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC).

Table 1

	Centre 1 – Consultant Breast Radiologists						
	1	2	3	4	5	6	7
Total Procedures	276	37	501	81	116	207	131
Procedures where core biopsy not attempted	0 (0%)	0 (0%)	0 (0%)	2 (2.5%)	0 (0%)	0 (0%)	1 (0.8%)
Procedures with successful visualisation of SLN	269 (97.5%)	33 (89.2%)	457 (91.2%)	59 (72.8%)	94 (81%)	187 (90.3%)	118 (90.1%)
Procedures without SLN tissue retrieved	1 (0.4%)	1 (2.7%)	64 (12.8%)	2 (2.5%)	10 (8.6%)	18 (8.7%)	34 (26%)
Successful retrieval of lymphoid tissue in those with visualised SLN	268 (99.6%)	32 (97%)	393 (86%)	57 (96.6%)	84 (89.4%)	169 (90.4%)	84 (71.2%)
Total successful visualisation and core biopsy of SLN	268 (97.1%)	32 (86.5%)	393 (78.4%)	57 (70.4%)	84 (72.4%)	169 (81.6%)	84 (64.1%)

Table 2

Centre	Prevalence of LN metastases	TP	FP	TN	FN	Sensitivity	Specificity	PPV	NPV
1	22%	84	2	635	95	46.9% (39.4-55.5%)	99.7 % (98.95-100%)	97.7% (91.9-99.7%)	87.0% (84.3-89.3%)
2	27%	31	3	153	28	52.5% (39.1-65.7%)	98.1% (94.5-99.6%)	91.2% (76.3-98.1%)	84.5% (78.4-89.5%)
3	35%	13	0	52	15	46.4% (27.5-66.1%)	100% (93.2-100%)	100% (75.3-100%)	86.9% (82.4-90.3%)
4	29%	5	1	26	6	45.5% (16.7-76.6%)	96.3% (81.0-99.9%)	77.6% (31.3-96.3%)	86.2 % (78.4 -91.5%)



Table 3

		Volume of disease at the end of surgical treatment (microbubble/ CEUS false negative core biopsy)		
Centre 1	All Patients	Micrometastasis/ ITC	Low (<2 LN macrometastases)	High (2 or more LN macrometastases)
Total number of patients	816	37	46	12
		15 (41%) ALND	34 (74%) ALND	12 (100%) ALND
Median age in years (range)	61 (30-94)	61 (42-89)	55 (32-90)	53 (36-69)
<b>Receptor Status</b>				
ER positive	707 (87%)	32 (86%)	40 (87%)	11 (92%)
ER unknown	4 (0.4%)	0	0	0
Her-2 positive	79 (10%)	3 (8%)	2 (4%)	1 (8%)
Her-2 not recorded	13 (1.6%)	0	2 (4%)	0
ER-/PR-/Her-2-	72 (10%)	4 (10.8%)	4 (9%)	0
ER-/PR-/HER-2+	28 (3%)	1 (2.7%)	1 (2%)	1 (8%)
<b>Invasive Tumour Size</b>				
DCIS + microinvasion	5 (0.6%)	0	0	0
T1	472 (58%)	19 (51%)	17 (37%)	4 (33%)
T2	199 (24%)	10 (27%)	18 (39%)	5 (42%)
T3	29 (4%)	5 (14%)	3 (7%)	1 (8%)
Multifocal	111 (14%)	3 (8%)	8 (17%)	2 (17%)
Unknown	0	0	0	0
<b>Tumour Grade</b>				
Grade 1	180 (22%)	6 (16%)	7 (15%)	4 (33%)
Grade 2	392 (48%)	20 (54%)	21 (46%)	6 (50%)
Grade 3	209 (26%)	11 (30%)	14 (30%)	1 (8%)
Mixed grade	29 (4%)	0	4 (9%)	1 (8%)
Unknown	1 (0.1%)	0	0	0
<b>Tumour Type</b>				
IDC	660 (81%)	23 (62%)	38 (83%)	11 (92%)
ILC	103 (13%)	11 (30%)	6 (13%)	1 (8%)
Other	29 (4%)	1 (3%)	1 (2%)	0
Mixed	17 (2%)	2 (5%)	1 (2%)	0
Unknown	2 (0.2%)	0	0	0

Table 4

		Volume of disease at the end of surgical treatment (microbubble/ CEUS false negative core biopsy)		
Centre 2	All Patients	Micrometastasis/ ITC	Low (<2 LN macrometastases)	High (2 or more LN macrometastases)
Total number of patients	215	10	11	7
		2 (20%) ALND	10 (91%) ALND	7 (100%) ALND
Median age in years (range)	64 (31-93)	63.5 (37-93)	62 (38-91)	55 (47-72)
<b>Receptor Status</b>				
ER positive	176 (82%)	10 (100%)	8 (73%)	6 (86%)
ER unknown	6 (3%)	0	1 (9%)	1 (14%)
Her-2 positive	29 (13%)	1 (10%)	0	1 (14%)
Her-2 not recorded	8 (4%)	0	1 (9%)	1 (14%)
ER-/PR-/Her-2-	23 (11%)	0	1 (9%)	0
ER-/PR-/HER-2+	10 (5%)	0	0	0
<b>Invasive Tumour Size</b>				
DCIS + microinvasion	0	0	0	0
T1	118 (55%)	2 (20%)	6 (55%)	2 (29%)
T2	71 (33%)	6 (60%)	4 (36%)	5 (71%)
T3	9 (4%)	0	0	0
Multifocal	17 (8%)	2 (20%)	0	0
Unknown	0	0	1 (9%)	0
<b>Tumour Grade</b>				
Grade 1	35(16%)	1 (10%)	0	2 (29%)
Grade 2	102 (47%)	6 (60%)	5 (45.5%)	1 (14%)
Grade 3	68 (32%)	2 (20%)	4 (36%)	4 (57%)
Mixed grade	4 (2%)	0	0	0
Unknown	6 (3%)	0	2 (18%)	0
<b>Tumour Type</b>				
IDC	160 (74%)	7 (70%)	8 (73%)	5 (71%)
ILC	24 (11%)	2 (20%)	1 (9%)	0
Other	16 (7%)	0	0	0
Mixed	13 (6%)	1 (10%)	1 (9%)	2 (29%)
Unknown	2 (1%)	0	1 (9%)	0

1 **Enhanced pre-operative axillary staging using intradermal microbubbles**  
2 **and contrast-enhanced ultrasound to detect and biopsy sentinel lymph**  
3 **nodes in breast cancer: A potential replacement for axillary surgery**  
4  
5  
6  
7

8 **Short title:** Microbubble and CEUS core biopsy of sentinel nodes in breast  
9 cancer  
10  
11  
12  
13  
14  
15

16 **Abstract**

17 **Objectives**

18 To compare the experience of 4 UK Centres in the use of intradermal  
19 microbubbles and contrast enhanced ultrasound (CEUS) to pre-operatively  
20 identify and biopsy sentinel lymph nodes (SLN) in patients with breast cancer.  
21  
22  
23  
24  
25  
26  
27

28 **Methods**

29 In all centres, breast cancer patients had a microbubble/ CEUS SLN core biopsy  
30 prior to axillary surgery and patients in Centres 1 and 2 had a normal grey-scale  
31 axillary ultrasound. Data was collected between 2010 and 2016; 1361 from  
32 Centre 1 (prospective, sequential), 376 from Centre 2 (retrospective, sequential),  
33 122 from Centre 3 (retrospective, selected) and 48 from Centre 4 (prospective,  
34 selected).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 **Results**

46 SLN were successfully core biopsied in 80% (Centre 1), 79.6% (Centre 2), 77.5%  
47 (Centre 3) and 88% (Centre 4). The sensitivities to identify all SLN metastases  
48 were 46.9% (95% CI 39.4-55.1), 52.5% (95% CI 39.1-65.7), 46.4% (95% CI 27.5-  
49 66.1) and 45.5% (95% CI 16.7-76.6) respectively. Specificities; 99.7% (95% CI  
50 98.9-100), 98.1% (95% CI 94.5-99.6), 100% (95% CI 93.2-100%) and 96.3%  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

(95% CI 81-99.9) respectively. Negative predictive values; 87.0% (95% CI 84.3-89.3), 84.5% (95% CI 78.4-89.5), 86.9% (95% CI 82.4-90.3) and 86.2% (95% CI 78.4-91.5) respectively. At Centres 1 and 2, 12/730 (1.6%) and 7/181 (4%) respectively of patients with a benign microbubble/ CEUS SLN core biopsy had 2 or more lymph node (LN) macrometastases found at the end of primary surgical treatment.

### **Conclusions**

The identification and biopsy of SLN using CEUS is a reproducible technique.

### **Advances in knowledge**

In the era of axillary conservation, microbubble/ CEUS SLN core biopsy has the potential to succeed surgical staging of the axilla.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1 Sentinel lymph node excision (SLNE) is the axillary staging method of choice for  
2 breast cancer patients with normal axillary lymph nodes (LN) on grey-scale  
3 ultrasound or a benign biopsy of morphologically abnormal LN (1,2). Although  
4 SLNE has less reported morbidity than axillary lymph node dissection (ALND) it  
5 remains a surgical procedure performed under general anaesthesia with  
6 recognised immediate complications such as infection (11%) and long term  
7 problems with sensory loss (11%) and arm lymphoedema (5%) at 12 months (3)  
8 The operation is also reliant upon 2 tracers (radioactive isotope and blue dye) to  
9 locate SLN and maintain a false negative rate of 6% (4). The blue dye carries a  
10 hazard of anaphylaxis (0.9%) (5) and there are logistical challenges in obtaining  
11 medical grade radioisotopes.  
12

13 Recently, the practice of removing all malignant axillary LN to achieve local  
14 control has been challenged by the results of a trial where patients with SLN  
15 metastases were randomized to a completion ALND or no further axillary  
16 surgery (6). The local recurrence rate in the axilla was low with no difference  
17 between the groups despite the fact that 27.3% of patients in the ALND arm had  
18 further lymph node metastases retrieved (6). These results emphasize the role  
19 of modern adjuvant treatment in preventing [loco-regional](#) disease recurrence  
20 and have led to the introduction of conservative surgical management of the  
21 axilla for patients with up to 2 SLN macrometastases (7).  
22

23 In patients with breast cancer, SLN can be identified and percutaneously  
24 biopsied in the clinic using intradermally injected microbubbles and contrast-  
25 enhanced ultrasound (CEUS) (8, 9, 10). The technique was originally described in  
26 a swine melanoma model (11) and following a trial period, was incorporated  
27

1 into routine practice at Centre 1 for all invasive breast cancer patients with a  
2 normal grey-scale axillary ultrasound (12,13) to aid treatment planning such as  
3 the selection of appropriate axillary surgery (14), the initiation of neo-adjuvant  
4 systemic therapy (15) and reconstructive decisions for patients who may  
5 consequently benefit from post-mastectomy radiotherapy (16). The  
6 identification of SLN using microbubbles and CEUS was previously validated  
7 against the surgical detection of SLN using blue dye and radioisotope and  
8 concordance was found in 93% of cases that had a core biopsy (13). Other  
9 Centres in the UK also adopted the microbubble/ CEUS SLN biopsy technique for  
10 use in their own practice.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 Previous work from Centre 1 has shown that patients with invasive breast  
24 cancer and a normal grey-scale ultrasound and benign microbubble/ CEUS SLN  
25 core biopsy are unlikely to have extensive metastatic axillary disease that is both  
26 grey-scale ultrasound occult and missed on SLN core biopsy (14). These results  
27 suggest that complete radiological staging of the axilla might be feasible.  
28  
29  
30  
31  
32  
33  
34

35 We therefore aimed to assess the reproducibility of the microbubble/ CEUS SLN  
36 core biopsy procedure by comparing the test accuracy of Centre 1 with a smaller  
37 population of consecutive patients from Centre 2 as well as 2 other UK centres  
38 using patients, which were selected on the basis of tumour clinico-pathological  
39 features. We also examined the technical performance of individual radiologists  
40 at Centre 1. Lastly, we measured the volume of axillary metastases in patients  
41 from Centres 1 and 2 to determine the proportion of patients with a false  
42 negative microbubble/ CEUS SLN core biopsy who had 2 or more axillary LN  
43 macrometastases at the end of primary surgical treatment.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Methods

### *Study Design*

Data was collected from 4 Breast Units across the UK (figure 1). In Centres 1 and 2, patients were included with newly diagnosed breast cancer and an initial grey-scale axillary ultrasound +/- biopsy of indeterminate/ abnormal LN (17) and only those with normal axillary imaging/ benign pathology proceeded to have a microbubble/ CEUS SLN core biopsy. At Centre 3, patients with newly diagnosed grade 2 or 3 invasive breast cancer and a normal grey-scale axillary ultrasound or indeterminate LN that were inadequate (not enough cells to make a diagnosis) or benign on cytology were included. At Centre 4, patients with newly diagnosed T2 or larger tumours, grade 2 or 3 invasive breast cancer with grey-scale ultrasound indeterminate axillary LN were included. Centres 3 and 4 selected patients at the discretion of the multi-disciplinary team. In all centres, pregnant patients and those with locally advanced/ metastatic disease were excluded. At Centre 1, 1361 consecutive patients were identified from a prospective database of patients having microbubble/ CEUS guided SLN core biopsy between December 2010 and November 2016. A proportion of this data (570 patients) has been previously analysed and published ([14reference blinded for review](#)). Retrospective data was collected on 376 consecutive patients from Centre 2 between July 2010 and July 2014. Retrospective data was collected on 122 selected patients from Centre 3 between July 2013 and December 2015. Prospective data was collected on 48 selected patients from Centre 4 between June 2013 and February 2015.

1 *Identification and biopsy of SLN using intradermal microbubbles and CEUS*

2  
3 Following a diagnosis of breast cancer, patients had an injection of ultrasound  
4 contrast agent (Sonovue, BRACCO Imaging S.p.A, Italy). 0.2ml of ultrasound  
5 contrast agent was injected intra-dermally in the periareolar, upper outer  
6 quadrant position using a 26G needle with a 1 ml tuberculin syringe at Centre 1,  
7 Centre 2 and Centre 3. Centre 4 used a 24G needle with a 1 ml tuberculin syringe.  
8 Lymphatic channels were visualised on contrast pulse sequencing and tracked  
9 into the axilla. Areas of contrast accumulation were also imaged with grey scale  
10 or live dual images (figure 2). Further injections of contrast up to 1ml and 3  
11 consecutive injections together with injection site massage for 10-30 seconds  
12 were performed if progress of contrast through the lymphatic vessels was slow/  
13 not immediately evident. Once identified, the SLN was biopsied using a core  
14 biopsy technique, Centres 1 and 2 (Achieve automatic biopsy device, Carefusion,  
15 San Diego, California, USA. 2-3x 14-16G), Centre 3 (Trucut biopsy device, San  
16 Diego, California, USA. 3-4x 14G) and Centre 4 (Trucut biopsy device. 3-4x 18G).  
17 Microbubble/ CEUS guided SLN core biopsies were performed by consultant  
18 breast radiologists; 7 at Centre 1, 4 at Centre 2, 2 at Centre 3 and a single  
19 consultant breast radiologist at Centre 4. Three of the 7 radiologists at Centre 1  
20 also worked at Centre 2. Very rarely 2 SLN were visualised and both were  
21 biopsied at Centre 1, Centre 2 and Centre 3. At Centre 4, only the first enhancing  
22 SLN was biopsied. Centre 1 ultrasound examinations were performed with a  
23 Sequoia 512 Acuson (Siemens Medical Systems, Issaquah, Wash, USA) or LOGIQ  
24 9 (GE Healthcare, Fairfield, CT, USA) using a linear transducer operating at 4.5 to  
25 15MHz. Centre 2 ultrasound examinations were performed with a LOGIQ 9 using



1 a linear transducer operating at 4.5 to 15MHz, Centre 3 ultrasound examinations  
2 were performed with a Siemens Antaris or Acuson S1000 (Siemens Medical  
3 Systems, Issaquah, Wash, USA) and Centre 4 ultrasound examinations were  
4 performed with a Toshiba Aplio 500 (Toshiba Medical Systems Europe B.V.,  
5 Zilverstraat 1, 2718RP, Zoetermeer, The Netherlands) using a mid-range linear  
6 probe (PLT-704SBT Linear). All ultrasound machines provided: conventional  
7 grey-scale, pulse-inversion harmonic grey scale, contrast specific sonographic  
8 imaging with live dual images of tissue only and contrast agent images. In order  
9 to reduce microbubble destruction, low mechanical index values were applied  
10 (MI: 0.1-0.4). Each centre performed histological analysis on biopsy samples.  
11 Centres 1 and 2 also used pancytokeratin (MNF116) immunohistochemical  
12 staining for patients with a lobular phenotype.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

### 31 *Surgical Management of the axilla*

32 Initially, at Centre 1 and Centre 2 if the microbubble/ CEUS SLN core biopsy  
33 contained malignant cells, patients were advised to have an ipsilateral ALND.  
34  
35 Later in the study period (Centre 1 and Centre 2) and throughout the study  
36 period at Centre 3 and Centre 4, to avoid surgical overtreatment, patients were  
37 recommended to have ALND only if a macrometastasis (>2mm) was seen in the  
38 core biopsy specimen and those with core biopsy isolated tumour cells (ITC) or  
39 micrometastases (<2mm) had SLNE. Patients also had SLNE if the core biopsy  
40 identified indeterminate cellular changes, normal lymphoid tissue, inadequate  
41 tissue sampling or the patient declined primary ALND. Before surgery, patients  
42 had an injection of radioactive isotope (Nanocoll, G.E Healthcare, Chicago, USA),  
43 between 20 and 40mbeq in the peri-areolar, upper outer quadrant position.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Later that day or the following day, patients underwent surgical resection. After  
2 anaesthetic induction, patients received a 2ml injection of blue dye (Bleu Patente  
3 V 2.5%, Guerbet, France) sub-dermally in the periareolar, upper outer quadrant  
4 position. A gamma probe (Navigator GPS, RMD Instruments, Watertown, USA)  
5 was used to identify SLN. All SLN within the axilla were excised and sent for local  
6 histological analysis including immunohistochemical staining for patients with a  
7 lobular phenotype at Centres 1 and 2. The total volume of axillary metastases at  
8 the end of surgical treatment was determined for each patient using the  
9 following scoring system: 1 (LN containing a macrometastasis), 0.5 (LN  
10 containing a micrometastasis) and 0.2 (LN containing isolated tumour cells)(12).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

### 26 *Statistics*

27  
28 The proportion of patients who had SLN identified was calculated. Using  
29 contingency tables, the sensitivity and specificity of a successful (LN tissue  
30 retrieved) CEUS guided SLN core biopsy, as the index test to identify SLN  
31 metastases in breast cancer patients with invasive disease was determined with  
32 axillary surgery (SLNE/ ALND) as the reference standard. The estimated values  
33 were calculated along with corresponding 95% confidence intervals (CI)  
34 illustrating the uncertainty in the results. All CI were calculated using the exact  
35 binomial method. The prevalence of SLN metastases in each population was  
36 derived from the reference standard. For centres 1 and 2, positive and negative  
37 predictive values were calculated directly, as they both employed consecutive  
38 enrolment so the study prevalence will be representative of the population of  
39 interest. For centres 3 and 4, which only included a non-randomly selected  
40 subset of eligible patients, Bayesian methods were used to calculate positive and  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 negative predictive values, using the prevalence from centre 1 (the largest centre  
2 with consecutive enrolment). The Mann-Whitney test was used to compare the  
3 volume of axillary metastases between patients with a false negative and true  
4 positive microbubble/ CEUS SLN biopsy. All analysis was completed using Stata  
5 software (Stata version 13.1; Stata Corp LP, College Station, Tx, USA).  
6  
7  
8  
9

### 10 11 12 *Ethics*

13  
14  
15  
16 Kent Research Ethics Committee, UK approved the original trials at  
17 Centre 1 (reference numbers: 04/Q1801/25 and 11/H1101/1) and the  
18 Medicines and Healthcare Products Regulatory Agency (MHRA) sanctioned the  
19 use of microbubbles by intraparenchymal injection (Eudract Number: 2004-  
20 002423-41). All participating Centres were NHS Trusts and the New Procedures  
21 Committees at Centre 1, Centre 2, and Centre 3 and the Breast Services  
22 Committee at Centre 4 each approved the use of intradermal microbubbles and  
23 CEUS to identify and biopsy SLN in patients with breast cancer at their  
24 institution.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Results

### Visualisation and targeted core biopsy of SLN using intradermal microbubbles and CEUS

A total of 1361 consecutive patients were identified from a prospectively maintained database at Centre 1. Complete data was not available for 12 patients and 1 patient had a fine needle aspiration biopsy (FNAB) rather than a core biopsy. Sentinel lymph nodes were clearly visualised by CEUS in 1216 of the 1348 patients (90%). In the 132 cases where the intradermally injected microbubbles failed to traffic through lymphatics to identify SLN, 25 (19%) had previous surgery on the breast/ axilla, a post breast biopsy haematoma or malignant involvement of the nipple areolar complex. A successful SLN core biopsy was achieved in 1083 of the 1348 patients (80%). The performance statistics of the 7 Centre 1 radiologists are presented in Table 1.

A total of 376 patients were identified retrospectively at Centre 2. Complete data was not available for 15 patients, 3 patients had a FNAB and 15 patients did not have the microbubbles/ CEUS procedure. Sentinel lymph nodes were clearly visualised by CEUS in 290 of 343 patients (84.5%). A successful core biopsy of SLN was achieved in 273 of 343 patients (79.6%).

At Centre 3, 121 patients were identified retrospectively. Complete data was not available for one patient. Sentinel lymph nodes were clearly visualised by CEUS in 109 of the 120 patients (90.8%). A successful core biopsy of SLN was achieved in 93 of 120 patients (77.5%). Prospective data was collected on 48 patients at Centre 4. There were technical problems with the initial 5 patients, as the lymphatics could not be visualised. However, once the settings on the ultrasound

1 machine were optimised, SLN were clearly visualised and successfully core  
2 biopsied in 38 of 43 patients (88%). In all centres, there were no allergic  
3 reactions following the administration of contrast agent and only one significant  
4 bleeding complication (Centre 1) of a large haematoma after SLN core biopsy  
5 which was evacuated 2 weeks later at the time of primary surgical treatment.  
6  
7  
8  
9

10  
11  
12  
13 **Identification of SLN metastases in patients with invasive breast cancer and**  
14 **a successful microbubble/ CEUS SLN core biopsy that underwent primary**  
15 **surgical treatment**  
16  
17  
18  
19

20  
21 In patients with a successful microbubble/ CEUS SLN core biopsy, the following  
22 were excluded from final analysis; those with pre-invasive disease (DCIS), those  
23 who had primary systemic or endocrine therapy and patients who did not  
24 proceed with axillary surgery because of choice or inability to tolerate a general  
25 anaesthetic. At Centres 1 and 2, patients with un-biopsied grey-scale abnormal  
26 LN were excluded and 6 patients at Centre 1 were also excluded because of  
27 incomplete surgical data.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 At Centre 1, 816 of the 1083 patients with a successful microbubble/ CEUS SLN  
39 core biopsy went on to have primary surgical treatment. [Evidence of a core  
40 biopsy tract was seen in the excised SLN of 656 \(80%\) patients, 127 \(16%\)  
41 patients did not have a core biopsy tract visualized in excised SLN, 2 \(0.2%\)  
42 patients had a core biopsy tract seen in non-SLN and data regarding the presence  
43 of a core biopsy tract was not recorded for 31 \(3.8%\) patients.](#) For the other  
44 centres; 215 of 273 patients (Centre 2), 80 of 93 patients (Centre 3) and 38 of 38  
45 patients (Centre 4) with a successful microbubble/ CEUS SLN core biopsy went  
46 on to have primary surgical treatment. [Comprehensive data documenting](#)  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

[evidence of previous biopsy in excised SLN was not available for Centres 2, 3 and](#)

[4.](#) For each centre, the accuracy of a successful microbubble/ CEUS SLN core biopsy to identify SLN metastases in patients with invasive breast cancer is presented in Table 2.

**Volume of axillary metastases at the end of primary surgical treatment in patients with invasive breast cancer, radiologically normal axillary LN and a successful microbubble/ CEUS SLN core biopsy in Centres 1 and 2**

At Centre 1, 95 patients had a false negative microbubble/ CEUS SLN core biopsy with metastases found in the excised SLN. [Of these, evidence of a core biopsy tract was seen in the excised SLN of 69 \(72.6%\) patients, 23 \(24.2%\) patients did not have a core biopsy tract visualised in excised SLN and data regarding the presence of a core biopsy tract was not recorded for 3 \(3.2%\) patients.](#) Sixty-one patients (64%) [with a false negative microbubble/ CEUS SLN core biopsy result](#) went on to have a completion ALND and 34 (36%) had no further axillary surgery (axillary conservation). At the end of primary surgical treatment, 12 patients (12.6%) were found to have 2 or more LN macrometastases and 82 patients (87.4%) had less than 2 LN macrometastases. Of the 730 patients with an initial benign microbubble/ CEUS SLN core biopsy, only 12 (1.6%) had 2 or more LN macrometastases found at the end of primary surgical treatment (Table 3). Eighty-four patients at Centre 1 had a true positive microbubble/ CEUS SLN core biopsy and 81 (96%) had an ALND whereas 3 (4%) had a SLNE. Of these 84 patients, 42 (50%) had 2 or more axillary LN macrometastases at the end of primary surgical treatment.

1 At Centre 2, 28 patients had a false negative microbubble/ CEUS SLN core biopsy  
2 with metastases found in the excised SLN. Nineteen patients (68%) went on to  
3 have a completion ALND and 9 (32%) had axillary conservation. At the end of  
4 primary surgical treatment, 7 patients (25%) were found to have 2 or more  
5 axillary macrometastases and 22 patients (75%) had a malignant axillary LN  
6 score less than 2 or had ITC in multiple LN (one patient with ITC in 13 LN). Of the  
7 181 patients with an initial benign microbubble/ CEUS SLN core biopsy, 7 (4%)  
8 had 2 or more axillary macrometastases found at the end of primary surgical  
9 treatment (Table 4). Thirty-one patients at Centre 2 had a true positive  
10 microbubble/ CEUS SLN core biopsy and 30 (97%) had an ALND whereas 1 (3%)  
11 had a SLNE. Of these 31 patients, 14 (45%) had 2 or more axillary LN  
12 macrometastases at the end of primary surgical treatment.

13 For both centres, at the end of surgical treatment the difference between the  
14 final malignant LN score of false negative versus true positive microbubble/  
15 CEUS SLN core biopsies was statistically significant (figure 3).

## 16 Discussion

17 Despite the use of different ultrasound machines and variations in the methods  
18 and patient selection, the identification and core biopsy of SLN using intradermal  
19 microbubbles and CEUS in patients with breast cancer is a reproducible  
20 technique across multiple centres. Overall, the visualisation of SLN across the 4  
21 centres ranged from 84.5% to 90.8% and a successful core biopsy from 77.5% to  
22 88%. ~~The sensitivities of a microbubble/ CEUS core biopsy as a test to identify  
23 SLN metastases in patients with invasive breast cancer at Centres 2, 3 and 4  
24 were within the 95% confidence intervals of Centre 1. The factors that~~

1 appeared to adversely affect the visualisation of SLN at Centre 1 included  
2 previous surgery and disease involvement of the nipple areolar complex.

3  
4 There is undoubtedly a learning curve associated with the procedure and  
5 familiarity with the equipment is important as demonstrated by the data from  
6 Centre 4 where they experienced technical problems with the first 5 patients.  
7  
8 The performance of Centre 1 radiologists also ~~highlight~~ illustrate the distinct  
9 competencies of the 2 components of the procedure, namely SLN identification  
10 and SLN core biopsy. Even though identical ultrasound equipment and methods  
11 were used, the percentage of procedures with successful visualisation of SLN  
12 varied from 97% to 73%, which suggests either that not all radiologists at Centre  
13 1 received adequate procedural training before performing the test or some  
14 found it difficult to visualise microbubbles trafficking through lymphatic  
15 channels. Six of the 7 radiologists were fairly consistent in their ability to  
16 successfully core biopsy visualised SLN, but 1 obviously struggled and only  
17 successfully retrieved lymphoid tissue in 70% of cases. Anecdotally, most  
18 radiologists accustomed to the procedure recommend that novices observe 3  
19 cases, then perform 10 cases supervised before undertaking 30 independent  
20 procedures with an audit of their results. Once proficient, the whole procedural  
21 time is 15 to 30 minutes.

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48 There is scope to improve the technology of CEUS. In swine models, LN  
49 metastases can be identified as areas devoid of contrast agent (11) and in a  
50 recent study of breast cancer patients, the sensitivity of CEUS as a test to identify  
51 SLN metastases using only enhancement patterns (no biopsy) was 81.8% (18).  
52  
53 Innovations such as ultrafast ultrasound (19), super resolution imaging (20) and  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



improved lymphatic microbubble transit (21) may improve the ability of clinicians to visualise SLN and achieve a reliable standard.

~~The sensitivities of a microbubble/ CEUS core biopsy as a test to identify SLN metastases in patients with invasive breast cancer and a normal grey-scale axillary ultrasound/ benign axillary LN biopsy is consistently around 50% with at Centres 2, 3 and 4 were within the 95% confidence intervals of Centre 1. The sensitivity of the technique as a test to identify SLN metastases is consistently around 50%. As grey-scale axillary ultrasound can usefully identify approximately 50% of LN metastases (22), the addition of a microbubble/ CEUS SLN core biopsy substantially increases the overall detection rate for metastatic axillary LN. Consequently, the negative predictive value of the test is high and <5% of patients (Centres 1 and 2) with a normal grey-scale ultrasound and benign microbubble/ CEUS SLN core biopsy had 2 or more LN macrometastases detected by axillary surgery.~~

~~but because of the low prevalence of LN metastases in the studied patient groups, the negative predictive value of the test is high.~~

We have previously speculated that the technique has a high false negative rate because the core biopsy fails to pick up small metastatic deposits in SLN (13) and this may be the reason why very few patients with a false negative microbubble/ CEUS core biopsy at Centres 1 and 2 had two or more axillary LN macrometastases found at the end of primary surgical treatment. Yet, retrieving more LN tissue with a vacuum-assisted biopsy technique does not appear to appreciably increase the sensitivity of microbubbles and CEUS (23).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Alternatively, it is usual for only one SLN to be visualised and biopsied with CEUS but the median number of SLN retrieved with a surgical excision is 2 (3). [In this series, 80% of surgically excised LN at Centre 1 showed evidence of a previous core biopsy and this proportion dropped to 72.6% in patients with a false negative benign microbubble/ CEUS SLN core biopsy.](#) This raises the possibility that in the false negative cases, the second or subsequent SLN contained the metastases and perhaps more than one SLN should be actively sought with the microbubbles/ CEUS procedure.

When compared to Centre 1, a higher proportion of Centre 2 patients with a benign microbubble/ CEUS SLN core biopsy had two or more axillary macrometastases found at the end of surgical treatment (1.8 vs 4%). This may be related to the smaller patient sample size in Centre 2 or the higher prevalence of LN metastases in patients with a successful microbubble/ CEUS SLN core biopsy in Centre 2 (27% vs 22% at Centre 1). As the patient and clinicopathologic features of the tumours were similar in both centres, the greater prevalence of LN metastases at Centre 2 may be a consequence of a lower initial metastatic LN detection rate with grey-scale axillary ultrasound.

The difference in the volume of axillary disease between patients with a false negative and true positive microbubble/ CEUS SLN core biopsy at the end of primary surgical treatment was statistically significant in Centres 1 and 2. When compared to the false negative groups in Centres 1 and 2, more patients with a true positive microbubble/ CEUS SLN core biopsy had complete axillary surgery (ALND) rather than axillary conservation (SLNE). The retrieval of a greater number of LN in the true positive groups from Centres 1 and 2 may therefore have increased the total number of axillary LN metastases found at the end of

1 surgical treatment and influenced the final metastatic score. However, in a  
2 previous publication from Centre 1 ([14reference blinded for review](#)) comparing  
3 only patients with complete axillary surgery (ALND), the difference in the  
4 volume of axillary disease between those with a false negative and true positive  
5 microbubble/ CEUS SLN core biopsy remained stastically significant.  
6  
7

8  
9  
10  
11 [An argument against using CEUS to biopsy SLN in routine practice is that](#)  
12 [patients with a biopsy containing malignant cells are committed to a primary](#)  
13 [ALND for what may be a low burden of axillary disease. This can be mitigated](#)  
14 [against by offering SLNE to patients with micrometastases in the core biopsy](#)  
15 [specimen. It should also be noted that 50% of patients at Centre 1 and 45% of](#)  
16 [patients at Centre 2 with a true positive microbubble/ CEUS SLN core biopsy had](#)  
17 [2 or more axillary macrometastases found at the end of surgical treatment and](#)  
18 [therefore using the test for patients who are not eligible for axillary conservation](#)  
19 [\(7\) is beneficial.](#)  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

## 38 **Conclusion**

39  
40  
41 [In patients with breast cancer, the identification and biopsy of SLN using](#)  
42 [intradermally injected microbubbles and CEUS is a reproducible technique. The](#)  
43 [results of the American College of Surgeons Oncology group Z0011 trial \(6\) have](#)  
44 [changed practice by showing that loco-regional control of axillary metastases is](#)  
45 [not solely dependant upon surgical excision and residual disease can be treated](#)  
46 [with adjuvant therapy. In addition, anatomic staging of breast cancer is likely to](#)  
47 [become less relevant to treatment decisions as tumour genomic and molecular](#)  
48 [assays are better understood \(24\). Based on recent information obtained from](#)  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Centre 1, a SLNE costs 3.6x the cost of a microbubble/ CEUS SLN core biopsy  
2 (£671.63 vs £189 respectively). Omitting axillary surgery will improve theatre  
3 utilisation (potentially allowing more cases to be added to a list) as well as  
4 reducing the anaesthetic time for each patient. In the era of axillary conservation,  
5 molecular medicine and dwindling resources, ~~which provides useful information~~  
6 ~~that can influence treatment decisions. In the era of axillary conservation, the~~  
7 combination of grey-scale axillary ultrasound and a microbubble/ CEUS SLN core  
8 biopsy has the potential to succeed surgical staging of the axilla. Further work  
9 now needs to be undertaken to refine the procedure with protocols, standard  
10 setting and training.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Veronesi U, Paganelli G, Viale G, et al. A randomised comparison of sentinel-node biopsy with routine axillary dissection in breast cancer. *N Engl J Med* 2003;349:546-553.
2. CG80 Early and locally advanced breast cancer: full guideline. United Kingdom, National Institute for Health and Clinical Excellence, February 2009. (Accessed July 2016 at <https://www.nice.org.uk/guidance/cg80/resources/early-and-locally-advanced-breast-cancer-diagnosis-and-treatment-975682170565>).
3. Mansel RE, Fallowfield L, Kissin M, Goyal A, Newcombe RG, Dixon J M et al. Randomized Multicenter Trial of Sentinel Node Biopsy Versus Standard Axillary Treatment in Operable Breast Cancer: The Almanac Trial. *J Nat Can Inst* 2006; 98: 599-609.
4. Pesek S, Ashikaga T, Krag L E, Krag D. The false-negative rate of sentinel node biopsy in patients with breast cancer: a meta-analysis. *World J Surg* 2012; 36: 2239-2251.
5. Barthelmes L, Goyal A, Newcombe RG, McNeill F, Mansel RE. Adverse reactions to patent blue V dye – The NEW START and ALMANAC experience. *Eur J Surg Oncol* 2010; 36: 399-403.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
6. Giuliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PW, Blumencranz PW, et al. Axillary Dissection vs No Axillary Dissection in Women With Invasive Breast Cancer and Sentinel Node Metastasis. JAMA 2011; 305: 569-575.
7. Association of Breast Surgery Consensus Statement Management of the Malignant Axilla in Early Breast Cancer, March 2015. (Accessed June 2017 at [http://www.associationofbreastsurgery.org.uk/media/50934/axilla\\_abs\\_consensus\\_statement\\_16\\_3\\_15.pdf](http://www.associationofbreastsurgery.org.uk/media/50934/axilla_abs_consensus_statement_16_3_15.pdf)).
8. Omoto K, Matsunaga H, Take N, Hozumi Y, Takehara M, Omoto Y et al. Sentinel Node detection method using contrast-enhanced ultrasonography with sonazoid in breast cancer: preliminary clinical study. Ultrasound Med Biol 2009; 35: 1249-1256.
9. Sever A, Jones S, Cox K, Weeks J, Mills P, Jones P. Preoperative localization of sentinel lymph nodes using intradermal microbubbles and contrast-enhanced ultrasonography in patients with breast cancer. Br J Surg 2009; 96:1295-1299.
10. Rautiainen S, Sudah M, Joukainen S, Sironen R, Vanninen R and Sutela A. Contrast-enhanced ultrasound-guided axillary lymph node core biopsy: Diagnostic accuracy in preoperative staging of invasive breast cancer. Eur J Radiol 2015; 84: 2130-2136.
11. Goldberg BB, Merton DA, Liu JB et al. Sentinel lymph nodes in a swine model with melanoma: contrast-enhanced lymphatic US. Radiology 2004;230:324-330.

12. Sever A, Broillet A, Schneider M, Cox K, Jones S et al. Dynamic visualisation of lymphatic channels and sentinel lymph nodes using intradermal microbubbles and contrast enhanced ultrasound in a swine model and patients with breast cancer. *J Ultrasound Med* 2010; 29: 1699-704.
13. Cox K, Sever A, Jones S, Weeks J, Mills P, Devalia H et al. Validation of a technique using microbubbles and contrast enhanced ultrasound (CEUS) to biopsy sentinel lymph nodes (SLN) in pre-operative breast cancer patients with a normal grey-scale axillary ultrasound. *Eur J Surg Oncol* 2013; 39: 760-765.
14. Cox K, Weeks J, Mills P, Chalmers R, Devalia H, Fish D, Sever A. Contrast-Enhanced Ultrasound Biopsy of Sentinel Lymph Nodes in Patients with Breast Cancer: Implications for Axillary Metastases and Conservation. *Ann Surg Oncol* 2016; 23: 58-64.
15. Kilbride KE, Lee MC, Nees AV, Cimmino VM, Diehl KM, Sabel MS, Hayes DF, Schott AF, Klee CG, Chang AE, Newman LA. Axillary staging prior to neoadjuvant chemotherapy for breast cancer: predictors of recurrence. *Ann Surg Oncol* 2008; 15: 3252-3258.
16. Su YL, Li SH, Chen YY, Chen HC, Tay Y, Huang CH, Chou FF, Wu SC, Rau KM. Post-mastectomy radiotherapy benefits subgroups of breast cancer patients with T1-2 tumor and 1-3 axillary lymph node(s) metastasis. *Radiol Oncol* 2014; 48 (3): 314-322.
17. Yang WT, Metreweli C, Lam PK, Chang J. Benign and malignant breast masses and axillary nodes: evaluation with echo enhanced color power Doppler Ultrasound. *Radiology* 2001; 220:795-802.

18. Xie F, Zhang D, Cheng L, Yu L, Yang L, Tong F et al. Intradermal microbubbles and contrast-enhanced ultrasound (CEUS) is a feasible approach for sentinel lymph identification in early-stage breast cancer. *World J Surg Oncol* 2015; 13: 319.
19. Tanter M, Fink M. Ultrafast imaging in biomedical ultrasound. *IEEE Trans Ultrason Ferroelectr Freq Control* 2014; 61: 102-119.
20. Christensen-Jeffries K, Browning RJ, Tang M-X, Dunsby CW; Eckersley RJ, In Vivo Acoustic Super-Resolution and Super-Resolved Velocity Mapping Using Microbubbles. *IEEE TRANSACTIONS ON MEDICAL IMAGING* 2015; 34: 433-440
21. Gorce J-M, Arditi M, Schneider M. Influence of bubble size distribution on the echogenicity of ultrasound contrast agents. *Investigative Radiology* 2000; 35: 661-671.
22. [Diepstraten SC, Sever AR, Buckens CF, Veldhuis WB, van Dalen T, van den Bosch MA, Mali WP and Verkooijen HM. Value of preoperative ultrasound-guided axillary lymph node dissection in breast cancer: a systematic review and meta-analysis. \*Ann Surg Oncol\* 2014; 21: 51-59.](#)
23. [Britton P, Willsher P, Taylor K, Kilburn-Toppin F, Provenzano E, Forouhi P, Benson J, Agrawal A, Forman JR and Wallis MG. Microbubble detection and ultrasound-guided vacuum-assisted biopsy of axillary lymph nodes in patients with breast cancer. \*Clin Radiol\* 2017; 72: 772-779.](#)
24. [Donovan CA, Giuliano AE. Evolution of the Staging System in Breast Cancer. \*Ann Surg Oncol\* 2017; 24: 3469-3470.](#)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure Captions**

**Figure 1**

Diagram showing the selection and flow of participants from each centre through the study. Lymph node (LN), Sentinel Lymph Node (SLN), Contrast enhanced ultrasound (CEUS), Ductal Carcinoma in situ (DCIS),

1 isolated tumour cells (ITC), surgical sentinel lymph node biopsy (SLNB),  
2 axillary lymph node dissection (ALND) and POSNOC – (POsitive Sentinel  
3 NODe: adjuvant therapy alone versus adjuvant therapy plus Clearance or  
4 axillary radiotherapy. A randomised controlled trial of axillary treatment  
5 in women with early stage breast cancer who have metastases in one or  
6 two sentinel nodes).  
7  
8  
9  
10  
11  
12  
13  
14  
15

## 16 **Figure 2**

17  
18 A. Ultrasound contrast pulse sequencing image of a SLN (white arrow)  
19 after injection of intradermal microbubbles (between 0.2 and 1 ml  
20 injected using a 26G needle with 1ml tuberculin syringe) into the UOQ  
21 periareolar area of the breast. B. Grey scale ultrasound image of the same  
22 SLN as visualised in A. Images provided by Centre 3.  
23  
24  
25  
26  
27  
28  
29  
30  
31

## 32 **Figure 3**

33  
34 A. Volume of axillary disease at the end of primary surgical treatment for  
35 individual patients at Centres 1 and 2 who had a false negative (FN) or  
36 true positive (TP) microbubble/ CEUS core biopsy of SLN. Axillary lymph  
37 node dissection (ALND) and sentinel lymph node excision (SLNE). B. The  
38 total volume of axillary metastases at the end of surgical treatment was  
39 determined using a scoring system (isolated tumour cells = 0.2, each LN  
40 micrometastasis = 0.5 and each LN macrometastasis = 1). The Mann-  
41 Whitney test was used to compare the volume of axillary metastases  
42 between patients with a false negative and true positive microbubble/  
43 CEUS SLN biopsy in both centres.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1                   **Table 1**

2  
3                   Performance statistics of 7 consultant breast radiologists at Centre 1  
4  
5                   (Total 1349 of 1361 procedures). In 12 cases the data was incomplete and  
6  
7                   the name of the radiologist was not recorded. For radiologist no.1, one  
8  
9                   successful procedure was a fine needle aspiration biopsy rather than a  
10  
11                   core biopsy.  
12  
13  
14  
15

16  
17  
18                   **Table 2**

19  
20                   Test accuracy of CEUS guided SLN biopsy using intradermally injected  
21  
22                   microbubbles in women with normal (centres 1 and 2) or indeterminate  
23  
24                   results from previous grey scale ultrasound (with or without previous  
25  
26                   biopsy). Reference standard is Sentinel Lymph node excision (SLNE) or  
27  
28                   axillary lymph node dissection (ALND). Positive predictive value (PPV)  
29  
30                   and negative predictive value (NPV) were calculated directly for centres 1  
31  
32                   and 2 who employed consecutive recruitment, and using Bayesian  
33  
34                   methods with 22% prevalence at centres 3 and 4 as these were not a  
35  
36                   consecutively or randomly selected group. True positive (TP), false  
37  
38                   positive (FP), true negative (TN) and false negative (FN).  
39  
40  
41  
42  
43  
44  
45  
46  
47

48                   **Table 3**

49  
50                   Age and clinicopathological characteristics of all patients at Centre 1 (first  
51  
52                   column) with a successful microbubble/CEUS SLN core biopsy before  
53  
54                   primary surgical treatment and Centre 1 patients with false negative  
55  
56                   microbubble/ CEUS SLN core biopsies sub-divided into micrometastases  
57  
58  
59  
60  
61  
62  
63  
64  
65

( $<2\text{mm}$ )/ isolated tumour cells (ITC), low volume metastases and high volume metastases identified at the end of primary surgical treatment. Data are expressed as  $n$  (%). Estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (Her2), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC).

**Table 4**

Age and clinicopathological characteristics of all patients at Centre 2 (first column) with a successful microbubble/CEUS SLN core biopsy before primary surgical treatment and Centre 2 patients with false negative microbubble/ CEUS SLN core biopsies sub-divided into micrometastases ( $<2\text{mm}$ )/ isolated tumour cells (ITC), low volume metastases and high volume metastases identified at the end of primary surgical treatment. Data are expressed as  $n$  (%). Estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (Her2), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC).

Table 1

	Centre 1 – Consultant Breast Radiologists						
	1	2	3	4	5	6	7
Total Procedures	276	37	501	81	116	207	131
Procedures where core biopsy not attempted	0 (0%)	0 (0%)	0 (0%)	2 (2.5%)	0 (0%)	0 (0%)	1 (0.8%)
Procedures with successful visualisation of SLN	269 (97.5%)	33 (89.2%)	457 (91.2%)	59 (72.8%)	94 (81%)	187 (90.3%)	118 (90.1%)
Procedures without SLN tissue retrieved	1 (0.4%)	1 (2.7%)	64 (12.8%)	2 (2.5%)	10 (8.6%)	18 (8.7%)	34 (26%)
Successful retrieval of lymphoid tissue in those with visualised SLN	268 (99.6%)	32 (97%)	393 (86%)	57 (96.6%)	84 (89.4%)	169 (90.4%)	84 (71.2%)
Total successful visualisation and	268 (97.1%)	32	393 (78.4%)	57	84	169	84

core biopsy of SLN		(86.5%)		(70.4%)	(72.4%)	(81.6%)	(64.1%)
--------------------	--	---------	--	---------	---------	---------	---------

Table 2

Centre	Prevalence of LN metastases	TP	FP	TN	FN	Sensitivity	Specificity	PPV	NPV
1	22%	84	2	635	95	46.9% (39.4-55.5%)	99.7% (98.95-100%)	97.7% (91.9-99.7%)	87.0% (84.3-89.3%)
2	27%	31	3	153	28	52.5% (39.1-65.7%)	98.1% (94.5-99.6%)	91.2% (76.3-98.1%)	84.5% (78.4-89.5%)
3	35%	13	0	52	15	46.4% (27.5-66.1%)	100% (93.2-100%)	100% (75.3-100%)	86.9% (82.4-90.3%)
4	29%	5	1	26	6	45.5% (16.7-76.6%)	96.3% (81.0-99.9%)	77.6% (31.3-96.3%)	86.2% (78.4-91.5%)

Table 3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Centre 1	All Patients	Volume of disease at the end of surgical treatment (microbubble/ CEUS false negative core biopsy)		
		Micrometastasis/ ITC	Low (<2 LN macrometastases)	High (2 or more LN macrometastases)
Total number of patients	816	37	46	12
		15 (41%) ALND	34 (74%) ALND	12 (100%) ALND
Median age in years (range)	61 (30-94)	61 (42-89)	55 (32-90)	53 (36-69)
<b>Receptor Status</b>				
ER positive	707 (87%)	32 (86%)	40 (87%)	11 (92%)
ER unknown	4 (0.4%)	0	0	0
Her-2 positive	79 (10%)	3 (8%)	2 (4%)	1 (8%)
Her-2 not recorded	13 (1.6%)	0	2 (4%)	0
ER-/PR-/Her-2-	72 (10%)	4 (10.8%)	4 (9%)	0
ER-/PR-/HER-2+	28 (3%)	1 (2.7%)	1 (2%)	1 (8%)
<b>Invasive Tumour Size</b>				
DCIS + microinvasion	5 (0.6%)	0	0	0
T1	472 (58%)	19 (51%)	17 (37%)	4 (33%)
T2	199 (24%)	10 (27%)	18 (39%)	5 (42%)
T3	29 (4%)	5 (14%)	3 (7%)	1 (8%)
Multifocal	111 (14%)	3 (8%)	8 (17%)	2 (17%)
Unknown	0	0	0	0
<b>Tumour Grade</b>				
Grade 1	180 (22%)	6 (16%)	7 (15%)	4 (33%)
Grade 2	392 (48%)	20 (54%)	21 (46%)	6 (50%)
Grade 3	209 (26%)	11 (30%)	14 (30%)	1 (8%)
Mixed grade	29 (4%)	0	4 (9%)	1 (8%)

Unknown	1 (0.1%)	0	0	0
<b>Tumour Type</b>				
IDC	660 (81%)	23 (62%)	38 (83%)	11 (92%)
ILC	103 (13%)	11 (30%)	6 (13%)	1 (8%)
Other	29 (4%)	1 (3%)	1 (2%)	0
Mixed	17 (2%)	2 (5%)	1 (2%)	0
Unknown	2 (0.2%)	0	0	0

Table 4

		Volume of disease at the end of surgical treatment (microbubble/ CEUS false negative core biopsy)		
Centre 2	All Patients	Micrometastasis/ ITC	Low (<2 LN macrometastases)	High (2 or more LN macrometastases)
Total number of patients	215	10	11	7
		2 (20%) ALND	10 (91%) ALND	7 (100%) ALND
Median age in years (range)	64 (31-93)	63.5 (37-93)	62 (38-91)	55 (47-72)
<b>Receptor Status</b>				
ER positive	176 (82%)	10 (100%)	8 (73%)	6 (86%)
ER unknown	6 (3%)	0	1 (9%)	1 (14%)
Her-2 positive	29 (13%)	1 (10%)	0	1 (14%)
Her-2 not recorded	8 (4%)	0	1 (9%)	1 (14%)
ER-/PR-/Her-2-	23 (11%)	0	1 (9%)	0
ER-/PR-/HER-2+	10 (5%)	0	0	0
<b>Invasive Tumour Size</b>				
DCIS + microinvasion	0	0	0	0
T1	118 (55%)	2 (20%)	6 (55%)	2 (29%)
T2	71 (33%)	6 (60%)	4 (36%)	5 (71%)
T3	9 (4%)	0	0	0
Multifocal	17 (8%)	2 (20%)	0	0
Unknown	0	0	1 (9%)	0
<b>Tumour Grade</b>				
Grade 1	35(16%)	1 (10%)	0	2 (29%)
Grade 2	102 (47%)	6 (60%)	5 (45.5%)	1 (14%)
Grade 3	68 (32%)	2 (20%)	4 (36%)	4 (57%)



1	Mixed grade	4 (2%)	0	0	0
2	Unknown	6 (3%)	0	2 (18%)	0
3	<b>Tumour Type</b>				
4	IDC	160 (74%)	7 (70%)	8 (73%)	5 (71%)
5	ILC	24 (11%)	2 (20%)	1 (9%)	0
6	Other	16 (7%)	0	0	0
7	Mixed	13 (6%)	1 (10%)	1 (9%)	2 (29%)
8	Unknown	2 (1%)	0	1 (9%)	0
9					
10					
11					
12					
13					
14					
15					
16					
17					
18					
19					
20					
21					
22					
23					
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
51					
52					
53					
54					
55					
56					
57					
58					
59					
60					
61					
62					
63					
64					
65					

Figure 1




Figure 2

Figure 2






Figure 3

A



B

Centre	no. of true positives	Median LN score (IQR)	no. of false negatives	Median LN score (IQR)	P-value
1	84	1.75 (1 – 3)	95	1 (0.5 – 1)	<0.0005
2	31	1.5 (1 – 2)	28	1 (0.5 – 2)	0.01