

1 **Title**

2 **Pathogenesis of HIV-1 and *Mycobacterium tuberculosis* co-infection**

3 **Authors**

4 Lucy CK Bell and Mahdad Noursadeghi

5 **Author affiliations**

6 Division of Infection & Immunity, Cruciform Building, University College London, WC1E 6BT, London, UK

7 Correspondence to M.N. m.noursadeghi@ucl.ac.uk

8 Glossary

| Term | Definition |
|-----------------------------------|--|
| Acid-fast bacilli (AFB) | Bacteria which are resistant to decolourisation during laboratory staining procedures, which is a recognised property of <i>Mycobacteria</i> . This arises due to the high mycolic acid content of the bacterial cell wall. Several diagnostic tests for TB rely on this property, including Ziehl-Neelsen staining. |
| Bacillary load | The measurable quantity of bacteria within a host organism or sample. |
| Efferocytosis | The process by which dead or dying cells are cleared by phagocytosis. |
| Extra-pulmonary | Anatomical locations beyond the thoracic cavity or lung. |
| Granulomatous pathology | Chronic inflammatory foci within tissues, primarily made up of a core of activated macrophages surrounded by CD4 ⁺ T cells. |
| Giant cells | Multinucleated cells derived from macrophages, typically found within granulomatous inflammation. |
| HIV-1 long terminal repeat | Repetitive non-coding sequences at each end of the HIV-1 proviral DNA, which are formed during reverse transcription and play important roles in integration and regulation of viral gene expression. |
| Immunoregulation | Mechanisms by which the immune system self-regulates via negative feedback loops, e.g. the production of immunosuppressive cytokines. |
| Independent risk factor | A variable that improves the prediction of outcome in a statistical model which already includes other variables. |
| Inflammasome | Multimeric molecular complexes formed during innate immune signalling which activate caspase enzymes, control maturation of the pro-inflammatory cytokines IL-1 β and IL-18, and may lead to cell death via pyroptosis. |
| Immunosenescence | The observable decline in immune function associated with ageing. |
| Immunodominant | The antigenic epitopes most commonly targeted by the adaptive immune response. |
| Lung apices | The upper lobe of each lung. |
| Mycobacteraemia | Circulation of mycobacteria in the bloodstream, identified on culture of blood. |
| Necrotic granuloma | Granulomatous inflammation with a core of dead cells. |
| Phagosome | A cytoplasmic vesicle formed as a result of the cellular uptake of particles >0.75 μ m in diameter. |
| Pleural effusion | An accumulation of fluid in the pleural cavity, the anatomical compartment which surrounds the lungs. This can arise due to a range of causes, one of which is infections such as TB. |

| | |
|-----------------------------------|---|
| Pro-inflammatory cytokines | Extracellular signalling molecules secreted chiefly by immune cells, which specific cell-surface receptors to trigger inflammatory processes. |
| Pulmonary cavitation | Formation of large airspaces in the lung parenchyma due to tissue destruction. |
| Pyroptosis | A specific cell death pathway triggered activation of caspase 1. |
| Resting T cells | T cells which have not been activated by binding of their cognate antigen to the T cell receptor or stimulation by mitogens. |
| Quasispecies | A genetically heterogenous population arising from a process of mutation and selection. |
| Sentinel cell | Tissue resident cell that initiates a host immune response. |
| Serodiscordant couples | A sexual partnership in which one partner is HIV-1 infected and the other is not. |
| Sympatric speciation | The evolutionary process by which one species adapts to another with which it overlaps geographically. |
| Tuberculin skin test | Intradermal injection of a standardised preparation of purified protein derivative of killed and homogenised <i>M. tuberculosis</i> . |
| Th17 cells | CD4 positive T helper cell subset that produces interleukin (IL) 17 on stimulation, which in turn has a canonical role in augmenting neutrophil responses to infection. |
| Viral rebound | Development of a detectable plasma viral load in a HIV-1 positive individual following a period of virological suppression, typically associated with an interruption in ART or the development of drug resistance. |

10 **Abstract**

11 Co-infection with *Mycobacterium tuberculosis* is the leading cause of death in HIV-1 infected individuals. It has
12 long been known that HIV-1 infection alters the course of *M. tuberculosis* infection and substantially increases
13 the risk of active tuberculosis (TB). It has also become clear that TB increases levels of HIV-1 replication,
14 propagation and genetic diversity. Therefore, co-infection provides reciprocal advantages to both pathogens.
15 In this Review, we describe the epidemiological associations between the two pathogens, selected interactions
16 of each pathogen with the host and our current understanding of how they affect the pathogenesis of TB and
17 HIV-1/AIDS in co-infected individuals. We evaluate the mechanisms and consequences of HIV-1 depletion of
18 T cells on immune responses to *M. tuberculosis*. We also discuss the effect of HIV-1 infection on the control
19 of *M. tuberculosis* by macrophages through phagocytosis, autophagy and cell death, and we propose models
20 by which dysregulated inflammatory responses drive the pathogenesis of TB and HIV-1/AIDS.

21 **Key points**

- 22 – There were 1.14 million new cases of HIV-1/TB co-infection and 400,000 deaths that were attributed to
23 co-infection in 2015.
- 24 – The risk of TB increases by 2-5 fold in early HIV-1 infection and by more than 20-fold in advanced HIV-1
25 disease. Approximately 4-fold increased risk of TB persists in HIV-1 infected patients treated with
26 antiretroviral therapy.
- 27 – HIV-1 infects CD4⁺ T cells and macrophages. *Mycobacterium tuberculosis* primarily infects macrophages,
28 which require CD4⁺ T cells to augment intracellular clearance of microbial pathogens. Hence the depletion
29 of CD4⁺ T cells that is associated with HIV-1 infection is thought to have a major role in the increased risk
30 of TB in HIV-1 infected people.
- 31 – Co-infection of HIV-1 and *M. tuberculosis* at the level of individual macrophages may also occur, but has
32 not been demonstrated in vivo. This is important because experimental models show that HIV-1 infection
33 of macrophages can attenuate phagocytosis and intracellular killing by the autophagy pathway.
- 34 – Progressive HIV-1 disease and TB are both characterised by chronic inflammation driven by the failure to
35 clear either pathogen. The chronic nature of these responses may undermine host protection by promoting
36 an immunoregulatory phenotype that is characterised by attenuated T cell responses.
- 37 – Advanced HIV-1 infection is associated with reduced immunopathology of TB co-infection, but the
38 introduction of antiretroviral therapy can exacerbate the immunopathology of TB, giving rise to immune
39 reconstitution inflammatory syndrome (IRIS). This reflects recovery of innate immune inflammatory
40 responses to *M. tuberculosis*, which may be exacerbated by the recirculation of *M. tuberculosis* reactive
41 T cells and failure of the normal homeostatic control of inflammatory responses.
- 42 – The proinflammatory response to *M. tuberculosis* may exacerbate HIV-1/AIDS disease progression by
43 increasing virus propagation through increased transcription and cell-cell transmission.

44

45 Introduction

46 The retrovirus HIV-1, which causes Acquired Immunodeficiency Syndrome (AIDS), transmitted to humans from
47 primates in the 20th century¹. The causative agent of tuberculosis (TB), *Mycobacterium tuberculosis*, is an
48 unencapsulated acid-fast bacillus which has been a pathogen of humans for millennia². HIV/AIDS and TB are
49 each among the 10 leading causes of death world-wide. The interaction between these two pathogens
50 substantially contributes to this high incidence of mortality. Estimates for the incidence of HIV-1 and *M.*
51 *tuberculosis* infection in 2015 highlights the scale of these epidemics (**Table 1**)^{3,4}. The geographical
52 convergence of the HIV/TB 'syndemics' in Africa and in Eastern Europe, as well as the demographic
53 convergence in particular at-risk groups, such as prisoners⁵ and miners in Southern Africa⁶, further exacerbate
54 the burden of co-infection morbidity and mortality. The impact of TB is also increasing with the emergence of
55 multi-drug resistant (MDR)-TB (**Table 1**), for which HIV-1 is an independent risk factor⁷. Moreover, the rates
56 of undiagnosed TB in HIV-1 positive individuals as revealed by post-mortem studies⁸, suggest that the disease
57 burden that is associated with co-infection has been underestimated.

58 An increased risk of TB throughout the course of HIV-1 disease has been established through epidemiological
59 studies^{9,10} (**Figure 1a**). HIV-1 co-infection also influences the clinical phenotype of TB. HIV-1 positive patients
60 with CD4⁺ T cell counts in the normal range present with classic symptoms of pulmonary TB, but disease that
61 is restricted to the lung apices is less frequent, whereas pleural effusions and lymph node disease are more
62 likely¹¹. In advanced AIDS, *M. tuberculosis* frequently causes disseminated extra-pulmonary disease and
63 mycobacteraemia^{12,13}.

64 TB may also exacerbate HIV-1 disease and AIDS progression (**Figure 1b**). Active TB is associated with higher
65 HIV-1 viral loads in the blood¹⁴ and cerebrospinal fluid (CSF)¹⁵, and an increased genetic heterogeneity of the
66 viral quasispecies¹⁶. This is likely to be driven by increases in HIV-1 replication at the site of co-infection^{17,18}.
67 In addition, *M. tuberculosis* co-infection may contribute to higher levels of systemic immune activation that is
68 associated with HIV-1 disease progression^{19,20}, even in the context of latent TB infection (LTBI)²¹. The
69 importance of these phenotypes in driving HIV-1 disease progression is unknown, but based on our
70 understanding of how high viral loads²² and immune activation²³ can drive immunosuppression, co-infection
71 with *M. tuberculosis* may accelerate AIDS progression. Accordingly, TB has been associated with an increased
72 incidence of additional opportunistic infections²⁴, and may be an independent risk factor for progression to
73 AIDS²⁵. This hypothesis is supported by the observation that an episode of successfully treated TB is
74 associated with a four-fold increase in all-cause mortality associated with HIV HIV²⁶.

75 Understanding the mechanisms that underpin HIV-1 and TB co-infection is paramount to identifying the best
76 approaches to overcome the global burden of disease that is caused by these pathogens. In this Review, we
77 describe the specific host-pathogen interactions which have emerged as common features of both organisms.
78 In particular, their ability to infect macrophages, induce type 1 interferon (IFN) responses and the role of chronic
79 or dysregulated inflammation in the pathogenesis of disease. We then describe our current understanding of
80 the mechanisms by which each organism influences the pathogenesis of the co-infecting pathogen, and
81 consider the future challenges for research in this field.

82 HIV-1 infection

83 Cellular targets of HIV-1

84 CD4⁺ T cells are considered the primary target cells for HIV-1. Macrophages are also infected by HIV-1 and
85 may have an important role in HIV/AIDS pathogenesis. CD4⁺ T cells and macrophages are also thought to be
86 crucial for host defence against *M. tuberculosis*²⁷ (Figure 2). Given that macrophages are the primary
87 intracellular niche for *M. tuberculosis*, their permissivity to HIV-1 raises the possibility that the two pathogens
88 can co-infect individual cells. HIV-1 infection of monocyte-derived macrophages *in vitro* is well described
89 despite the fact that macrophages express SAMHD1, an endonuclease that is active in non-dividing cells.
90 SAMHD1 has dNTPase activity and therefore depletes the cell of nucleotides that are required for DNA
91 synthesis, thereby restricting HIV-1 reverse transcription²⁸. The simian immunodeficiency virus (SIV)
92 accessory protein Vpx (which is absent in HIV-1) degrades SAMHD1 and makes human myeloid cells more
93 permissive to retroviral infection. Interestingly, Vpx deficient SIV still infects myeloid cells in non-human
94 primates²⁹, suggesting that Vpx is not necessary for infection of myeloid cells *in vivo*. Nonetheless, the
95 observation that HIV-1 has not evolved to counteract SAMHD1 restriction had led to the hypothesis that HIV-1
96 infection of macrophages does not occur *in vivo*. However, sub-populations of human macrophages enter a
97 G1-like state in which SAMHD1 activity is downregulated, making the cells substantially more permissive to
98 HIV-1³⁰. These data provide a mechanism by which a retrovirus that cannot counteract SAMHD1 restriction is
99 able to infect non-dividing myeloid cells. A subset of human alveolar macrophages that are infected with HIV-
100 1 *in vivo*, have been identified by RNA fluorescence *in situ* hybridisation, suggesting that the virus in these
101 cells may be transcriptionally active³¹. Furthermore, sustained HIV-1 replication in the absence of T cells has
102 been demonstrated in non-human primates and in humanised mouse models^{32,33}.

103 Long-lived memory CD4⁺ T cells are thought to be the dominant cell type in which HIV-1 can establish latency
104 and evade clearance by antiretroviral therapy (ART). Proviral DNA can also be detected in alveolar
105 macrophages in patients on ART³⁴. Moreover, viral rebound following ART treatment interruption has been
106 described in humanized myeloid-only mice, indicating that macrophages may also act as a long term viral
107 reservoir³⁵. Importantly, the viral reservoir during ART is not completely latent, but exhibits low levels of HIV-1
108 replication³⁶. This may occur in 'sanctuary sites' where drugs do not reach sufficient concentrations. Lymphoid
109 tissue has been proposed as one such site^{37,38}. Macrophage populations within the central nervous system
110 may be another³⁹. Therefore, HIV-1 replication may still influence *M. tuberculosis* infection even in patients
111 receiving effective ART, particularly if this occurs within macrophages.

112 Induction and evasion of type 1 IFN responses

113 Type 1 IFNs represent the canonical innate immune response to viral infections⁴⁰. This response coincides
114 with the peak in viraemia that follows primary HIV-1 infection⁴¹. In various models, HIV-1 has been shown to
115 activate innate IFN responses through pattern recognition receptors (PRRs), including toll-like receptor (TLR)-
116 7⁴², RIG-I⁴³ and DDX3⁴⁴ RNA sensors, as well as cGAS⁴⁵, PQBP1⁴⁶ and IFI16⁴⁷ DNA sensors. This response
117 is thought to contribute to suppression of primary HIV-1 viraemia⁴⁸, albeit without achieving complete
118 suppression owing to the range of counter measures that are employed by viruses to overcome IFN-inducible
119 antiviral host proteins⁴⁹. Interestingly, HIV-1 infection and replication in macrophages, representing key tissue
120 resident innate immune sentinel cells, fails to induce innate IFN responses as a result of interactions between
121 the viral capsid and host proteins that shield the nascent DNA products of viral reverse transcription from host
122 DNA sensors⁵⁰. Macrophages successfully restrict HIV-1 in response to type 1 IFN⁵¹. Therefore, the ability of

123 the virus to evade innate immune detection in macrophages may have an important role in establishing
124 persistent HIV-1 infection.

125 **HIV-1 infection as a chronic inflammatory disease**

126 HIV-1 infection is increasingly considered a chronic inflammatory disease that leads to immunodeficiency.
127 Inflammatory markers are elevated throughout the asymptomatic phase of infection⁵² and correlate with the
128 rate of progression to AIDS²³. The same phenotype is not evident in non-pathogenic SIV infection of
129 non-human primates and in HIV-1 infected children who do not progress to immunodeficiency⁵³.

130 Several mechanisms are thought to lead to the chronic immune activation that is associated with HIV-1
131 infection⁵². Chief among these is the translocation of microbial products from the gastrointestinal lumen into
132 the bloodstream, following massive T cell depletion in gastrointestinal-associated lymphoid tissue (GALT)
133 during primary HIV-1 infection. This is thought to be caused by lytic infection of GALT Th17 cells⁵⁴, which are
134 particularly permissive to retroviral infection⁵⁵. Non-productive HIV-1 infection of resting T cells by cell-cell
135 spread in lymphoid tissue may also contribute to T cell depletion and immune activation. In this model,
136 incomplete DNA products of reverse transcription are recognised by the cytosolic DNA sensor IFI16, leading
137 to activation of the inflammasome, leading to cell death by pyroptosis and the release of proinflammatory
138 IL1 β ^{56,57}, thereby linking CD4⁺ T cell depletion and chronic inflammation^{58,59}. In contrast to this inflammatory
139 mechanism of T cell death, HIV-1 proviral DNA integration may activate DNA damage pathways that can cause
140 T cell apoptosis⁶⁰. Chronic immune activation may also lead to premature immune senescence or
141 compensatory immunoregulation^{61–63}. Research efforts have focussed on the role of chronic type 1 IFN
142 responses in persistent HIV-1 infection. In a non-human primate model of pathogenic HIV-1 infection, chronic
143 IFN stimulation caused IFN desensitisation and consequently, reduced expression of IFN-dependent antiviral
144 factors, leading to increased viral replication and further T cell depletion⁴⁸.

145 The strong link between HIV-1 infection and chronic inflammation raises the question of whether the virus
146 gains an advantage by driving this phenotype. One which, co-infection with *M. tuberculosis* may be expected
147 to compound as it also induces chronic inflammation. Inflammatory signalling may benefit HIV-1 by directly
148 stimulating viral replication (discussed below). In addition, the recruitment of leukocytes could potentially
149 provide HIV-1 with a source of target cells to infect. This may be particularly important as HIV-1 infects new
150 cells most efficiently by direct cell-cell transmission via a 'virological synapse' actively orchestrated by the
151 virus⁶⁴. Cell-cell transmission also shields the virus from neutralising antibodies⁶⁵, leads to more rapid viral
152 gene expression⁶⁶ and enables the virus to overcome cell tropism barriers, such as infection of macrophages
153 by non-macrophage tropic viruses⁶⁷.

154 [Au: Please insert a brief summary paragraph (2-3 sentences) to conclude this section and to refocus the
155 reader back to HIV/TB co-infection.]

156

157 **M. tuberculosis infection**

158 **Infection of macrophages**

159 *M. tuberculosis* is a facultative intracellular pathogen of macrophages. Once inside, *M. tuberculosis* quickly
160 adapts to the environment within the phagosome through transcriptional reprogramming in order to upregulate
161 iron scavenging mechanisms, switch to anaerobic respiratory pathways, and to use cholesterol as a carbon
162 source and aspartate as a nitrogen source⁶⁸. Within macrophages, *M. tuberculosis*-containing phagosomes

163 fail to undergo the normal process of maturation and acidification that is associated with phagolysosomal fusion
164 and by which phagosomal cargo is usually degraded. Multiple *M. tuberculosis* virulence factors are thought to
165 contribute to this phenotype⁶⁹.

166 Macrophage cell death is a key feature of granulomatous pathology in TB. Central to this process is the
167 bacterial ESX-1 secretion system and the secreted effector molecule ESAT6. These are encoded by the RD1
168 locus which is deleted in the live attenuated *Mycobacterium bovis*, bacille Calmette–Guérin (BCG) vaccine.
169 ESX-1 is required for *M. tuberculosis* to escape from phagosomes into the host cell cytoplasm, and for
170 triggering cell death pathways⁷⁰. ESAT6 can also trigger macrophage cell death through apoptosis⁷¹. In
171 addition, ESAT6 has been reported to activate the inflammasome⁷², suggesting that *M. tuberculosis* may cause
172 cell death through inflammasome/caspase 1-mediated pyroptosis. The triggering of cell death promotes
173 bacterial dissemination through efferocytosis⁷³ and by releasing bacteria into the extracellular space for onward
174 transmission to new hosts.

175 **Induction of innate immune type 1 IFN responses**

176 Mycobacterial lipids, lipoproteins and nucleic acids trigger a range of innate immune responses when they are
177 sensed by host PRRs within macrophages^{74,75}. These responses are thought to be crucial for further immune
178 cell recruitment and for the production of antimicrobial peptides⁷⁶. Innate immune responses to *M. tuberculosis*
179 include type 1 IFN responses that have conventionally been associated with antiviral responses⁷⁷. The
180 observation that type 1 IFNs are immunosuppressive in chronic viral infections has led to studies to determine
181 whether type 1 IFNs counteract immunoprotective IFN γ -dependent or IL1 β -dependent mechanisms of
182 *M. tuberculosis* clearance⁷⁸. Induction of type 1 IFN responses is principally mediated through the recognition
183 of *M. tuberculosis* nucleic acids by the cytosolic DNA sensor cGAS⁷⁵. This is dependent on ESX-1-mediated
184 *M. tuberculosis* phagosomal escape, which also drives inflammasome maturation of IL1 β ⁷⁹. Interestingly, the
185 levels of effector molecules (ESAT6) that are secreted by the ESX-1 system determines the outcome of host
186 cellular responses polarised towards either type 1 IFNs or IL1 β). *M. tuberculosis* strains that are more virulent
187 have been found to produce more ESAT6 and more type 1 IFNs⁸⁰.

188 **Chronic inflammation to promote transmission**

189 Similar to HIV-1 infection, chronic inflammation is the hallmark of TB pathology. Immunopathogenesis may
190 even be more important in this case because, unlike HIV-1, *M. tuberculosis* is an obligate pathogen. Its ability
191 to escape the intracellular niche, cause pulmonary cavitation and induce coughing through chronic
192 inflammation within airways, are necessary for its dispersal between individuals. Matrix metalloproteinase
193 (MMP)-1 has a crucial role in pulmonary cavitation that is associated with *M. tuberculosis* infection⁸¹. This
194 protein belongs to a family of host proteinases that degrade the extracellular matrix. MMPs are produced by
195 macrophages, epithelial cells and fibroblasts in response to pro-inflammatory cytokines. In this context, chronic
196 pro-inflammatory T cells may contribute to the pathology in response to persistent *M. tuberculosis*. The
197 observation that virulent *M. tuberculosis* strains have highly conserved immunodominant T cell epitopes
198 suggests that *M. tuberculosis* does not rely on antigenic variation to evade protective immunity^{82,83}. In addition,
199 it raises the possibility that the conservation of these immunodominant responses may be beneficial to the
200 pathogen. Thereby, *M. tuberculosis* may commandeer T cell responses to promote immunopathology and
201 consequently, its transmission.

HIV/TB co-infection

HIV-1 depletion of *M. tuberculosis* reactive T cells

Notwithstanding the hypothesis presented above that T cell responses in TB may contribute to pathogenesis of disease, they have long been thought to have an important role in immunological protection against *M. tuberculosis* by promoting intracellular bacterial killing or restriction (Figure 2). Genetic deficiencies in IL12 signalling (which is required for Th1 cell differentiation), or IFN γ signalling (representing the canonical product of Th1 responses), give rise to Mendelian susceptibility to mycobacterial disease (MSMD)⁸⁴. HIV-1 co-infection further highlights the importance of T cell mediated immunity. Substantially increased risk of TB and its extrapulmonary dissemination is strongly correlated with CD4⁺ T cell depletion in HIV-1 infected individuals. T cell depletion is evident in peripheral blood, in the respiratory tract and at the site of tuberculin skin test (TST) challenge⁸⁵⁻⁸⁷. Assuming that CD4⁺ T cell protection against *M. tuberculosis* is conferred by the pro-inflammatory cytokines that they produce, it is notable that the proportions of polyfunctional *M. tuberculosis* reactive T cells, which produce the pro-inflammatory cytokines IFN γ , tumour necrosis factor (TNF) and interleukin (IL)2 are also depleted in HIV-1 infected individuals⁸⁶. Hence, HIV-1 depletes T cell populations that are likely to be functionally important for protection against TB. Transcriptional profiling of biopsies taken from the site of the TST challenge in humans confirmed that T cell recruitment and IFN γ activity were both substantially reduced in HIV-1/TB co-infected patients, with blood CD4⁺ T cell counts of <200 per mL, indicative of advanced HIV-1 disease⁸⁵.

HIV-1 infected T cells may also contribute to the increased risk of TB in early HIV-1 disease before substantial depletion of peripheral blood CD4 counts⁸⁷. HIV-1 DNA was detected more frequently in *M. tuberculosis* specific T cells. These T cells produced high levels of IL2 which made them more permissive to HIV-1 infection. In comparison to the total memory T cell population or memory T cells that specifically recognise human cytomegalovirus, *M. tuberculosis* specific T cells were preferentially depleted in early HIV-1 infection. Taken together these data suggest that the depletion of these cells may be a direct result of HIV-1 infection. Transcriptional profiling of the TST challenge site biopsies in HIV-1/TB co-infected patients with blood CD4⁺ T cell counts >200 /mL also revealed less T cell recruitment at the site of the antigenic challenge, compared to HIV-1 negative patients with active TB⁸⁵. However, the functional significance of the reduced T cell recruitment observed in this study is currently unknown, as comparable levels of IFN γ inducible gene expression were found in the HIV-1 positive and negative groups. Hence, IFN γ activity as a surrogate of robust CD4 T cell responses to mycobacterial antigens was preserved in early HIV-1 disease, despite previous reports of preferential depletion of *M. tuberculosis* specific T cells. These data suggest that increased risk of TB in HIV-1 infected patients is not solely mediated by T cell depletion.

Of the other T cell populations that may contribute to HIV-1-associated TB, Th17 and Th22 cells are the most plausible candidates. A functional role for these T cell populations in immunological protection against TB is primarily based on data obtained from experiments in mice and by their role in the recruitment of phagocytic cells including macrophages⁸⁸⁻⁹⁰. The depletion of these T cell subtypes during primary HIV-1 infection^{54,91} may therefore contribute to differences in the immune response to *M. tuberculosis* in HIV-1 co-infected patients compared to HIV negative patients. Another T cell population that becomes depleted in HIV-1 infection are mucosal associated invariant T (MAIT) cells⁹². These are CD8⁺ innate lymphoid cells which recognise bacterial metabolites of vitamin B that are presented by a non-polymorphic MHC-like molecule, MR1⁹². MAIT cells are activated by *M. tuberculosis* and are enriched at the site of TB disease⁹³. Therefore, their depletion in HIV-1

243 infection may attenuate a component of host immune responses to *M. tuberculosis*. MAIT cells are not infected
244 by HIV-1. Their depletion is thought to be caused indirectly by immune activation. Importantly, by comparison
245 to the general population, the risk of active TB remains higher in HIV-1 infected patients even after becoming
246 established on effective ART⁹⁴. In this context, the failure of ART to restore the T cell repertoire, including MAIT
247 cells^{95,96}, may also be a significant factor in the persistently elevated risk of TB.

248 **HIV-1 inhibition of phagocytosis and autophagy in macrophages**

249 *M. tuberculosis* has evolved to survive and grow within macrophages. Nonetheless, *M. tuberculosis*
250 phagocytosis by macrophages is thought to restrict mycobacterial growth. The best evidence for this is comes
251 from experiments in the zebrafish *Mycobacterium marinum* model in which bacillary uptake by macrophages
252 is elegantly visualised. In this model, macrophage depletion, delayed macrophage recruitment or necrotic
253 macrophage cell death are all associated with increased microbial burden^{73,97,98}. Once the bacteria are
254 phagocytosed, phagolysosomal fusion that would lead to bacterial killing is inhibited by *M. tuberculosis*. This
255 may be overcome by the autophagy pathway^{99,100}, and by inducible production of bacteriocidal nitric oxide or
256 a range of antimicrobial peptides, all of which are generally upregulated by the action of IFN γ .

257 HIV-1 infection has been reported to inhibit macrophage phagocytosis dependent on diverse cell surface
258 receptors and mediated by HIV-1 infection of the affected cell¹⁰¹. In this study, the HIV-1 accessory protein Nef
259 was found to be both necessary and sufficient to inhibit phagocytosis by directly interacting with adapter protein
260 AP-1 to inhibit the recruitment of recycling endosomes that are required for phagosome biogenesis¹⁰¹. It is also
261 possible that these effects may be mediated indirectly by the action of circulating virus-free accessory proteins
262 on uninfected macrophages, which can be detected *in vivo*^{102,103}. Moreover, impaired phagocytosis has also
263 been observed in HIV-infected alveolar macrophages *ex vivo*³¹. Interestingly, HIV-1 Nef also inhibits the
264 autophagy pathway by blocking the maturation of autophagosomes through a direct interaction with the
265 autophagy regulator, Beclin-1¹⁰⁴. This inhibition was found to protect nascent virion assembly from autophagic
266 degradation. Consistent with these effects of HIV-1 infection on phagocytosis and autophagy (Figure 2), HIV-1
267 co-infection in macrophages that are infected with *M. tuberculosis* has been associated with increased
268 mycobacterial growth¹⁰⁵. Interestingly, however, vitamin D treatment of co-infected macrophages was reported
269 to restrict both *M. tuberculosis* and HIV-1 replication by an autophagy-dependent mechanism, suggesting that
270 the inhibition of autophagy by HIV-1 is easily overcome by the action of vitamin D¹⁰⁶. Vitamin D deficiency is
271 undoubtedly prevalent amongst populations at greatest risk of co-infection¹⁰⁷. Therefore, if HIV-1 inhibition of
272 autophagy is an important determinant for increased risk of TB, vitamin D supplementation may substantially
273 reduce the incidence of TB disease in HIV-1 infected patients while at the same time supporting immune
274 control of the virus. However, this hypothesis has yet to be tested in clinical trials.

275 **Macrophage cell death and tissue necrosis in HIV/TB co-infection**

276 HIV-1 does not cause macrophage cell death⁵¹. A number of reports suggest that HIV-1 proteins reduce
277 *M. tuberculosis*-associated macrophage apoptosis; potentially by HIV-1 Nef inhibition of TNF responses to
278 *M. tuberculosis*^{108–110}. Cellular apoptosis has been considered as a mechanism for limiting intracellular
279 *M. tuberculosis* growth. Therefore, by inhibiting apoptosis, HIV-1 infection may compromise *M. tuberculosis*
280 restriction. However, live cell imaging data has recently contradicted this hypothesis by demonstrating that cell
281 death is associated with *M. tuberculosis* growth rather than restriction¹¹¹. A key observation in HIV-1 infected
282 patients with pulmonary TB is the presence of fewer necrotic granuloma and less pulmonary cavitation.
283 Interestingly, in active TB MMP1 levels are significantly lower in respiratory tract samples from HIV-1 infected

284 patients with severe CD4⁺ T cell depletion, compared to HIV-1 negative patients¹¹². These observations have
285 largely supported the hypothesis that T cell responses to *M. tuberculosis* contribute substantially to cellular
286 necrosis and tissue damage. In agreement with this cellular necrosis and tissue damage are reduced in
287 patients with AIDS, but enhanced host cell viability in advanced HIV-1 disease does not achieve better
288 *M. tuberculosis* control. Consistent with the reduction in pulmonary cavitation, HIV-1 co-infected patients may
289 transmit less *M. tuberculosis*¹¹³. Nonetheless, at the population level, an increased incidence of active TB in
290 HIV-1 infected patients ultimately promotes the onward transmission of *M. tuberculosis*.

291 **Immunopathology of *M. tuberculosis* in HIV-1 infected patients**

292 The most direct evidence to support the hypothesis that there is a reduction of *M. tuberculosis*
293 immunopathology in patients with AIDS, is the phenomenon of TB immune reconstitution inflammatory
294 response syndrome (TB-IRIS). TB-IRIS is the development of increased inflammatory pathology in patients
295 following the commencement of ART, and may manifest by either a worsening of known TB disease, or
296 'unmasking' of previously asymptomatic *M. tuberculosis* infection¹¹⁴. TB-IRIS occurs in ~15% of HIV-1 infected
297 patients starting ART¹¹⁵ and is most commonly found in patients with very low peripheral blood CD4⁺ T cell
298 counts and evidence of a high *M. tuberculosis* bacillary load before starting ART. The pathological features of
299 TB-IRIS include systemic responses such as fever and increased acute neutrophilic inflammation at the site
300 of *M. tuberculosis* infection. Comparisons of peripheral blood transcriptional profiles in cohorts of HIV/TB
301 co-infected patients with and without TB-IRIS revealed that in cases of TB-IRIS, there were increases in the
302 expression of IFN, MyD88 and inflammasome-dependent innate immune responses¹¹⁶. These data suggest
303 that TB-IRIS may be caused by the recovery of innate immune responses to *M. tuberculosis*, and presumably
304 failure of immunoregulation that would ordinarily control pathogenic innate inflammatory responses. By
305 inference these data suggest that HIV-1 may downregulate innate immune and immunoregulatory responses
306 to *M. tuberculosis*, as well as classical Th1 responses. With the notable exception of type 1 IFNs, the wide
307 repertoire of pro-inflammatory transcriptional innate immune responses at the site of TST challenge, were
308 found to be lower in patients with active TB and advanced HIV-1 co-infection compared to HIV-1 negative
309 controls⁸⁵. By contrast, patients presenting with unmasking TB-IRIS had substantially higher pro-inflammatory
310 transcriptional responses to the TST compared to HIV-1 negative patients with active TB, consistent with
311 exaggerated inflammatory responses. In this study, unmasking TB-IRIS was associated with features that are
312 associated with Th2 responses and increased granulocyte colony stimulating factor (CSF3) expression that is
313 known to augment neutrophil responses⁸⁵.

314 How HIV-1 might inhibit innate immune responses to *M. tuberculosis* remains unclear. Both pathogens can
315 infect macrophages, which are widely recognised to generate pro-inflammatory innate immune responses.
316 Attenuated innate immune responses to prototypic stimuli such as lipopolysaccharide (LPS) by alveolar
317 macrophages from HIV-infected individuals has been reported^{117,118}. The HIV-1 accessory proteins Nef, Vpu
318 and Vpr have each been reported to inhibit innate immune intracellular signalling pathways¹¹⁹. Consistent with
319 these reports, HIV-1 infection of macrophages attenuated activation of the canonical NFκB pathway in
320 response to LPS¹²⁰. However, genome-wide transcriptional responses to LPS were largely preserved. In the
321 same experimental model of macrophages that were infected with HIV-1, pro-inflammatory innate immune
322 responses to *M. tuberculosis* co-infection were also preserved¹²¹⁻¹²³. Instead, HIV-1 infection was associated
323 with attenuated immunoregulatory IL10 expression, leading to exaggerated pro-inflammatory responses at
324 subsequent time points¹²¹. These data suggest that any HIV-1 associated inhibition of innate immune

325 pro-inflammatory responses to *M. tuberculosis* by HIV-1 does not arise because of co-infection at the cellular
326 level.

327 Indirect effects on macrophage innate immune responses may also result from HIV-1 modulation of T cells,
328 for example, by alterations in the cytokine milieu that acts on uninfected macrophages. Therefore, an
329 alternative hypothesis may be that pathogenic innate immune responses to *M. tuberculosis* are amplified by
330 T helper cells. In such a model, the immunopathogenesis of TB-IRIS may be driven by the recirculation of
331 *M. tuberculosis* reactive T cells, leading to the recovery of innate immune inflammatory responses and
332 compounded by high bacterial loads which have accumulated in the immunosuppressed patient providing a
333 higher dose for stimulation of innate immune responses¹¹⁴. In this context, HIV-1 attenuation of IL10 responses
334 in macrophages may represent foci of deficient immunoregulation that leads to pathological inflammation. In
335 this regard, we found that deficient IL-10 responses persist in infected macrophages in the presence of
336 antiretrovirals in vitro¹²¹, and others have demonstrated the persistence of dysregulated phagocyte phenotypes
337 after antiretroviral treatment, such as increased TLR-2 expression¹²⁴ and a dysregulation of complement
338 pathways¹²⁵. Hence, TB-IRIS may be caused by a combination of high bacterial burden, T cell recovery and
339 failure of immune regulation in HIV-1 infected macrophages (Figure 3). Interestingly, TST challenge
340 experiments revealed attenuated IL10 responses in HIV-1 infected patients with CD4⁺ T cells >200 /mL⁸⁵.
341 Therefore, an increased risk of active TB in early HIV-1 disease may also partly reflect a propensity for
342 immunopathology as a result of inadequate IL10 regulation.

343 **Effects of TB on HIV-1 replication**

344 Increased HIV-1 viral loads in the lungs of co-infected patients with pulmonary TB is well established^{17,18,126}.
345 This is commonly associated with increased viral load in peripheral blood¹⁴. Whether the increase in circulating
346 virus arises from replication in the lung alone or is also due to increased systemic virus replication is not known.
347 As most HIV-1 replication occurs in activated T cells, their recruitment to sites of granulomatous inflammation
348 in TB may facilitate rapid virus propagation through the accumulation of tightly packed permissive cells and
349 hence cell-cell transmission. However, the host immune response to *M. tuberculosis* also increases HIV-1
350 transcription. The HIV-1 long terminal repeat (LTR) includes binding sites for several host transcription factors
351 which are activated by innate immune and cytokine signalling pathways. These include the NFκB, AP1,
352 CCAAT/enhancer binding protein (C/EBP), CREB/ATF and NFAT families of transcription factors¹²⁷. Innate
353 immune activation by *M. tuberculosis* or mycobacterial products increased HIV-1 transcription and replication
354 in myeloid cell lines, through the action of C/EBP, NFκB and NFAT5¹²⁸⁻¹³⁰.

355 Experimental data on the effects of *M. tuberculosis* infection on HIV-1 transcription in macrophages are
356 inconclusive. In macrophages that were infected with HIV-1, co-infection with BCG caused a dose dependent
357 suppression of virus production. This was attributed to the C/EBP binding motif in the viral LTR and associated
358 with production of a type 1 IFN inducible inhibitory isoform of C/EBPβ, leading to a model in which
359 mycobacterial induction of type 1 IFNs may restrict HIV-1 transcription¹³¹. These data are somewhat
360 inconsistent with the current view of ESX-1-dependent induction of type 1 IFN responses by *M. tuberculosis*⁷⁹
361 given that BCG lacks ESX-1¹³². Interestingly, alveolar macrophages from healthy lung tissue express high
362 levels of inhibitory C/EBPβ, but this is strongly downregulated in cells that are isolated from the site of
363 pulmonary TB¹³¹, suggesting that the pro-inflammatory milieu that are present in active TB granuloma
364 overcomes type 1 IFN-mediated inhibition of virus transcription. Direct evidence for this hypothesis was shown
365 in HIV-1 infected macrophages where co-infection with *M. tuberculosis* led to an initial decrease in HIV-1

366 transcription followed by a substantial increase¹²¹. The increase in virus transcription was co-incident with
367 sustained pro-inflammatory responses as a result of HIV-1 attenuation of early IL10 regulatory responses, and
368 complementation of deficient IL10 responses reversed the increase in viral transcription in *M. tuberculosis*
369 co-infected macrophages¹²¹. IL10 has been reported to inhibit HIV-1 transcription by STAT3-dependent
370 induction of inhibitory C/EBP β and by inhibition of cyclin-T1 that is required for HIV-1 Tat-dependent
371 transactivation of viral transcription^{133–135}. Conversely, it has long been established that some canonical
372 pro-inflammatory cytokines, for example, TNF, IL6 and IL1 β , individually or synergistically upregulate HIV-1
373 transcription^{136–138}. Notably, IFN γ (which is enriched within TB granuloma) shows substantial overlap with the
374 antiviral effects of type 1 IFNs⁵¹. Despite this, IFN γ may still act synergistically with TNF to promote virus
375 transcription¹³⁹.

376 Increased HIV-1 replication in macrophages as a result of pro-inflammatory responses to *M. tuberculosis* may
377 also be accompanied by increased macrophage permissivity to nascent HIV-1 infection. Macrophages had
378 previously been considered as terminally differentiated cells that are unable to replicate. Mouse experiments
379 have demonstrated macrophage replication in response to inflammatory stimuli, for example modelled by
380 helminth infection¹⁴⁰ and for maintenance of embryologically derived resident populations^{141–143}. Recent
381 findings show that polyploid giant cells within granuloma arise from macrophages that enter cell cycle but do
382 not complete cytokinesis¹⁴⁴. In addition, macrophages that enter the cell cycle but arrest in a G1-like state,
383 downregulate SAMHD1 activity to allow DNA synthesis, and consequently become more permissive to HIV-1
384 by facilitating reverse transcription³⁰. Taken together, the pro-inflammatory cytokine response to *M.*
385 *tuberculosis* (facilitated by HIV-1 attenuation of immune regulation), recruitment and activation of T cells, and
386 modulation of macrophage cell cycle phenotype act in concert to enhance virus replication and propagation
387 (Figure 4). Hence these co-infecting pathogens successfully cooperate to usurp host defences to their own
388 advantage.

389 **Future perspectives**

390 In resource-poor settings where the highest incidence of HIV-1 and *M. tuberculosis* co-infection is found, ART
391 has substantially reduced the incidence of co-infection and improved clinical outcomes¹⁴⁵. In the pre-ART era,
392 the majority of co-infection was evident in people with advanced HIV-1 disease. Assuming that ART
393 programmes continue to grow in these settings, the majority of morbidity and mortality that is associated with
394 HIV/TB co-infection may be caused by the increased risk of TB in patients with early HIV-1 infection prior to
395 ART initiation; TB-IRIS arising in patients starting ART; and the residual increased risk of active TB in patients
396 on ART. Therefore, ongoing research that focusses on the mechanisms by which the two pathogens interact
397 in these circumstances, and identification of possible therapeutic targets, is necessary. Specific priorities
398 include being able to (1) stratify the risks of active TB in order to inform optimal use of systematic screening
399 for active TB or strategies for preventative therapy; (2) to identify opportunities for host directed therapies that
400 treat or reduce the risk of TB-IRIS, or population level interventions to reduce the risk of active TB, such as
401 vitamin D supplementation; and (3) to evaluate the effects of active TB or LTBI on subclinical viral replication
402 and diversification that may promote HIV-1 drug resistance and persistence.

403 The application of whole-genome sequencing and single genome amplification¹⁴⁶ will offer greater depths of
404 resolution to explore the effect of *M. tuberculosis* co-infection on HIV-1 diversity. Likewise, whole-genome
405 sequencing of *M. tuberculosis* is expected to offer more insight into transmission chains¹⁴⁷ in order to assess

406 the impact of HIV-1 on the spread of *M. tuberculosis*, and whether HIV-1 influences sympatric speciation of
407 *M. tuberculosis*¹⁴⁸. Interestingly, a recent whole-genome sequencing study of *M. tuberculosis* strains in HIV-1
408 infected and uninfected individuals suggested virus-induced changes to bacterial evolution that disrupt the
409 unusually high degree of epitope conservation in *M. tuberculosis*¹⁴⁹.

410 HIV-1, and for the most part *M. tuberculosis*, are exclusively human pathogens, often rendering conventional
411 small models of disease misrepresentative. The recent developments in humanised mouse models for HIV-1
412 may also represent a significant opportunity to study co-infection. Likewise, the use of non-human primates in
413 which the combination of blood sampling and functional imaging offers the opportunity for detailed and
414 well-controlled longitudinal experiments to study co-infection. The prospect of human TB challenge models
415 are also emerging¹⁵⁰. In addition to the primary goal of developing tools to evaluate *M. tuberculosis* vaccine
416 efficacy, these would offer unprecedented opportunities to identify correlates of protection and to study
417 pathogenesis in co-infected individuals, particularly in the context of ART. These models could enable the
418 assessment of variables other than T cell depletion during co-infection, and explore the effects of chronic
419 inflammation, ageing, smoking, obesity and diabetes. Ultimately, the study of HIV/TB co-infection will also be
420 crucial in the development of effective vaccines for these important pathogens, based on our understanding
421 of how they undermine host defences through their interactions.

422 **Acknowledgements**

423 LCKB was funded by a Medical Research Council Doctoral Training Award through the University College
424 London MB PhD programme. MN is supported by a Wellcome Trust Investigator Award and National Institute
425 of Health Research Biomedical Research Centre Funding to University College Hospitals NHS Foundation
426 Trust and University College London.

427 **Competing interests statement**

428 The authors declare no competing interests.

429 **Publisher's note**

430 Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional
431 affiliations.

432 **Author biographies**

433 Lucy C. K. Bell completed medical training at the University of Cambridge and University College London
434 (UCL), during which she completed a PhD, studying the impact of HIV-1 co-infection on immune responses to
435 TB. She is now an academic clinical trainee at Guys' & St Thomas' Hospitals and King's College London, and
436 maintains active research interests in HIV-1, TB and transcriptomics.

437 Mahdad Noursadeghi is a Clinician Scientist and leads a research group in Infection and Immunity at University
438 College London, focusing on HIV-1 and Tuberculosis. They model innate immune host-pathogen interactions
439 in human macrophages, and use molecular profiling of challenge experiments in humans and sampling of
440 tissues at the site of disease in order to understand host-pathogen interactions in vivo.

441 Table of contents blurb

Text

442 Co-infection with *Mycobacterium tuberculosis* is the leading cause of death in HIV-1 infected individuals. In
443 this Review, Bell and Noursadeghi describe the epidemiological associations between the two pathogens,
444 selected interactions of each pathogen with the host and our current understanding of how they affect the
445 pathogenesis of tuberculosis and HIV-1/AIDS in co-infected individuals.

446 **Subject categories**

447 Biological sciences / Immunology / Infectious diseases / HIV infections

448 [URI /631/250/255/1901]

449 Biological sciences / Immunology / Infectious diseases / Tuberculosis

450 [URI /631/250/255/1856]

451 Biological sciences / Microbiology / Virology / Viral pathogenesis

452 [URI /631/326/596/2555]

453 Biological sciences / Microbiology / Bacteria / Bacterial pathogenesis

454 [URI /631/326/41/2531]

455 Biological sciences / Immunology / Immunological disorders / Immunological deficiency syndromes / HIV
456 infections

457 [URI /631/250/249/1570/1901]

458 Health sciences / Pathogenesis / Immunopathogenesis

459 [URI /692/420/2780]

460 **Tables**461 **Table 1: HIV-1, active tuberculosis (TB) and HIV-1/TB co-infection in 2015^{3,4}.**

| Global burden of disease | |
|--|-----------------------------|
| New HIV-1 infections | 2.1 million |
| Individuals living with HIV-1 | 36.7 million |
| New cases of active TB | 10.4 million |
| New cases of MDR-TB | 480,000 |
| Individuals with latent TB infection (LTBI) | 1 in 4 individuals globally |
| Mortality | |
| Deaths attributed to HIV-1 | 1.1 million |
| Deaths attributed to TB | 1.8 million |
| TB case fatality rate | 17% |
| Burden of co-infection | |
| Cases of active TB among individuals with HIV-1 | 1.14 million |
| Deaths attributed to HIV-1/TB co-infection | 400,000 |

462

463 **Figure legends**464 **Figure 1. HIV-1/TB co-infection increases the risk of active TB and HIV-1 disease progression.**

465 **(A)** The risk of active tuberculosis (TB) increases to 2-5 fold above baseline soon after an individual is infected
466 with HIV-1 during the early and chronic phases of infection. As HIV-1 progresses and causes severe
467 immunodeficiency, the risk of TB is further increased to at least 20-fold greater than the general population.
468 TB risk is accrued with longer times spent at low blood CD4⁺ T cell counts. Moreover, antiretroviral therapy
469 (ART) for HIV-1 does not fully restore the risk to baseline. There remains >4 fold increased rates of active TB
470 even once CD4⁺ T cell counts have reconstituted. **(B)** Incident *Mycobacterium tuberculosis* co-infection in
471 HIV-1 infected people increases HIV-1 replication and consequently viral diversity. It may also potentiate
472 chronic immune activation, accelerating the progression of HIV-1 disease.

473 **Figure 2. HIV-1 Nef may compromise host control of *Mycobacterium tuberculosis* by inhibition of
474 bacterial phagocytosis and autophagic clearance of phagosomal cargo.**

475 Macrophage control of *M. tuberculosis* is thought to be mediated by bacterial phagocytosis and the clearance
476 of *M. tuberculosis* containing phagosomes that fail to undergo phagolysosomal fusion via the autophagy
477 pathway. HIV-1 co-infection can undermine this mechanism of host defence at multiple levels. *M. tuberculosis*
478 clearance by this pathway is also dependent on vitamin D and augmented by Th1 responses through the action
479 of IFN γ . Deficiency of IFN γ responses owing to depletion of *M. tuberculosis* reactive Th1 cells in progressive
480 HIV-1 disease is thought to be the canonical mechanism by which HIV-1 compromises host defence against
481 *M. tuberculosis*. In addition, the HIV-1 Nef accessory protein reduces macrophage phagocytic capacity by
482 inhibiting AP-1-mediated endosomal recycling that is needed to form nascent phagosomes. Autophagosome
483 assembly is increased in HIV-1 infected macrophages, but their maturation and clearance function by fusion
484 with lysosomes is attenuated by the interaction of HIV-1 Nef with the autophagy related gene, Beclin-1.
485 Interestingly, vitamin D supplementation may overcome this inhibition of autophagosome maturation to
486 improve both *M. tuberculosis* clearance, and HIV-1 restriction by autophagy.

487 **Figure 3. High bacillary burden, T cell recovery and HIV-1-induced failure of immunoregulation drive
488 TB immune reconstitution inflammatory syndrome (TB-IRIS).**

489 Exaggerated pro-inflammatory responses that are normally derived from innate immune activation of myeloid
490 cells are the dominant feature of TB-IRIS, and T cell derived IFN γ responses are known to augment innate
491 immune responses by macrophages (A). Therefore the most likely model for exaggerated macrophage derived
492 inflammation in TB-IRIS is the combined effects of high bacillary burden in immunocompromised patients
493 before the onset of antiretroviral therapy (ART) **(B)** and recirculation of *Mycobacterium tuberculosis* reactive T
494 cells after ART initiation **(C)**. Although ART effectively blocks cell-cell propagation of the virus, the treatment
495 does not clear integrated HIV-1 provirus in macrophages which continue to express HIV-1 proteins. **(D)** In this
496 context HIV-1 inhibition of macrophage IL10 responses to *M. tuberculosis* may also contribute to exaggerated
497 inflammatory responses by a failure of immunoregulation **(D)**. CC, chemokines.

498 **Figure 4. Tuberculosis increases HIV-1 replication and propagation through innate immune signalling
499 pathways, proinflammatory cytokines and failure of immunoregulation.**

500 **(A)** Innate immune signalling pathways in macrophages can increase HIV-1 transcription through activation of
501 NF κ B, C/EBP, CREB/ATF and NFAT transcription factors. The host cell response to innate immune activation
502 by *Mycobacterium tuberculosis* leads to the production of a range of proinflammatory cytokines and
503 chemokines. These drive the local recruitment of T cells as part of a prototypic cell mediated immune response
504 **(B)**. The accumulation of activated T cells provides a population of cells permissive to HIV-1 and allows for

Figure legends

505 rapid virus propagation by direct cell-cell spread **(C)**. The pro-inflammatory cytokines also serve to promote
506 transactivation of virus replication through the action of NFkB and NFAT transcription factors **(D)**. HIV-1
507 attenuation of IL10 responses to *M. tuberculosis* favours the virus by reducing IL10 mediated inhibition of HIV-
508 1 transcription via C/EBP β and by promoting pro-inflammatory responses through a failure of
509 immunoregulation **(E)**. Although *M. tuberculosis* induces type1 IFN responses in macrophages, which would
510 be expected to promote an antiviral state, any autocrine or paracrine inhibition of HIV-1 replication is transient
511 **(F)**. Recent data has emerged to show that *M. tuberculosis* causes macrophage polyploidy through activation
512 of the cell cycle coupled to cytokinesis failure **(G)**. G1-like macrophages are more permissive to HIV-1.
513 Whether, *M. tuberculosis* induction of multinucleated giant cells further increases the HIV-1 permissive host
514 cellular niche, merits further investigation. CC, chemokines.
515

516 **References**

-
- 517 1. Korber, B. *et al.* Timing the ancestor of the HIV-1 pandemic strains. *Science* **288**, 1789–1796 (2000).
- 518 2. Russell, D. G. Who puts the tubercle in tuberculosis? *Nat. Rev. Microbiol.* **5**, 39–47 (2007).
- 519 3. UNAIDS. *Global AIDS update 2016*. (2016).
- 520 4. World Health Organisation. *Global tuberculosis report 2016*. (2016).
- 521 5. Dolan, K. *et al.* Global burden of HIV, viral hepatitis, and tuberculosis in prisoners and detainees. *Lancet*
- 522 *Lond. Engl.* **388**, 1089–1102 (2016).
- 523 6. Corbett, E. L. *et al.* Risk factors for pulmonary mycobacterial disease in South African gold miners. A
- 524 case-control study. *Am. J. Respir. Crit. Care Med.* **159**, 94–99 (1999).
- 525 7. Mesfin, Y. M., Hailemariam, D., Biadgign, S. & Kibret, K. T. Association between HIV/AIDS and Multi-
- 526 Drug Resistance Tuberculosis: A Systematic Review and Meta-Analysis. *PLoS ONE* **9**, (2014).
- 527 8. Gupta, R. K., Lucas, S. B., Fielding, K. L. & Lawn, S. D. Prevalence of tuberculosis in post-mortem
- 528 studies of HIV-infected adults and children in resource-limited settings: a systematic review and meta-
- 529 analysis. *AIDS Lond. Engl.* **29**, 1987–2002 (2015).
- 530 **Metanalysis including greater than 3200 autopsies from low and middle income countries which**
- 531 **estimated that TB was the cause of death in 37.2% of HIV-1 infected individuals.**
- 532 9. Sonnenberg, P. *et al.* How soon after infection with HIV does the risk of tuberculosis start to increase? A
- 533 retrospective cohort study in South African gold miners. *J. Infect. Dis.* **191**, 150–158 (2005).
- 534 10. Getahun, H., Gunneberg, C., Granich, R. & Nunn, P. HIV infection-associated tuberculosis: the
- 535 epidemiology and the response. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* **50 Suppl 3**, S201-207
- 536 (2010).
- 537 11. Lawn, S. D., Evans, A. J., Sedgwick, P. M. & Acheampong, J. W. Pulmonary tuberculosis: radiological
- 538 features in west Africans coinfecting with HIV. *Br. J. Radiol.* **72**, 339–344 (1999).
- 539 12. Naing, C., Mak, J. W., Maung, M., Wong, S. F. & Kassim, A. I. B. M. Meta-analysis: the association
- 540 between HIV infection and extrapulmonary tuberculosis. *Lung* **191**, 27–34 (2013).
- 541 13. Gilks, C. F. *et al.* Extrapulmonary and disseminated tuberculosis in HIV-1-seropositive patients
- 542 presenting to the acute medical services in Nairobi. *AIDS Lond. Engl.* **4**, 981–985 (1990).

References

- 543 14. Goletti, D. *et al.* Effect of Mycobacterium tuberculosis on HIV replication. Role of immune activation. *J.*
544 *Immunol. Baltim. Md* 1950 **157**, 1271–1278 (1996).
- 545 15. Marais, S., Meintjes, G., Lesosky, M., Wilkinson, K. A. & Wilkinson, R. J. Interleukin-17 mediated
546 differences in the pathogenesis of HIV-1-associated tuberculous and cryptococcal meningitis. *AIDS*
547 *Lond. Engl.* **30**, 395–404 (2016).
- 548 16. Collins, K. R. *et al.* Human immunodeficiency virus type 1 (HIV-1) quasispecies at the sites of
549 Mycobacterium tuberculosis infection contribute to systemic HIV-1 heterogeneity. *J. Virol.* **76**, 1697–1706
550 (2002).
- 551 17. Lawn, S. D. *et al.* Anatomically compartmentalized human immunodeficiency virus replication in HLA-
552 DR+ cells and CD14+ macrophages at the site of pleural tuberculosis coinfection. *J. Infect. Dis.* **184**,
553 1127–1133 (2001).
- 554 18. Nakata, K. *et al.* Mycobacterium tuberculosis enhances human immunodeficiency virus-1 replication in
555 the lung. *Am. J. Respir. Crit. Care Med.* **155**, 996–1003 (1997).
- 556 19. Toossi, Z. *et al.* Systemic immune activation and microbial translocation in dual HIV/TB infected subjects.
557 *J. Infect. Dis.* (2013). doi:10.1093/infdis/jit092
- 558 20. Meng, Q. *et al.* Immune Activation at Sites of HIV/TB Co-Infection Contributes to the Pathogenesis of
559 HIV-1 Disease. *PloS One* **11**, e0166954 (2016).
- 560 21. Sullivan, Z. A., Wong, E. B., Ndung'u, T., Kasprovicz, V. O. & Bishai, W. R. Latent and Active
561 Tuberculosis Infection Increase Immune Activation in Individuals Co-Infected with HIV. *EBioMedicine* **2**,
562 334–340 (2015).
- 563 22. Mellors, J. W. *et al.* Plasma viral load and CD4+ lymphocytes as prognostic markers of HIV-1 infection.
564 *Ann. Intern. Med.* **126**, 946–954 (1997).
- 565 23. Deeks, S. G. *et al.* Immune activation set point during early HIV infection predicts subsequent CD4+ T-
566 cell changes independent of viral load. *Blood* **104**, 942–947 (2004).
- 567 24. Whalen, C. *et al.* Accelerated course of human immunodeficiency virus infection after tuberculosis. *Am.*
568 *J. Respir. Crit. Care Med.* **151**, 129–135 (1995).
- 569 25. Badri, M., Ehrlich, R., Wood, R., Pulerwitz, T. & Maartens, G. Association between tuberculosis and HIV
570 disease progression in a high tuberculosis prevalence area. *Int. J. Tuberc. Lung Dis. Off. J. Int. Union*
571 *Tuberc. Lung Dis.* **5**, 225–232 (2001).

References

- 572 26. Kabali, C. *et al.* Increased mortality associated with treated active tuberculosis in HIV-infected adults in
573 Tanzania. *Tuberc. Edinb. Scotl.* **93**, 461–466 (2013).
- 574 27. Nunes-Alves, C. *et al.* In search of a new paradigm for protective immunity to TB. *Nat. Rev. Microbiol.*
575 **12**, 289–299 (2014).
- 576 28. Lahouassa, H. *et al.* SAMHD1 restricts the replication of human immunodeficiency virus type 1 by
577 depleting the intracellular pool of deoxynucleoside triphosphates. *Nat. Immunol.* **13**, 223–228 (2012).
- 578 29. Calantone, N. *et al.* Tissue myeloid cells in SIV-infected primates acquire viral DNA through phagocytosis
579 of infected T cells. *Immunity* **41**, 493–502 (2014).
- 580 30. Mlcochova, P. *et al.* A G1-like state allows HIV-1 to bypass SAMHD1 restriction in macrophages. *EMBO*
581 *J.* e201696025 (2017). doi:10.15252/embj.201696025
- 582 **HIV-1 was able to productively infect macrophages in non-replicative cell cycle during which SAMHD1**
583 **is inactivated, providing a niche which HIV-1 has exploited without a countermeasure for SAMHD1**
584 **restriction.**
- 585 31. Jambo, K. C. *et al.* Small alveolar macrophages are infected preferentially by HIV and exhibit impaired
586 phagocytic function. *Mucosal Immunol.* (2014). doi:10.1038/mi.2013.127
- 587 **Direct evidence was identified for productive HIV-1 infection in up to 5% of alveolar macrophages in**
588 **bronchoalveolar lavage specimens by RNA fluorescence in situ hybridisation, and further**
589 **assessments showed that these cells exhibited impaired phagocytosis of reported beads.**
- 590 32. Honeycutt, J. B. *et al.* Macrophages sustain HIV replication in vivo independently of T cells. *J. Clin.*
591 *Invest.* (2016). doi:10.1172/JCI84456
- 592 **HIV-1 was able to sustain long term productive infection in vivo, in a mouse model with human myeloid**
593 **cells , but without T cells, indicating that macrophages are sufficient to support chronic HIV-1 infection.**
- 594 33. Igarashi, T. *et al.* Macrophage are the principal reservoir and sustain high virus loads in rhesus macaques
595 after the depletion of CD4+ T cells by a highly pathogenic simian immunodeficiency virus/HIV type 1
596 chimera (SHIV): Implications for HIV-1 infections of humans. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 658–663
597 (2001).
- 598 34. Cribbs, S. K., Lennox, J., Caliendo, A. M., Brown, L. A. & Guidot, D. M. Healthy HIV-1-infected individuals
599 on highly active antiretroviral therapy harbor HIV-1 in their alveolar macrophages. *AIDS Res. Hum.*
600 *Retroviruses* **31**, 64–70 (2015).
- 601 35. Honeycutt, J. B. *et al.* HIV persistence in tissue macrophages of humanized myeloid-only mice during
602 antiretroviral therapy. *Nat. Med.* **23**, 638–643 (2017).

References

- 603 36. Churchill, M. J., Deeks, S. G., Margolis, D. M., Siliciano, R. F. & Swanstrom, R. HIV reservoirs: what,
604 where and how to target them. *Nat. Rev. Microbiol.* **14**, 55–60 (2016).
- 605 37. Lorenzo-Redondo, R. *et al.* Persistent HIV-1 replication maintains the tissue reservoir during therapy.
606 *Nature* **530**, 51–56 (2016).
- 607 38. Boritz, E. A. *et al.* Multiple Origins of Virus Persistence during Natural Control of HIV Infection. *Cell* **166**,
608 1004–1015 (2016).
- 609 39. Churchill, M. J. *et al.* Use of laser capture microdissection to detect integrated HIV-1 DNA in
610 macrophages and astrocytes from autopsy brain tissues. *J. Neurovirol.* **12**, 146–152 (2006).
- 611 40. McNab, F., Mayer-Barber, K., Sher, A., Wack, A. & O’Garra, A. Type I interferons in infectious disease.
612 *Nat. Rev. Immunol.* **15**, 87–103 (2015).
- 613 41. Stacey, A. R. *et al.* Induction of a striking systemic cytokine cascade prior to peak viremia in acute human
614 immunodeficiency virus type 1 infection, in contrast to more modest and delayed responses in acute
615 hepatitis B and C virus infections. *J. Virol.* **83**, 3719–3733 (2009).
- 616 42. Beignon, A.-S. *et al.* Endocytosis of HIV-1 activates plasmacytoid dendritic cells via Toll-like receptor–
617 viral RNA interactions. *J. Clin. Invest.* **115**, 3265–3275 (2005).
- 618 43. Berg, R. K. *et al.* Genomic HIV RNA induces innate immune responses through RIG-I-dependent sensing
619 of secondary-structured RNA. *PLoS One* **7**, e29291 (2012).
- 620 44. Gringhuis, S. I. *et al.* HIV-1 blocks the signaling adaptor MAVS to evade antiviral host defense after
621 sensing of abortive HIV-1 RNA by the host helicase DDX3. *Nat. Immunol.* **18**, 225–235 (2017).
- 622 45. Gao, D. *et al.* Cyclic GMP-AMP Synthase Is an Innate Immune Sensor of HIV and Other Retroviruses.
623 *Science* **341**, 903–906 (2013).
- 624 46. Yoh, S. M. *et al.* PQBP1 Is a Proximal Sensor of the cGAS-Dependent Innate Response to HIV-1. *Cell*
625 **161**, 1293–1305 (2015).
- 626 47. Jakobsen, M. R. *et al.* IFI16 senses DNA forms of the lentiviral replication cycle and controls HIV-1
627 replication. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E4571–4580 (2013).
- 628 48. Sandler, N. G. *et al.* Type I interferon responses in rhesus macaques prevent SIV infection and slow
629 disease progression. *Nature* **511**, 601–605 (2014).

References

- 630 49. Doyle, T., Goujon, C. & Malim, M. H. HIV-1 and interferons: who's interfering with whom? *Nat. Rev.*
631 *Microbiol.* **13**, 403–413 (2015).
- 632 50. Rasaiyaah, J. *et al.* HIV-1 evades innate immune recognition through specific cofactor recruitment.
633 *Nature* **503**, 402–405 (2013).
- 634 51. Tsang, J. *et al.* HIV-1 infection of macrophages is dependent on evasion of innate immune cellular
635 activation. *AIDS Lond. Engl.* **23**, 2255–2263 (2009).
- 636 52. Douek, D. C., Roederer, M. & Koup, R. A. Emerging concepts in the immunopathogenesis of AIDS.
637 *Annu. Rev. Med.* **60**, 471–484 (2009).
- 638 53. Muenchhoff, M. *et al.* Nonprogressing HIV-infected children share fundamental immunological features
639 of nonpathogenic SIV infection. *Sci. Transl. Med.* **8**, 358ra125 (2016).
- 640 54. Brenchley, J. M. *et al.* Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral
641 infections. *Blood* **112**, 2826–2835 (2008).
- 642 55. Stieh, D. J. *et al.* Th17 Cells Are Preferentially Infected Very Early after Vaginal Transmission of SIV in
643 Macaques. *Cell Host Microbe* **19**, 529–540 (2016).
- 644 56. Doitsh, G. *et al.* Cell death by pyroptosis drives CD4 T-cell depletion in HIV-1 infection. *Nature advance*
645 **online publication**, (2013).
- 646 57. Monroe, K. M. *et al.* IFI16 DNA sensor is required for death of lymphoid CD4 T cells abortively infected
647 with HIV. *Science* **343**, 428–432 (2014).
- 648 58. Galloway, N. L. K. *et al.* Cell-to-Cell Transmission of HIV-1 Is Required to Trigger Pyroptotic Death of
649 Lymphoid-Tissue-Derived CD4 T Cells. *Cell Rep.* **12**, 1555–1563 (2015).
- 650 59. Doitsh, G. & Greene, W. C. Dissecting How CD4 T Cells Are Lost During HIV Infection. *Cell Host Microbe*
651 **19**, 280–291 (2016).
- 652 60. Cooper, A. *et al.* HIV-1 causes CD4 cell death through DNA-dependent protein kinase during viral
653 integration. *Nature* **498**, 376–379 (2013).
- 654 61. Deeks, S. G. HIV Infection, Inflammation, Immunosenescence, and Aging. *Annu. Rev. Med.* **62**, 141–
655 155 (2011).
- 656 62. Khaitan, A. & Unutmaz, D. Revisiting Immune Exhaustion During HIV Infection. *Curr. HIV/AIDS Rep.* **8**,
657 4–11 (2011).

References

- 658 63. Beyer, M. *et al.* Tumor-necrosis factor impairs CD4(+) T cell-mediated immunological control in chronic
659 viral infection. *Nat. Immunol.* **17**, 593–603 (2016).
- 660 **Evidence from HIV-1 infected patients and the chronic LCMV mouse infection model indicates that**
661 **persistently elevated TNF levels can inhibit helper T cell function by upregulating expression of**
662 **inhibitory molecules such as PD1, suggesting a mechanism by which chronic inflammation can lead**
663 **to immunodeficiency.**
- 664 64. Jolly, C., Kashefi, K., Hollinshead, M. & Sattentau, Q. J. HIV-1 cell to cell transfer across an Env-induced,
665 actin-dependent synapse. *J. Exp. Med.* **199**, 283–293 (2004).
- 666 65. Jolly, C. Cell-to-cell transmission of retroviruses: Innate immunity and interferon-induced restriction
667 factors. *Virology* **411**, 251–259 (2011).
- 668 66. Boullé, M. *et al.* HIV Cell-to-Cell Spread Results in Earlier Onset of Viral Gene Expression by Multiple
669 Infections per Cell. *PLoS Pathog.* **12**, e1005964 (2016).
- 670 67. Baxter, A. E. *et al.* Macrophage Infection via Selective Capture of HIV-1-Infected CD4+ T Cells. *Cell Host*
671 *Microbe* **16**, 711–721 (2014).
- 672 68. Schnappinger, D. *et al.* Transcriptional Adaptation of Mycobacterium tuberculosis within Macrophages.
673 *J. Exp. Med.* **198**, 693–704 (2003).
- 674 **Transcriptional profiling of *M. tuberculosis* isolated from phagosomes compared to bacteria grown in**
675 **broth culture revealed that in response to phagosomal uptake and the effects of host IFN γ or inducible**
676 **nitric oxide synthase, the bacteria upregulate iron scavenging systems, expression of dormancy**
677 **related genes and genes that support anaerobic respiration.**
- 678 69. Peddireddy, V., Doddam, S. N. & Ahmed, N. Mycobacterial Dormancy Systems and Host Responses in
679 Tuberculosis. *Front. Immunol.* **8**, (2017).
- 680 70. Simeone, R. *et al.* Phagosomal Rupture by Mycobacterium tuberculosis Results in Toxicity and Host Cell
681 Death. *PLoS Pathog.* **8**, (2012).
- 682 71. Volkman, H. E. *et al.* Tuberculous granuloma induction via interaction of a bacterial secreted protein with
683 host epithelium. *Science* **327**, 466–469 (2010).
- 684 72. Mishra, B. B. *et al.* Mycobacterium tuberculosis protein ESAT-6 is a potent activator of the NLRP3/ASC
685 inflammasome. *Cell. Microbiol.* **12**, 1046–1063 (2010).
- 686 73. Davis, J. M. & Ramakrishnan, L. The Role of the Granuloma in Expansion and Dissemination of Early
687 Tuberculous Infection. *Cell* **136**, 37–49 (2009).

References

- 688 74. Kleinnijenhuis, J., Oosting, M., Joosten, L. A. B., Netea, M. G. & Van Crevel, R. Innate immune
689 recognition of Mycobacterium tuberculosis. *Clin. Dev. Immunol.* **2011**, 405310 (2011).
- 690 75. Watson, R. O. *et al.* The Cytosolic Sensor cGAS Detects Mycobacterium tuberculosis DNA to Induce
691 Type I Interferons and Activate Autophagy. *Cell Host Microbe* **17**, 811–819 (2015).
- 692 76. Liu, P. T. *et al.* Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response.
693 *Science* **311**, 1770–1773 (2006).
- 694 77. Berry, M. P. R. *et al.* An interferon-inducible neutrophil-driven blood transcriptional signature in human
695 tuberculosis. *Nature* **466**, 973–977 (2010).
- 696 78. Mayer-Barber, K. D. *et al.* Host-directed therapy of tuberculosis based on interleukin-1 and type I
697 interferon crosstalk. *Nature* **511**, 99–103 (2014).
- 698 79. Manzanillo, P. S., Shiloh, M. U., Portnoy, D. A. & Cox, J. S. Mycobacterium Tuberculosis Activates the
699 DNA-Dependent Cytosolic Surveillance Pathway within Macrophages. *Cell Host Microbe* **11**, 469–480
700 (2012).
- 701 80. Wassermann, R. *et al.* Mycobacterium tuberculosis Differentially Activates cGAS- and Inflammasome-
702 Dependent Intracellular Immune Responses through ESX-1. *Cell Host Microbe* **17**, 799–810 (2015).
- 703 **Inflammasome activation leading to secretion of active IL1 β and induction of type 1 IFNs by**
704 **macrophages infected with *M. tuberculosis* is dependent on the mycobacterial ESX1 secretion system,**
705 **and specific targeting of EsxA secretion attenuated induction of IFNs but not activation of the**
706 **inflammasome.**
- 707 81. Elkington, P. *et al.* MMP-1 drives immunopathology in human tuberculosis and transgenic mice. *J. Clin.*
708 *Invest.* (2011). doi:10.1172/JCI45666
- 709 82. Comas, I. *et al.* Human T cell epitopes of Mycobacterium tuberculosis are evolutionarily hyperconserved.
710 *Nat. Genet.* **42**, 498–503 (2010).
- 711 83. Coscolla, M. *et al.* M. tuberculosis T Cell Epitope Analysis Reveals Paucity of Antigenic Variation and
712 Identifies Rare Variable TB Antigens. *Cell Host Microbe* **18**, 538–548 (2015).
- 713 84. Al-Muhsen, S. & Casanova, J.-L. The genetic heterogeneity of mendelian susceptibility to mycobacterial
714 diseases. *J. Allergy Clin. Immunol.* **122**, 1043-1051; quiz 1052-1053 (2008).
- 715 85. Bell, L. C. K. *et al.* In Vivo Molecular Dissection of the Effects of HIV-1 in Active Tuberculosis. *PLOS*
716 *Pathog* **12**, e1005469 (2016).

References

- 717 **Genome-wide transcriptional profiling of biopsies from the site of the tuberculin skin test to make**
718 **molecular and systems level assessments of human immune responses to a standardised**
719 **mycobacterial challenge revealed deficient IL10 responses in HIV-1 infected patients before severe**
720 **immunodeficiency, preserved type 1 IFN responses in HIV-1 infected patients with severe**
721 **immunodeficiency, and exaggerated Th2 responses during unmasking TB-IRIS after antiretroviral**
722 **therapy.**
- 723 86. Kalsdorf, B. *et al.* HIV-1 infection impairs the bronchoalveolar T-cell response to mycobacteria. *Am. J.*
724 *Respir. Crit. Care Med.* **180**, 1262–1270 (2009).
- 725 87. Geldmacher, C. *et al.* Preferential infection and depletion of Mycobacterium tuberculosis-specific CD4 T
726 cells after HIV-1 infection. *J. Exp. Med.* **207**, 2869–2881 (2010).
- 727 **Preferential depletion of *M. tuberculosis* reactive CD4 T cells which produced more IL2, were more**
728 **permissive to HIV-1 infection, suggested that HIV-1 targeting of these cells may contribute to increased**
729 **risk of TB in early HIV-1 infection, before generalised T cell depletion.**
- 730 88. Cruz, A. *et al.* Pathological role of interleukin 17 in mice subjected to repeated BCG vaccination after
731 infection with Mycobacterium tuberculosis. *J. Exp. Med.* **207**, 1609–1616 (2010).
- 732 89. Nandi, B. & Behar, S. M. Regulation of neutrophils by interferon- γ limits lung inflammation during
733 tuberculosis infection. *J. Exp. Med.* **208**, 2251–2262 (2011).
- 734 90. Treerat, P. *et al.* Novel role for IL-22 in protection during chronic Mycobacterium tuberculosis HN878
735 infection. *Mucosal Immunol.* **10**, 1069–1081 (2017).
- 736 91. Kim, C. J. *et al.* A role for mucosal IL-22 production and Th22 cells in HIV-associated mucosal
737 immunopathogenesis. *Mucosal Immunol.* **5**, 670–680 (2012).
- 738 92. Saeidi, A. *et al.* Functional role of mucosal-associated invariant T cells in HIV infection. *J. Leukoc. Biol.*
739 **100**, 305–314 (2016).
- 740 93. Jiang, J. *et al.* Mucosal-associated invariant T-cell function is modulated by programmed death-1
741 signaling in patients with active tuberculosis. *Am. J. Respir. Crit. Care Med.* **190**, 329–339 (2014).
- 742 94. Gupta, A., Wood, R., Kaplan, R., Bekker, L.-G. & Lawn, S. D. Tuberculosis incidence rates during 8 years
743 of follow-up of an antiretroviral treatment cohort in South Africa: comparison with rates in the community.
744 *PloS One* **7**, e34156 (2012).
- 745 95. Heather, J. M. *et al.* Dynamic Perturbations of the T-Cell Receptor Repertoire in Chronic HIV Infection
746 and following Antiretroviral Therapy. *Front. Immunol.* **6**, (2016).

References

- 747 **Comprehensive TCR repertoire analysis by next generation sequencing of samples of HIV-1 infected**
748 **patients showed incomplete reconstitution of the T cell repertoire 3 months after effective antiretroviral**
749 **therapy.**
- 750 96. Cosgrove, C. *et al.* Early and nonreversible decrease of CD161⁺ /MAIT cells in HIV infection. *Blood*
751 **121**, 951–961 (2013).
- 752 97. Cambier, C. J. *et al.* Mycobacteria manipulate macrophage recruitment through coordinated use of
753 membrane lipids. *Nature* **505**, 218–222 (2014).
- 754 98. Roca, F. J. & Ramakrishnan, L. TNF Dually Mediates Resistance and Susceptibility to Mycobacteria via
755 Mitochondrial Reactive Oxygen Species. *Cell* **153**, 521–534 (2013).
- 756 99. Alonso, S., Pethe, K., Russell, D. G. & Purdy, G. E. Lysosomal killing of Mycobacterium mediated by
757 ubiquitin-derived peptides is enhanced by autophagy. *Proc. Natl. Acad. Sci.* **104**, 6031–6036 (2007).
- 758 100. Gutierrez, M. G. *et al.* Autophagy Is a Defense Mechanism Inhibiting BCG and Mycobacterium
759 tuberculosis Survival in Infected Macrophages. *Cell* **119**, 753–766 (2004).
- 760 101. Mazzolini, J. *et al.* Inhibition of phagocytosis in HIV-1-infected macrophages relies on Nef-dependent
761 alteration of focal delivery of recycling compartments. *Blood* **115**, 4226–4236 (2010).
- 762 **Inhibition of macrophage phagocytic uptake by the effect of HIV-1 Nef accessory protein on AP-1**
763 **mediated recycling of endosomes required for phagosome formation.**
- 764 102. Toossi, Z., Liu, S., Wu, M., Mayanja-Kizza, H. & Hirsch, C. S. Short Communication: Circulating Plasma
765 HIV-1 Viral Protein R in Dual HIV-1/Tuberculosis Infection. *AIDS Res. Hum. Retroviruses* **30**, 644–647
766 (2014).
- 767 103. Fujii, Y., Otake, K., Tashiro, M. & Adachi, A. Soluble Nef antigen of HIV-1 is cytotoxic for human CD4⁺
768 T cells. *FEBS Lett.* **393**, 93–96 (1996).
- 769 104. Kyei, G. B. *et al.* Autophagy pathway intersects with HIV-1 biosynthesis and regulates viral yields in
770 macrophages. *J. Cell Biol.* **186**, 255–268 (2009).
- 771 **Induction of autophagosome formation in HIV-1 infected macrophages, wherein HIV-1 Nef mediated**
772 **autophagosome maturation supports HIV-1 replication but counteracts autophagic degradation of the**
773 **virus and might be expected to inhibit autophagic clearance of mycobacteria.**
- 774 105. Pathak, S., Wentzel-Larsen, T. & Asjo, B. Effects of in vitro HIV-1 infection on mycobacterial growth in
775 peripheral blood monocyte-derived macrophages. *Infect. Immun.* **78**, 4022–4032 (2010).

References

- 776 106. Campbell, G. R. & Spector, S. A. Vitamin D Inhibits Human Immunodeficiency Virus Type 1 and
777 Mycobacterium tuberculosis Infection in Macrophages through the Induction of Autophagy. *PLoS Pathog.*
778 **8**, (2012).
- 779 107. Martineau, A. R. *et al.* Reciprocal seasonal variation in vitamin D status and tuberculosis notifications in
780 Cape Town, South Africa. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19013–19017 (2011).
- 781 108. Patel, N. R. *et al.* HIV impairs TNF-alpha mediated macrophage apoptotic response to Mycobacterium
782 tuberculosis. *J. Immunol. Baltim. Md 1950* **179**, 6973–6980 (2007).
- 783 109. Patel, N. R., Swan, K., Li, X., Tachado, S. D. & Koziel, H. Impaired M. tuberculosis-mediated apoptosis
784 in alveolar macrophages from HIV+ persons: potential role of IL-10 and BCL-3. *J. Leukoc. Biol.* **86**, 53–
785 60 (2009).
- 786 110. Kumawat, K., Pathak, S. K., Spetz, A.-L., Kundu, M. & Basu, J. Exogenous Nef is an inhibitor of
787 Mycobacterium tuberculosis-induced tumor necrosis factor-alpha production and macrophage apoptosis.
788 *J. Biol. Chem.* **285**, 12629–12637 (2010).
- 789 111. Mahamed, D. *et al.* Intracellular growth of Mycobacterium tuberculosis after macrophage cell death leads
790 to serial killing of host cells. *eLife* **6**, e22028 (2017).
- 791 112. Walker, N. F. *et al.* Doxycycline and HIV infection suppress tuberculosis-induced matrix
792 metalloproteinases. *Am. J. Respir. Crit. Care Med.* **185**, 989–997 (2012).
- 793 113. Huang, C.-C. *et al.* The effect of HIV-related immunosuppression on the risk of tuberculosis transmission
794 to household contacts. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* **58**, 765–774 (2014).
- 795 114. Bell, L. C. K., Breen, R., Miller, R. F., Noursadeghi, M. & Lipman, M. Paradoxical reactions and immune
796 reconstitution inflammatory syndrome in tuberculosis. *Int. J. Infect. Dis. IJID Off. Publ. Int. Soc. Infect.*
797 *Dis.* **32**, 39–45 (2015).
- 798 115. Müller, M. *et al.* Immune reconstitution inflammatory syndrome in patients starting antiretroviral therapy
799 for HIV infection: a systematic review and meta-analysis. *Lancet Infect. Dis.* **10**, 251–261 (2010).
- 800 116. Lai, R. P. J. *et al.* HIV–tuberculosis-associated immune reconstitution inflammatory syndrome is
801 characterized by Toll-like receptor and inflammasome signalling. *Nat. Commun.* **6**, 8451 (2015).
- 802 **Pro-inflammatory responses in TB-IRIS cases were enriched for innate immune MyD88 and**
803 **inflammasome mediated pathways in myeloid cells suggesting that unregulated recovery of these**
804 **pathways after antiretroviral therapy may be responsible for the pathogenesis of IRIS.**

References

- 805 117. Tachado, S. D. *et al.* MyD88-dependent TLR4 signaling is selectively impaired in alveolar macrophages
806 from asymptomatic HIV+ persons. *Blood* **115**, 3606–3615 (2010).
- 807 118. Tachado, S. D., Li, X., Swan, K., Patel, N. & Koziel, H. Constitutive activation of phosphatidylinositol 3-
808 kinase signaling pathway down-regulates TLR4-mediated tumor necrosis factor-alpha release in alveolar
809 macrophages from asymptomatic HIV-positive persons in vitro. *J. Biol. Chem.* **283**, 33191–33198 (2008).
- 810 119. Noursadeghi, M., Katz, D. R. & Miller, R. F. HIV-1 infection of mononuclear phagocytic cells: the case
811 for bacterial innate immune deficiency in AIDS. *Lancet Infect. Dis.* **6**, 794–804 (2006).
- 812 120. Noursadeghi, M. *et al.* Genome-wide innate immune responses in HIV-1-infected macrophages are
813 preserved despite attenuation of the NF-kappa B activation pathway. *J. Immunol. Baltim. Md 1950* **182**,
814 319–328 (2009).
- 815 121. Tomlinson, G. S. *et al.* HIV-1 Infection of Macrophages Dysregulates Innate Immune Responses to
816 *Mycobacterium tuberculosis* by Inhibition of Interleukin-10. *J. Infect. Dis.* **209**, 1055–1065 (2014).
- 817 **HIV-1 infected macrophages exhibited selective depletion of IL10 responses to *M. tuberculosis***
818 **co-infection, leading to a failure of immunoregulation, exaggerated proinflammatory responses and**
819 **increased HIV-1 replication.**
- 820 122. Maddocks, S. *et al.* Gene expression in HIV-1/*Mycobacterium tuberculosis* co-infected macrophages is
821 dominated by *M. tuberculosis*. *Tuberc. Edinb. Scotl.* **89**, 285–293 (2009).
- 822 123. Pathak, S., Wentzel-Larsen, T. & Asjö, B. Effects of in vitro HIV-1 infection on mycobacterial growth in
823 peripheral blood monocyte-derived macrophages. *Infect. Immun.* **78**, 4022–4032 (2010).
- 824 124. Tan, D. B. A. *et al.* TLR2-induced cytokine responses may characterize HIV-infected patients
825 experiencing mycobacterial immune restoration disease. *AIDS Lond. Engl.* **25**, 1455–1460 (2011).
- 826 125. Tran, H. T. T. *et al.* Modulation of the complement system in monocytes contributes to tuberculosis-
827 associated immune reconstitution inflammatory syndrome. *AIDS Lond. Engl.* **27**, 1725–1734 (2013).
- 828 126. Toossi, Z. *et al.* Increased replication of HIV-1 at sites of *Mycobacterium tuberculosis* infection: potential
829 mechanisms of viral activation. *J. Acquir. Immune. Defic. Syndr.* **28**, 1–8 (2001).
- 830 127. Van Lint, C., Bouchat, S. & Marcello, A. HIV-1 transcription and latency: an update. *Retrovirology* **10**, 67
831 (2013).

References


- 832 128. Zhang, Y., Nakata, K., Weiden, M. & Rom, W. N. Mycobacterium tuberculosis enhances human
833 immunodeficiency virus-1 replication by transcriptional activation at the long terminal repeat. *J. Clin.*
834 *Invest.* **95**, 2324–2331 (1995).
- 835 129. Henderson, A. J., Zou, X. & Calame, K. L. C/EBP proteins activate transcription from the human
836 immunodeficiency virus type 1 long terminal repeat in macrophages/monocytes. *J. Virol.* **69**, 5337–5344
837 (1995).
- 838 130. Ranjbar, S., Jasenosky, L. D., Chow, N. & Goldfeld, A. E. Regulation of Mycobacterium tuberculosis-
839 Dependent HIV-1 Transcription Reveals a New Role for NFAT5 in the Toll-Like Receptor Pathway. *PLoS*
840 *Pathog* **8**, e1002620 (2012).
- 841 131. Honda, Y. *et al.* Type I Interferon Induces Inhibitory 16-kD CCAAT/ Enhancer Binding Protein (C/EBP) β ,
842 Repressing the HIV-1 Long Terminal Repeat in Macrophages: Pulmonary Tuberculosis Alters C/EBP
843 Expression, Enhancing HIV-1 Replication. *J. Exp. Med.* **188**, 1255–1265 (1998).
- 844 132. Gröschel, M. I., Sayes, F., Simeone, R., Majlessi, L. & Brosch, R. ESX secretion systems: mycobacterial
845 evolution to counter host immunity. *Nat. Rev. Microbiol.* **14**, 677–691 (2016).
- 846 133. Wang, Y. & Rice, A. P. Interleukin-10 inhibits HIV-1 LTR-directed gene expression in human
847 macrophages through the induction of cyclin T1 proteolysis. *Virology* **352**, 485–492 (2006).
- 848 134. Kootstra, N. A., van 't Wout, A., Huisman, H. G., Miedema, F. & Schuitemaker, H. Interference of
849 interleukin-10 with human immunodeficiency virus type 1 replication in primary monocyte-derived
850 macrophages. *J. Virol.* **68**, 6967–6975 (1994).
- 851 135. Tanaka, N. *et al.* Interleukin-10 Induces Inhibitory C/EBP β through STAT-3 and Represses HIV-1
852 Transcription in Macrophages. *Am. J. Respir. Cell Mol. Biol.* **33**, 406–411 (2005).
- 853 136. Duh, E. J., Maury, W. J., Folks, T. M., Fauci, A. S. & Rabson, A. B. Tumor necrosis factor alpha activates
854 human immunodeficiency virus type 1 through induction of nuclear factor binding to the NF-kappa B sites
855 in the long terminal repeat. *Proc. Natl. Acad. Sci. U. S. A.* **86**, 5974–5978 (1989).
- 856 137. Poli, G., Kinter, A. L. & Fauci, A. S. Interleukin 1 induces expression of the human immunodeficiency
857 virus alone and in synergy with interleukin 6 in chronically infected U1 cells: inhibition of inductive effects
858 by the interleukin 1 receptor antagonist. *Proc. Natl. Acad. Sci. U. S. A.* **91**, 108–112 (1994).

References

- 859 138. Poli, G. *et al.* Interleukin 6 induces human immunodeficiency virus expression in infected monocytic cells
860 alone and in synergy with tumor necrosis factor alpha by transcriptional and post-transcriptional
861 mechanisms. *J. Exp. Med.* **172**, 151–158 (1990).
- 862 139. Han, X., Becker, K., Degen, H. J., Jablonowski, H. & Strohmeyer, G. Synergistic stimulatory effects of
863 tumour necrosis factor alpha and interferon gamma on replication of human immunodeficiency virus type
864 1 and on apoptosis of HIV-1-infected host cells. *Eur. J. Clin. Invest.* **26**, 286–292 (1996).
- 865 140. Jenkins, S. J. *et al.* Local macrophage proliferation, rather than recruitment from the blood, is a signature
866 of TH2 inflammation. *Science* **332**, 1284–1288 (2011).
- 867 141. Aziz, A., Soucie, E., Sarrazin, S. & Sieweke, M. H. MafB/c-Maf deficiency enables self-renewal of
868 differentiated functional macrophages. *Science* **326**, 867–871 (2009).
- 869 142. Schulz, C. *et al.* A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science*
870 **336**, 86–90 (2012).
- 871 143. Hashimoto, D. *et al.* Tissue-Resident Macrophages Self-Maintain Locally throughout Adult Life with
872 Minimal Contribution from Circulating Monocytes. *Immunity* **38**, 792–804 (2013).
- 873 144. Herrtwich, L. *et al.* DNA Damage Signaling Instructs Polyploid Macrophage Fate in Granulomas. *Cell*
874 **167**, 1264–1280.e18 (2016).
- 875 145. Lawn, S. D., Kranzer, K. & Wood, R. Antiretroviral Therapy for Control of the HIV-associated Tuberculosis
876 Epidemic in Resource-Limited Settings. *Clin. Chest Med.* **30**, 685–699 (2009).
- 877 146. Smyth, R. P. & Negroni, M. A step forward understanding HIV-1 diversity. *Retrovirology* **13**, 27 (2016).
- 878 147. Walker, T. M. *et al.* Assessment of Mycobacterium tuberculosis transmission in Oxfordshire, UK, 2007-
879 12, with whole pathogen genome sequences: an observational study. *Lancet Respir. Med.* **2**, 285–292
880 (2014).
- 881 148. Fenner, L. *et al.* HIV infection disrupts the sympatric host-pathogen relationship in human tuberculosis.
882 *PLoS Genet.* **9**, e1003318 (2013).
- 883 149. Koch, A. S. *et al.* The Influence of HIV on the Evolution of Mycobacterium tuberculosis. *Mol. Biol. Evol.*
884 **34**, 1654–1668 (2017).
- 885 150. Kaufmann, S. H. E. *et al.* TB biomarkers, TB correlates and human challenge models: New tools for
886 improving assessment of new TB vaccines. *Tuberc. Edinb. Scotl.* **99 Suppl 1**, S8–S11 (2016).

Figure 1

A



B




Figure 2




Figure 3




Figure 4

