

Alternate Healthy Eating Index 2010, Dietary inflammatory index and risk of mortality: Results from the Whitehall II cohort study and meta-analysis of previous DII and mortality studies.

Nitin Shivappa^{1,2,3}, James R. Hebert^{1,2,3}, Mika Kivimaki⁴, Tasnime Akbaraly^{4,5}

¹ Cancer Prevention and Control Program, University of South Carolina, Columbia, SC 29208, USA

² Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA

³ Connecting Health Innovations LLC, Columbia, SC 29201, USA

⁴ University College London, Department of Epidemiology and Public Health, London, United Kingdom

⁵ MMDN, University Montpellier, EPHE, INSERM, U1198, Montpellier, F-34095 France

Address correspondence and reprint requests to: Dr. Nitin Shivappa, Cancer Prevention and Control Program, University of South Carolina, 915 Greene Street, Suite 241, Columbia, SC 29208, Telephone: (803) 576-5666, Fax: (803) 576-5624, e-mail: shivappa@mailbox.sc.edu

Running Title: Association between DII, AHEI-2010 and mortality risk in Whitehall II Study

Abstract:

We aimed to examine the association between the Alternative Healthy Eating Index updated in 2010 (AHEI - 2010), the Dietary Inflammatory Index (DIITM) and risk of mortality in the Whitehall II study. We also conducted a meta-analysis on the DII-based results from previous studies to summarize the overall evidence. Data on dietary behavior assessed by self-administered repeated food frequency questionnaires and on mortality status were available for 7627 participants from the Whitehall II cohort. Cox proportional hazards regression models were performed to assess the association between cumulative average of AHEI-2010 and DII scores and mortality risk. During 22 years of follow-up, 1001 participants died (450 from cancer, 264 from cardiovascular disease). Both AHEI-2010 and DII were associated with all-cause mortality. After adjustment for a large range of potential confounders, for each increment of dietary indices z-score, hazard ratio (HR)=0.82, 95 % CI: 0.76-0.88 for AHEI-2010 and HR= 1.18, 95 % CI: 1.08-1.29 for DII were observed. Significant associations were also observed with cardiovascular and cancer mortality risk. For DII, a meta-analysis (using fixed effects) from this and 4 previous studies showed a positive association of DII score with all-cause (HR=1.04, 95 % CI: 1.03-1.05, n=28,891 deaths), cardiovascular (HR=1.05, 95 % CI: 1.03-1.07, n=10,424 deaths) and cancer mortality (HR=1.05, 95 % CI: 1.03-1.07, n=8269). The present study confirms the validity to assess overall diet through AHEI-2010 and DII in the Whitehall II cohort and highlights the importance of considering diet indices related to inflammation when evaluating all cause, cardiovascular and cancer mortality risk.

Keywords: Dietary indices; mortality risk; Whitehall II cohort Study; meta-analysis;

INTRODUCTION

Various studies have been conducted exploring diet and dietary components in relation to, both from all-cause and cause-specific mortality⁽¹⁾. There has recently been a shift in the focus from individual nutrient and food studies to dietary pattern studies because dietary components are consumed in combination and are correlated with one another⁽²⁾. With the dietary pattern approach overall quality of takes into account the complexity of the diet and the potentially synergistic or antagonistic effects of individual dietary components⁽³⁾.

Chronic inflammation is known to be associated with a variety of chronic health conditions including arthritis, diverticulitis, cardiovascular disease, diabetes^(4; 5; 6), and common epithelial cancers, with colorectal cancer^(7; 8; 9) being the most extensively studied. Dietary factors can contribute to an individual's underlying state of chronic inflammation^(10; 11) and there is growing evidence that specific dietary components influence inflammation^(12; 13; 14), potentially influencing risk of all-cause, cancer and cardiovascular mortality^(15; 16; 17; 18). We previously showed in the Whitehall II cohort that a high score in Alternative Healthy Eating Index (AHEI) score was associated both with inflammatory markers and mortality risk.

The Dietary Inflammatory Index (DII) was developed to characterize an individual's diet on a continuum from maximally anti- to pro-inflammatory⁽¹⁹⁾. Thus far, the DII has been found to be associated with C-reactive protein (CRP)^(20; 21), interleukin-6^(22; 23; 24), and tumor necrosis factor (TNF)-alpha⁽²³⁾. Until now, the DII has been applied to mortality outcomes only in four cohort studies and no overall evaluation of this evidence exists. Since the creation of the AHEI in 2002 which was based on foods and nutrients predictive of chronic disease risk, substantial evidence has emerged to support a role of additional dietary factors in the development of chronic disease and a new measure of diet quality that incorporates current scientific evidence on diet and health has been proposed: the AHEI-2010⁽²⁵⁾. In this analysis, we assessed the association between the DII, the AHEI-2010 and all-cause, cardiovascular, and cancer mortality risk in a population-based prospective cohort of British men and women participating in the Whitehall II cohort. A secondary objective was to summarize the overall evidence of the association between DII and mortality from the Whitehall II study and all previous studies using meta-analysis.

METHODS

Study population

Participants of the Whitehall II study were London-based office staff, aged 35–55 years, who worked in 20 civil service departments at study inception ⁽²⁶⁾. Baseline screening (phase 1: 1985–1988, n = 10,308) consisted of a clinical examination and a self-administered questionnaire. Subsequent phases of data collection alternated between a clinical examination along with a questionnaire survey (Phase 3: 1991/93, n = 8815; Phase 5: 1997/99, n=7263; Phase 7: 2002/04, n=6943; Phase 9: 2007/09) and a postal questionnaire alone (Phases 2, 4, 8 and 10). After the study was described to each participant, written informed consent was obtained. The University College London Ethics Committee approved the study. Phase 3 (1991–1993) was considered the baseline for the purpose of this study, because it was the first assessment of the AHEI-2010 and DII. Analyses were carried out on the 7627 participants alive at phase 3 with information on vital status and with complete data on dietary indices and covariates at phase 3.

Dietary assessment

Dietary intake at phase 3 was assessed with the use of a semi-quantitative food-frequency questionnaire (FFQ) with 127 food items, as described previously ^(27; 28). The validity and the reliability of the FFQ in terms of nutrient and food consumption have been documented in detail both in our cohort and in another independent UK cohort ^(28; 29). The selected frequency category for each food item was converted to a daily intake. Nutrient intakes were computed by multiplication of the consumption frequency for each food by its nutrient content (for specified portions), and then the nutrient contributions from all foods were summed. Frequency of consumption for multivitamin supplements was also collected. Nutrient values were calculated with the use of a computerized system developed for the Whitehall II dietary data.

Alternative Healthy Eating Index 2010 (AHEI- 2010)

Scoring criteria for AHEI-2010 are described in detail elsewhere ⁽²⁵⁾. The AHEI-2010 is based on 11 components: six components for which the highest intakes were supposed to be ideal (vegetables, fruit, whole grains, nuts and legumes, long chain omega-3 fats (docosahexaenoic acid and eicosapentaenoic acid), and

polyunsaturated fatty acids), one component for which moderate intake was supposed to be ideal (alcohol), and four components for which avoidance or lowest intake were supposed to be ideal (sugar sweetened drinks and fruit juice, red and processed meat, trans fat, and sodium). Each component is given a minimal score of 0 and a maximal score of 10, with intermediate values scored proportionally, and has the potential to contribute 0-10 points to the total score. All the component scores are summed to obtain a total AHEI-2010 score, which ranges from 0 to 110, with a higher score representing a healthier diet (see online supplementary material for the distribution of the baseline score).

Dietary Inflammatory Index (DII)

The DII is based on literature published through 2010 linking diet to inflammation. Developing the DII involved reviewing and scoring nearly 2000 scientific articles on diet and six inflammatory markers [i.e., CRP, interleukin (IL)-1 β , IL-4, IL-6, IL-10, and TNF- α] based on cell culture and laboratory animal experiments, and cross-sectional, longitudinal and intervention trials in humans. Individual intakes of food parameters on which the DII is based have then been compared to a world standard database of dietary intake based on datasets from 11 different regions worldwide. A complete description of the DII is available elsewhere. Briefly, to calculate DII scores for participants in this study, the dietary data at phase 3 were first linked to the world database that provided a robust estimate of a mean and standard deviation for each parameter⁽¹⁹⁾. These then become the multipliers to express an individual's exposure relative to the "standard global mean" as a z-score. This is achieved by subtracting the "standard global mean" from the amount reported and dividing this value by the standard deviation. To minimize the effect of "right skewing," we converted this value to a centered percentile score. The centered percentile score for each food parameter for each individual was then multiplied by the respective food parameter-specific inflammatory effect score, which was derived from the literature review, in order to obtain a food parameter-specific DII score for an individual. All of the food parameter-specific DII scores were then summed to create the overall DII score for each participant in the study⁽¹⁹⁾. A description of validation of the DII, including comparing hs-CRP values to DII derived from both dietary recalls and a structured questionnaire similar to an FFQ, is available elsewhere⁽¹⁹⁾. More negative values represent more

anti-inflammatory diets. The DII score when calculated from all 45 food parameters could range from a global minimum of -8.87 (maximally anti-inflammatory) to a global maximum $+7.98$ (maximally pro-inflammatory), though the typical observed range in most studies where DII is calculated from an average of 27 food parameters is from about -5.5 to $+5.5$ ^(24; 30; 31). Details regarding the construct validation of the DII have been detailed elsewhere ⁽¹⁹⁾. In the present study, a total of 27 of 45 food parameters were available from the FFQ and were used to calculate the DII. These include energy, carbohydrate, protein, total fat, alcohol, fiber, cholesterol, saturated fat, MUFA, PUFA, omega-3, omega-6, trans-fat, niacin, thiamin, riboflavin, vitamin B12, B6, iron, magnesium, zinc, selenium, vitamin A, vitamin C, vitamin D, vitamin E, folic acid. Steps involved in the DII calculation are described in Online Supplementary Figure A. DII scores were calculated from nutrients derived only from dietary items.

We compute the AHEI-2010 and the DII scores from each FFQ administered in phase 3 (1991-93), phase 5 (1997-99) and phase 7 (2002-2004). To reduce measurement errors and to represent long-term dietary intake, we calculated the cumulative average of AHEI-2010 and DII.

Mortality assessment

The Whitehall II study is linked to the National Health Services death and electronic patient records with the use of the National Health Services identification number assigned to all British citizens. A total of 10,297 participants (99.9%) were successfully traced and have been followed through these registries. Mortality data, which included the date and cause of death, were available through the National Health Services Central Registry until 28 February 2015. Death certificates were coded with the use of the 9th or 10th revision of the International Classification of Diseases (ICD) (<http://www.who.int/classifications/icd/en/>). We analyzed all-cause mortality and mortality from specific causes, such as CVD (ICD-9 codes 390.0–458.9 and ICD-10 codes I00–I99) and cancer (ICD-9 140.0–209.9 and ICD-10 C00–C97). Non cancer/non-CVD mortality includes all remaining deaths that are classified neither as cancer nor CVD, and included deaths from diseases of the

respiratory system (most common other cause of death), digestive system, or nervous system; injuries; poisoning; and external causes.

Covariates

Sociodemographic variables consisted of age, sex, ethnicity (White/South Asian / Black), marital status (married or cohabiting versus living alone) and occupational position with the use of current (or last, for retired participants) British civil service employment grade, defined on the basis of salary and grouped into 3 categories: high (senior administrators) / intermediate (executives, professionals, and technical staff) / low (clerical and office support staff) grades.

Health behaviors included in the analysis were smoking (current / former / nonsmoker), total energy intake estimated from the food frequency questionnaire (in kcal/day), alcohol intake patterns [none : 0 glass per day, moderate: <2 (3) glasses per day for women (men), and heavy consumption: ≥ 2 (3) glasses per day for women (men)] and physical activity assessed via questionnaire data and categorized into 3 groups (high, intermediate, low) according to frequency of participation in “vigorous” (e.g., running, hard swimming, playing squash), “moderately energetic” (e.g., dancing, cycling, leisurely swimming), and “mildly energetic” physical activity⁽³²⁾. Health status was ascertained with the use of a number of measures: prevalence of cardiovascular diseases on the basis of clinically verified events, which included non-fatal myocardial infarction, definite angina, self-reported stroke or transient ischemic attack; hypertension (systolic or diastolic blood pressure ≥ 140 or ≥ 90 mm Hg, respectively, or use of hypertensive drugs); concentration of high density lipoprotein (HDL) cholesterol (in mmol/L), use of lipid-lowering drugs; type 2 diabetes (diagnosed with the use of the World Health Organization (WHO) definition); body mass index (BMI; in kg/m^2) and self-reported longstanding illness (participants responded to a question asking if they had a longstanding illness: yes/no).

Covariates were obtained from the baseline questionnaire (phase 3) and updated every 5 years (at phase 5 and phase 7). All variables (other than sex and ethnicity) included time-varying covariates.

Statistical analysis

Characteristics of participants as a function of mortality status were first described and tested using chi-square tests for categorical variables and Student t-test for continuous variables. Next, comparison of characteristics associated with DII and AHEI-2010 tertiles were performed using chi-square tests for categorical variables and ANOVA for continuous variables. Associations between AHEI-2010 and DII tertiles were assessed using chi-square tests.

The main analyses consisted of exploring the association between the risk of mortality and the cumulative average of AHEI-2010 and DII scores in the Whitehall II cohort. Stratified proportional Cox hazard models analyses were used to calculate hazard ratios (HRs) and 95% confidence intervals (95% CIs) for death. In these analyses the AHEI-2010 and DII were first considered as continuous variables by using z -score. Then AHEI-2010 and DII were categorized into tertiles, with the lowest tertile serving as the reference group for both indices. We calculated a test for trend across the tertiles of the AHEI-2010 and DII score by treating the categories as an ordinal variable in a proportional hazards model. We used three multivariable models, In model 1 we adjusted age, sex and ethnicity; in model 2 we additionally adjusted for occupational grade, living alone, smoking habits, alcohol consumption (only for DII), physical activity; and model 3 was further adjusted for health status factors including body mass index (BMI), antecedent of CVD, use of lipids lowering drugs, HDL cholesterol, hypertension, type 2 diabetes and longstanding illness. These time-varying covariates were chosen *a priori* as they previously had been shown to be strong risk factors for mortality in this cohort. To minimize the effect of missingness, we replaced missing data on these covariates with the last valid values. These analyses of the association between DII, AHEI-2010 and overall mortality were repeated for cause-specific mortality including cardiovascular and cancer mortality. Stratified analyses also were carried out by tertiles of energy intake. Analyses were performed using SAS statistical software, version 9.4.

A second analysis consisted for the DII of meta-analysis results from Whitehall II and all four previous studies. To date, there were four other studies ^(33; 34; 35; 36) that have been published on the DII and mortality. We synthesised this evidence by pooling effect estimates from those studies and the current study. As the number of studies was modest ⁽³⁷⁾ fixed-effects meta-analysis was used. For comparison, we repeated the analysis using an

alternative approach, random-effects meta-analysis. We used continuous DII score as the exposure because results for that variable were available from all five studies. Mortality analyses were restricted to deaths from all-causes, cancer and cardiovascular disease. Heterogeneity in study-specific estimates was evaluated using I^2 statistics. All meta-analyses were performed using Stata, version MP 13.1 with the "metan" command (Online Supplementary Material Table F).

RESULTS

Analyses from the Whitehall II study were carried out on the 7627 participants alive at phase 3 with information on vital status and with data on dietary indices and covariates as described in the Flow chart mapping the selection of participants (Online supplementary Figure B). Characteristics of these participants as a function of mortality status are detailed in Table 1. Distribution of AHEI-2010 and DII score are presented in Online Supplementary Material- Table A. DII had a mean \pm SD value of 0.37 ± 1.41 , with a range from -3.28 (most anti-inflammatory score) to +4.54 (most pro-inflammatory score). AHEI had a mean \pm SD value of 48.7 ± 10.0 , with a range from 15.5 (least healthy diet) to 85.5 (healthiest diet). Participants in the highest tertile of AHEI-2010 (reflecting a healthier diet) compared to those in the lowest tertile (reflecting an unhealthier diet) were less likely to be men, white with a low socioeconomic position and more likely to be younger, with higher occupational grade, to be married, and to have healthy behaviors (smoking habits, alcohol consumption and physical activity). Their average mean of total energy intake was also higher. Regarding health factors, the group of participants in the highest tertile of AHEI-2010 were less likely to have hypertension, they showed higher means of HDL cholesterol and lower means of BMI and. Similar pattern of socio-demographic, health behavior and health status factors associated with DII tertiles were observed.

The correlation coefficient between AHEI-2010 and DII was -0.41. In Table 2 we show that 54.4% of participants in tertile 1 of DII; i.e., with the most anti-inflammatory diets, are in tertile 3 of AHEI-10 (corresponding to the healthiest diet) and 51.9 % of participants in tertile 3 of DII i.e., with the most pro-inflammatory diets, are in tertile 1 of AHEI (corresponding to the unhealthiest diet). However, there is some

discordance, 18.1% of participants in the most anti-inflammatory group (tertile 1) of DII are in the unhealthiest diet group for AHEI-2010

In total, 1001 total deaths were identified during the 22 years of follow-up; 450 cancer deaths, 254 cardiovascular deaths. As illustrated in Figure 1 and Figure 2, both AHEI-2010 and DII were associated with all-cause mortality. Whatever the model, for each SD increment of AHEI-2010 (9.8 points) a 20% reduced risk of mortality was observed, and for DII, one increment of 1SD (1.3 points) was associated with an increased by 20% of mortality hazard ratio. Results of the analyses in which AHEI-2010 and DII scores were categorized in tertiles are concordant with these findings (results are presented in Online Supplementary Material- Table C). In sex-, age- and ethnicity-adjusted model, participants in the 3rd and 2nd tertiles of AHEI-2010 showed, respectively, 30% and 40% reduced risk of all-cause mortality risk over the 22 years of follow-up. Further adjustment for health behaviors (occupational grade, living alone, smoking habits, physical activity and total energy intake) in model 2 and health status factors (body mass index, antecedent of CVD, use of lipids lowering drugs, HDL cholesterol, hypertension, type 2 diabetes and long-standing illness) in model 3) did not substantially attenuate the association. Online Supplementary Material- Table C shows that cumulative average of AHEI score tertiles remained significantly associated with mortality risk after full adjustment (Model 3) ($HR_{\text{Tertile3 versus Tertile1}} = 0.73$, 95% CI: 0.69 to 0.87; $HR_{\text{Tertile 2 versus Tertile1}} = 0.79$, 95% CI: 0.66 to 0.93 ; $P_{\text{trend}} < 0.001$). Regarding DII, participants in tertile 3 (most pro-inflammatory group) had 47% higher risk of all-cause mortality compared to participants in tertile 1 (most anti-inflammatory group) ($HR_{\text{Tertile3 versus Tertile1}} = 1.47$, 95% CI: 1.19 to 1.82; $P_{\text{trend}} < 0.001$). Regarding the important differences in total energy intake average means across DII tertiles, we carried out an additional analysis to describe the DII –all cause mortality association across total energy intake tertiles. Results (Online Supplementary Material- Table B) showed that the DII was significantly associated with all-cause mortality only among participants in the lowest tertile of total energy intake ($HR_{\text{Tertile3 versus Tertile1}} = 1.72$, 95% CI: 1.08 to 2.72; $P_{\text{trend}} = 0.02$).

Over the 22 years of follow-up, 264 death caused by cardiovascular diseases occurred. Results of the association between cumulative average of AHEI-2010 and DII scores and risk of cardiovascular mortality are

detailed in figures 1 and 2 and in Online Supplementary Material- Table D. Higher AHEI-2010 score was consistently associated with lower risk of cardiovascular mortality whatever the level of adjustment considered (in full adjusted model, HR=0.80, 95% CI: 0.69 to 0.93). Similar results were observed when tertiles of AHEI-2010 and DII were considered (AHEI-2010HR_{Tertile3 versus Tertile1} = 0.68, 95% CI: 0.48 to 0.95; P_{trend} =0.03); and (DIIHR_{Tertile3 versus Tertile1} = 1.46, 95% CI: 1.00 to 2.13; P_{trend} 0.05).

Regarding cancer mortality, 450 death occurred over the follow-up. Figures 1 shows that AHEI-2010 was significantly associated with lower risk of cancer mortality (HR=0.87, 95% CI=0.78 to 0.98). Online Supplementary Material- Table E confirms that in sex-, age- and ethnicity-adjusted models, participants in higher tertiles of AHEI-2010 showed a lower risk of cancer mortality compared to those in the bottom tertile. However further adjustment for health behavior and health status factors substantially attenuate the association. Participants with high DII score showed a significant higher risk of cancer mortality and this association remains statistically significant whatever the way DII was analysed and the level of adjustment (Figure 2 and Online Supplementary Material- Table E).

To our knowledge four other previous studies investigated the DII –mortality risk (all cause, and caused by cardiovascular diseases or by cancer). A full description of these studies is provided in Table 3. DII for the IWHS in the US was calculated from FFQ that was adapted from the 126-item instrument developed by Willett et al, in total there were 27 food parameters that were used for DII calculation and this study had information on supplement use⁽³⁵⁾. DII for SMC in Sweden was calculated using a 96-item food frequency questionnaire from 27 food parameters without supplements⁽³⁶⁾. In the NHANES III study in the US, DII was calculated using 24 hour recall without supplements from 27 food parameters⁽³⁴⁾. In the SUVIMAX study conducted in France, DII was calculated using multiple 24-hour records from 37 food parameters⁽³³⁾. We combine these results with those from the Whitehall II study. We then conducted a meta-analysis on data from the total of 91,520 participants across the five studies. Figures 3, 4 and 5 show positive associations between DII score and all-cause, cardiovascular and cancer mortality, respectively. The pooled hazard ratios from fixed and random effect models were 1.04 (95 % CI: 1.03 to 1.05) and 1.05 (95 % CI: 1.03 to 1.07) for all cause

mortality (28,891 deaths), 1.05 (95 % CI: 1.03 to 1.07) and 1.05 (95 % CI: 1.03 to 1.08) for cardiovascular mortality (10,424 deaths); and 1.05 (95 % CI: 1.03 to 1.07) and 1.06 (95 % CI: 1.02 to 1.09) for cancer mortality (8269 deaths). Thus, no differences were observed in summary estimates between the two methods of meta-analysis.

DISCUSSION

In this prospective cohort study of 7627 British men and women, adherence to healthy diet assessed by the AHEI-2010 was associated with lower risk of all cause, cardiovascular mortality and in a lesser extent to cancer mortality while participants with higher DII– reflecting a consumption of a pro-inflammatory diet– showed higher risk of all-cause, cardiovascular and cancer mortality. Meta-analyses of the DII –mortality risk reported **in this study along** with those from previous cohort studies, consisting of a total of 91,520 participants, confirmed a higher risk of overall, cardiovascular and cancer mortality in individuals with a high DII score.

We observed concordance between the two dietary indices, DII and AHEI-2010. More than 50% of the participants in the healthiest group of AHEI-2010 were in the most anti-inflammatory group and similar results were observed for the least healthy and most pro-inflammatory groups of AHEI-2010 and DII, respectively. This is along expected lines because components, such as vegetables and fruits, which contribute to healthier AHEI-2010 scores, also contribute to anti-inflammatory DII scores. Both adherence to diet high in AHEI-2010 score and diet low in DII scores constitutes healthy diet. We did observe some discordance between the two indices. The importance of dietary features common to all: higher intakes of whole grains, vegetables, fruit and plant-based components which contribute a range of phytochemicals with potential to impact disease risk through different mechanisms working at different stages of cancer initiation and development. Differences between different indices and outcomes may reflect how other items such as fruit juice, green leafy vegetables and low-fat dairy were considered differently by the different indices. The DII is based on the idea that many disease-determining mechanisms operate in a pro-inflammatory environment ⁽³¹⁾, whereas the AHEI 2010 is based on foods and nutrients predictive of chronic disease risk ⁽²⁵⁾. The most anti-inflammatory diets are likely to include foods that feature prominently in other ‘high-quality diets’.

AHEI-2010 was found to predict all cause, cardiovascular and cancer mortality. These results fall in line with a recent meta-analysis conducted by Onvani et al. ⁽³⁸⁾ investigating how diet quality indices as assessed by Healthy Eating, Alternative Healthy Eating Index relate to all-cause and specific causes of mortality (cardiovascular mortality, cancer mortality). Conversely previous studies examining the DII-mortality association produced mixed findings. In the Swedish Mammography study, DII was positively associated with all-cause and digestive cancer mortality but not with overall cancer mortality ⁽³⁶⁾. In the Iowa Women's Health Study, increasing DII was associated with greater risk of all-cause, CVD, digestive tract cancer, coronary heart disease and chronic obstructive pulmonary disease mortalities ⁽²⁴⁾. Among digestive-tract cancers, colorectal cancer (CRC) is known to be strongly related to inflammation ^(39; 40; 41; 42) and represents the majority of digestive-tract cancers ^(20; 43). The DII has been shown to be associated with various digestive tract cancers including CRC ^(31; 44; 45; 46; 47; 48) and with CVDs ^(49; 50; 51). In a couple of other studies, the DII was shown to be associated with increased prostate cancer mortality ^(33; 52). Our meta-analysis including all data available confirm an association between DII, all-cause mortality, CVD mortality and overall cancer mortality. Participants with higher DII score showed lower mean of total energy intake compared to participants with high DII score. Conversely participants in the highest tertile of AHEI-2010 were more likely to have higher total energy intake. One possibility is that the overall higher intake of food (contributing to higher overall energy intake) would encompass many anti-inflammatory dietary components amongst WHI participants who consume higher amounts of food in general.

Although the mechanisms of how a healthy diet reduces mortality are not known, a plausible pathway is that a pro-inflammatory diet increasing systemic inflammation also increases insulin resistance ^(53; 54). Higher consumption of food items such as meat and butter and lower consumption of food items like vegetables and fruits has been shown to increase systemic inflammation by increasing levels of high-sensitivity CRP, E-selectin and soluble vascular cell adhesion molecule-1 ⁽⁵³⁾, which are then responsible for increasing insulin resistance ⁽⁵⁴⁾. Insulin resistance caused by increasing circulating levels of insulin, triglycerides, and non-esterified fatty acids ^(55; 56) is associated with an increased risk of chronic diseases, such as cardiovascular

diseases, which can be fatal. On the other hand, the cardio-protective role of higher scores on the AHEI-2010 could be explained by other biological mediators such as arrhythmia, thrombosis, and insulin resistance ⁽⁵⁷⁾.

Our study has several strengths. First, it is population-based and employs a prospective design. It also, benefits from complete ascertainment of deaths through the National Health Services Central Registry until 31 August 2014, and detailed information on diet. This study also had a long follow-up for the outcomes studied. Although DII has been used to predict mortality in a European; i.e., Swedish cohort ⁽³⁶⁾, this is the first time the DII has been used in a cohort study in Britain whose population and dietary habits are more diverse than in Sweden ^(58; 59). The main limitation of this study was that information on diet was self-reported, which can lead to a potential misclassification of the exposure. Classification errors in our prospective study, however, were non-differential with respect to the occurrence of death and most likely led to an attenuation of the results. Dietary data were available only at one-time point. Participants' dietary habit might have changed during the follow-up period. However, previous studies have reported that dietary pattern classification is moderately stable over long periods of time during adulthood ^(60; 61; 62; 63; 64; 65). Another important limitation of the study includes non-availability of information on the remaining 18 food parameters for DII calculation, Britain with a diverse population which include several Asians availability of information on food parameters like spices would have modified the results.

In conclusion, diet indices –the AHEI-2010 and DII- that possibly reflect pro / anti -inflammatory properties of diet are associated with long-term all-cause, cardiovascular and cancer mortality in the UK Whitehall II study. Future steps might include investigating how these dietary indices are related to chronic inflammation and their biomarkers.

Acknowledgements

Study was supported by grants from the UK Medical Research Council (MRC, K013351); the British Heart Foundation; the British Health and Safety Executive; the British Department of Health; the National Heart, Lung, and Blood Institute (R01HL036310); the National Institute on Aging, NIH (R01AG013196 and R01AG034454); and the Agency for Health Care Policy and Research (grant HS06516). Drs. Shivappa and

Hébert were supported by grant number R44DK103377 from the United States National Institute of Diabetes and Digestive and Kidney Diseases. Prof. Kivimaki was supported by the MRC and NordForsk.

The authors have declared no conflicts of interest.

Dr. James R. Hébert owns controlling interest in Connecting Health Innovations LLC (CHI), a company planning to license the right to his invention of the dietary inflammatory index (DII) from the University of South Carolina in order to develop computer and smart phone applications for patient counseling and dietary intervention in clinical settings. Dr. Nitin Shivappa is an employee of CHI. The subject matter of this paper will not have any direct bearing on that work, nor has that activity exerted any influence on this project.

We thank all of the participating civil service departments and their welfare, personnel, and establishment officers; the British Occupational Health and Safety Agency; the British Council of Civil Service Unions; all participating civil servants in the Whitehall II study; and all members of the Whitehall II study team. The Whitehall II Study team comprises research scientists, statisticians, study coordinators, nurses, data managers, administrative assistants and data entry staff, who make the study possible.

Author contribution: M.K. designed and conducted the study; N. S. calculated the DII scores and wrote the first draft of the manuscript; T.A. performed analyses and worked on the methods and results sections of the manuscript. N. S., J. R. H., M.K. and T.A. provided suggestions and revised the manuscript.

All the authors approved the final version of the manuscript.

Data Sharing: Whitehall II data, protocols, and other metadata are available to the scientific community. Please refer to the Whitehall II data sharing policy at <https://www.ucl.ac.uk/whitehallII/data-sharing>.

REFERENCES

1. Schwingshackl L, Hoffmann G (2015) Diet quality as assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension score, and health outcomes: a systematic review and meta-analysis of cohort studies. *J Acad Nutr Diet* **115**, 780-800.e785. doi: 710.1016/j.jand.2014.1012.1009. Epub 2015 Feb 1011.
2. Ocke MC (2013) Evaluation of methodologies for assessing the overall diet: dietary quality scores and dietary pattern analysis. *Proc Nutr Soc* **72**, 191-199. doi: 110.1017/S0029665113000013. Epub 0029665113002013 Jan 0029665113000030.
3. Jacobs DR, Jr., Steffen LM (2003) Nutrients, foods, and dietary patterns as exposures in research: a framework for food synergy. *Am J Clin Nutr* **78**, 508S-513S.
4. Ouchi N, Parker JL, Lugus JJ *et al.* (2011) Adipokines in inflammation and metabolic disease. *Nature reviews Immunology* **11**, 85-97.
5. Calabro P, Golia E, Yeh ET (2009) CRP and the risk of atherosclerotic events. *Seminars in immunopathology* **31**, 79-94.

6. Miller MA, Cappuccio FP (2007) Inflammation, sleep, obesity and cardiovascular disease. *Current vascular pharmacology* **5**, 93-102.
7. Chung Y-C, Chang Y-F (2003) Serum interleukin-6 levels reflect the disease status of colorectal cancer. *Journal of Surgical Oncology* **83**, 222-226.
8. Terzic J, Grivennikov S, Karin E *et al.* (2010) Inflammation and colon cancer. *Gastroenterology* **138**, 2101-2114 e2105.
9. Toriola AT, Cheng TY, Neuhauser ML *et al.* (2013) Biomarkers of inflammation are associated with colorectal cancer risk in women but are not suitable as early detection markers. *Int J Cancer* **132**, 2648-2658.
10. Lin OS (2009) Acquired risk factors for colorectal cancer. *Methods in molecular biology* **472**, 361-372.
11. Wogan GN, Hecht SS, Felton JS *et al.* (2004) Environmental and chemical carcinogenesis. *Seminars in cancer biology* **14**, 473-486.
12. de Mello VD, Schwab U, Kolehmainen M *et al.* (2011) A diet high in fatty fish, bilberries and wholegrain products improves markers of endothelial function and inflammation in individuals with impaired glucose metabolism in a randomised controlled trial: the Sysdimet study. *Diabetologia* **54**, 2755-2767.
13. Khoo J, Piantadosi C, Duncan R *et al.* (2011) Comparing effects of a low-energy diet and a high-protein low-fat diet on sexual and endothelial function, urinary tract symptoms, and inflammation in obese diabetic men. *J Sex Med* **8**, 2868-2875.
14. Luciano M, Mottus R, Starr JM *et al.* (2012) Depressive symptoms and diet: their effects on prospective inflammation levels in the elderly. *Brain Behav Immun* **26**, 717-720.
15. Chang AR, Lazo M, Appel LJ *et al.* (2014) High dietary phosphorus intake is associated with all-cause mortality: results from NHANES III. *Am J Clin Nutr* **99**, 320-327.
16. Cheung CL, Sahni S, Cheung BM *et al.* (2014) Vitamin K intake and mortality in people with chronic kidney disease from NHANES III. *Clin Nutr*.
17. Cohen HW, Hailpern SM, Alderman MH (2008) Sodium intake and mortality follow-up in the Third National Health and Nutrition Examination Survey (NHANES III). *J Gen Intern Med* **23**, 1297-1302.
18. Deng X, Song Y, Manson JE *et al.* (2013) Magnesium, vitamin D status and mortality: results from US National Health and Nutrition Examination Survey (NHANES) 2001 to 2006 and NHANES III. *BMC Med* **11**, 187.
19. Shivappa N, Steck SE, Hurley TG *et al.* (2014) Designing and developing a literature-derived, population-based dietary inflammatory index. *Public health nutrition* **17**, 1689-1696.
20. Shivappa N, Steck SE, Hurley TG *et al.* (2014) A population-based dietary inflammatory index predicts levels of C-reactive protein in the Seasonal Variation of Blood Cholesterol Study (SEASONS). *Public health nutrition* **17**, 1825-1833.
21. Wirth MD, Burch J, Shivappa N *et al.* (2014) Association of a dietary inflammatory index with inflammatory indices and metabolic syndrome among police officers. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine* **56**, 986-989.
22. Shivappa N, Hebert JR, Rietzschel ER *et al.* (2015) Associations between dietary inflammatory index and inflammatory markers in the Asklepios Study. *The British journal of nutrition* **113**, 665-671.
23. Tabung FK, Steck SE, Zhang J *et al.* (2015) Construct validation of the dietary inflammatory index among postmenopausal women. *Annals of epidemiology* **25**, 398-405.
24. Wood LG, Shivappa N, Berthon BS *et al.* (2015) Dietary inflammatory index is related to asthma risk, lung function and systemic inflammation in asthma. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology* **45**, 177-183.
25. Chiuve SE, Fung TT, Rimm EB *et al.* (2012) Alternative dietary indices both strongly predict risk of chronic disease. *The Journal of nutrition* **142**, 1009-1018.
26. Marmot M, Brunner E (2005) Cohort Profile: the Whitehall II study. *Int J Epidemiol* **34**, 251-256.
27. Akbaraly TN, Ferrie JE, Berr C *et al.* (2011) Alternative Healthy Eating Index and mortality over 18 y of follow-up: results from the Whitehall II cohort. *Am J Clin Nutr*.
28. Brunner E, Stallone D, Juneja M *et al.* (2001) Dietary assessment in Whitehall II: comparison of 7 d diet diary and food-frequency questionnaire and validity against biomarkers. *The British journal of nutrition* **86**, 405-414.
29. Bingham SA, Gill C, Welch A *et al.* (1997) Validation of dietary assessment methods in the UK arm of EPIC using weighed records, and 24-hour urinary nitrogen and potassium and serum vitamin C and carotenoids as biomarkers. *Int J Epidemiol* **26 Suppl 1**, S137-151.
30. Shivappa N, Bosetti C, Zucchetto A *et al.* (2014) Association between dietary inflammatory index and prostate cancer among Italian men. *The British journal of nutrition*, 1-6.

31. Shivappa N, Prizment AE, Blair CK *et al.* (2014) Dietary inflammatory index and risk of colorectal cancer in the Iowa Women's Health Study. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* **23**, 2383-2392.
32. Sabia S, Dugravot A, Kivimaki M *et al.* (2012) Effect of intensity and type of physical activity on mortality: results from the Whitehall II cohort study. *American journal of public health* **102**, 698-704.
33. Graffouillere L, Deschasaux M, Mariotti F *et al.* (2016) Prospective association between the Dietary Inflammatory Index and mortality: modulation by antioxidant supplementation in the SU.VI.MAX randomized controlled trial. *Am J Clin Nutr* **103**, 878-885.
34. Shivappa N, Steck SE, Hussey JR *et al.* (2015) Inflammatory potential of diet and all-cause, cardiovascular, and cancer mortality in National Health and Nutrition Examination Survey III Study. *European journal of nutrition*.
35. Shivappa N, Blair CK, Prizment AE *et al.* (2015) Association between inflammatory potential of diet and mortality in the Iowa Women's Health study. *European journal of nutrition*.
36. Shivappa N, Harris H, Wolk A *et al.* (2015) Association between inflammatory potential of diet and mortality among women in the Swedish Mammography Cohort. *European journal of nutrition*.
37. Guolo A, Varin C (2015) Random-effects meta-analysis: the number of studies matters. *Statistical methods in medical research*.
38. Onvani S, Haghighatdoost F, Surkan PJ *et al.* (2016) Adherence to the Healthy Eating Index and Alternative Healthy Eating Index dietary patterns and mortality from all causes, cardiovascular disease and cancer: a meta-analysis of observational studies. *Journal of human nutrition and dietetics : the official journal of the British Dietetic Association*.
39. Erlinger TP, Platz EA, Rifai N *et al.* (2004) C-reactive protein and the risk of incident colorectal cancer.[see comment]. *JAMA* **291**, 585-590.
40. Gunter MJ, Stolzenberg-Solomon R, Cross AJ *et al.* (2006) A prospective study of serum C-reactive protein and colorectal cancer risk in men. *Cancer Research* **66**, 2483-2487.
41. Otani T, Iwasaki M, Sasazuki S *et al.* (2006) Plasma C-reactive protein and risk of colorectal cancer in a nested case-control study: Japan Public Health Center-based prospective study. *Cancer Epidemiology, Biomarkers & Prevention* **15**, 690-695.
42. Nikiteas NI, Tzanakis N, Gazouli M *et al.* (2005) Serum IL-6, TNFalpha and CRP levels in Greek colorectal cancer patients: prognostic implications. *World Journal of Gastroenterology* **11**, 1639-1643.
43. Herszenyi L, Tulassay Z (2010) Epidemiology of gastrointestinal and liver tumors. *European review for medical and pharmacological sciences* **14**, 249-258.
44. Shivappa N, Bosetti C, Zucchetto A *et al.* (2014) Dietary inflammatory index and risk of pancreatic cancer in an Italian case-control study. *The British journal of nutrition*, 1-7.
45. Lu Y, Shivappa N, Lin Y *et al.* (2015) Diet-related inflammation and oesophageal cancer by histological type: a nationwide case-control study in Sweden. *European journal of nutrition*.
46. Shivappa N, Hebert JR, Rashidkhani B (2015) Dietary Inflammatory Index and Risk of Esophageal Squamous Cell Cancer in a Case-Control Study from Iran. *Nutrition and cancer*, 1-7.
47. Shivappa N, Zucchetto A, Serraino D *et al.* (2015) Dietary inflammatory index and risk of esophageal squamous cell cancer in a case-control study from Italy. *Cancer causes & control : CCC* **26**, 1439-1447.
48. Tabung FK, Steck SE, Ma Y *et al.* (2015) The association between dietary inflammatory index and risk of colorectal cancer among postmenopausal women: results from the Women's Health Initiative. *Cancer causes & control : CCC* **26**, 399-408.
49. O'Neil A, Shivappa N, Jacka FN *et al.* (2015) Pro-inflammatory dietary intake as a risk factor for CVD in men: a 5-year longitudinal study. *The British journal of nutrition*, 1-9.
50. Ramallal R, Toledo E, Martinez-Gonzalez MA *et al.* (2015) Dietary Inflammatory Index and Incidence of Cardiovascular Disease in the SUN Cohort. *PLoS one* **10**, e0135221.
51. Garcia-Arellano A, Ramallal R, Ruiz-Canela M *et al.* (2015) Dietary Inflammatory Index and Incidence of Cardiovascular Disease in the PREDIMED Study. *Nutrients* **7**, 4124-4138.
52. Zucchetto A, Gini A, Shivappa N *et al.* (2016) Dietary inflammatory index and prostate cancer survival. *Int J Cancer* **139**, 2398-2404.
53. Esmailzadeh A, Kimiagar M, Mehrabi Y *et al.* (2007) Dietary Patterns and Markers of Systemic Inflammation among Iranian Women. *The Journal of nutrition* **137**, 992-998.
54. Festa A, D'Agostino R, Howard G *et al.* (2000) Chronic Subclinical Inflammation as Part of the Insulin Resistance Syndrome: The Insulin Resistance Atherosclerosis Study (IRAS). *Circulation* **102**, 42-47.

55. Bruce WR, Wolever TM, Giacca A (2000) Mechanisms linking diet and colorectal cancer: the possible role of insulin resistance. *Nutrition and cancer* **37**, 19-26.
56. Bruce WR, Giacca A, Medline A (2000) Possible mechanisms relating diet and risk of colon cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* **9**, 1271-1279.
57. Neelakantan N, Naidoo N, Koh WP *et al.* (2016) The Alternative Healthy Eating Index Is Associated with a Lower Risk of Fatal and Nonfatal Acute Myocardial Infarction in a Chinese Adult Population. *J Nutr* **146**, 1379-1386.
58. Murakami K, Livingstone MBE (2016) Energy density of meals and snacks in the British diet in relation to overall diet quality, BMI and waist circumference: findings from the National Diet and Nutrition Survey. *Brit J Nutr* **116**, 1479-1489.
59. Lundberg-Hallén N, Öhrvik V (2015) Key foods in Sweden: Identifying high priority foods for future food composition analysis. *Journal of Food Composition and Analysis* **37**, 51-57.
60. Jain M, Howe GR, Harrison L *et al.* (1989) A study of repeatability of dietary data over a seven-year period. *American journal of epidemiology* **129**, 422-429.
61. Jensen OM, Wahrendorf J, Rosenqvist A *et al.* (1984) The reliability of questionnaire-derived historical dietary information and temporal stability of food habits in individuals. *American journal of epidemiology* **120**, 281-290.
62. Lindsted KD, Kuzma JW (1989) Long-term (24-year) recall reliability in cancer cases and controls using a 21-item food frequency questionnaire. *Nutrition and cancer* **12**, 135-149.
63. Mursu J, Steffen LM, Meyer KA *et al.* (2013) Diet quality indexes and mortality in postmenopausal women: the Iowa Women's Health Study. *Am J Clin Nutr* **98**, 444-453.
64. Sijtsma FP, Meyer KA, Steffen LM *et al.* (2012) Longitudinal trends in diet and effects of sex, race, and education on dietary quality score change: the Coronary Artery Risk Development in Young Adults study. *Am J Clin Nutr* **95**, 580-586.
65. Thompson FE, Metzner HL, Lamphiear DE *et al.* (1990) Characteristics of individuals and long term reproducibility of dietary reports: the Tecumseh Diet Methodology Study. *J Clin Epidemiol* **43**, 1169-1178.

TABLES

Table 1 : Characteristics of the 7627 Whitehall II participants according to survival status over 22-y of follow-up.

Characteristics at baseline		All N= 7627	Alive N= 6626	Deceased N= 1001	P value
		(%) or m± sd	%	%	
Socio demographic					
Sex	Men	5308 (69.6)	69.7	68.8	0.57
	Women	2319 (30.4)	30.3	31.2	
Age, years		50.1 ± 6.0	49.5 ± 5.9	53.7 ± 5.8	
Ethnicity	White	6968 (91.4)	91.5	90.1	0.29
	South Asian	405 (5.3)	5.2	5.9	
	Black	254 (3.3)	3.2	4.0	
Occupational Grade	Low	1233 (16.2)	15.3	22.0	<0.001
	Intermediate	3460 (45.4)	45.7	43.2	
	High	2934 (38.5)	39.0	34.9	
Marital status	Married	5856 (76.8)	77.1	74.9	0.13
	cohabited				
	Living alone	1771 (23.2)	22.9	25.1	
Health behaviours					
Total energy intake, kcal/day		2094 ± 634	2096 ± 624	2081 ± 695	0.52
Smoking Status	Non smoker	3844 (50.4)	51.7	41.8	<0.001
	Ex-smoker	2636 (36.6)	34.7	32.9	
	Current smoker	1147 (15.0)	13.6	24.4	
Alcohol intake	None	1460 (19.1)	18.6	22.9	<0.001
	Moderate	4849 (63.6)	64.5	57.2	
	Heavy	1318 (17.3)	16.9	19.9	
Physical activity	Low	1538 (20.2)	19.5	24.5	<0.001
	Intermediate	2156 (28.3)	28.2	28.5	
	High	3933 (51.6)	52.2	47.0	
Health Status					
Body mass index, kg/m ²		25.3 ± 3.7	25.2 ± 3.6	26.0 ± 4.2	
Hypertension	No	6168 (80.9)	82.3	71.2	<0.001
	Yes	1459 (19.1)	17.7	28.9	
Type 2 diabetes	No	7390(96.9)	97.3	93.9	<0.001
	Yes	237 (3.1)	2.7	6.1	
Antecedent of CVD	No	7390 (96.9)	97.5	92.8	<0.001
	Yes	237 (3.1)	2.5	7.2	
HDL cholesterol, mmol/L		1.43 ± 0.41	1.44 ±0.41	1.38 ±0.42	<0.001
Use of lipids lowering drugs	No	7571 (99.3)	99.3	99.3	0.88
	Yes	56 (0.73)	0.74	0.70	
Longstanding illness	No	5053 (66.2)	67.3	59.3	<0.001
	Yes	2574 (33.7)	32.7	40.7	
Dietary exposure					
AHEI-2010	T1	2640 (34.6)	33.5	41.8	<0.001
	T2	2332 (30.6)	30.9	28.1	
	T3	2655 (34.8)	35.5	30.2	

DII	T1	2542 (33.3)	33.6	31.4	<0.001
	T2	2543 (33.3)	33.8	30.3	
	T3	2542 (33.3)	32.6	38.4	

CVD: Cardiovascular disease; AHEI-2010: Alternative Healthy Eating Index 2010; DII: dietary inflammatory Index; T : Tertile

Occupational position with the use of current (or last for retired participants) British civil service employment grade, was defined on the basis of salary and grouped into 3 categories: high (senior administrators) / intermediate (executives, professionals, and technical staff) / low (clerical and office support staff) grades.

Alcohol intake categories were defined as none : 0 glass per day ; moderate:< 2 (3) glasses per day for women (men); and heavy consumption : ≥ 2 (3) glasses per day for women (men).

Physical activity was categorized into 3 groups (high, intermediate, low) according to frequency of participation in “vigorous” (eg, running, hard swimming, playing squash), “moderately energetic” (eg, dancing, cycling, leisurely swimming), and “mildly energetic” physical activity.

Table 2: Relationship between Alternative Healthy Eating Index 2010 (AHEI-2010) and Dietary Inflammatory Index (DII)

AHEI-2010 tertiles	DII tertiles		
	Tertile 3	Tertile 2	Tertile 1
	(Most pro-inflammatory)		(Most anti-inflammatory)
Tertile 1 (Unhealthy diet)	1319 (51.9)	860 (33.8)	461 (18.1)
Tertile 2	809 (31.8)	825 (32.4)	698 (27.5)
Tertile 3 (Healthiest diet)	414 (16.2)	858 (33.7)	1383 (54.4)

Table 3: Characteristics of included studies									
	Sample size	Follow up (years)	Dietary Assessment tool used	Number of food parameters used for DII calculation	Mean and standard deviation of DII	DII included information from supplements	Assessment of mortality	Total number of deaths	Adjustment
Whitehall II	7622	22	127 item FFQ	27	0.37 ±1.41	No	Deaths were identified by linkage to the National Health Services death and electronic patient records with the use of the National Health Services identification number assigned to all British citizens	1001	adjusted for age, sex, ethnicity, occupational grade, living alone, smoking habits, alcohol consumption, physical activity, total energy intake, body mass index, antecedent of CVD, use of lipids lowering drugs, HDL cholesterol, hypertension, type 2 diabetes and longstanding illness
NHANES III	12366	18	24-Hour recalls	27	0.73±2.20	No	death were identified through	2795	adjustment was made for age, BMI,

							linkage to National Death Index records through December 31, 2006, by the National Center for Health Statistics		smoking status, sex, race/ethnicity, diabetes status, hypertension status, physical activity, medical history of CVD, and poverty index (PI).
Swedish Mammography Cohort	33747	15	96 item FFQ	27	0.64 ± 1.45	No	death were identified through linkage to the Swedish Cause of Death Registry at National Board of Health and Welfare.	7095	Adjusted for age, energy intake, BMI , education, smoking status, physical activity, and alcohol intake
Iowa Women's Health Study	37525	24	adapted from the 126-item FFQ developed by Willett and group at Harvard	27	-0.87± 2.02	Yes	Deaths were identified through the State Health Registry of Iowa or the National Death Index for women who did not respond to the last	17793	Adjusted for age , BMI , smoking status, pack-years of smoking, HRT use , education, prevalent diabetes, prevalent hypertension,

							follow-up questionnaire (2004) or who emigrated from Iowa.		prevalent heart disease, prevalent cancer, total energy intake
SUVIMAX	8089	13	24-Hour dietary records	37	0.6 ± 1.7 for women and 0.8 ± 1.5 for men	No	Death were identified from relatives or physicians during follow-up. At the end of follow-up, vital status of all subjects of the cohort and causes of death were verified with the national death registry (CépiDC).	207	Adjusted for age , sex, intervention group of the initial SUVIMAX trial, number of 24-h dietary records, BMI, physical activity, smoking status, educational level, family history of cancer in first-degree relatives, family history of CVD in first-degree relatives, energy intake without alcohol, and alcohol intake