



van der Nelson, H. A., Draycott, T., Siassakos, D., Yau, C. W. H., & Hatswell, A. J. (2017). Carbetocin versus oxytocin for prevention of postpartum haemorrhage at caesarean section in the United Kingdom: An economic impact analysis. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, *210*, 286-291. https://doi.org/10.1016/j.ejogrb.2017.01.004

Peer reviewed version

License (if available): CC BY-NC-ND

Link to published version (if available): 10.1016/j.ejogrb.2017.01.004

Link to publication record in Explore Bristol Research PDF-document

University of Bristol - Explore Bristol Research General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: http://www.bristol.ac.uk/pure/about/ebr-terms

Table 4: Overall cost as a result of using either oxytocin or carbetocin for prevention of postpartum haemorrhage at caesarean section

	Oxytocin	Carbetocin	Change with Carbetocin
PPH 500- 1000ml	65	43	-22
events			
PPH 1000- 1500ml	13	9	-5
events			
PPH >1500ml events	10	6	-3
PPH Events	88	58	-30
	Oxytocin	Carbetocin	Change with carbetocin
PPH Events	£105,227	£69,242	-£35,985
PPH prophylaxis	£3,600	£26,460	£22,860
PPH re-treatment	£3,600	£2,250	-£1,350
Ante-Natal	£124,312	£124,312	£0
Recovery	£1,774,419	£1,761,637	-£12,783
Follow up	£102,351	£102,090	-£261
Total cost	£2,113,508	£2,085,990	-£27,518