

1 **Children's compliance with wrist worn accelerometry within a cluster randomised controlled**  
2 **trial: Findings from The Healthy Lifestyles Programme (HeLP)**

3

4

5 **Running heading:** Children's Accelerometry Compliance within an RCT

6

7

8

9

10

11

12

13

14

15

16

17

18

Accepted

19 **Abstract**

20 **Purpose:** To assess children's compliance with wrist worn accelerometry during a randomised  
21 control trial and to examine whether compliance differed by allocated condition or gender.

22 **Method:** 886 children within the Healthy Lifestyles Programme (HeLP) trial were randomly allocated  
23 to wear a GENEActiv accelerometer at baseline and 18 month follow up. Compliance with minimum  
24 wear time criteria ( $\geq 10$  hours for 3 week, 1 weekend day) was obtained for both time points. Chi-  
25 squared tests were used to determine associations between compliance, group allocation and gender.

26 **Results:** At baseline, 851 children had useable data, 830 (97.5%) met the minimum wear time criteria,  
27 631 (74.1%) had data for 7 days at 24 hours/day. At follow up, 789 children had useable data, 745  
28 (94.4%) met the minimum wear time criteria, 528 (67%) children had complete data. Compliance did  
29 not differ by gender (baseline;  $X_2 = 1.66$ ,  $p = 0.2$ , follow up;  $X_2 = 0.76$ ,  $p = 0.4$ ) or by group at follow  
30 up ( $X_2 = 2.35$ ,  $p = 0.13$ ).

31 **Conclusion:** The use of wrist worn accelerometers and robust trial procedures resulted in high  
32 compliance at two time points regardless of group allocation, demonstrating the feasibility of using  
33 precise physical activity monitors to measure intervention effectiveness.

34 **Trial registration:** ISRCTN 15811706

35

36

37

38

39

## 40 **Background**

41 Assessing children's physical activity (PA) using accelerometry is now common place in cohort  
42 studies (1, 2, 3) and randomised control trials.(4, 5) However, researchers still face challenges  
43 regarding choice of minimum wear time criteria and participant compliance (i.e. those who meet or  
44 exceed the minimum wear criteria), which can substantially affect interpretation of results. Setting a  
45 high wear time threshold for inclusion in data analysis tends to improve the precision of PA estimates  
46 (6) but often substantially increases the number of data files that have to be excluded from analyses  
47 due to missing data. This can result in selection bias, as the sample retained may differ on exposure to  
48 the intervention in a clinical trial, the outcome variable or other important covariates.(7, 8) Hence it is  
49 desirable to maximise both the retained sample size and the accelerometer wear time period. Recent  
50 developments in the design of activity monitors and wear protocols have sought to address these two  
51 methodological challenges.(9)

52  
53 Evidence demonstrates that the use of a waterproof, wrist worn accelerometer, designed to be discrete  
54 and minimise discomfort, can reduce periods of non-wear in adults (10) with similar high compliance  
55 demonstrated in small samples of children. (11) This, in turn, reduces the need for statistical  
56 imputation methods, assumptions regarding missing values (7, 12) and the associated risks of  
57 selection bias and misclassification of PA. In addition, there is evidence that implementing a 24 hour  
58 wear protocol, albeit with waist worn devices, rather than the more commonly used 'waking hours  
59 only' protocol can also increase wear time compliance. (9) It would be expected that the combination  
60 of increasing both comfort/convenience/waterproofing and manipulating the wear time protocol  
61 should yield higher compliance, in turn leading to more precise estimates of PA across the entire  
62 week.

63  
64 Despite evidence of high compliance at single measurement points (11) with wrist worn devices,  
65 evidence regarding compliance over multiple measure periods is limited. Assessing compliance over  
66 multiple measures is of particular importance in determining the effectiveness of behavioural

67 interventions in randomised trials. Low compliance with the pre specified accelerometer wear time  
68 can exacerbate any loss to follow up; participants may complete all other trial outcomes but are  
69 treated as lost to follow up for PA outcomes due to their failure to meet the minimum accelerometer  
70 wear time criteria. Howie and colleagues (13) demonstrated a reduction from 89% to 79% compliance  
71 with wrist worn devices over 4 measurement points using 10 hours for four days criteria. The authors  
72 did not take into consideration the compliance with extended wear time that may be achievable with  
73 wrist worn devices, nor whether compliance differed across intervention arm. Despite randomisation  
74 prior to participants being allocated to intervention or control groups, if non-compliance with wear  
75 time is systematically different by group allocation, bias may be introduced.

76

77 The aim of this study was to examine children's compliance with a wrist worn accelerometer at  
78 baseline and 18 month follow up within a cluster randomised controlled trial, (5) using both  
79 traditional and extended wear time criteria. Secondly, the study aimed to examine whether compliance  
80 with follow up measures differed by group allocation (i.e. intervention vs. control groups), and  
81 whether compliance was associated with gender.

82

### 83 **Methods**

#### 84 *Participants*

85 Data from the present study were obtained as part of the Healthy Lifestyles Programme (HeLP), a  
86 school based, cluster randomised control trial of a novel obesity prevention programme. (5) The trial  
87 involved 32 schools and 53 classes of Year 5 children (aged 9-10 years) across Devon, UK. One Year  
88 5 class from each participating school was randomly selected to receive an accelerometer at baseline  
89 (n=886). Data were collected in two phases, with 16 schools in each cohort. Baseline physical activity  
90 data were collected in October 2012 and 2013 for Cohort 1 and Cohort 2 respectively. Schools were  
91 then randomised to receive HeLP (5) or to the control arm (usual practice). Full details of recruitment  
92 and study procedures are provided elsewhere (5, 14)

93 (<http://clahrc-peninsula.nihr.ac.uk/research/help-the-healthy-lifestyles-programme>). Follow up PA  
94 data were collected 18 months post baseline, in June 2014 and 2015 for Cohorts 1 and 2, respectively.  
95 Ethical approval for the trial was obtained from the Peninsula College of Medicine and Dentistry in  
96 March 2012 (reference number 12/03/140).

#### 97 *Physical activity measurement*

98 Physical activity was assessed using a GENEActiv (ActivInsights Ltd, Kimbolton, UK) tri-axial  
99 accelerometer, measuring 43mm x 40mm x 13mm. The GENEActiv was attached to a polyurethane  
100 strap, and worn at the wrist, like a watch. It can measure between +/- 8g at a rate of up to 100Hz.  
101 During the present study, data were collected at a rate of 85.7Hz.

#### 102 *Anthropometric measures*

103 Children's height was measured using a SECA (hamburg, Germany) stadiometer and recorded to the  
104 nearest 0.1cm. Weight was measured using a Tanita Body Composition Analyser SC-330 (U.K ltd.,  
105 Middlesex UK) and recorded to the nearest 0.1kg. BMI sds were calculated using the Cole (15) BMI  
106 reference curves for children. Waist circumference was measured 4cm above the umbilicus using a  
107 flexible (non-elastic) tape measure.

#### 108 *Protocol*

109 Prior to distributing the monitors, parents received a reminder letter about the date the GENEActiv  
110 would be given to children and the date of removal. Monitors were distributed by HeLP co-ordinators  
111 to small groups (~10 per group) of children at a time. Participants were informed about the monitor  
112 placement and were asked to wear the monitors on their non-dominant wrist, continuously for a  
113 period of eight days, which included one familiarisation day. During these sessions, each child was  
114 provided with an information pack, including reminder sheets to display at home, and letters to  
115 distribute to sport coaches to prevent removal during extracurricular activities, alongside dates for

116 monitor collection. The devices were collected by HeLP co-ordinators, with follow up visits made to  
117 collect any not returned on the planned collection day.

118 Anthropometric data was collected in a private room by two trained and blinded assessors during a  
119 specifically designed lesson relating to measurement (14).

#### 120 *Data analysis*

121 Data were downloaded using GENEActiv PC software version 1.4 and analysed using the GGIR  
122 software (16, 17, 18) package for R (*cran.r-project.org*). Data processing included auto calibration  
123 using local gravity as a reference (16) and the detection of abnormally high values (16, 19). The raw  
124 values from each axis are used to create a vector magnitude ( $\sqrt{x^2 + y^2 + z^2} - 1g$ ) with negative  
125 values rounded to zero (20) creating the Euclidean Norm minus one (ENMO; measured in milli-g(mg)  
126 units) as reported elsewhere (16, 20.) Data were averaged over 1 second epochs, with the first and  
127 final 30 minutes removed from analysis, in order to minimise inclusion of spurious data at the  
128 beginning and end of data capture. Non-wear time was apparent if the standard deviation of two axes  
129 was less than 13mg and the value range was less than 50mg. Non-wear was assessed over 60 minute  
130 windows, using moving increments of 15 minutes. (2, 16) Time spent in different PA intensities were  
131 estimated using published accelerometer cut-points. (21)

132 Compliance was established for the minimum wear criteria of  $\geq 10$  hours for  $\geq 3$  week and 1 weekend  
133 day (22) at baseline and 18 month follow up. For data collected at baseline, a compliance matrix was  
134 created to report the number and percentage of children meeting multiple valid hours / day  
135 combinations. In order to assess compliance with valid hour/day combinations at the 18 month follow  
136 up, a further compliance matrix was created which only included those children who met the  
137 minimum wear time criteria at baseline, allowing a more thorough examination of any potential  
138 impact of non-compliance on the overall loss to follow up within the trial. Compliance with minimum  
139 wear criteria at baseline and follow up was also reported by gender for both time-points.

140 Pearson's Chi-squared test was used to assess whether compliance with minimum wear time was  
141 associated with group allocation (intervention vs. control) at the 18 month follow up. Only  
142 participants who met the minimum wear time criteria at baseline were included in this analysis.  
143 Sensitivity analysis using all available data from the 18 month follow up, irrespective of baseline  
144 compliance was also undertaken.

## 145 **Results**

146 Characteristics of the 886 participants (423 male; 463 female) allocated to receive accelerometry at  
147 baseline and 18 month follow up are outlined in table 1.

### 148 *Baseline*

149 Of the 886 participants; 851 had useable data (n = 409 male); missing data (n=35) were a result of  
150 monitor failure (including calibration error), or participant absence during the measurement period;  
151 shown in figure 1. Of those with useable data, 830 (97.5%) children met the minimum wear time  
152 criteria of  $\geq 10$  hours for 3 weekdays and 1 weekend day. Table 2 shows the number and percentage of  
153 children complying with varying combinations of valid hours / days. When split by gender, 96.8%  
154 (396/409) of males and 98.1% (434/442) of females met the minimum wear time criteria, there was no  
155 significant association between gender and compliance at baseline ( $X_2 = 1.66$ ,  $p = 0.2$ ).

156 Table 1. Anthropometric and physical activity characteristics at baseline and 18 months

	Baseline			18 month follow up		
	Mean (SD)			Mean (SD)		
	Whole cohort	Intervention	control	Whole cohort	Intervention	control
n	886	428	458	861	412	449
Gender (n male)	423	208	215	411	200	211
age (years)	9.7 (0.3)	9.8 (0.3)	9.7 (0.3)	11.3 (0.3)	11.3 (0.3)	11.3 (0.3)
height (cm)	138.3 (6.8)	138.7 (6.9)	137.8 (6.7)	147.7 (7.6)	148.2 (7.7)	147.2 (7.6)
weight (kg)	33.6 (7.5)	34.3 (8.1)	33.0 (7.0)	40.5 (9.8)	41.3 (10.4)	39.8 (9.1)
BMI sds <sup>a</sup>	0.19 (1.2)	0.27 (1.2)	0.11 (1.1)	0.19 (1.2)	0.27 (1.3)	0.12 (1.2)
waist circumference (cm)	61.0 (7.4)	61.7 (7.8)	60.4 (7.0)	64.1 (8.4)	64.6 (9.0)	63.7 (7.9)
<b>Physical Activity characteristics <sup>b</sup></b>						
n	830	408	422	745	359	386
ENMO (mg)	49.3 (11.1)	49.0 (11.3)	49.6 (10.9)	51.8 (13.4)	52.1 (14.0)	51.5 (13.0)
Total PA <sup>c</sup> (minutes)	183.9 (35.7)	182.7 (36.7)	185.0 (34.7)	198.9 (42.0)	199.7 (43.9)	198.1 (40.2)
Light PA (minutes)	130.3 (24.4)	129.4 (24.7)	131.1 (24.2)	141.3 (27.4)	141.7 (27.8)	141.1 (27.1)
Moderate PA (minutes)	40.2 (11.7)	40.0 (12.1)	40.4 (11.4)	43.8 (14.8)	44.3 (16.2)	43.5 (13.4)



Vigorous PA (minutes)	13.42 (6.2)	13.3 (6.2)	13.5 (6.2)	13.6 (7.5)	13.7 (7.7)	13.5 (7.4)
MVPA (minutes)	53.6 (16.5)	53.3 (16.8)	53.9 (16.2)	57.5 (20.9)	58.0 (22.3)	57.0 (19.4)

---

*PA - physical activity; <sup>a</sup>BMI sds calculated using Standard Deviation Scores were derived for body mass index (BMI), based on the UK 1990 BMI reference curves for children [15] <sup>b</sup>Physical activity characteristics for those who met the minimum inclusion criteria .<sup>c</sup> Total physical activity includes time in light, moderate and vigorous PA.*

157

158

159

160

161

162

163

164

165

166 Table 2. Number (percentage) of children achieving different wear time combinations (days / hours) at baseline

	8 hours	10 hours	12 hours	14 hours	16 hours	18 hours	20 hours	22 hours	24 hours
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
7 days	812 (95.4)	807 (94.8)	803 (94.4)	800 (94.0)	789 (92.7)	759 (89.2)	751 (88.2)	742 (87.2)	631 (74.1)
6 days	819 (96.2)	818 (96.1)	815 (95.8)	813 (95.5)	808 (94.9)	801 (94.1)	797 (93.7)	791 (92.9)	765 (89.9)
5 days	826 (97.1)	825 (96.9)	824 (96.8)	822 (96.6)	819 (96.2)	814 (95.7)	813 (95.5)	808 (94.9)	802 (94.2)
4 days	840 (98.7)	832 (97.8)	830 (97.5)	830 (97.5)	825 (96.9)	817 (96.0)	816 (95.9)	814 (95.7)	811 (95.3)
3 days	847 (99.5)	843 (99.0)	839 (98.6)	839 (98.6)	835 (98.1)	828 (97.3)	826 (97.1)	824 (96.8)	822 (96.6)
2 days	849 (99.8)	848 (99.6)	847 (99.5)	846 (99.4)	840 (98.7)	838 (98.5)	836 (98.2)	833 (97.9)	829 (97.4)
1 day	851 (100)	851 (100)	851 (100)	851 (100)	850 (99.9)	844 (99.2)	843 (99.1)	842 (98.9)	839 (98.6)

167

168

169 *Follow up*

170 At follow up (18 months), 25 children had moved out of area, resulting in 861 children potentially  
171 available for follow up measures. Their characteristics are presented in Table 1. Of these 861 children,  
172 789 (91.6%) had useable accelerometer data, and 745 (94.4%) met the minimum wear time criteria;  
173 528 children (67%) achieved 24 hours for 7 days. When split by gender, 93.7% of males (356/380)  
174 and 95.1% (389/409) of females met the minimum wear criteria, no significant difference in  
175 compliance between gender was apparent at follow up ( $X_2=0.76$ ,  $p = 0.38$ ).

176 When considering only those participants who had valid baseline data ( $n=830$ ), 746 were potentially  
177 available for follow up; of the 84 children who were not available for follow up measures, 22 had  
178 moved out of area, 12 children were absent during the testing period, 4 children failed to return the  
179 device, 45 monitors failed or had calibration error and 1 child developed a rash and stopped wearing  
180 the accelerometer. Of the original 886 children randomised to participate in physical activity data  
181 collection, 705 (79.5%) met the minimum wear time criteria at both baseline and follow up. Analysis  
182 by gender showed 79.9% (338/423) of males and 79.2% (367/463) of females meeting the minimum  
183 wear criteria at both time points. Table 3 presents compliance for combinations of days and hours at  
184 18 months for only those children who had valid baseline data.

185 For the 830 participants who had valid baseline data and were followed up at 18 months ( $n=705$ ),  
186 Pearson's chi-squared showed no evidence of a statistical association between allocated group and  
187 compliance with minimum accelerometer wear time criteria at follow up; 6.8 % ( $n=25$ ) in the  
188 intervention arm and 4.2 % ( $n=16$ ) of children in the control arm did not meet minimum valid day  
189 criteria ( $X_2 = 2.35$ ,  $p = 0.13$ ). Sensitivity analysis using all available data from the 18 month follow  
190 up ( $n=789$ ) also showed no association between group allocation and compliance ( $X_2 = 1.24$ ,  $p =$   
191 0.27).

192

193 Table 3. Number (%) of children achieving wear time combinations (days / hours) at follow up\*

	8 hours	10 hours	12 hours	14 hours	16 hours	18 hours	20 hours	22 hours	24 hours
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
7 days	669 (89.7)	663 (88.9)	660 (88.5)	651 (87.3)	642 (86.1)	607 (81.4)	592 (79.4)	575 (77.1)	499 (66.9)
6 days	690 (92.5)	687 (92.1)	683 (91.6)	675 (90.5)	663 (88.9)	650 (87.1)	646 (86.6)	632 (84.7)	613 (82.2)
5 days	707 (94.7)	702 (94.1)	697 (93.4)	693 (92.9)	686 (92.0)	675 (90.5)	669 (89.7)	664 (89.0)	654 (87.7)
4 days	721 (96.7)	719 (96.4)	714 (95.7)	713 (95.6)	706 (94.6)	696 (93.3)	689 (92.4)	680 (91.2)	674 (90.3)
3 days	731 (98.0)	729 (97.7)	725 (97.2)	721 (96.6)	715 (95.8)	707 (94.8)	703 (94.2)	696 (93.3)	691 (92.6)
2 days	744 (99.7)	744 (99.7)	742 (99.5)	740 (99.2)	734 (98.4)	729 (97.7)	726 (97.3)	717(96.1)	709 (95.0)
1 day	746 (100)	746 (100)	746 (100)	745 (99.9)	741 (99.3)	737 (98.8)	735 (89.5)	733 (98.3)	726 (97.3)

194 \*only children with valid wear time at baseline are included in table 3.

195

196

197

198

## 199 Discussion

200 The primary aim of this study was to examine children's compliance with accelerometer wear time  
201 over two measurement points during a randomised controlled trial. The secondary aim was to assess  
202 whether compliance differed by group (intervention/control) allocation and gender. The results  
203 demonstrate high compliance with wrist worn accelerometry at both baseline (97.5%) and 18 month  
204 follow up (94.4%), with equally high compliance demonstrated by both males and females. Moreover,  
205 high rates of compliance were also apparent when assessing whether minimum wear time criteria was  
206 met at both time points; 705/886 (82.8%) children had  $\geq 10$  hours of wear time for  $\geq 4$  days (including  
207 1 weekend day) at both baseline and 18 months. Chi squared tests showed no association between  
208 gender and compliance (males vs females) at baseline and follow up. Nor were there associations with  
209 group allocation (intervention v control) and compliance with minimum wear time at 18 month follow  
210 up. It appears, therefore, that constant wear, avoiding having to remember to put on or activate an  
211 accelerometer, using a watch-like wrist-worn device is acceptable to children and, consequently,  
212 facilitates reliable data collection.

213

214 These findings demonstrate that high levels of compliance at multiple time points can be obtained by  
215 combining the use of wrist worn, waterproof accelerometers and a 24 hour wear time protocol within  
216 a cluster randomised controlled trial in 9-11 year old children. Beyond meeting minimum wear time  
217 requirements, extended periods of wear can be achieved; providing more accurate estimates of PA, as  
218 the possibility of under or over estimating PA based only on capturing small portions of the day (23)  
219 or a limited number of days (24) is reduced. This may be particularly important in children's activity  
220 measurement due to the variation in their activity over the day.(25) In addition, very few children  
221 were lost due to non-wear, further reducing the impact of missing data and possible selection bias.  
222 Baseline compliance in the present study is slightly higher than rates previously reported in samples  
223 of children with wrist worn devices; Fairclough et al. (11) reported 89% compliance with  $\geq 10$  hours  
224 for 3 week days and 1 weekend day. Additionally, compliance in the HeLP trial compares favourably  
225 to data collected with similar populations within large cohort studies using waist worn monitors. (1, 3,  
226 26, 27, 28). For example the Millennium cohort study reported 67% of children complying with  $\geq 10$

227 hours on at least 2 days (3); compliance rates within HeLP for the same criteria were 99.6%. It is  
228 likely that higher compliance rates within HeLP are due to monitor placement and trial procedures, as  
229 many studies using waist worn devices employ a 'waking time only' protocol (9).

230 Direct comparison of compliance rates between studies is challenging as the method of detecting non-  
231 wear can affect estimates of compliance even when the definition of compliance is the same. In the  
232 present study the method of detecting non-wear is based on non-wear algorithms that use the raw  
233 acceleration values from all three accelerometer axis (16). Arguably this method is more likely to  
234 accurately classify non-wear compared to methods based simply on extended periods of consecutive  
235 '0' counts (29). It is not clear whether the latter method leads to a greater under or overestimate of  
236 non-wear compared to methods based on raw acceleration.

237 Assessing the compliance at 18 month follow up using two methods a) as an independent time point  
238 and b) by considering rate of compliance at follow up with only those children who had provided  
239 'valid' baseline data, allows for a more in depth view of how non-compliance with accelerometer  
240 wear may impact on loss to follow up within large scale trials of behavioural interventions. When the  
241 18 month follow up time point is treated independently, compliance with minimum wear is similar to  
242 that observed at baseline (94.4%). Yet considering the rate of compliance across both time points  
243 provides important information for planning future trials; these results indicate that high compliance  
244 with minimum wear can be achieved at both baseline and follow up, with 79.5% of the original  
245 sample having valid data at both time points. Results indicate that the percentage of participants  
246 treated as 'lost' due to accelerometer non-compliance is low when using a combination of wrist worn  
247 devices, a 24 hour wear protocol and comprehensive trial procedures. These results are encouraging  
248 for future trials as previous studies reported a large drop in compliance with a minimum wear time of  
249  $\geq 10$  hours for  $\geq 3$  days between two time points (from 75% to 56%) when using waist worn devices.  
250 (30)

251

252 Previously, trials of behavioural interventions assessing physical activity with accelerometers have  
253 reported lower compliance in the control group, (31) risking systematic missing data and selection  
254 bias. Results from the HeLP trial show that it is possible to achieve very high compliance in both

255 allocated groups (intervention and control); possibly due to the cluster nature of the trial. In turn this  
256 allows greater sensitivity to detect potential intervention effects (32) and limited loss to follow up due  
257 to missing data. As a result, more precise physical activity estimates are possible, as is the capacity to  
258 detect small changes. Consequently, future studies may benefit from a reduction in required  
259 recruitment targets. (32)

260

261 Estimates of PA are reported to differ by gender in childhood, with males accumulating more MVPA  
262 than females (27). It is important to ensure that any observed differences are a result of actual  
263 behaviour rather than a consequence of systematic error resulting from differences in wear time  
264 compliance between genders. The present study demonstrated no association between gender and  
265 compliance with minimum wear time criteria at either measurement point. However it is important  
266 that differences in compliance are assessed prior to concluding whether are behavioural differences  
267 exist. (32).

268

269 Whilst providing important findings regarding compliance, the limitations of this study that arise from  
270 device failure should be highlighted. A substantial number of participants' data were lost as a result of  
271 device failure; this was particularly noticeable at the 18 month follow up assessment, where data from  
272 48 participants were not able to be recovered from the device, increasing missing data at follow up.  
273 The device failure appeared to be a result of battery failure over time; future studies should take into  
274 consideration the life span of these devices during the study design and procurement phases.

275

276 Using a combination of a waterproof, wrist worn accelerometer and a 24 hour wear protocol means  
277 no conclusions can be made as to which factor or which combination of factors were most important  
278 in increasing compliance; previously Tudor-Locke et al. (9) demonstrated that increased compliance  
279 and wear time with waist-worn devices can be increased using a 24 hour protocol, rather than a wake  
280 time only protocol. Alternatively, Fairclough et al. (11) demonstrated higher compliance with wrist  
281 placement rather than waist-worn devices. It is clear, however, that combination of the two  
282 approaches and robust trial protocols provide the best compliance with accelerometer wear time.

283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307

## **Conclusion**

High compliance with accelerometer wear time protocols can be achieved with children participating in a cluster randomised controlled trial at both baseline and follow up and does not differ by group (intervention/control) allocation. Constant wear of waterproof, wrist worn accelerometers alongside robust trial procedures should be utilised in physical activity research to minimise the number of children with missing data at follow up through non-compliance.

## **Acknowledgements**

This research was supported by a grant PHR project 10/3010/01 from the National Institute for Health Research (NIHR) Public health research programme. In addition, this research was supported and funded by the NIHR Collaboration for Leadership in Applied Health Research and Care South West Peninsula (PenCLAHRC). The views expressed are those of the authors and not necessarily those of the NHS, NIHR or the Department of Health.

The authors would like to acknowledge the Healthy lifestyle co-ordinators for facilitating data collection; Jackie Minton, Helen Eke, Chloe Thomas and Camilla McHugh from the University of Exeter Medical School, and Alison Hurst, Data Manager for the Healthy Lifestyles programme based in the University of Exeter Medical school.



308 **References**

- 309 1. Booth JN, Leary SD, Joinson C, Ness AR, Tomporowski, PD, Boyle JM, Reilly, JJ. Associations  
310 between objectively measured physical activity and academic attainment in adolescents from a  
311 UK cohort. *Br J sports Med.* 2013; doi:10.1136/bjsports-2013-092334
- 312 2. Da Silva ICM, Van Hees VT, Ramires V et al. Physical activity levels in three Brazilian birth  
313 cohorts as assessed with raw triaxial wrist accelerometry. *Int J Epidemiol.* 2014; 43: 1959-1968.  
314 doi: 10.1093/ije/dyu203
- 315 3. Griffiths LJ, Cortina-Borja M, Sera F, et al. How active are our children? Findings from the  
316 Millennium Cohort Study. *BMJ Open.* 2013;3:e002893. doi: 10.1136/bmjopen-2013-002893
- 317 4. Jago R, Edwards MJ, Sebire SJ, et al. Bristol girls dance project (BGDP): protocol for a cluster  
318 randomised controlled trial of an after-school dance programme to increase physical activity  
319 among 11–12 year old girls. *BMC Public Health* . 2013; 13:1003
- 320 5. Wyatt K, Lloyd J, Abraham C, et al. The Healthy Lifestyles Programme (HeLP), a novel school-  
321 based intervention to prevent obesity in school children: study protocol for a randomised  
322 controlled trial. *Trials.* 2013; 14:95
- 323 6. Herrmann SD, Barreira TV, Kang M, Ainsworth BE. Impact of accelerometer wear time on  
324 physical activity data: a NHANES semisimulation data approach. *Br J Sports Med.* 2012; 10: 742-  
325 749. doi:10.1136/bjsports-2012-091410
- 326 7. Toftager M, Kristensen PL, Oliver M et al. Accelerometer data reduction in adolescents: effects  
327 on sample retention and bias. *Int J Behav Nutr Phys Act.* 2013;10:140. doi: 10.1186/1479-5868-  
328 10-140
- 329 8. Kang M, Rowe DA, Barreira TV, Robinson TS, Mahar MT. Individual information-centred  
330 approach for handling physical activity missing data. *Res Q Exerc Sport.* 2009; 80: 131 – 137.
- 331 9. Tudor-Locke C, Barreira TV, Schuna Jr JM, et al. Improving wear time compliance with a 24-  
332 hour waist worn accelerometer protocol in the international study of childhood obesity lifestyle  
333 and environment (ISCOLE). *Int J behave Nutr Phys Act.* 2015; 12:11 doi 10.1186/s12966-015-  
334 0172-x

- 335 10. Sabia S, Van Hees VT, Shipley MJ et al. Associations between questionnaire- and accelerometer-  
336 assessed physical activity: the role of sociodemographic factors. *Am J Epidemiol.* 2014; 179: 781  
337 – 790.
- 338 11. Fairclough SJ, Noonan R, Rowlands AV, Van Hees V, Knowles Z, Boddy LM. Wear compliance  
339 and activity in children wearing wrist and hip mounted accelerometers. *Med Sci Sports Exerc.*  
340 2016; 48: 245-53.
- 341 12. Montori VM, Guyatt GH. Intention-to-treat principle. *CMAJ.* 2001; 165:1339 – 1341.
- 342 13. Howie EK, McVeigh JA, Straker LM. Comparison of Compliance and Intervention Outcomes  
343 Between Hip- and Wrist-Worn Accelerometers During a Randomized Crossover Trial of an  
344 Active Video Games Intervention in Children. *J Phys Act Health.* 2016; 13: 964-969.
- 345 14. Lloyd J, Creanor S, Price L, et al. Trial baseline characteristics of a cluster randomised controlled  
346 trial of a school-located obesity prevention programme; the Healthy Lifestyles Programme  
347 (HeLP) trial. *BMC Public Health.* 2017; 17: 291. doi: 10.1186/s12889-017-4196-9.
- 348 15. Cole TJ, Freeman JV, Preece MA. British 1990 growth reference centiles for weight, height, body  
349 mass index and head circumference fitted by maximum penalized likelihood. *StatMed.*  
350 1998;17:407–29.
- 351 16. Van Hees VT, Fang Z, Zhao JH, Heywood J, Mirkes E, Sabia S. Package ‘GGIR’: Raw  
352 Accelerometer Data Analysis. <https://cran.r-project.org/web/packages/GGIR/index.html>
- 353 17. Van Hees VT, Gorzelniak L, Leon ECD, et al. Separating Movement and Gravity Components in  
354 an Acceleration Signal and Implications for the Assessment of Human Daily Physical Activity.  
355 *Plos ONE.* 2013;8: e61691. doi:10.1371/journal.pone.0061691
- 356 18. van Hees VT, Fang Z, Langford J, et al. .Autocalibration of accelerometer data for free-living  
357 physical activity assessment using local gravity and temperature: an evaluation on four continents.  
358 *J Appl Physiol.* 2014; 117: 738-744.
- 359 19. Rowlands AV, Yates T, Davis M, Khunti K, Edwardson CL. Raw Accelerometer Data Analysis  
360 with GGIR R-package: Does Accelerometer Brand Matter? *Med Sci Sports Exerc.* 2016; 48:  
361 1935-1941.

- 362 20. Hildebrand M, van Hees VT, Hansen BH, Ekelun U. Age group comparability of raw  
363 accelerometer output from wrist- and hip-worn monitors. *Med Sci Sports Exerc.* 2014; 46: 1816-  
364 24.
- 365 21. Phillips LRS, Parfitt G, Rowlands A. Calibration of the GENEActiv accelerometer for assessment of  
366 physical activity intensity in children. *J Sci Med Sport.* 2013; 16: 124 – 128.  
367 doi:10.1016/j.jsams.2012.05.013
- 368 22. Trost, SG, Mciver KL, Pate RR. Conducting Accelerometer-Based Activity Assessments in Field-  
369 Based Research. *Med Sci Sports Exerc.* 2005;37,Suppl: S531–S543.
- 370 23. Catellier DJ, Hannan PJ, Murray DM. Imputation of missing data when measuring physical  
371 activity by accelerometry. *Med Sci Sports Exerc.* 2005; 37, Suppl: s555 – s562.  
372 doi:10.1249/01.mss.0000185651.59486.4e
- 373 24. Trost, SG, Pate RR , Freedson PS, Sallis J, Taylor WC. Using objective physical activity  
374 measures with youth: How many days of monitoring are needed? *Med Sci Sports Exerc.* 2000;  
375 32:426–431
- 376 25. Jago R, Anderson CB, Baranowski T, Watson K. Adolescent patterns of physical activity,  
377 differences by gender, day and time of day. *Am J Pre Med.*2005; 28: 447 – 452,  
378 doi:10.1016/j.amepre.2005.02.007
- 379 26. Colley R, Garriguet D, Janssens I, Craig C, Clarke J, Tremblay MS. Physical activity of Canadian  
380 children and youth: accelerometer results from the 2007 to 2009 Canadian Health Measures  
381 Survey. *Health Rep.* 2011; 21:63 – 64.
- 382 27. Esliger D, Hall J. Accelerometry in children. In Craig R, Mindell J, Hirani V, editors. *Health*  
383 *Survey for England: Physical activity and Fitness.* Leeds: The NHS Information Centre for health  
384 and social care; 2008.p 159-173
- 385 28. Troiano R, Berrigan D, Dodd K, Masse LC, Tilert T, McDowell M. Physical activity in the  
386 United States measured by accelerometer. *Med Sci Sports Exerc.*2008; 40: 181-8.  
387 doi:10.1249/mss.0b013e31815a51b3
- 388 29. Rowlands AV, Esliger DW, Eady J, Eston RG. Empirical evidence to inform decisions  
389 regarding identification of non-wear periods from accelerometer habitual physical activity

- 390 data. In: Bacquet, G. and Bethoin, S. *Children and Exercise XXV*. 2010. Routledge, London,  
391 219-222.
- 392 30. Audrey S, Bell S, Hughes R, Campbell, R. Adolescent perspectives on wearing accelerometers to  
393 measure physical activity in population-based trials. *Euro J Public Health*. 2012;23:475-480  
394 doi:10.1093/eurpub/cks081
- 395 31. Magnusson KT, Sigurgeirsson I, Sveinsson T, Johannsson E. Assessment of a two-year school-  
396 based physical activity intervention among 7 – 9 year old children. *Int J behave Nutr Phys Act*.  
397 2011; 8: 138 – 149. doi: 10.1186/1479-5868-8-138
- 398 32. Rennie KL, Wareham NJ. The assessment of physical activity in individuals and populations:  
399 why try to be more precise about how physical activity is assessed? *Int J Obes Relat Metab*  
400 *Disord*.1998; suppl 2: s30 – s38.
- 401
- 402
- 403
- 404
- 405
- 406
- 407
- 408
- 409
- 410
- 411
- 412