



## University of Dundee

### Accuracy of alcohol and breast cancer risk information on Drinkaware's website

Larsen, John; Wallace, Paul; Sim, Fiona; Chick, Jonathan; Jarvis, Sarah; Lidington, Iona; Neidle, Stephen; Ogden, Graham; Owens, Lynn

*Published in:*  
Drug and Alcohol Review

*DOI:*  
[10.1111/dar.12676](https://doi.org/10.1111/dar.12676)

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

#### *Citation for published version (APA):*

Larsen, J., Wallace, P., Sim, F., Chick, J., Jarvis, S., Lidington, I., ... Owens, L. (2018). Accuracy of alcohol and breast cancer risk information on Drinkaware's website. *Drug and Alcohol Review*. DOI: 10.1111/dar.12676

#### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## COMMENTARY

## Accuracy of alcohol and breast cancer risk information on Drinkaware's website

JOHN LARSEN<sup>1</sup> , PAUL WALLACE<sup>2</sup>, FIONA SIM<sup>3</sup>, JONATHAN CHICK<sup>4</sup>, SARAH JARVIS<sup>3</sup>, IONA LIDINGTON<sup>5</sup>, STEPHEN NEIDLE<sup>2</sup>, GRAHAM OGDEN<sup>6</sup> & LYNN OWENS<sup>7</sup>

<sup>1</sup>Drinkaware—Research and Impact, London, UK, <sup>2</sup>University College, London, UK, <sup>3</sup>National Health Service England, London, UK, <sup>4</sup>Edinburgh Napier University, Edinburgh, UK, <sup>5</sup>Kingston Primary Care Trust, Kingston, UK, <sup>6</sup>Dental Hospital and School, University of Dundee, Dundee, UK, and <sup>7</sup>Alcohol Services/Hepatology, Royal Liverpool and Broadgreen University Hospitals National Health Service Trust, Liverpool, UK

### Abstract

*A recent paper in Drug and Alcohol Review analysed the information on cancer disseminated by 27 alcohol industry funded organisations. The independent UK alcohol education charity Drinkaware was among the organisations whose information was studied, and based on the analysis claims were made of misrepresentation of evidence about the alcohol-related risk of cancer and alcohol industry influence. This commentary challenges the validity of these findings in respect to the evidence relating to the Drinkaware information, as the analysis is found to be misrepresenting the information by both disregarding the wider information content provided and the order and prominence with which alcohol-related cancer risk is presented. Furthermore, it is argued that the public has a right to be provided with relevant evidence-based information about cancer risk. It is critical that Drinkaware's important public health function is not compromised by unjustified allegations of inaccuracy and by unwarranted attacks on its independence and integrity. [Larsen J, Wallace P, Sim F, Chick J, Jarvis S, Lidington I, Neidle S, Ogden G, Owens L. Accuracy of alcohol and breast cancer risk information on Drinkaware's website. Drug Alcohol Rev 2018]*

**Key words:** alcohol, cancer risk, alcohol education, risk communication, public health.

As medical and scientific advisors to and responsible for research within the independent UK alcohol education charity Drinkaware, we write in response to the paper by Professor Mark Petticrew *et al.* 'How alcohol industry organisations mislead the public about alcohol and cancer' published online in *Drug and Alcohol Review* on 7 September 2017 [1]. This paper reports on a study designed to analyse the information on cancer disseminated by alcohol industry funded organisations, most commonly Social Aspects Public Relations Organisations. It reportedly aimed to determine the extent to which the organisations selected for study fully and accurately communicate the scientific evidence on cancer.

The authors report the findings of a qualitative analysis of information related to alcohol and cancer

risk obtained from the websites and documents of 27 organisations. The analysis is reported to have involved extraction of data independently by two authors, who coded and analysed data using documentary analysis methods. They report their method as comprising: 'reading and understanding meanings of individual texts to identify subthemes; identification of thematic clusters of nodes; triangulation between documents and organisations; checking for reliability/validity and the use of representative examples.'

The authors claim that the analysis demonstrated that most of the organisations included in the study were disseminating misrepresentations of the evidence about the association between alcohol and cancer, and

---

John Larsen PhD, Director of Evidence and Impact, Paul Wallace MD, PhD, Emeritus Professor of Primary Care, Fiona Sim MD, GP and Senior Clinical Adviser, Jonathan Chick MBChB, DSc, FRCPsych, Consultant Psychiatrist and Honorary Professor, Sarah Jarvis MD, GP, Iona Lidington MSc, Associate Director of Public Health, Stephen Neidle, PhD, DSc, Emeritus Professor of Chemical Biology, Graham Ogden MD, PhD, Professor of Oral Surgery, Lynn Owens PhD, Nurse Consultant. Correspondence to Dr John Larsen, Evidence and Impact, Drinkaware, Salisbury House, London Wall, London EC2M 5QQ, UK. Tel.: +442077669907; E-mail: jlarsen@drinkaware.co.uk

Received 7 December 2017; accepted for publication 18 January 2018.

that strategies of ‘denial/omission’, ‘distortion’ and ‘distraction’ were employed. The paper concluded that the alcohol industry appears to be engaged in extensive misrepresentation of evidence about the alcohol-related risk of cancer through its influence on the organisations studied.

Drinkaware was included among the 27 organisations considered. We address in this commentary the claims made in the paper specifically concerning Drinkaware, as we are only involved with information content provided by this organisation.

Looking at the substance of the analysis, material from Drinkaware is explicitly mentioned in relation to the ‘distraction’ strategy as an example of seeking to minimise the role of alcohol by pointing to a wide range of other risk factors, presenting alcohol as ‘just one risk among many’. The one quote from the Drinkaware website that is presented in the manuscript is the following:

*‘For example, the fact that you are female is a risk factor in developing breast cancer. We also know breast cancer is age-related so you’re more likely to develop it as you get older and that you’re more prone to breast cancer if it is part of your family history. These are all factors beyond our control. We also know that risk is related to the ‘hormone environment’ that women experience during the course of early pregnancy, child birth and breastfeeding which all exert a protective effect.’*

Petticrew *et al.* claim that providing this information is misleading because it emphasises potential moderating factors ‘without acknowledging the clear independent risk of alcohol consumption’. However, all the statements made in this paragraph have long been accepted as mainstream opinion in the breast cancer medical and scientific community and there is overwhelming evidence of their correctness. As medical and scientific advisors we stand by the importance of including this information, and it is our opinion that without this paragraph we would have failed to provide women with a complete picture of the risks of alcohol and breast cancer, making us guilty of precisely the crimes of omission of which Petticrew *et al.* accuse other bodies.

Furthermore, in using only this quote the authors fail to make any reference to the comprehensive provision of information in the preceding paragraphs on the same Drinkaware webpage: ‘Alcohol and breast cancer’ (provided in Supplementary Table 2 published together with the Petticrew *et al.* paper). This webpage clearly, comprehensively and unequivocally sets out the evidence linking alcohol to the causation of breast cancer and its sources. The first section on the webpage presents the research evidence: ‘It is clear from a number of large scale studies that there is a link between alcohol consumption and cancer. Globally,

one in five (21.6%) of all alcohol-related deaths are due to cancer. Breast cancer is the most common cancer among women’. Below this, three key points are emphasised: ‘[t]here is good evidence to suggest that alcohol increases the risk of developing breast cancer’; ‘[o]f course drinking alcohol does not mean you will automatically get breast cancer, it does mean your risk of developing it will be increased’ and ‘[h]ow much you drink over your lifetime is what increases the risk’. When presenting further details of the research evidence it is highlighted that ‘[a] number of studies have found that a woman’s risk of breast cancer increases by 7–12% for every 10 g of alcohol per day’. Throughout, the relevant primary sources for these evidence claims are clearly referenced. Other webpages on the Drinkaware website address the link between alcohol and several other cancers, with which we note Petticrew *et al.* do not find fault.

Having reviewed very carefully the page singled out for criticism by Petticrew *et al.*, we stand by its content and refute any misrepresentation of the clear independent risk of alcohol consumption in relation to breast cancer, either intentional or unintentional. Therefore, such criticism is, in our considered opinion, wholly unjustified and unprofessional.

Furthermore, we categorically refute the authors’ inference that information on alcohol and breast cancer provided by Drinkaware was influenced in any way by the alcohol industry. Drinkaware was established in 2007 as an independent alcohol education charity. It is run by an independent Board of Trustees, and all medical and health information provided by Drinkaware is developed with guidance from and approval of an independent Medical Advisory Panel of senior medical and scientific experts. Drinkaware is acknowledged by UK government agencies to be a major provider of evidence-based public information on alcohol in the UK, with more than 9 million individual website visitors each year [2] and in excess of 400,000 people have downloaded the Drinkaware ‘Track and Calculate Units’ app [3].

It is vital that Drinkaware’s important public health function is not compromised by unjustified allegations of inaccuracy and by entirely unwarranted attacks on its independence and integrity. We therefore expect Petticrew *et al.* to address the inaccuracies in their paper, which we have highlighted in this commentary.

## Acknowledgements

The lead author is employed within the UK alcohol education charity Drinkaware, and the co-authors are

members of the independent Medical Advisory Panel of Drinkaware. Drinkaware is primarily funded by voluntary and unrestricted donations from UK alcohol producers, retailers and supermarkets, based on an agreement between the UK government and the alcohol industry. The funders of Drinkaware have no involvement in or influence over the activities of the charity which is governed by an independent Board of trustees.

## References

- [1] Petticrew M, Maani Hessari N, Knai C, Weiderpass E. How alcohol industry organisations mislead the public about alcohol and cancer. *Drug Alcohol Rev* 2017. <https://doi.org/10.1111/dar.12596> [Epub ahead of print].
- [2] Facts & Impacts: Drinkaware Impact Report 2015. Available at: [https://www.drinkaware.co.uk/media/1666/drinkaware\\_2015\\_impact\\_report.pdf](https://www.drinkaware.co.uk/media/1666/drinkaware_2015_impact_report.pdf) (accessed 23 November 2017).
- [3] Attwood S, Parke H, Larsen J, Morton KL. Using a mobile health application to reduce alcohol consumption: a mixed-methods evaluation of the drinkaware track & calculate units application. *BMC Public Health* 2017;17:394. <https://doi.org/10.1186/s12889-017-4358-9>.