



This is a repository copy of *The DNases of pathogenic Lancefield streptococci*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/127831/>

Version: Accepted Version

Article:

Remington, A. and Turner, C.E. orcid.org/0000-0002-4458-9748 (2018) The DNases of pathogenic Lancefield streptococci. *Microbiology*, 164 (3). pp. 242-250. ISSN 1350-0872

<https://doi.org/10.1099/mic.0.000612>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 **The Deoxyribonucleases of Pathogenic Lancefield Streptococci**

2 Alex Remington and Claire E Turner

3 Department of Molecular Biology & Biotechnology, The Florey Institute, University of Sheffield,
4 Sheffield, UK.

5

6

7 **Corresponding author:**

8 Dr Claire Turner

9 Department of Molecular Biology & Biotechnology

10 Firth Court

11 Western Bank

12 Sheffield

13 S10 2TN

14 c.e.turner@sheffield.ac.uk

15

16 **Keywords:** *Streptococcus pyogenes*, DNases, neutrophil extracellular traps

17

18 **Abstract**

19 Deoxyribonucleases (DNases) are abundant among the pathogenic streptococci, with most species
20 harbouring genes for at least one. Despite their prevalence, however, the role for these extracellular
21 enzymes is still relatively unclear. The DNases of the Lancefield group A *Streptococcus*, *S. pyogenes*
22 are the best characterised, with a total of eight DNase genes identified so far. Six are known to be
23 associated with integrated prophages. Two are chromosomally encoded, and one of these is cell-wall
24 anchored. Homologues of both prophage-associated and chromosomally encoded *S. pyogenes*
25 DNases have been identified in other streptococcal species, as well as other unique DNases. A major
26 role identified for streptococcal DNases appears to be in the destruction of extracellular traps
27 produced by immune cells, such as neutrophils, to ensnare bacteria and kill them. These traps are
28 composed primarily of DNA which can be degraded by the secreted and cell wall anchored
29 streptococcal DNases. DNases can also reduce TLR-9 signalling to dampen the immune response and
30 produce cytotoxic deoxyadenosine to limit phagocytosis. Upper respiratory tract infection models of
31 *S. pyogenes* have identified a role for DNases in potentiating infection and transmission, possibly by
32 limiting the immune response or through some other unknown mechanism. Streptococcal DNases
33 may also be involved in interacting with other microbial communities through communication,
34 bacterial killing and disruption of competitive biofilms, or control of their own biofilm production. The
35 contribution of DNases to pathogenesis may therefore be wide ranging and extend beyond direct
36 interference with the host immune response.

37 Introduction

38 A number of clinically significant eukaryotic and prokaryotic microorganisms produce
39 deoxyribonucleases (DNases) as virulence factors. These include Gram-positive bacterial pathogens
40 such as *Staphylococcus aureus* and streptococcal species such as *Streptococcus pyogenes* [1, 2]. Gram-
41 negative pathogens such as *Helicobacter pylori* and *Shigella flexneri*, in addition to members of the
42 genera *Salmonella* and *Yersinia*, also implement these enzymes in a similar manner [3, 4]. Further, the
43 opportunistic fungal pathogens *Candida albicans* and *Cryptococcus neoformans* and the malarial
44 parasites of the genus *Plasmodium* are also known to utilise DNases in pathogenesis [5-7].

45 The genus *Streptococcus* comprises a multitude of obligate and opportunistic pathogens of both
46 humans and animals [8]. A classification system identified by Rebecca Lancefield in the 1930s is still
47 used to classify the beta-haemolytic streptococci based on their type specific carbohydrate antigen
48 [8]. *S. pyogenes*, the Lancefield Group A *Streptococcus* (GAS) is a major human pathogen responsible
49 for a diversity of clinical manifestations and considerable global disease burden exceeding 700 million
50 infections per annum [9]. Clinical manifestations include superficial infections such as pharyngitis,
51 non-bullous impetigo and scarlet fever in addition to potentially lethal invasive manifestations such
52 as streptococcal toxic shock syndrome, necrotising fasciitis and puerperal sepsis [10]. Furthermore, *S.*
53 *pyogenes* is associated with serious post-infectious sequelae, notably post-streptococcal
54 glomerulonephritis and rheumatic fever [11]. *Streptococcus agalactiae*, the Lancefield group B
55 *Streptococcus* (GBS) is another major human pathogen and, although present as a commensal in the
56 gastrointestinal and genitourinary tract, it is a leading cause of neonatal morbidity and mortality
57 worldwide, often associated with neonatal meningitis, pneumonia and sepsis [12]. *Streptococcus*
58 *dysgalactiae* sbsp. *equisimilis* (SDSE) can carry the Lancefield group C or G antigens and has only
59 relatively recently been recognised as a major human pathogen, increasing in incidence and
60 prevalence [13]. The spectrum of symptoms associated with SDSE infection are similar to those
61 observed with *S. pyogenes*, indeed the two species share many virulence factors and significant DNA
62 sequence similarity [13]. Other Lancefield group C streptococci include *Streptococcus equi* subsp. *equi*

63 (hereafter referred to as *S. equi*) and *S. equi* subsp. *zoepidemicus* (hereafter referred to as *S.*
64 *zoepidemicus*). *S. equi* is almost exclusively a pathogen of horses and is believed to be descended
65 from an ancestral strain of *S. zoepidemicus*, which will readily colonise and infect humans in addition
66 to a vast spectrum of domestic and livestock animals. Both species share in excess of 80% DNA
67 sequence similarity with *S. pyogenes* [14].

68 The ability for Lancefield streptococci to cause a wide range of disease may be due to an extensive
69 arsenal of virulence factors. Some of these factors, which include DNases, are associated with mobile
70 genetic elements and can transmit between strains and even species. The function of DNases during
71 pathogenesis is still relatively unclear and the potential for DNases to be virulence factors has only
72 recently been explored.

73 **DNase history & nomenclature**

74 Historically, it was believed that *S. pyogenes* produced only four DNases and these were serologically
75 classified as DNase A, DNase B, DNase C and DNase D. Anti-DNase B titres have been used as a
76 serological biomarker of streptococcal infection and post-streptococcal immune sequelae since at
77 least the 1970s [15, 16]. However, the identity of DNase B would not be truly known for over a decade
78 when it was demonstrated that the chromosomally encoded DNase *spdB* or mitogenic factor (*mf*; then
79 thought to be a streptococcal superantigen) was in fact DNase B [17]. It would be many years
80 subsequent to the initial use of DNase B in the clinical laboratory before it was established that these
81 enzymes could contribute to the pathogenesis of *S. pyogenes* [2]. DNase C is now known to be Spd3,
82 and DNase D has been identified as Sda2 (SdaD2) [2], however, the identity of DNase A in the original
83 serological system is not currently known.

84 Since their discovery, the classification, nomenclature and role of the DNases found in streptococci
85 has been a confusing topic. With advances in molecular biology and the application of whole genome
86 sequencing, it is now apparent that a number of variants exist for the majority of DNases and there is
87 homology between DNases of different streptococcal species.

88 Genetic identification and classification of streptococcal DNases

89 *DNases of the Lancefield group A Streptococcus*

90 The DNases of group A *Streptococcus* are by far the best characterised and currently eight have been
91 identified; *spnA*, *spdB*, *sda1*, *sda2*, *spd1*, *spd3*, *spd4* and *sdn*. Both *spnA* and *spdB* are encoded on the
92 chromosome and have been shown to be common to all *S. pyogenes* isolates tested, existing as
93 different alleles related to the *emm* genotype of the isolate [18, 19]. We confirmed the presence of
94 both *spnA* and *spdB* in all available completed *S. pyogenes* genomes (NCBI, n=54) representing 25
95 different *emm* genotypes. SpnA is the only *S. pyogenes* DNase to be identified that is cell wall anchored
96 via an LPXTG motif [19].

97 The other six *S. pyogenes* DNases are associated with prophages or prophage-like elements (Table 1).
98 *S. pyogenes* has a close evolutionary relationship with temperate bacteriophage as most strains are
99 polylysogenised, and prophage and prophage-like-elements account for ~10% of the *S. pyogenes*
100 genome [20, 21]. Bacteriophages are transmissible between hosts, carrying genes for bacterial
101 virulence factors; the streptococcal superantigens, DNases and a secreted phospholipase. Different
102 bacteriophages may carry the same virulence factor and each factor may exist as a different allele, as
103 has been shown for the streptococcal superantigens [22]. By mining the 54 available genomes of *S.*
104 *pyogenes* we identified that at least two different DNA sequence alleles exist for each of the six
105 prophage associated DNases (Table 1). The most prolific prophage-associated DNase genes were *spd1*
106 and *spd3*. The other four prophage-associated DNase genes were restricted to isolates belonging to
107 only 2-5 different *emm* genotypes, which may be reflective of host-specificity or functionality of the
108 associated prophage. Twelve isolates did not carry any prophage-associated DNases and they
109 belonged to the *emm* genotypes *emm44*, 59, 71, 82, 83, 89 and 101. Although other *emm89* isolates
110 did carry prophage-associated DNases, two *emm89* isolates (H293 and MGAS23530) have been shown
111 not to have any prophage elements integrated into their chromosomes, and therefore only carry
112 chromosomal DNase genes [23, 24].

113 *DNases of other streptococci*

114 Literature search and BLAST analyses identified DNases in other streptococcal species that are similar
115 to those found in *S. pyogenes*. Despite the prevalence of *spd1* and *spd3* among *S. pyogenes* isolates,
116 we only identified three potential homologues of either of these genes (Figure 1). Homologues of *spd1*
117 were identified by BLAST analysis (sharing $\geq 90\%$ identity over 75% or more of the sequence length) in
118 isolates of *S. iniae* and *S. porcinus*, pathogens of fish and swine respectively, although they may also
119 cause disease in humans. A third gene, similar to *spd1* was identified in *S. zooepidemicus* MGCS1056,
120 and is one of three inferred DNases in this genome termed *sdzA* (the *spd1* homologue), *sdzB* and *sdzD*
121 [25]. The gene *sdzA* was unique to MGCS1056 but the other two DNases were identified in the other
122 three *S. zooepidemicus* genomes and the two genomes of the closely related equine pathogen *S. equi*.
123 Alleles of the *sdzD* gene cluster with *sda* genes of *S. pyogenes* and SDSE, suggesting they are
124 homologous. Although in *S. pyogenes*, *sda* is associated with prophages, no such elements were
125 identified associated with the *sda*-like genes of *S. zooepidemicus* [25], *S. equi* or SDSE. This was also
126 the case for the third DNase gene of *S. zooepidemicus* and *S. equi*, *sdzB*, although it shares some
127 similarity to the *S. pyogenes* chromosomally encoded *spdB* as well as the prophage associated *spd4*.
128 Recently, two further DNases were identified in *S. zooepidemicus* that are cell wall anchored, termed
129 ENuc and 5Nuc [26]. The *enuc* alleles identified in both *S. zooepidemicus* and *S. equi* clustered with
130 other identified cell surface anchored DNases (Figure 1) that include *spnA*-like alleles from SDSE. The
131 SDSE *spnA*-like alleles all have a cell wall anchor motif, except *spnA.5* carried by the SDSE strain
132 GGS_124, which has a truncation mutation resulting in the loss of the far C-terminal region including
133 the LPXTG anchor motif. Other cell surface anchored DNases that have been identified include the *S.*
134 *sanguinis* cell wall anchored nuclease, SWAN and the *S. suis* SsnA. *S. suis* does not have a Lancefield
135 antigen and is a pathogen of swine, but can cause severe zoonotic infection in humans. SsnA of *S. suis*
136 has been previously identified as a functional DNase [27, 28] along with EndAsuis, although EndAsuis
137 is membrane anchored and shows homology to *endA* of *S. pneumoniae* which may play a role in
138 competence [29]. The gene *endAsuis* did not show any homology to other streptococcal DNases

139 analysed (Figure 1) although similar DNases with a role in competence may exist in other streptococcal
140 species. The *nucA* gene of *S. agalactiae* also appeared unrelated to any other identified streptococcal
141 DNases, but has confirmed DNase activity [30]. It seems likely that other DNases exist in *S. agalactiae*
142 but have yet to be identified.

143 We did identify two prophage-associated DNases genes in SDSE. One was in strain GGS_124 and
144 associated with a prophage element that shares ~90% identity to prophage 315.3 from *S. pyogenes*
145 *emm3* genome MGAS315 [31]. Although in *S. pyogenes*, the prophage 315.3 is associated with the
146 DNase *spd4*, the gene found in GGS_124 (SDSE167_1285, SDSE *sdn*) is 100% identical to a different
147 prophage-associated *S. pyogenes* DNase, *sdn.5* (Figure 1). Another SDSE strain, 167 also has a
148 prophage-like element associated with a DNase gene [32]. The prophage is most closely related to a
149 prophage-element found in *emm1* NCTC8198 *S. pyogenes*, although this prophage is not associated
150 with any virulence factors, the prophage element in the SDSE strain 167 is associated with an *sda*-like
151 gene (SDEG_1103, SDSE *sda2*). However, in this strain, the gene carries a mutation that would truncate
152 the protein. These findings suggest an exchange of prophages and associated virulence factors
153 between *S. pyogenes* and SDSE.

154

155 **The Role of DNases**

156 *Immune evasion*

157 Originally it was thought that DNases facilitated dissemination of streptococci through tissue planes
158 in the human host by liquefying purulent exudate produced during infection [33]. It has also been
159 speculated that in reducing the viscosity of the microenvironment, DNases expedite transmission of
160 progeny phage particles between bacterial hosts, potentially conferring a selection advantage to both
161 bacteriophage and bacterium [34]. Although this may still be the case, a recently described role for
162 DNases is in the evasive strategy implemented by *S. pyogenes* to prevent neutrophil activation and
163 degradation of neutrophil extracellular traps (NETs) (Figure 2) [35]. NETs are composed of chromatin,

164 histones, proteolytic enzymes and other peptides, and produced by neutrophils on degranulation
165 whereupon they bind to invading microorganisms by charge interaction [36]. Once entrapped,
166 secreted cationic antimicrobial peptides attack the offending agent and neutralise virulence factors.
167 Similar extracellular trap structures have been described in association with mast cells [37] and
168 eosinophils [38], all of which can be degraded by DNases [39].

169 By secreting DNases, such as Sda1, *S. pyogenes* is able to escape these bactericidal traps by degrading
170 their chromatin backbone, thus surviving and spreading (Figure 2) [2, 35]. The ENuc and 5Nuc DNases
171 of *S. zooepidemicus* also have the capacity to degrade NETs, both synergistically and alone, and
172 enabled *S. zooepidemicus* to spread systemically in a murine model of infection [26]. The *S.*
173 *agalactiae* DNase NucA is also able to degrade NETs and its loss results in reduced virulence [30].

174 Cell wall anchored DNases have also been associated with NET degradation. The first description of a
175 cell wall located DNase was the discovery of SsnA of *S. suis* [40]. *S. pyogenes*, further to secreting
176 extracellular DNases, is also able to implement the cell wall anchored DNase SpnA to escape these
177 traps [41]. The SWAN (Sanguinis cell wall anchored nuclease) of *S. sanguinis*, an opportunistic
178 periodontal pathogen, has been shown to degrade NETs [42]. *S. pneumoniae* also produces a cell wall
179 located nuclease, EndA, which is capable of degrading NETs [43] and a homologue of this enzyme,
180 EndAsuis, can be found in *S. suis*. EndAsuis is reported to increase survival in NETs and is produced in
181 addition to the aforementioned secreted nuclease [29]. Although it is unclear why streptococci might
182 implement both cell-wall anchored and secreted DNases, they may provide necessary localised DNase
183 activity in the immediate environment as well as more wide-spread activity [44]. *S. pyogenes* remained
184 attenuated following deletion of the cell-wall anchored DNase SpnA, despite complementation with a
185 secreted form of the enzyme [44]. SpnA may also have an additional role in pathogenesis that is not
186 related to its enzymatic activity [44] and this may well extend to other streptococcal cell wall anchored
187 DNases.

188 Another method by which *S. pyogenes* is able to evade innate immunity is by degrading its own nucleic
189 acids. Indeed, depolymerisation of bacterial DNA by DNases has been shown to prevent killing of *S.*
190 *pyogenes* by reducing TLR-9 signalling and subsequent recognition of un-methylated CpG-rich DNA by
191 macrophages [45]. The ENuc and 5Nuc DNases of *S. zooepidemicus* too have a dual action against the
192 innate immune system. Both nucleases degrade extracellular traps but also possess 5'-nucleotidase
193 activity and produce cytotoxic deoxyadenosine as a substrate, that impedes phagocytosis by
194 macrophages [26]. A similar mechanism was also identified in *S. pyogenes* whereby a cell wall
195 anchored 5' nucleotidase (S5nA) acted synergistically with the DNase SpnA to cleave NETs and
196 generate the cytotoxic deoxyadenosine [46]. S5Na and SpnA are closely related to 5Nuc and ENuc of
197 *S. zooepidemicus*, respectively (Figure 1). Other streptococcal species also express 5' nucleotidases
198 and so similar mechanisms may exist for DNase synergy.

199 *Nutrient scavenging*

200 Elimination of DNA during infection may also have an indirect impact on pathogenesis, serving more
201 than one purpose. For certain strains of *S. pyogenes*, nucleic acid derivatives are essential for growth.
202 Indeed, efforts to standardise a laboratory method for sulphonamide sensitivity testing in the 1940s
203 were hindered by this necessity [47]. Scavenging nucleic acids during infection or colonisation by
204 implementing may therefore provide nutrients during both colonisation and infection. In addition,
205 extracellular trap formation by neutrophils and mast cells ultimately leads to death of these
206 phagocytes [39], and could provide further nutrients for the bacteria. This may also explain why *S.*
207 *pyogenes* and other streptococci possess both secreted and cell-wall- anchored DNases, to retain
208 DNase activity close to the bacterium for nutrition.

209 *Role of DNases during infection*

210 Experimental infection data obtained using a genotype *emm1* strain with three DNases revealed that
211 sequential inactivation of these genes, most importantly SdaD2, significantly impeded the capacity of
212 the strain to establish pharyngeal infection in cynomolgus macaques [2]. Similarly, the acquisition of

213 the Spd1 DNase by ST15 *emm3* *S. pyogenes* was associated with increased nasal and airborne shedding
214 in a murine nasopharyngeal infection model [48]. While not inherently more invasive, nor more lethal,
215 the *emm3* strains that had acquired Spd1 were found to be overrepresented in an upsurge of disease,
216 and their emergence was coincident with a dramatic but transient spike in invasive *emm3* disease in
217 the United Kingdom [49]. The exact role DNase play during upper-respiratory tract infection has yet
218 to be elucidated but these findings support the potential for DNases to contribute directly to infection.
219 DNases have also been shown to contribute to the disease progression in murine skin and soft tissue
220 infections [2,19,35] and the *Galleria mellonella* model of invasive disease [46], which may be due to
221 their role in preventing NET-mediated killing as well as some other as yet un-identified role [46].

222 *Bacterial competition and communication*

223 Spd1 is also reported to have ribonuclease (RNase) activity [50]. A number of secreted eukaryotic
224 RNases are known to be bactericidal; human RNases, such as the eosinophil cationic protein (also
225 known as RNase 3) and the keratinocyte-derived RNase 7 have been shown to play an important role
226 in innate immunity and defence against both Gram-positive and Gram-negative pathogens, by
227 attacking the bacterial cell wall [51-53]. It may be the case, therefore, that the duality of some of the
228 streptococcal DNases in their ability to also degrade RNA, may serve a similar offensive purpose.
229 Microbial RNases have also been reported to have the capacity to damage eukaryotic cells both
230 directly and indirectly of their ribonucleolytic activity, targeting various cellular components, leading
231 to altered gene expression, cellular dysfunction and cell death [54].

232 The contact-dependent growth inhibition (CDI) toxin of *Yersinia kristensenii* was recently identified as
233 a novel bacterial RNase of the RNase A superfamily with a key role in bacterial competition and growth
234 [55]. The RNase activity of streptococcal DNases may therefore also serve to mediate cell-cell
235 interactions within and between bacterial species, coordinating microbial communities such as those
236 observed in non-sterile sites and biofilm.

237 It is also possible that these enzymes may be used by streptococci to compete with commensal
238 microorganisms in non-sterile sites, such as in the nasopharynx or on the skin. Indeed, nucleic acids
239 are a fundamental component of many microbial biofilms [56], and streptococcal DNases may be able
240 to effectively eliminate biofilms formed by other bacteria or regulate the formation of its own biofilm.

241 The EndA nuclease of *S. pneumoniae* plays a role in immune evasion, virulence and competence [43,
242 57]. EndA degrades double-stranded DNA to single-stranded DNA during transformation for the
243 purposes of uptake and recombination [57, 58]. Although no such mechanism has been described in
244 *S. pyogenes*, it is possible that streptococcal DNases could reduce the potential for competing bacterial
245 cells in the environment to be transformed by degrading extracellular DNA.

246 **Summary**

247 The prevalence of DNases suggests an important role in the biology of many streptococci, particularly
248 *S. pyogenes*, with at least two being found in all strains tested. With the increased use of whole
249 genome sequencing it may be that more DNases are identified and it will be important to maintain a
250 consistent classification system across streptococcal species, similar to that proposed for the
251 streptococcal superantigens [22].

252 It has been demonstrated experimentally that acquisition of prophage-associated DNases does not
253 necessarily increase the virulence of a strain, and the genetic background of the bacterial host may
254 play a role [59]. Although this could also be dependent on the types of virulence assays used and the
255 sensitivity of both *in-vitro* and *in-vivo* disease models which are required to build a more complete
256 picture of how DNases function. Indeed, DNase production has been shown previously to require
257 interaction with eukaryotic cells or induction by other external triggers [34].

258 Both chromosomally-encoded and prophage-associated DNases have also been shown to be under
259 the control of the extensive regulatory systems used by streptococci. This includes the control of
260 virulence system (CovR/S or CsrR/S), which negatively regulates *sdaD2* but positively regulates *spdB*

261 in M1 strains [60] and Rgg, which negatively controls *spdB* and *spd3* [61]. Other regulators such as
262 PerR, Ihk/Irr and CodY have also been shown to influence expression of DNases [62-64]. Further work
263 is required to fully understand the complex regulation of DNase expression, which could also be
264 influenced by genotype and associated prophage.

265 Interestingly, Walker et al demonstrated that *sda1* (*sdaD2*) expression during disease is essential for
266 *emm1* strains and places a selective pressure upon CovR/S to mutate, not only to de-repress *sda1*
267 expression but to down-regulate the protease SpeB which degrades Sda1 [65]. Sda1 can therefore
268 influence the infection potential of isolates not only through direct means of protection against NETs,
269 but also by indirectly promoting the development of 'hyper-virulent' CovR/S mutant strains.

270 The contribution of DNases to bacterial colonisation and infection may be extensive. The main focus
271 of DNase research so far has been on the destruction of NETs, however there are other potential roles
272 for DNases that may facilitate infection and warrant further research.

273 **Funding**

274 This work was supported by funding from the Florey Institute, University of Sheffield.

275

276 **Conflict of Interest**

277 The authors declare no conflict of interest.

278

279 **Abbreviations**

280 *Streptococcus dysgalactiae subsp equisimilis* (SDSE)

281 Group B *Streptococcus* (GBS)

282 Group A *Streptococcus* (GAS)

283 Neutrophil extracellular traps (NETs)

284

285

286 **References**

287

288 1. **Berends ET, Horswill AR, Haste NM, Monestier M, Nizet V et al.** Nuclease expression by
289 *Staphylococcus aureus* facilitates escape from neutrophil extracellular traps. *J Innate Immun*
290 2010;2(6):576-586.

291 2. **Sumby P, Barbian KD, Gardner DJ, Whitney AR, Welty DM et al.** Extracellular
292 deoxyribonuclease made by group A *Streptococcus* assists pathogenesis by enhancing evasion of the
293 innate immune response. *Proc Natl Acad Sci U S A* 2005;102(5):1679-1684.

294 3. **Kim JM, Choe MH, Asaithambi K, Song JY, Lee YS et al.** *Helicobacter pylori* HP0425 Targets
295 the Nucleus with DNase I-Like Activity. *Helicobacter* 2016;21(3):218-225.

296 4. **Weinrauch Y, Drujan D, Shapiro SD, Weiss J, Zychlinsky A.** Neutrophil elastase targets
297 virulence factors of enterobacteria. *Nature* 2002;417(6884):91-94.

298 5. **Sanchez M, Colom F.** Extracellular DNase activity of *Cryptococcus neoformans* and
299 *Cryptococcus gattii*. *Rev Iberoam Micol* 2010;27(1):10-13.

300 6. **Genton B, Reed ZH.** Asexual blood-stage malaria vaccine development: facing the challenges.
301 *Curr Opin Infect Dis* 2007;20(5):467-475.

302 7. **de Paula Menezes R, de Melo Riceto EB, Borges AS, de Brito Roder DV, dos Santos Pedroso**
303 **R.** Evaluation of virulence factors of *Candida albicans* isolated from HIV-positive individuals using
304 HAART. *Arch Oral Biol* 2016;66:61-65.

305 8. **Facklam R.** What happened to the streptococci: overview of taxonomic and nomenclature
306 changes. *Clin Microbiol Rev* 2002;15(4):613-630.

307 9. **Carapetis JR, Steer AC, Mulholland EK, Weber M.** The global burden of group A streptococcal
308 diseases. *Lancet Infect Dis* 2005;5(11):685-694.

309 10. **Walker MJ, Barnett TC, McArthur JD, Cole JN, Gillen CM et al.** Disease manifestations and
310 pathogenic mechanisms of Group A *Streptococcus*. *Clin Microbiol Rev* 2014;27(2):264-301.

311 11. **Cunningham MW.** Pathogenesis of group A streptococcal infections. *Clin Microbiol Rev*
312 2000;13(3):470-511.

313 12. **Melin P.** Neonatal group B streptococcal disease: from pathogenesis to preventive strategies.
314 *Clin Microbiol Infect* 2011;17(9):1294-1303.

315 13. **Brandt CM, Spellerberg B.** Human infections due to *Streptococcus dysgalactiae* subspecies
316 *equisimilis*. *Clin Infect Dis* 2009;49(5):766-772.

317 14. **Holden MT, Heather Z, Paillot R, Steward KF, Webb K et al.** Genomic evidence for the
318 evolution of *Streptococcus equi*: host restriction, increased virulence, and genetic exchange with
319 human pathogens. *PLoS Pathog* 2009;5(3):e1000346.

320 15. **Shulman ST, Ayoub EM, Victorica BE, Gessner IH, Tamer DF et al.** Differences in antibody
321 response to streptococcal antigens in children with rheumatic and non-rheumatic mitral valve disease.
322 *Circulation* 1974;50(6):1244-1251.

323 16. **Kaplan EL, Anthony BF, Chapman SS, Ayoub EM, Wannamaker LW.** The influence of the site
324 of infection on the immune response to group A streptococci. *J Clin Invest* 1970;49(7):1405-1414.

325 17. **Sriskandan S, Unnikrishnan M, Krausz T, Cohen J.** Mitogenic factor (MF) is the major DNase
326 of serotype M89 *Streptococcus pyogenes*. *Microbiology* 2000;146 (Pt 11):2785-2792.

327 18. **Matsumoto M, Sakae K, Hashikawa S, Torii K, Hasegawa T et al.** Close correlation of
328 streptococcal DNase B (*sdaB*) alleles with emm genotypes in *Streptococcus pyogenes*. *Microbiol*
329 *Immunol* 2005;49(10):925-929.

330 19. **Hasegawa T, Minami M, Okamoto A, Tatsuno I, Isaka M et al.** Characterization of a virulence-
331 associated and cell-wall-located DNase of *Streptococcus pyogenes*. *Microbiology* 2010;156(Pt 1):184-
332 190.

333 20. **Banks DJ, Beres SB, Musser JM.** The fundamental contribution of phages to GAS evolution,
334 genome diversification and strain emergence. *Trends Microbiol* 2002;10(11):515-521.

335 21. **Beres SB, Musser JM.** Contribution of exogenous genetic elements to the group A
336 *Streptococcus* metagenome. *PLoS One* 2007;2(8):e800.

- 337 22. **Commons RJ, Smeesters PR, Proft T, Fraser JD, Robins-Browne R et al.** Streptococcal
338 superantigens: categorization and clinical associations. *Trends Mol Med* 2014;20(1):48-62.
- 339 23. **Beres SB, Kachroo P, Nasser W, Olsen RJ, Zhu L et al.** Transcriptome remodeling contributes
340 to epidemic disease caused by the human pathogen *Streptococcus pyogenes*. *MBio* 2016;7(3).
- 341 24. **Turner CE, Abbott J, Lamagni T, Holden MT, David S et al.** Emergence of a new highly
342 successful acapsular group A *Streptococcus* clade of genotype *emm89* in the United Kingdom. *MBio*
343 2015;6(4):e00622.
- 344 25. **Beres SB, Sesso R, Pinto SW, Hoe NP, Porcella SF et al.** Genome sequence of a Lancefield
345 group C *Streptococcus zooepidemicus* strain causing epidemic nephritis: new information about an old
346 disease. *PLoS One* 2008;3(8):e3026.
- 347 26. **Ma F, Guo X, Fan H.** Extracellular nucleases of *Streptococcus equi* subsp. *zooepidemicus*
348 degrade neutrophil extracellular traps and impair macrophage activity of the host. *Appl Environ*
349 *Microbiol* 2017;83(2).
- 350 27. **Haas B, Bonifait L, Vaillancourt K, Charette SJ, Gottschalk M et al.** Characterization of DNase
351 activity and gene in *Streptococcus suis* and evidence for a role as virulence factor. *BMC Res Notes*
352 2014;7:424.
- 353 28. **de Buhr N, Neumann A, Jerjomiceva N, von Kockritz-Blickwede M, Baums CG.** *Streptococcus*
354 *suis* DNase SsnA contributes to degradation of neutrophil extracellular traps (NETs) and evasion of
355 NET-mediated antimicrobial activity. *Microbiology* 2014;160(Pt 2):385-395.
- 356 29. **de Buhr N, Stehr M, Neumann A, Naim HY, Valentin-Weigand P et al.** Identification of a novel
357 DNase of *Streptococcus suis* (EndAsuis) important for neutrophil extracellular trap degradation during
358 exponential growth. *Microbiology* 2015;161(Pt 4):838-850.
- 359 30. **Derre-Bobillot A, Cortes-Perez NG, Yamamoto Y, Kharrat P, Couve E et al.** Nuclease A
360 (Gbs0661), an extracellular nuclease of *Streptococcus agalactiae*, attacks the neutrophil extracellular
361 traps and is needed for full virulence. *Mol Microbiol* 2013;89(3):518-531.
- 362 31. **Suzuki H, Lefebure T, Hubisz MJ, Pavinski Bitar P, Lang P et al.** Comparative genomic analysis
363 of the *Streptococcus dysgalactiae* species group: gene content, molecular adaptation, and promoter
364 evolution. *Genome Biol Evol* 2011;3:168-185.
- 365 32. **Watanabe S, Kirikae T, Miyoshi-Akiyama T.** Complete genome sequence of *Streptococcus*
366 *dysgalactiae* subsp. *equisimilis* 167 carrying Lancefield group C antigen and comparative genomics of
367 *S. dysgalactiae* subsp. *equisimilis* strains. *Genome Biol Evol* 2013;5(9):1644-1651.
- 368 33. **Sherry S, Johnson A, Tillett WS.** The action of streptococcal desoxyribose nuclease
369 (streptodornase) *in vitro* and on purulent pleural exudations of patients. *J Clin Invest* 1949;28(5 Pt
370 2):1094-1104.
- 371 34. **Broudy TB, Pancholi V, Fischetti VA.** The *in vitro* interaction of *Streptococcus pyogenes* with
372 human pharyngeal cells induces a phage-encoded extracellular DNase. *Infect Immun* 2002;70(6):2805-
373 2811.
- 374 35. **Buchanan JT, Simpson AJ, Aziz RK, Liu GY, Kristian SA et al.** DNase expression allows the
375 pathogen group A *Streptococcus* to escape killing in neutrophil extracellular traps. *Curr Biol*
376 2006;16(4):396-400.
- 377 36. **Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y et al.** Neutrophil extracellular
378 traps kill bacteria. *Science* 2004;303(5663):1532-1535.
- 379 37. **von Kockritz-Blickwede M, Goldmann O, Thulin P, Heinemann K, Norrby-Teglund A et al.**
380 Phagocytosis-independent antimicrobial activity of mast cells by means of extracellular trap
381 formation. *Blood* 2008;111(6):3070-3080.
- 382 38. **Yousefi S, Gold JA, Andina N, Lee JJ, Kelly AM et al.** Catapult-like release of mitochondrial
383 DNA by eosinophils contributes to antibacterial defense. *Nat Med* 2008;14(9):949-953.
- 384 39. **von Kockritz-Blickwede M, Nizet V.** Innate immunity turned inside-out: antimicrobial defense
385 by phagocyte extracellular traps. *J Mol Med (Berl)* 2009;87(8):775-783.
- 386 40. **Fontaine MC, Perez-Casal J, Willson PJ.** Investigation of a novel DNase of *Streptococcus suis*
387 serotype 2. *Infect Immun* 2004;72(2):774-781.

- 388 41. **Chang A, Khemlani A, Kang H, Proft T.** Functional analysis of *Streptococcus pyogenes* nuclease
389 A (SpnA), a novel group A streptococcal virulence factor. *Mol Microbiol* 2011;79(6):1629-1642.
- 390 42. **Morita C, Sumioka R, Nakata M, Okahashi N, Wada S et al.** Cell wall-anchored nuclease of
391 *Streptococcus sanguinis* contributes to escape from neutrophil extracellular trap-mediated
392 bacteriocidal activity. *PLoS One* 2014;9(8):e103125.
- 393 43. **Beiter K, Wartha F, Albiger B, Normark S, Zychlinsky A et al.** An endonuclease allows
394 *Streptococcus pneumoniae* to escape from neutrophil extracellular traps. *Curr Biol* 2006;16(4):401-
395 407.
- 396 44. **Uchiyama S, Andreoni F, Schuepbach RA, Nizet V, Zinkernagel AS.** DNase Sda1 allows invasive
397 M1T1 Group A *Streptococcus* to prevent TLR9-dependent recognition. *PLoS Pathog*
398 2012;8(6):e1002736.
- 399 45. **Zheng L, Khemlani A, Lorenz N, Loh JM, Langley RJ et al.** Streptococcal 5'-Nucleotidase A
400 (S5nA), a novel *Streptococcus pyogenes* virulence factor that facilitates immune evasion. *J Biol Chem*
401 2015;290(52):31126-31137.
- 402 46. **Chalmers C, Khemlani AHJ, Sohn CR, Loh JMS, Tsai CJ et al.** Streptococcus pyogenes nuclease
403 A (SpnA) mediated virulence does not exclusively depend on nuclease activity. *J Microbiol Immunol*
404 *Infect* 2017.
- 405 47. **Wilson AT.** Loss of Group Carbohydrate during Mouse Passages of a Group a Hemolytic
406 *Streptococcus*. *J Exp Med* 1945;81(6):593-596.
- 407 48. **Afshar B, Turner CE, Lamagni TL, Smith KC, Al-Shahib A et al.** Enhanced nasopharyngeal
408 infection and shedding associated with an epidemic lineage of *emm3* group A *Streptococcus*. *Virulence*
409 2017:1-11.
- 410 49. **Al-Shahib A, Underwood A, Afshar B, Turner CE, Lamagni T et al.** Emergence of a novel
411 lineage containing a prophage in *emm/M3* group A *Streptococcus* associated with upsurge in invasive
412 disease in the UK. *Microb Genom* 2016;2(6):e000059.
- 413 50. **Korczyńska JE, Turkenburg JP, Taylor EJ.** The structural characterization of a prophage-
414 encoded extracellular DNase from *Streptococcus pyogenes*. *Nucleic Acids Res* 2012;40(2):928-938.
- 415 51. **Harder J, Schroder JM.** RNase 7, a novel innate immune defense antimicrobial protein of
416 healthy human skin. *J Biol Chem* 2002;277(48):46779-46784.
- 417 52. **Rosenberg HF.** RNase A ribonucleases and host defense: an evolving story. *J Leukoc Biol*
418 2008;83(5):1079-1087.
- 419 53. **Torrent M, Badia M, Moussaoui M, Sanchez D, Nogues MV et al.** Comparison of human
420 RNase 3 and RNase 7 bactericidal action at the Gram-negative and Gram-positive bacterial cell wall.
421 *FEBS J* 2010;277(7):1713-1725.
- 422 54. **Makarov AA, Ilinskaya ON.** Cytotoxic ribonucleases: molecular weapons and their targets.
423 *FEBS Lett* 2003;540(1-3):15-20.
- 424 55. **Batot G, Michalska K, Ekberg G, Irimpan EM, Joachimiak G et al.** The CDI toxin of *Yersinia*
425 *kristensenii* is a novel bacterial member of the RNase A superfamily. *Nucleic Acids Res*
426 2017;45(9):5013-5025.
- 427 56. **Whitchurch CB, Tolker-Nielsen T, Ragas PC, Mattick JS.** Extracellular DNA required for
428 bacterial biofilm formation. *Science* 2002;295(5559):1487.
- 429 57. **Zhu L, Kuang Z, Wilson BA, Lau GW.** Competence-independent activity of pneumococcal EndA
430 mediates degradation of extracellular DNA and NETs and is important for virulence. *PLoS One*
431 2013;8(7):e70363.
- 432 58. **Lacks S, Greenberg B.** A deoxyribonuclease of *Diplococcus pneumoniae* specific for
433 methylated DNA. *J Biol Chem* 1975;250(11):4060-4066.
- 434 59. **Venturini C, Ong CL, Gillen CM, Ben-Zakour NL, Maamary PG et al.** Acquisition of the Sda1-
435 encoding bacteriophage does not enhance virulence of the serotype M1 *Streptococcus pyogenes*
436 strain SF370. *Infect Immun* 2013;81(6):2062-2069.

- 437 60. **Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM.** Genome-wide analysis of group a
438 streptococci reveals a mutation that modulates global phenotype and disease specificity. *PLoS Pathog*
439 2006;2(1):e5.
- 440 61. **Anbalagan S, Chaussee MS.** Transcriptional regulation of a bacteriophage encoded
441 extracellular DNase (Spd-3) by Rgg in *Streptococcus pyogenes*. *PLoS One* 2013;8(4):e61312.
- 442 62. **Voyich JM, Sturdevant DE, Braughton KR, Kobayashi SD, Lei B et al.** Genome-wide protective
443 response used by group A *Streptococcus* to evade destruction by human polymorphonuclear
444 leukocytes. *Proc Natl Acad Sci U S A* 2003;100(4):1996-2001.
- 445 63. **McDowell EJ, Callegari EA, Malke H, Chaussee MS.** CodY-mediated regulation of
446 *Streptococcus pyogenes* exoproteins. *BMC Microbiol* 2012;12:114.
- 447 64. **Wang CH, Chiang-Ni C, Kuo HT, Zheng PX, Tsou CC et al.** Peroxide responsive regulator PerR
448 of group A *Streptococcus* is required for the expression of phage-associated DNase Sda1 under
449 oxidative stress. *PLoS One* 2013;8(12):e81882.
- 450 65. **Walker MJ, Barnett TC, McArthur JD, Cole JN, Gillen CM et al.** Disease manifestations and
451 pathogenic mechanisms of Group A *Streptococcus*. *Clin Microbiol Rev* 2014;27(2):264-301.
- 452

Table 1. Prophage-associated DNases in *S. pyogenes* and their alleles.


Gene	Allele*	Isolate	Locus
<i>sda1</i>	<i>sda1.1</i>	MGAS8232 (M18)	spyM18_1746
	<i>sda1.2</i>	MGAS10394 (M6), JRS4 (M6)	M6_Spy1339, SPYJRS4_1267
<i>sda2</i>	<i>sda2.1</i> (<i>sdaD2</i>)	MGAS5005 (M1), A20 (M1), M1_476 (M1), HKU360 (M12), MGAS2096 (M12)	M5005_Spy1415, A20_1463, M1GAS476_1494, SpyOHK_02660, MGAS2096_Spy1441
	<i>sda2.2</i>	MGAS9429 (M12)	MGAS9429_Spy1417
<i>spd1</i>	<i>spd1.1</i>	SF370 (M1), MGAS10270 (M2), MGAS10750 (M4), MEW427 (M4), Manfredo (M5), MGAS9429 (M12), MGAS2096 (M12), HKU360 (M12), HKU488 (M12), MGAS8232 (M18), M23ND (M23), STAB120304 (M75), STAB14018 (M75), NGAS743 (M87), JMUB1235 (M89), MGAS27061 (M89), MGAS11027 (M89), NGAS322 (M114)	SPy0712, MGAS10270_Spy0598, MGAS10750_Spy0622, AWM58_02815, SpyM51263, MGAS9429_Spy0594, MGAS2096_Spy0602, SPYOHK_01985, HKU488_01495, spyM18_0779, FE90_0223, B5D85_03105, AYM92_03010, DI45_06730, JMUB1235_0583, MGAS27061_0582, MGAS11027_0597, SD89_06900
	<i>spd1.2</i>	MGAS10394 (M6), JRS4 (M6), MGAS6180 (M28), M28PF1 (M28), STAB9014 (M28), MEW123 (M28), STAB10015 (M28)	M6_Spy1195, SPYJRS4_1111, M28_Spy0968, ABO05_04560, VT08_04870, AWM59_04435, VU19_04860
	<i>spd1.3</i>	Alab49 (M53), AP53 (M53)	SPYALAB49_001168, AUQ45_1179
<i>spd3</i>	<i>spd3.1</i>	SF370 (M1), MGAS5005 (M1), M1_476 (M1), A20 (M1), 5448 (M1), AP1 (M1), NCTC8198 (M1), MGAS10270 (M2), MGAS10750 (M4), MEW427 (M4), Manfredo (M5), GUR (M11), HKU488 (M12), HSC5 (M14), NZ131 (M49), Alab49 (M53), AP53 (M53), STAB13021 (M66), STAB14018 (M75), STAB120304 (M75), STAB090229 (M75), NGAS743 (M87)	SPy_1436, M5005_Spy1169, M1GAS476_1231, A20_1204, SP5448_03755, SPAP1_02890, ERS445054_01298, MGAS10270_Spy0852, MGAS10750_Spy0888, AWM58_03950, SpyM50534, B2G65_01915, HKU488_01108, L897_05810, Spy49_1455, SPYALAB49_001299, AUQ45_1308, AXK13_07360, AYM92_04205, B5D85_04335, B4W66_03985, DI45_04190
	<i>spd3.2</i>	M23ND (M23)	FE90_0649
	<i>spd3.3</i>	MGAS8232 (M18)	spyM18_1446
	<i>spd3.4</i>	MGAS10394 (M6)	M6_Spy1541
	<i>spd4</i>	<i>spd4.1</i>	MGAS315 (M3), SSI-1 (M3), STAB902 (M3)
<i>spd4.2</i>		Manfredo (M5)	SpyM50691
<i>sdn</i>	<i>sdn.1</i>	MGAS315 (M3)	SpyM3_1409
	<i>sdn.2</i>	SSI-1 (M3), STAB902 (M3)	SPs0455, STAB902_02580
	<i>sdn.3</i>	MGAS10394 (M6)	M6_Spy0067
	<i>sdn.4</i>	NGAS743 (M87), MGAS11027 (M89)	DI45_06360, MGAS11027_0659
	<i>sdn.5</i>	STAB90229 (M75)	B4W66_07530

* Allele based on nucleotide sequence of the entire coding region.

455 **Figure Legends**

456 **Figure 1. Phylogenetic analysis of streptococcal DNases.** Full length coding regions of each identified
457 and potential DNase gene were aligned using MUSCLE and a neighbour-joining tree created. Bootstrap
458 values greater than 80% are shown on branches. Multiple alleles were compared for all prophage-
459 associated *S. pyogenes* DNase genes (red) but only single representative alleles for the chromosomal
460 DNases *spnA*, *spnB* and *s5nA* are shown (blue). Alleles for *Streptococcus dysgalactiae subsp equisimilis*
461 (SDSE) DNase genes were determined from five completed genomes; AC-2713 (NC_019042.1),
462 GGS_124 (AP010935.1), RE378 (AP011114.1), 167 (AP012976.1), ATCC12394 (CP002215.1). Alleles
463 for *Streptococcus zooepidemicus* DNase genes (green) were determined from four completed
464 genomes; H70 (FM204884.1), CY (CP006770.1), ATCC35246 (CP002904.1), MGCS10565 (CP001129.1).
465 Alleles for *Streptococcus equi* DNase genes (green) were determined from two completed genomes;
466 ATCC39806 (CP021972.1), 4047 (FM204883.1). Other DNase genes comprise *endAsuis* (SSU1009) and
467 *ssnA* (SSU1760) from *S. suis* strain P1/7 (AM946016.1), *swan* (SSA_1750) from *S. sanguinis* SK36
468 (CP000387.1), *spnA* (K710_1281) from *S. iniae* SF1 (CP005941.1), *spd1* (STRPO_1639) from *S. porcinus*
469 str. Jelinkova 176 (AEUU02000001.1), group B *Streptococcus* (GBS) *nuc* (gbs0661) from *S. agalactiae*
470 NEM316 (AL732656.1).

471 **Figure 2. Streptococcal DNases can degrade neutrophil extracellular traps (NETs).** NETs are
472 composed primarily of DNA (blue strands) associated with histones and other antibacterial factors
473 (yellow circles). Bacteria can be ensnared in the DNA traps and killed by the associated factors (left-
474 hand figure). Streptococcal secreted and cell-anchored DNases (indicated as scissors) degrade NETs,
475 allowing the bacteria to escape prevent killing (Right-hand figure).




Surface anchored DNases

476

477

Bacteria are trapped in DNA released from neutrophils, and killed by associated factors



Streptococcal DNases degrade DNA traps freeing the bacteria and preventing killing

