



MSC 46A12

**FOURIER-BESSEL'S TRANSFORM OF A GENERALIZED FUNCTION  
VANISHING OUTSIDE A BOUNDED SURFACE****E.L. Shishkina**

Voronezh State University,

Universitetskaya pl., 1, Voronezh, Russia, e-mail: [ilina\\_dico@mail.ru](mailto:ilina_dico@mail.ru)

**Key words:** Fourier-Bessel's transform, generalized functions, Schwartz's function, Euclidian space, linear functionals.

Let  $\mathbb{R}_n^+$  denote an Euclidean space of points  $x = (x_1, \dots, x_n)$ ,  $x_1 > 0, \dots, x_n > 0$  and the multiindex  $\gamma = (\gamma_1, \dots, \gamma_n)$  runs through fixed positive numbers. The space  $S_{ev}(\mathbb{R}_n^+) = S_{ev}$  is the subspace of the Schwartz function space that consists of functions  $\varphi(x)$  even in each variable  $x_1, \dots, x_n$ . The space of linear continuous functionals, whose regular representatives are generated by the linear weighted form

$$(f, \varphi)_\gamma = \int_{\mathbb{R}_n^+} f(x)\varphi(x)x^\gamma dx, \quad x^\gamma = \prod_{i=1}^n x_i^{\gamma_i},$$

is called the distribution space over  $S_{ev}$  and is denoted by  $S'_{ev}(\mathbb{R}_n^+) = S'_{ev}$ .

The Fourier-Bessel transform is denoted by formula

$$F_B[f](\xi) = \widehat{f}(\xi) = \int_{\mathbb{R}_n^+} f(x)\mathbf{j}_\gamma(x, \xi)x^\gamma dx,$$

where  $\mathbf{j}_\gamma(x, \xi) = \prod_{i=1}^n j_{\frac{\gamma_i-1}{2}}(x_i \xi_i)$ ,  $j_\nu(t) = \Gamma(\nu + 1) \left(\frac{2}{t}\right)^\nu J_\nu(t)$ ,  $t \in \mathbb{R}_1$ ,  $J_\nu(t)$  is Bessel functions of the first kind. Spaces  $S_{ev}$  and  $S'_{ev}$  are invariant to Fourier-Bessel transform (see [1]).

For the generalized function  $f \in S'_{ev}$ , vanishing outside a bounded surface  $\Omega \subset \mathbb{R}_n^+$  the Fourier-Bessel transform is functional

$$(f(x), \mathbf{j}_\gamma(x, \xi))_\gamma = \int_{\Omega} f(\sigma)\mathbf{j}_\gamma(x, \xi)x^\gamma dx,$$

which acts as follows: function  $\mathbf{j}_\gamma(x, \xi)$  is replaced by a test function  $\varphi_0(x, \xi) = \mathbf{j}_\gamma(x, \xi)$  for  $x \in \Omega$  and  $\varphi_0(x, \xi) = 0$  for  $x \notin \Omega$ , then functional  $f$  applies to  $\varphi_0(x, \xi)$ . The number obtained is independent of choice of this function  $\varphi_0(x, \sigma)$ .

The Fourier-Bessel transform of any generalized function  $f \in S'_{ev}$  vanishing outside a bounded surface for any test function  $\psi(x) \in S_{ev}$  is denoted by formula

$$\int (f(x), \mathbf{j}_\gamma(x, \sigma))_\gamma \widehat{\psi}(\sigma)\sigma^\gamma d\sigma = \frac{2^{n-|\gamma|}}{\prod_{j=1}^n \Gamma^2\left(\frac{\gamma_j+1}{2}\right)} (f, \psi)_\gamma.$$



We shall introduce a singular generalized weighted function (compare with construction in [2] page 247)

$$(\delta_\gamma(r-a), \varphi)_\gamma = \int_{S_n^+(a)} \varphi(x) x^\sigma dS, \quad \varphi(x) \in S_{ev}.$$

The Fourier-Bessel transform of  $\delta_\gamma(r-R)$  is calculated according to the formula:

$$F_B[\delta_\gamma(r-R)](\xi) = \int_{S_n^+(R)} \mathbf{j}_\gamma(x, \xi) x^\gamma dS_R = R^{n+|\gamma|-1} |S_n^+(1)|_\gamma j_{\frac{n+|\gamma|-2}{2}}(R|\xi|). \quad (1)$$

Following [3] we introduce the operator  $\Delta_\gamma$ :

$$\Delta_\gamma = \sum_{i=1}^n B_{\gamma_i},$$

where  $B_{\gamma_i}$  is Bessel operator:

$$B_{\gamma_i} = \frac{\partial^2}{\partial x_i^2} + \frac{\gamma_i}{x_i} \frac{\partial}{\partial x_i}, \quad i = 1, \dots, n.$$

The formula (1) can be used for solving a problem

$$\frac{\partial^2 u}{\partial t^2} + \frac{\alpha}{t} \frac{\partial u}{\partial t} = \Delta_\gamma u(x, t), \quad (2)$$

$$u(x, 0) = 0, \quad u_t(x, 0) = \delta_\gamma(x). \quad (3)$$

For  $0 < n + |\gamma| - \alpha < 3$ ,  $|\gamma| = \gamma_1 + \dots + \gamma_n$  the solution of (2)-(3) is

$$u(x, t) = C_{\alpha, \gamma}(n) t^{1-n-|\gamma|} {}_2F_1 \left( \frac{n+|\gamma|-\alpha}{2}, \frac{n+|\gamma|-1}{2}, \frac{n+|\gamma|}{2}; \frac{|x|^2}{t^2} \right),$$

where

$$C_{\alpha, \gamma}(n) = 2^{n+|\gamma|-\alpha-2} |S_1^+(n)|_\gamma \frac{\Gamma\left(\frac{1-\alpha}{2}\right) \Gamma\left(\frac{n+|\gamma|-\alpha}{2}\right)}{\Gamma\left(\frac{3-n-|\gamma|}{2}\right)}.$$

### References

1. Lyakhov L.N. Multipliers of the mixed Fourier–Bessel transform // Tr. Mat. Inst. Steklova. – 1997. – 17, V.214. – P.234–249.
2. Gel'fand, I.M., Shilov, G.E. Generalized functions. Vol. I: Properties and operations / Boston: Academic Press, 1964.
3. Kipriyanov I.A. Singular Elliptic Boundary Value Problems / Moscow: Nauka, 1996.



## ПРЕОБРАЗОВАНИЕ ФУРЬЕ-БЕССЕЛЯ ОБОБЩЕННОЙ ФУНКЦИИ ИСЧЕЗАЮЩЕЙ ВНЕ ОГРАНИЧЕННОЙ ПОВЕРХНОСТИ

Е.Л. Шишкина

Воронежский государственный университет,  
пл. Университетская, 1, Воронеж, Россия, e-mail: [ilina\\_dico@mail.ru](mailto:ilina_dico@mail.ru)

**Ключевые слова:** преобразование Фурье-Бесселя, обобщенные функции, функция Шварца, евклидово пространство, линейные функционалы.