

From the Department of Pathology, Royal Veterinary and Agricultural University, Copenhagen, Denmark.

IMMUNOGLOBULIN LEVELS IN NON-ABORTED AND ABORTED FETUSES FROM DANISH HERDS OF CATTLE

By

H. Bielefeldt Ohmann

OHMANN, H. BIELEFELDT: *Immunoglobulin levels in non-aborted and aborted fetuses from Danish herds of cattle.* Acta vet. scand. 1981, 22, 428—434. — The applicability of serological tests in the diagnosis of intrauterine infections in Danish cattle was investigated. Fetuses from slaughter animals, experimentally infected and spontaneously aborted fetuses, premature and stillborn calves, were subjected to necropsy and histological studies, and to microbiological and serological examinations. The latter comprised rocket immunoelectrophoresis and radial immunodiffusion. In sera of the fetuses from slaughter animals, immunoglobulins were either not detectable or only present in very small quantities, whereas sera of fetuses and calves with pathological changes and/or verified infection contained considerable amounts of IgM and IgG. IgA was also detected in the latter group. The results corroborate the diagnostic importance of immunoglobulin determination in aborted fetuses and stillborn calves.

bovine fetuses; intrauterine infection; immunoglobulins; cattle.

Apparently, no overall picture of abortion as a contribution to reproductive failure in Danish cattle stock is available. This applies to the incidence as well as to the significance of various infectious and non-infectious causes.

From all over the world serological investigations have come widely into use as a contribution to establishing a diagnosis of fetal infection as a cause of abortion. It is assumed that no transplacental transfer of immunoglobulins to the bovine fetus will occur during a normal pregnancy (*Brambell 1970*). Further, it has been shown that normal precolostral calf serum contains only small amounts of immunoglobulins, if any (*Sawyer et al. 1973, Ellis et al. 1978*). However, following the first appearance

of immune reactivity around day 90 of gestation the fetus gradually gains competence to infectious and non-infectious antigens (Schultz 1973). Factors resulting in antigenic stimulation of the bovine fetus may be of significance for bovine abortion (Miller & Quinn 1975, Kirkbride *et al.* 1977, Ellis *et al.*). The objects of the present study were to assess the applicability of serological tests for purposes of diagnosis of intrauterine infections in Danish cattle and to establish the comparative value of results available from abroad.

MATERIALS AND METHODS

Fifty-three bovine fetuses were collected at an abattoir. Ages of fetuses were estimated from tabular data based on the relationship of age to crown-rump length and weight measurements (Hubbert *et al.* 1972). Blood was collected from the brachial vessels, the serum separated and stored at -20°C until investigated. From six fetuses, less than ten weeks, amniotic, pleural and abdominal fluids were collected and used alternatively to serum. Eight aborted fetuses, two prematurely delivered calves and three calves stillborn at normal term were submitted for autopsy. Blood was collected as above or from the heart. Alternatively, from two fetuses serosanguinous pleural fluid was used. Five fetuses, five to six months of gestational age, formed part in an experiment, in which four fetuses were infected experimentally in utero with bovine viral diarrhoea virus (BVDV) and one received a sham-inoculum (Ohmann *et al.* 1981 b). Tissue samples were collected for microbiological and morphological studies. The histological examinations were carried out on formalin-fixed specimens stained with haematoxylin-eosin. Tissue samples from all fetuses were examined for cytopathogenic viral agents by inoculating supernatants of ground tissue onto bovine kidney cell cultures. Following one subcultivation all samples were tested for the presence of noncytopathogenic BVDV by indirect immunofluorescence (Ohmann *et al.* 1981 a). From ten animals, including the five experimental fetuses, tissues were additionally examined for BVDV-antigen by immunocytochemical methods (Ohmann *et al.* 1981 a). Routine bacteriological examination was carried out on tissue samples from 35 fetuses.

Rocket immunoelectrophoresis for the detection of IgG and IgM was essentially performed as outlined by Dalsgaard *et al.* (1979), using a phosphate buffer, pH 7.0 (Olitzki 1959), rabbit-

anti-bovine IgG (DAKO Immunoglobulins, DK) and rabbit-anti-bovine IgM (Miles Lab. Ltd, Slough, UK). Electrophoresis was always done in parallel to a pooled sample of serum from clinically healthy, BVDV-free calves (lot NK 800807, State Veterinary Institute of Virus Research, DK). The radial immunodiffusion (RID) tests were performed with commercially prepared testkits specific for bovine IgA, IgG and IgM (Miles Lab. Ltd, Slough, UK). The diffusion-zone was read after 48 h. Serum samples were also tested for the presence of BVDV-specific antibodies (SA), (Jensen 1981, Ohmann 1981).

RESULTS AND DISCUSSION

As will appear from Table 1, IgG was detected in four (7.7 %) and IgM in six (11.5 %) of the fetuses from slaughter animals. As the RID was only carried out on selected samples in this group, the results are not generally representative. Except for one, all fetuses with detectable serum-immunoglobulins were older than 150 days gestational age. Only in two fetuses, eight and eight and a half to nine months respectively (Table 2), did the IgG concentration amount to that found in precolostral newborn calf serum (Sawyer *et al.* 1973). The specificity of the immunoglobulin was not determined. Considerable amounts of IgG and IgM were detected in all serum samples from aborted and congenitally BVDV-infected fetuses and calves. The concentrations were equal to or higher than that found in precolostral bovine calf serum (Sawyer *et al.*, Ohmann 1981). With concentrations of more than 2.76 mg/ml in the serum of three stillborn calves (Table 2), the level may even exceed that of postcolostral newborn calf serum (Sawyer *et al.*). The immunoglobulins are assumed to origin from a fetal production. This is corroborated by the simultaneous presence of IgG and IgM in the absence of IgA, i.e., a selective occurrence of the former two types. Also, the absence of detectable specificity of the immunoglobulins in most fetal sera, in the presence of specific reactivity in the maternal serum (Ohmann, unpubl. data), support this assumption. IgA was only found in congenitally infected individuals. Although only infrequently found in fetuses of ungulates (Silverstein *et al.* 1963, Ellis *et al.* 1978), IgA may be produced in response to a persistent infection, i.e., as a result of endogenous induction of a secondary response (Ohmann 1981).

Table 1. Results of the virological bacteriological and serological investigations of 71 fetuses and calves.

Gestational age in months	Number of fetuses investigated	Virological examination	Bacteriological examination	Rocket immunoelectrophoresis		Radial immunodiffusion			Pathological changes	BVDV-SA in serum
				IgG	IgM	IgA	IgG	IgM		
abattoir										
≤ 2	9	0/9 ^z	ND	0/9	0/9	ND	ND	ND	0/9	0/9
2-3	8	0/8	0/5	0/8	0/8	ND	ND	ND	0/8	0/8
3-4	8	0/8	0/2	0/8	1/8	0/1	0/1	0/1	0/8	0/8
4-5	14	0/14	0/6	0/14	0/14	ND	0/4	ND	0/14	0/14
5-6	9 ^d	0/9	0/7	1/9	0/9	0/5	1/9	0/4	0/9	0/9
6-7	3	0/3	0/1	0/3	2/3	0/3	0/3	0/3	0/3	0/3
7-8	2	0/2	0/1	2/2	2/2	0/2	1/2	0/2	0/2	0/2
8-9	1	0/1	0/1	1/1	1/1	0/1	1/1	0/1	0/1	0/1
aborted and infected										
5-6	4 ^b	4/4 ^x	0/4	4/4	4/4	1/4	4/4	4/4	4/4	2/4
6-7	2 ^a	2/2 ^x	1/2 ^c	2/2	2/2	ND	ND	ND	2/2	1/2
7½-8½	6 ^a	3/6 ^x	0/6	6/6	6/6	0/2	0/2	0/2	5/6	1/6
8½-9	2 ^y	2/2 ^x	1/2	2/2	2/2	1/2	2/2	2/2	2/2	0/2
9	3 ^q	1/3 ^x	ND	3/3	3/3	1/3	3/3	3/3	2/3	2/3

a aborted fetuses.

b experimentally BVDV-infected fetuses (Ohmann *et al.* 1981 b).

c the isolated bacterial flora was regarded as contamination.

d this group includes one sham-inoculated fetus (Ohmann *et al.* 1981 b).

q stillborn or dead immediately after birth without preceding colostrum intake.

x BVDV was isolated and/or the BVDV-antigen was demonstrated in all positive cases.

y premature birth; the calves were killed or died shortly after birth.

z number positive/number examined.

ND the test was not done.

The high immunoglobulin levels were correlated with isolation of BVDV or detection of viral antigen in tissues in 12 animals (70.6 %) and with pathological lesions in 15 animals (88.2 %). The microbiological findings are obviously not representative of the causes of abortion in Danish cattle herds. The results of the immunological investigations are nevertheless in accordance with data previously reported from elsewhere (Schultz *et al.* 1971, Horner *et al.* 1973, Sawyer *et al.*, Miller & Quinn 1975, Kirkbride *et al.* 1977, Ellis *et al.*).¹⁰ Thus, high serum-

Table 2. Results of the quantitative analysis, by radial immunodiffusion, for IgM, IgG and IgA in serum of control (abattoir)-, infected and aborted fetuses and newborn calves. Comparative data from the literature are included.

Gestational age in months	Number of objects investigated	IgM mg/100 ml		IgG mg/100 ml		IgA mg/100 ml	
		range	average	range	average	range	average
3—4	1	0	—	0	—	0	—
4—5	4	ND	—	0	—	ND	—
5—6	9	0	—	0—3.75 ^e	0.42	0	—
6—7	3	0	—	0	—	0	—
7—8	2	—	—	0—10 ^e	5	0	—
8—9	1	0	—	230	—	0	—
aborted and/or infected							
5—6	4 ^b	10—240	73	11—85	35	0—18 ^e	4.5
7½—8½	2 ^a	20—140	80	65—210	137	0	—
8½—9	2 ^y	20—25	22.5	30—265	147	0—10 ^e	5
9	3 ^q	36—110	69	276—1240	651	0—24 ^e	8
pre-colostral calves							
	100	6—19	11	10—27	16	ND	—
post-colostral calves							
	23	101—301	175	239—2400	755	ND	—
<i>(Sawyer et al. 1973)</i>							

ND the serum was not tested.

a aborted fetuses.

b experimentally BVDV-infected fetuses (*Ohmann et al. 1981 b*).

e only one positive in the group.

q stillborn or dead immediately after birth without preceding colostrum intake.

y premature birth; the calves were killed or died shortly after birth.

immunoglobulin concentrations in aborted fetuses and calves dead without preceding colostrum intake may be a weighty indication of fetal infection. It is concluded that screening of immunoglobulins in bovine fetuses may offer a valuable contribution to clarifying the causes of abortion among Danish cattle stock.

ACKNOWLEDGEMENTS

The author thanks Dr. K. J. Sørensen, the State Veterinary Institute for Virus Research, Lindholm, for performing the virus isolations and Dr. M. Holm Jensen, the State Veterinary Institute for Virus Research, for contributions to the investigations for neutralizing antibodies. The studies have been supported by the Bernhard Bang Memorial Fund and by the V. A. Goldschmidt's Fund.

REFERENCES

- Brambell, F. W. R.*: The Transmission of Passive Immunity from Mother to Young. American Elsevier Publ. Comp., New York 1970.
- Dalsgaard, K., E. Overby, J. J. Metzger & A. Basse*: Rapid method for screening of immunoglobulins in porcine fetuses, using rocket immunoelectrophoresis. Application of an interspecies reaction between human and porcine μ -chain. *Acta vet. scand.* 1979, 20, 313—320.
- Ellis, W. A., E. F. Logan & J. J. O'Brian*: Serum immunoglobulins in aborted and non-aborted bovine fetuses. *Clin. exp. Immunol.* 1978, 33, 136—141.
- Horner, G. W., R. H. Johnson, D. P. Dennett & W. R. Lane*: A serological study of bovine foetal immunoglobulins. *Aust. vet. J.* 1973, 49, 325—329.
- Hubbert, W. T., O. H. V. Stalheim & G. D. Booth*: Changes in organ weights and fluid volumes during growth of the bovine fetus. *Growth* 1972, 36, 217—233.
- Jensen, M. Holm*: Detection of antibodies against hog cholera virus and bovine viral diarrhoea virus in porcine serum. *Acta vet. scand.* 1981, 22, 85—98.
- Kirkbride, C. A., D. Martinovich & D. A. Woodhouse*: Immunoglobulins and lesions in aborted bovine foetuses. *N. Z. vet. J.* 1977, 25, 180—187.
- Miller, R. B. & P. J. Quinn*: Observations on abortion in cattle: a comparison of pathological, microbiological and immunological findings in aborted foetuses and foetuses collected at abattoirs. *Canad. J. comp. Med.* 1975, 39, 270—290.
- Ohmann, H. Bielefeldt*: I. The bovine Immune System. II. Bovine viral diarrhoea virus-infections. Ph.D. thesis, Copenhagen 1981.
- Ohmann, H. Bielefeldt, M. Holm Jensen, K. J. Sørensen & K. Dalsgaard*: Demonstration of bovine viral diarrhoea virus antigen in cryostat- and paraffin-sections of bovine tissues by the immunoperoxidase technique. *Acta path. microbiol. scand. Sect. C* 1981 a, 89, in press.
- Ohmann, H. Bielefeldt, M. Holm Jensen, K. J. Sørensen & K. Dalsgaard*: Experimental fetal infection with bovine viral diarrhoea virus (BVDV). In preparation, 1981 b.

- Olitzki, A. L.*: Studies on the antigenic structure of virulent and non-virulent Brucellae with the aid of agar gel precipitation technique. *Brit. J. exp. Pathol.* 1959, 40, 432—440.
- Sawyer, M., B. I. Osburn, H. D. Knight & J. W. Kendrick*: A quantitative serologic assay for diagnosing congenital infections of cattle. *Amer. J. vet. Res.* 1973, 34, 1281—1284.
- Schultz, R. D.*: Developmental aspects of the fetal bovine immune response: a review. *Cornell Vet.* 1973, 63, 507—535.
- Schultz, R. D., F. Confer & H. W. Dunne*: Occurrence of blood cells and serum proteins in bovine fetuses and calves. *Canad. J. comp. Med.* 1971, 35, 93—98.
- Silverstein, A. M., G. J. Thorbecke, K. L. Kraner & R. J. Lukes*: Foetal responses to antigenic stimulus. III. Gamma-globulin production in normal and stimulated foetal lambs. *J. Immunol.* 1963, 91, 384—395.

SAMMENDRAG

Immunoglobulin niveau hos ikke-aborterede og aborterede fostre fra danske kvægbesætninger.

Anvendeligheden af serologiske tests, som et bidrag til diagnostisering af intrauterine infektioner, blev undersøgt på et dansk kvægmateriale. Fostre fra slagtekvæg, eksperimentelt inficerede og spontant aborterede fostre samt præmature og dødfødte kalve blev underkastet makro- og mikroskopiske, mikrobiologiske og serologiske undersøgelser. Sidstnævnte omfattede raket immunoelektroforese og radial immundiffusion.

Immunoglobuliner var kun påviselige i små mængder hos ganske få fostre fra slagtekvæg, mens sera fra fostre og kalve med patologiske forandringer og/eller verificeret infektion indeholdt betydelige mængder IgM og IgG. IgA kunne også påvises i sera fra denne individgruppe. Resultaterne viser den diagnostiske betydning af en immunoglobulinbestemmelse for aborterede fostre og dødfødte kalve.

(Received July 13, 1981).

Reprints may be requested from: H. Bielefeldt Ohmann, the Department of Pathology, Royal Veterinary and Agricultural University, Bülowsvej 13, DK-1870 Copenhagen V, Denmark.