

FACTORIZATION OF THE IDENTITY THROUGH OPERATORS WITH LARGE DIAGONAL

NIELS JAKOB LAUSTSEN, RICHARD LECHNER, AND PAUL F.X. MÜLLER

ABSTRACT. Given a Banach space X with an unconditional basis, we consider the following question: does the identity operator on X factor through every operator on X with large diagonal relative to the unconditional basis? We show that on Gowers' unconditional Banach space, there exists an operator for which the answer to the question is negative. By contrast, for any operator on the mixed-norm Hardy spaces $H^p(H^q)$, where $1 \leq p, q < \infty$, with the bi-parameter Haar system, this problem always has a positive solution. The spaces L^p , $1 < p < \infty$, were treated first by Andrew [Studia Math. 1979].

1. INTRODUCTION

Let X be a Banach space. A *basis* for X will always mean a Schauder basis. We denote by I_X the identity operator on X , and write $\langle \cdot, \cdot \rangle$ for the bilinear duality pairing between X and its dual space X^* . By an *operator* on X , we understand a bounded and linear mapping from X into itself.

Suppose that X has a normalized basis $(b_n)_{n \in \mathbb{N}}$, and let $b_n^* \in X^*$ be the n^{th} coordinate functional. For an operator T on X , we say that:

- ▷ T has large diagonal if $\inf_{n \in \mathbb{N}} |\langle Tb_n, b_n^* \rangle| > 0$;
- ▷ T is diagonal if $\langle Tb_m, b_n^* \rangle = 0$ whenever $m, n \in \mathbb{N}$ are distinct;
- ▷ the identity operator on X factors through T if there are operators R and S on X such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{I_X} & X \\ R \downarrow & & \uparrow S \\ X & \xrightarrow{T} & X \end{array}$$

is commutative.

Suppose that the basis $(b_n)_{n \in \mathbb{N}}$ for X is unconditional. Then the diagonal operators on X correspond precisely to the elements of $\ell_\infty(\mathbb{N})$, and so for each operator T on X with large diagonal, there is a diagonal operator S on X such that $\langle STb_n, b_n^* \rangle = 1$ for each $n \in \mathbb{N}$. This observation naturally leads to the following question.

Question 1.1. *Can the identity operator on X be factored through each operator on X with large diagonal?*

In classical Banach spaces such as ℓ^p with the unit vector basis and L^p with the Haar basis, the answer to this question is known to be positive. These are the

Date: March 1, 2018.

2010 Mathematics Subject Classification. Primary: 30H10, 47A68, 47B37. Secondary: 46B15, 46B25, 47A53 .

Key words and phrases. Factorization of operators, mixed-norm Hardy spaces, Fredholm theory, Gowers-Maurey spaces.

The research of Lechner and Müller is supported by the Austrian Science Foundation (FWF) Pr.Nr. P23987 P22549.

theorems of Pełczyński [19] and Andrew [2], respectively; see also Johnson, Maurey, Schechtman and Tzafriri [10, Chapter 9].

The aim of the present paper is to establish the following two results.

- ▷ There exists a Banach space with an unconditional basis for which the answer to Question 1.1 is negative. This result relies heavily on the deep work of Gowers [7] and Gowers-Maurey [8].
- ▷ Question 1.1 has a positive answer for the mixed-norm Hardy space $H^p(H^q)$, where $1 \leq p, q < \infty$, with the bi-parameter Haar system as its unconditional basis. This conclusion can be viewed as a bi-parameter extension of Andrew's theorem [2] on the perturbability of the one-parameter Haar system in L^p .

The precise statements of these results, together with their proofs, are given in Sections 2 and 3–5, respectively.

Acknowledgements. It is our pleasure to thank Th. Schlumprecht for very informative conversations and for encouraging the collaboration between Lancaster and Linz. Special thanks are due to J. B. Cooper (Linz) for drawing our attention to the work of Andrew [2].

2. THE ANSWER TO QUESTION 1.1 IS NOT ALWAYS POSITIVE

The aim of this section is to establish the following result, which answers Question 1.1 in the negative.

Theorem 2.1. *There is an operator T on a Banach space X with an unconditional basis such that T has large diagonal, but the identity operator on X does not factor through T .*

The proof of Theorem 2.1 relies on two ingredients. The first of these is Fredholm theory, which we shall now recall the relevant parts of.

Given an operator T on a Banach space X , we set

$$\alpha(T) = \dim \ker T \in \mathbb{N}_0 \cup \{\infty\} \quad \text{and} \quad \beta(T) = \dim(X/T(X)) \in \mathbb{N}_0 \cup \{\infty\},$$

and we say that:

- ▷ T is an *upper semi-Fredholm operator* if $\alpha(T) < \infty$ and T has closed range;
- ▷ T is a *Fredholm operator* if $\alpha(T) < \infty$ and $\beta(T) < \infty$.

Note that the condition $\beta(T) < \infty$ implies that T has closed range (see, e.g., [4, Corollary 3.2.5]), so that each Fredholm operator is automatically upper semi-Fredholm. For an upper semi-Fredholm operator T , we define its *index* by

$$i(T) = \alpha(T) - \beta(T) \in \mathbb{Z} \cup \{-\infty\}.$$

The main property of the class of upper semi-Fredholm operators that we shall require is that it is stable under strictly singular perturbations in the following precise sense. Let T be an upper semi-Fredholm operator on a Banach space X , and suppose that S is an operator on X which is *strictly singular* in the sense that, for each $\varepsilon > 0$, every infinite-dimensional subspace of X contains a unit vector x such that $\|Sx\| \leq \varepsilon$. Then $T + S$ is an upper semi-Fredholm operator, and

$$i(T + S) = i(T).$$

A proof of this result can be found in [14, Proposition 2.c.10].

We shall require the following piece of notation in the proof of our next lemma. For an element x of a Banach space X and a functional $f \in X^*$, we write $x \otimes f$ for the rank-one operator on X defined by

$$(x \otimes f)y = \langle y, f \rangle x \quad (y \in X).$$

Lemma 2.2. *Let T be a diagonal upper semi-Fredholm operator on a Banach space with a basis. Then $\beta(T) = \alpha(T)$, so that T is a Fredholm operator with index 0.*

Proof. Let X be the Banach space on which T acts, and let $(b_n)_{n \in \mathbb{N}}$ be the basis for X with respect to which T is diagonal. Set $N = \{n \in \mathbb{N} : Tb_n = 0\}$. Since T is diagonal, we have $\ker T = \overline{\text{span}}\{b_n : n \in N\}$, and so the set N is finite, with $\alpha(T)$ elements. Consequently, we can define a projection of X onto $\ker T$ by $P_N = \sum_{n \in N} b_n \otimes b_n^*$. The fact that $\ker P_N = \overline{\text{span}}\{b_n : n \in \mathbb{N} \setminus N\}$ implies that $T(X) \subseteq \ker P_N$. Conversely, for each $n \in \mathbb{N} \setminus N$, we have $b_n = T(\langle Tb_n, b_n^* \rangle^{-1} b_n)$, so we conclude that $\ker P_N \subseteq T(X)$ because T has closed range. Hence

$$\beta(T) = \dim P_N(X) = \alpha(T) < \infty,$$

and the result follows. \square

The other main ingredient in the proof of Theorem 2.1 is the Banach space X_G which Gowers [7] created to solve Banach's hyperplane problem. This Banach space has subsequently been investigated in more detail by Gowers and Maurey [8, Section (5.1)]. Its main properties are as follows.

Theorem 2.3 (Gowers [7]; Gowers and Maurey [8]). *There is a Banach space X_G with an unconditional basis such that each operator on X_G is the sum of a diagonal operator and a strictly singular operator.*

Corollary 2.4. *Each upper semi-Fredholm operator on the Banach space X_G is a Fredholm operator of index 0.*

Proof. Let T be an upper semi-Fredholm operator on X_G . By Theorem 2.3, we can find a diagonal operator D and a strictly singular operator S on X_G such that $T = D + S$. The stability of the class of upper semi-Fredholm operators under strictly singular perturbations that we stated above implies that D is an upper semi-Fredholm operator with the same index as T , and hence the conclusion follows from Lemma 2.2. \square

Proof of Theorem 2.1. Let $X = X_G$ be the Banach space from Theorem 2.3, and let $(b_n)_{n \in \mathbb{N}}$ be the unconditional basis for X_G with respect to which each operator on X_G is the sum of a diagonal operator and a strictly singular operator. We may suppose that $(b_n)_{n \in \mathbb{N}}$ is normalized. Set

$$T = I_{X_G} + b_1 \otimes b_2^* + b_2 \otimes b_1^*.$$

Then T has large diagonal because $\langle Tb_n, b_n^* \rangle = 1$ for each $n \in \mathbb{N}$.

Assume towards a contradiction that $I_{X_G} = STR$ for some operators R and S on X_G . Then R is injective, and its range is complemented (because RST is a projection onto it), and it is thus closed, so that R is an upper semi-Fredholm operator with $\alpha(R) = 0$. This implies that R is a Fredholm operator of index 0 by Corollary 2.4, and hence R is invertible. Since ST is a left inverse of R , the uniqueness of the inverse shows that $R^{-1} = ST$, but this contradicts that the operator T is not injective (because $T(b_1 - b_2) = 0$). \square

As we have seen in the proof of Theorem 2.1, the identity operator need not factor through a Fredholm operator. If, however, we allow ourselves sums of two operators, then we can always factor the identity operator, as the following result shows.

Proposition 2.5. *Let T be a Fredholm operator on an infinite-dimensional Banach space X . Then there are operators R_1, R_2, S_1 , and S_2 on X such that*

$$I_X = S_1TR_1 + S_2TR_2.$$

Proof. Let $P = \sum_{j=1}^n x_j \otimes f_j$ be a projection of X onto the kernel of T , where $n \in \mathbb{N}$, $x_1, \dots, x_n \in X$, and $f_1, \dots, f_n \in X^*$, and let Q be a projection of X onto the range of T . Since this range is infinite-dimensional, we can find $y_1, \dots, y_n \in X$ and $g_1, \dots, g_n \in X^*$ such that $\langle Ty_j, g_k \rangle = \delta_{j,k}$ (the Kronecker delta) for each $j, k \in \{1, \dots, n\}$. The restriction $\tilde{T}: x \mapsto Tx, \ker P \rightarrow T(X)$, is invertible, so we may define an operator on X by $S_1 = J\tilde{T}^{-1}Q$, where $J: \ker P \rightarrow X$ is the inclusion. Set

$$R_1 = I_X - P, \quad R_2 = \sum_{j=1}^n y_j \otimes f_j, \quad \text{and} \quad S_2 = \sum_{k=1}^n x_k \otimes g_k.$$

Then, for each $z \in X$, we have

$$\begin{aligned} (S_1TR_1 + S_2TR_2)z &= J\tilde{T}^{-1}QT(z - Pz) + \sum_{j,k=1}^n \langle Ty_j, g_k \rangle \langle z, f_j \rangle x_k \\ &= (z - Pz) + Pz = z, \end{aligned}$$

from which the conclusion follows. \square

3. THE ANSWER TO QUESTION 1.1 IS POSITIVE IN MIXED-NORM HARDY SPACES

In many classical Banach spaces, the answer to Question 1.1 is known to be positive. This includes ℓ^p , $p \geq 1$, and L^p , $p > 1$, see Pełczyński [19] and Andrew [2], respectively. Closely related to this question is the work of Johnson, Maurey, Schechtman and Tzafriri [10, Chapter 9], in which they specify a criterion for an operator on a rearrangement invariant function space to be a factor of the identity.

We now turn to defining the mixed-norm Hardy spaces together with an unconditional basis, the bi-parameter Haar system. Let \mathcal{D} denote the collection of dyadic intervals given by

$$\mathcal{D} = \{[k2^{-n}, (k+1)2^{-n}] : n, k \in \mathbb{N}_0, 0 \leq k \leq 2^n - 1\}.$$

The dyadic intervals are nested, i.e. if $I, J \in \mathcal{D}$, then $I \cap J \in \{I, J, \emptyset\}$. For $I \in \mathcal{D}$ we let $|I|$ denote the length of the dyadic interval I . Let $I \in \mathcal{D}$ and $I \neq [0, 1)$, then \tilde{I} is the unique dyadic interval satisfying $\tilde{I} \supset I$ and $|\tilde{I}| = 2|I|$. Given $N_0 \in \mathbb{N}_0$ we define

$$\mathcal{D}_{N_0} = \{I \in \mathcal{D} : |I| = 2^{-N_0}\} \quad \text{and} \quad \mathcal{D}^{N_0} = \{I \in \mathcal{D} : |I| \geq 2^{-N_0}\}.$$

Let h_I be the L^∞ -normalized Haar function supported on $I \in \mathcal{D}$; that is, for $I = [a, b)$ and $c = (a+b)/2$, we have $h_I(x) = 1$ if $a \leq x < c$, $h_I(x) = -1$ if $c \leq x < b$, and $h_I(x) = 0$ otherwise. Moreover, let $\mathcal{R} = \{I \times J : I, J \in \mathcal{D}\}$ be the collection of dyadic rectangles contained in the unit square, and set

$$h_{I \times J}(x, y) = h_I(x)h_J(y), \quad (I \times J \in \mathcal{R}, x, y \in [0, 1)).$$

For $1 \leq p, q < \infty$, the *mixed-norm Hardy space* $H^p(H^q)$ is the completion of

$$\text{span}\{h_{I \times J} : I \times J \in \mathcal{R}\}$$

under the square function norm

$$\|f\|_{H^p(H^q)} = \left(\int_0^1 \left(\int_0^1 \left(\sum_{I \times J} |a_{I \times J}|^2 h_{I \times J}^2(x, y) \right)^{q/2} dy \right)^{p/q} dx \right)^{1/p}, \quad (3.1)$$

where $f = \sum_{I \times J} a_{I \times J} h_{I \times J}$. Then $(h_{I \times J})_{I \times J \in \mathcal{R}}$ is a 1-unconditional basis of $H^p(H^q)$, called the *bi-parameter Haar system*. We begin with the following facts:

- ▷ It is recorded by Capon [3] that the identity operator provides an isomorphism between $H^p(H^q)$ and $L^p(L^q)$, $1 < p, q < \infty$.
- ▷ Since the bi-parameter Haar system $\{h_{I \times J} : I \times J \in \mathcal{R}\}$ is an unconditional basis, we do not need to specify an ordering of its index set \mathcal{R} .

▷ This basis is L^∞ -normalized and not normalized in $H^p(H^q)$; we have

$$\|h_{I \times J}\|_{H^p(H^q)} = |I|^{1/p} |J|^{1/q}.$$

An operator $T : H^p(H^q) \rightarrow H^p(H^q)$ has large diagonal with respect to the L^∞ -normalized Haar system $\{h_{I \times J} : I \times J \in \mathcal{R}\}$ if and only if for some $\delta > 0$ we have that $|\langle Th_{I \times J}, h_{I \times J} \rangle| \geq \delta |I \times J|$ for all $I \times J \in \mathcal{R}$. The remaining sections of the paper are devoted to proving the following theorem.

Theorem 3.1. *Let $1 \leq p, q < \infty$ and $\delta > 0$, and let $T : H^p(H^q) \rightarrow H^p(H^q)$ be an operator satisfying*

$$|\langle Th_{I \times J}, h_{I \times J} \rangle| \geq \delta |I \times J| \quad \text{for all } I \times J \in \mathcal{R}.$$

Then the identity operator on $H^p(H^q)$ factors through T , that is, there are operators R and S such that the diagram

$$\begin{array}{ccc} H^p(H^q) & \xrightarrow{I_{H^p(H^q)}} & H^p(H^q) \\ R \downarrow & & \uparrow S \\ H^p(H^q) & \xrightarrow{T} & H^p(H^q) \end{array} \quad (3.2)$$

is commutative. Moreover, for any $\eta \in (0, 1]$ the operators R and S can be chosen such that $\|R\| \|S\| \leq (1 + \eta)/\delta$.

For related, local (finite dimensional, quantitative) factorization theorems in bi-parameter H^1 and BMO, see [18, 13]. Recently in [11], the second named author obtained local factorization results in mixed-norm Hardy and BMO spaces by combining methods of the present paper with techniques of [13]. Despite the fact that the constants in our theorem are independent of p and q , we remark that the passage to the non-separable limiting spaces (corresponding to $p = \infty$ or $q = \infty$) cannot be deduced routinely from the proof given below. The non-separable space SL^∞ consisting of functions with square function in L^∞ would be an example of such a limiting space. Factorization theorems in SL^∞ are treated by the second named author in [12].

The cornerstones upon which the constructions of the operators R, S in Theorem 3.1 rest are embeddings and projections onto a carefully chosen block basis of the bi-parameter Haar system in mixed-norm Hardy spaces.

4. CAPON'S LOCAL PRODUCT CONDITION AND ITS CONSEQUENCES

In this section, we treat embeddings and projections in $H^p(H^q)$. They are the main pillars of the construction underlying the proof of Theorem 3.1. We begin by listing some elementary and well known facts concerning $H^p(H^q)$ and its dual.

4.1. Basic facts and notation.

Let $1 \leq p, q < \infty$ and let $H^p(H^q)^*$ denote the dual space of $H^p(H^q)$, identified as a space of functions on $[0, 1]^2$. Then the duality pairing between $H^p(H^q)$ and $H^p(H^q)^*$ is given by

$$\langle f, g \rangle = \int_0^1 \int_0^1 f(x, y) g(x, y) \, dy \, dx.$$

Correspondingly, we have

$$\|g\|_{H^p(H^q)^*} = \sup_{\|f\|_{H^p(H^q)} \leq 1} |\langle f, g \rangle|.$$

Since $h_{I \times J}$, $I \times J \in \mathcal{R}$ is a 1-unconditional Schauder basis in $H^p(H^q)$, we may identify an element $g \in H^p(H^q)^*$ with the sequence $(\langle h_{I \times J}, g \rangle)_{I \times J}$. In the dual

space, the norm of $(\langle h_{I \times J}, g \rangle)_{I \times J}$ is equal to the norm of $(\langle h_{I \times J}, g \rangle)_{I \times J}$. See [14, Chapter 1].

If $1 < p, p', q, q' < \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$, $\frac{1}{q} + \frac{1}{q'} = 1$, it is recorded by Capon [3] that there is a constant $C_{p,q}$ such that for any finite linear combination f of Haar functions we have

$$C_{p,q}^{-1} \|f\|_{L^p(L^q)} \leq \|f\|_{H^p(H^q)} \leq C_{p,q} \|f\|_{L^p(L^q)}.$$

Consequently, the identity operator provides an isomorphism between $H^p(H^q)$ and $L^p(L^q)$, and the dual of $H^p(H^q)$ identifies with $H^{p'}(H^{q'})$. Capon's argument is based on the observation by Pisier that the UMD_p property of a Banach space does not depend on the value of $1 < p < \infty$. For a proof of Pisier's observation, we refer to [15] respectively [20, Chapter 5].

For the limiting cases we have $H^1(H^q)^* = \text{BMO}(H^{q'})$, $H^p(H^1)^* = H^{p'}(\text{BMO})$ and $H^1(H^1)^* = \text{BMO}(\text{BMO})$. See Maurey [16] and Müller [17].

Let $\{\mathcal{B}_R : R \in \mathcal{R}\}$ be a pairwise disjoint family, where each set \mathcal{B}_R is a finite collection of disjoint dyadic rectangles. Given a vector of scalars $\beta = (\beta_R : R \in \bigcup_{Q \in \mathcal{R}} \mathcal{B}_Q)$, we define

$$b_R^{(\beta)}(x, y) = \sum_{Q \in \mathcal{B}_R} \beta_Q h_Q(x, y), \quad x, y \in [0, 1) \quad (4.1)$$

and we call $\{b_R^{(\beta)} : R \in \mathcal{R}\}$ the *block basis generated by* $\{\mathcal{B}_R : R \in \mathcal{R}\}$ and $\beta = (\beta_R : R \in \bigcup_{Q \in \mathcal{R}} \mathcal{B}_Q)$. Now, let $1 \leq p, q < \infty$ be fixed. Note that $\{b_R^{(\beta)} : R \in \mathcal{R}\}$ is 1-unconditional in $H^p(H^q)$ since $\{h_R : R \in \mathcal{R}\}$ is 1-unconditional in $H^p(H^q)$, i.e.

$$\left\| \sum_{R \in \mathcal{R}} \gamma_R \alpha_R b_R^{(\beta)} \right\|_{H^p(H^q)} \leq \sup_{R \in \mathcal{R}} |\gamma_R| \left\| \sum_{R \in \mathcal{R}} \alpha_R b_R^{(\beta)} \right\|_{H^p(H^q)}, \quad (\gamma_R : R \in \mathcal{R}) \in \ell^\infty(\mathcal{R}),$$

whenever the series $\sum_{R \in \mathcal{R}} \alpha_R b_R^{(\beta)}$ converges. We say that the system $\{b_R^{(\beta)} : R \in \mathcal{R}\}$ is equivalent to the Haar system $\{h_R : R \in \mathcal{R}\}$ if the operator $B_\beta : H^p(H^q) \rightarrow H^p(H^q)$ given by

$$B_\beta(f) = \sum_{R \in \mathcal{R}} \frac{\langle f, h_R \rangle}{\|h_R\|_2^2} b_R^{(\beta)}, \quad f \in H^p(H^q),$$

is bounded and an isomorphism onto its range. In this case, whenever $C_1, C_2 > 0$ are constants such that

$$\frac{1}{C_1} \|f\|_{H^p(H^q)} \leq \|B_\beta f\|_{H^p(H^q)} \leq C_2 \|f\|_{H^p(H^q)}, \quad f \in H^p(H^q),$$

we say that $\{b_R^{(\beta)} : R \in \mathcal{R}\}$ is $C_1 C_2$ -equivalent to $\{h_R : R \in \mathcal{R}\}$.

If $\beta_R = 1$ for each $R \in \mathcal{R}$, then we write b_R instead of $b_R^{(\beta)}$ and B in place of B_β .

4.2. Uniform weak and weak* limits.

Let Γ denote the closed unit ball of $\ell^\infty(\mathcal{R})$, so that Γ consists of all families $\gamma = (\gamma_R : R \in \mathcal{R})$ of scalars with $|\gamma_R| \leq 1$ for each $R \in \mathcal{R}$. Given $\gamma \in \Gamma$, the 1-unconditionality of the bi-parameter Haar system implies that the definition

$$M_\gamma : h_R \mapsto \gamma_R h_R, \quad R \in \mathcal{R} \quad (4.2)$$

extends uniquely to an operator of norm $\sup_R |\gamma_R|$ on $H^p(H^q)$.

Lemma 4.1. *For $m \in \mathbb{N}$, let \mathcal{X}_m and \mathcal{Y}_m be non-empty, finite families of pairwise disjoint dyadic intervals, define $f_m = \sum_{I \in \mathcal{X}_m, J \in \mathcal{Y}_m} h_{I \times J}$, $X_m = \bigcup \mathcal{X}_m$, and $Y_m = \bigcup \mathcal{Y}_m$, and let $1 \leq p, q < \infty$. Then:*

- (i) $\|f_m\|_{H^p(H^q)} = |X_m|^{1/p} |Y_m|^{1/q}$ for all $m \in \mathbb{N}$;
- (ii) $\|f_m\|_{H^p(H^q)^*} = |X_m|^{1-1/p} |Y_m|^{1-1/q}$ for all $m \in \mathbb{N}$.

Suppose in addition that:

- ▷ $\mathcal{X}_m \cap \mathcal{X}_n = \emptyset$ or $\mathcal{Y}_m \cap \mathcal{Y}_n = \emptyset$ whenever $m, n \in \mathbb{N}$ are distinct;
- ▷ $X_m = X_n$ and $Y_m = Y_n$ for all $m, n \in \mathbb{N}$.

Then:

- (iii) the sequence $(|X_m|^{-1/p}|Y_m|^{-1/q}f_m)_{m \in \mathbb{N}}$ in $H^p(H^q)$ is isometrically equivalent to the unit vector basis of ℓ^2 ;
- (iv) for each $g \in H^p(H^q)^*$, $\sup_{\gamma \in \Gamma} |\langle M_\gamma f_m, g \rangle| \rightarrow 0$ as $m \rightarrow \infty$;
- (v) for each $g \in H^p(H^q)$, $\sup_{\gamma \in \Gamma} |\langle M_\gamma g, f_m \rangle| \rightarrow 0$ as $m \rightarrow \infty$.

Note that in (i), (iii), and (iv), we regard f_m as an element of $H^p(H^q)$, whereas in (ii) and (v), we regard it as an element of $H^p(H^q)^*$.

Proof. Set $\mathcal{B}_m = \{I \times J : I \in \mathcal{X}_m, J \in \mathcal{Y}_m\}$ for each $m \in \mathbb{N}$.

(i). This follows immediately from the definition of $\|\cdot\|_{H^p(H^q)}$.

(ii). For any $g = \sum_{K \times L \in \mathcal{B}_m} a_{K \times L} h_{K \times L} \in H^p(H^q)$ we obtain by Hölder's inequality that

$$\begin{aligned} |\langle f_m, g \rangle| &\leq \sum_{K \in \mathcal{X}_m} |K| \sum_{L \in \mathcal{Y}_m} |a_{K \times L}| |L| \leq |Y_m|^{1-1/q} \sum_{K \in \mathcal{X}_m} |K| \left(\sum_{L \in \mathcal{Y}_m} |a_{K \times L}|^q |L| \right)^{1/q} \\ &\leq |X_m|^{1-1/p} |Y_m|^{1-1/q} \left(\sum_{K \in \mathcal{X}_m} |K| \left(\sum_{L \in \mathcal{Y}_m} |a_{K \times L}|^q |L| \right)^{p/q} \right)^{1/p} \\ &= |X_m|^{1-1/p} |Y_m|^{1-1/q} \|g\|_{H^p(H^q)}, \end{aligned}$$

and thus we have proved $\|f_m\|_{H^p(H^q)^*} \leq |X_m|^{1-1/p} |Y_m|^{1-1/q}$. For the other inequality, recall from (i) that $\|f_m\|_{H^p(H^q)} = |X_m|^{1/p} |Y_m|^{1/q}$, thus

$$\langle f_m, f_m \rangle = |X_m| |Y_m| = |X_m|^{1-1/p} |Y_m|^{1-1/q} \|f_m\|_{H^p(H^q)}.$$

(iii). We observe that the first of the additional assumptions ensures that $\mathcal{B}_m \cap \mathcal{B}_n = \emptyset$ whenever $m, n \in \mathbb{N}$ are distinct. Set $X := X_m$ and $Y := Y_m$ for some (and hence all) $m \in \mathbb{N}$, and let $(c_m)_{m \in \mathbb{N}}$ be a sequence of scalars that vanishes eventually. Since

$$\sum_{R \in \mathcal{B}_m} \mathbb{1}_R(x, y) = \left(\sum_{I \in \mathcal{X}_m} \mathbb{1}_I(x) \right) \left(\sum_{J \in \mathcal{Y}_m} \mathbb{1}_J(y) \right) = \mathbb{1}_X(x) \mathbb{1}_Y(y)$$

for all $m \in \mathbb{N}$ and $x, y \in [0, 1)$, (3.1) implies that

$$\begin{aligned} \left\| \sum_m c_m f_m \right\|_{H^p(H^q)}^p &= \int_0^1 \left(\int_0^1 \left(\sum_m |c_m|^2 \mathbb{1}_X(x) \mathbb{1}_Y(y) \right)^{q/2} dy \right)^{p/q} dx \\ &= \left(\sum_m |c_m|^2 \right)^{p/2} |X| |Y|^{p/q}, \end{aligned}$$

from which the conclusion follows.

(iv). Let $g \in H^p(H^q)^*$ and $\varepsilon > 0$. For each $R \in \mathcal{R}$, we can choose a scalar β_R with $|\beta_R| = 1$ such that $\beta_R \langle h_R, g \rangle = |\langle h_R, g \rangle|$. Set $\beta = (\beta_R) \in \Gamma$. By (iii), the sequence $(f_m)_{m \in \mathbb{N}}$ converges weakly to 0, so we can find $m_0 \in \mathbb{N}$ such that $|\langle f_m, M_\beta^* g \rangle| \leq \varepsilon$ whenever $m \geq m_0$. Then, for each $\gamma = (\gamma_R) \in \Gamma$ and $m \geq m_0$ we have

$$\begin{aligned} |\langle M_\gamma f_m, g \rangle| &= \left| \sum_{R \in \mathcal{B}_m} \gamma_R \langle h_R, g \rangle \right| \leq \sum_{R \in \mathcal{B}_m} |\langle h_R, g \rangle| \\ &= \sum_{R \in \mathcal{B}_m} \beta_R \langle h_R, g \rangle = \langle M_\beta f_m, g \rangle \leq \varepsilon, \end{aligned}$$

as required.




FIGURE 1. For a dyadic rectangle $I \times J \in \mathcal{R}$, this figure depicts $\mathcal{B}_{I \times J} = \mathcal{X}_{I \times J} \times \mathcal{Y}_{I \times J}$ (the collection of the dark gray rectangles) contained in the unit square (the light gray area). Here, $\mathcal{X}_{I \times J} = \{K_0, K_1, K_2\}$. The dyadic rectangles in $K_i \times \mathcal{Y}_{I \times J}$ are connected by dotted lines.

(v). Given $g \in H^p(H^q)$ and $\varepsilon > 0$, we choose a finite subset \mathcal{F} of \mathcal{R} such that $\|g - Pg\|_{H^p(H^q)} \leq \varepsilon$, where $P : H^p(H^q) \rightarrow H^p(H^q)$ is the orthogonal projection given by $Pf = \sum_{R \in \mathcal{F}} \frac{\langle f, h_R \rangle}{|R|} h_R$. Since the sets \mathcal{B}_m , $m \in \mathbb{N}$, are pairwise disjoint and \mathcal{F} is finite, we can find $m_0 \in \mathbb{N}$ such that $(\bigcup_{m \geq m_0} \mathcal{B}_m) \cap \mathcal{F} = \emptyset$. Then, for each $m \geq m_0$ and $\gamma \in \Gamma$, we have $P^* f_m = 0$, and hence

$$\begin{aligned} |\langle M_\gamma g, f_m \rangle| &= |\langle M_\gamma g, (I - P)^* f_m \rangle| = |\langle M_\gamma (I - P)g, f_m \rangle| \\ &\leq \|M_\gamma\| \|g - Pg\|_{H^p(H^q)} \|f_m\|_{H^p(H^q)^*} \leq \varepsilon, \end{aligned}$$

where we have used that M_γ commutes with P , and that

$$\|f_m\|_{H^p(H^q)^*} = |X|^{1-1/p} |Y|^{1-1/q} \leq 1$$

by (ii). \square

4.3. Embeddings and projections.

For each $R \in \mathcal{R}$ let $\mathcal{X}_R, \mathcal{Y}_R \subset \mathcal{D}$ denote non-empty, finite collections of dyadic intervals that define the collection of dyadic rectangles \mathcal{B}_R by

$$\mathcal{B}_R = \{K \times L : K \in \mathcal{X}_R, L \in \mathcal{Y}_R\}, \quad R \in \mathcal{R}. \quad (4.3)$$

Now (4.1) assumes the following form, if $\beta_R = 1$ for each $R \in \mathcal{R}$:

$$b_R(x, y) = \left(\sum_{K \in \mathcal{X}_R} h_K(x) \right) \left(\sum_{L \in \mathcal{Y}_R} h_L(y) \right), \quad R \in \mathcal{R}; \quad (4.4)$$

see Figure 1.

Capon [3] discovered a condition for $\{\mathcal{B}_R : R \in \mathcal{R}\}$ which ensures that the block basis $\{b_R : R \in \mathcal{R}\}$ given by (4.4) is equivalent to the Haar system $\{h_R : R \in \mathcal{R}\}$ in $H^p(H^q)$, whenever $1 < p, q < \infty$ (see Theorem 4.2). The local product condition (P1)–(P4) has its roots in Capon's seminal work [3].

We now introduce some notation. For $R \in \mathcal{R}$ we set

$$X_R = \bigcup \{K : K \in \mathcal{X}_R\} \quad \text{and} \quad Y_R = \bigcup \{L : L \in \mathcal{Y}_R\}. \quad (4.5)$$

For each $I_0 \times J_0 \in \mathcal{R}$ we consider the following unions

$$X_{I_0} = \bigcup \{X_{I_0 \times J} : J \in \mathcal{D}\}, \quad Y_{J_0} = \bigcup \{Y_{I \times J_0} : I \in \mathcal{D}\}. \quad (4.6)$$




FIGURE 3. The dyadic rectangles $I \times J$, $I \times J_0$ and $I \times J_1$ in \mathcal{R} are such that $J_0 \cup J_1 = J$ and $J_0 \cap J_1 = \emptyset$. This figure depicts the collections $\mathcal{B}_{I \times J} = \mathcal{X}_{I \times J} \times \mathcal{Y}_{I \times J}$ in the top layer, and $\mathcal{B}_{I \times J_0} = \mathcal{X}_{I \times J_0} \times \mathcal{Y}_{I \times J_0}$ and $\mathcal{B}_{I \times J_1} = \mathcal{X}_{I \times J_1} \times \mathcal{Y}_{I \times J_1}$ in the bottom layer. Here, $\mathcal{X}_{I \times J} = \mathcal{X}_{I_0 \times J} = \mathcal{X}_{I_1 \times J} = \{K_0, K_1, K_2\}$. Each interval in $\mathcal{Y}_{I \times J}$ is split in two intervals, which are then placed into $\mathcal{Y}_{I \times J_0}$ and $\mathcal{Y}_{I \times J_1}$, respectively.

where the constants $C(p, q, C_X, C_Y) \rightarrow \infty$ in each of the cases $p \rightarrow 1$, $p \rightarrow \infty$, $q \rightarrow 1$ or $q \rightarrow \infty$.

The next theorem is our first major step towards proving Theorem 3.1. We show that the operator P is bounded on $H^p(H^q)$, $1 \leq p, q < \infty$ with an upper estimate for the norm independent of p or q . Specifically, Theorem 4.3 includes the cases $p = 1$ or $q = 1$.

Theorem 4.3. *Let $1 \leq p, q < \infty$, let $\{\mathcal{B}_R : R \in \mathcal{R}\}$ be a pairwise disjoint family which satisfies the local product condition (P1)–(P4) with constants C_X and C_Y , and let $\beta = (\beta_Q : Q \in \bigcup_{R \in \mathcal{R}} \mathcal{B}_R)$ be a family of scalars such that*

$$M := \sup_Q |\beta_Q| < \infty.$$

Then the operators $B_\beta, A_\beta : H^p(H^q) \rightarrow H^p(H^q)$ given by

$$B_\beta f = \sum_{R \in \mathcal{R}} \frac{\langle f, h_R \rangle}{\|h_R\|_2^2} b_R^{(\beta)} \quad \text{and} \quad A_\beta f = \sum_{R \in \mathcal{R}} \frac{\langle f, b_R^{(\beta)} \rangle}{\|b_R\|_2^2} h_R$$

satisfy the estimates

$$\begin{aligned} \|B_\beta f\|_{H^p(H^q)} &\leq M C_X^{1/p} C_Y^{1/q} \|f\|_{H^p(H^q)}, & f \in H^p(H^q), \\ \|A_\beta f\|_{H^p(H^q)} &\leq M C_X^{3+1/p} C_Y^{3+1/q} \|f\|_{H^p(H^q)}, & f \in H^p(H^q). \end{aligned} \quad (4.9)$$

If we additionally assume that

$$m := \inf_Q |\beta_Q| > 0,$$




FIGURE 4. In the figure, $\mathcal{X}_{I \times J_j} = \{K_0, K_1, K_2\}$, $0 \leq j \leq 3$, whereas $\mathcal{Y}_{I \times J_j}$ changes with each layer $0 \leq j \leq 3$. For $y_0 \in [0, 1)$, the light red vertical plane connects the lines $\ell = \{(x, y_0) : x \in [0, 1)\}$ in the four layers depicted in the figure.

and if we define the vector of scalars $\gamma = (\gamma_Q : Q \in \bigcup_{R \in \mathcal{R}} \mathcal{B}_R)$ by $\beta_Q \gamma_Q = 1$, then the diagram

$$\begin{array}{ccc}
 H^p(H^q) & \xrightarrow{I_{H^p(H^q)}} & H^p(H^q) \\
 & \searrow B_\beta & \nearrow A_\gamma \\
 & & H^p(H^q)
 \end{array} \tag{4.10}$$

is commutative, and the operator A_γ satisfies the estimate $\|A_\gamma\| \leq m^{-1} C_X^{3+1/p} C_Y^{3+1/q}$. Moreover, the composition $P_{\beta, \gamma} = B_\beta A_\gamma$ is the projection $P_{\beta, \gamma} : H^p(H^q) \rightarrow H^p(H^q)$ given by

$$P_{\beta, \gamma}(f) = \sum_{R \in \mathcal{R}} \frac{\langle f, b_R^{(\gamma)} \rangle}{\|b_R\|_2^2} b_R^{(\beta)}.$$

Consequently, the range of B_β is complemented (by $P_{\beta,\gamma}$), and B_β is an isomorphism onto its range. Finally, if $\beta_Q = \gamma_Q = 1$ for each Q , then $P_{\beta,\gamma}$ coincides with the orthogonal projection P defined by (4.8).

Before we proceed with the proof, we record some simple facts.

Lemma 4.4. *Let $\mathcal{B}_R = \mathcal{X}_R \times \mathcal{Y}_R \subset \mathcal{R}$, $R \in \mathcal{R}$ satisfy the conditions (P1) and (P3). Then*

$$C_X^{-1}C_Y^{-1}|R| \leq \|b_R\|_2^2 \leq C_X C_Y |R|, \quad R \in \mathcal{R}.$$

Proof. Let $R \in \mathcal{R}$ be fixed. By condition (P1) and (4.3), the collections \mathcal{X}_R and \mathcal{Y}_R each consist of pairwise disjoint dyadic intervals, thus, Lemma 4.1 (i) yields

$$\|b_R\|_2^2 = |X_R||Y_R|.$$

By (P3) and (4.7) we obtain

$$C_X^{-1}C_Y^{-1}|R| \leq |X_R||Y_R| \leq C_X C_Y |R|. \quad \square$$

Below we use Minkowski's inequality in various function spaces. For ease of reference, we include it in the form that we need it.

Lemma 4.5. *Let (Ω, μ) be a probability space.*

(i) *Let $1 \leq r < \infty$ and let $g_k \in L^r(\Omega)$ be real valued. Then*

$$\int_{\Omega} \left(\sum_k g_k^2 \right)^{r/2} d\mu \geq \left(\sum_k \left(\int_{\Omega} g_k d\mu \right)^2 \right)^{r/2}.$$

(ii) *Let $1 \leq r, s < \infty$ and let $g_{k,\ell} \in L^s(\Omega)$ be real valued. Then*

$$\int_{\Omega} \left(\sum_k \left(\sum_{\ell} g_{k,\ell}^2 \right)^{s/2} \right)^{r/s} d\mu \geq \left(\sum_k \left(\sum_{\ell} \left(\int_{\Omega} g_{k,\ell} d\mu \right)^2 \right)^{s/2} \right)^{r/s}.$$

Proof. First, we apply Minkowski's inequality (see e.g. [6, Corollary 5.4.2], [9, Theorem 202]) to the integral and the sum over ℓ :

$$\left(\sum_k \left(\sum_{\ell} \left(\int_{\Omega} g_{k,\ell} d\mu \right)^2 \right)^{s/2} \right)^{1/s} \leq \left(\sum_k \left(\int_{\Omega} \left(\sum_{\ell} g_{k,\ell}^2 \right)^{1/2} d\mu \right)^s \right)^{1/s}.$$

Secondly, applying Minkowski's inequality to the integral and the sum over k yields

$$\left(\sum_k \left(\int_{\Omega} \left(\sum_{\ell} g_{k,\ell}^2 \right)^{1/2} d\mu \right)^s \right)^{1/s} \leq \int_{\Omega} \left(\sum_k \left(\sum_{\ell} g_{k,\ell}^2 \right)^{s/2} \right)^{1/s} d\mu.$$

Finally, we obtain (ii) by Hölder's inequality.

The assertion (i) follows from (ii) by putting $s = 2$. \square

Lemma 4.6. *Assume that $(Z_I : I \in \mathcal{D})$ satisfies the following condition: For all $I, I_0, I_1 \in \mathcal{D}$ with $I_0 \cap I_1 = \emptyset$, $I_0 \cup I_1 \subset I$ we have that*

$$Z_{I_0} \cap Z_{I_1} = \emptyset \quad \text{and} \quad Z_{I_0} \cup Z_{I_1} \subset Z_I.$$

Let $0 < r < \infty$, $N_0 \in \mathbb{N}$ and $c_I \geq 0$ and define

$$f(z) = \left(\sum_{I \in \mathcal{D}^{N_0}} c_I \mathbb{1}_{Z_I}(z) \right)^r.$$

Then

$$\tilde{c}_I = \left(\sum_{E \supset I} c_E \right)^r - \left(\sum_{E \supsetneq I} c_E \right)^r$$

satisfies $\tilde{c}_I \geq 0$ and we obtain the identity

$$f(z) = \sum_{I \in \mathcal{D}^{N_0}} \tilde{c}_I \mathbb{1}_{Z_I}(z).$$

Proof. Observe that by telescoping and the tree structure of the sets $(Z_I : I \in \mathcal{D})$ we have that

$$\left(\sum_{I \in \mathcal{D}^{N_0}} c_I \mathbb{1}_{Z_I}(z) \right)^r = \sum_{I \in \mathcal{D}^{N_0}} \tilde{c}_I \mathbb{1}_{Z_I}(z).$$

The fact that $\tilde{c}_I \geq 0$ is self-evident. \square

Proof of Theorem 4.3. The proof will be split into three parts. In the first part, we will give the estimate for B_β , and in the second part, we will establish the estimate for A_β .

PART 1: THE ESTIMATE FOR B_β . We emphasize that our proof of the estimate for B_β only uses the conditions (P1)–(P3); specifically, we do not use (P4).

For $N_0 \in \mathbb{N}$ we define the collections of indices

$$\mathcal{R}_{N_0} = \{I_0 \times J_0 \in \mathcal{R} : I_0, J_0 \in \mathcal{D}_{N_0}\} \quad (4.11a)$$

and

$$\mathcal{R}^{N_0} = \{I_0 \times J_0 \in \mathcal{R} : I_0, J_0 \in \mathcal{D}^{N_0}\}. \quad (4.11b)$$

Let us assume that

$$f = \sum_{R \in \mathcal{R}^{N_0}} a_R h_R.$$

Then by (P1) and (4.3) we find that

$$\|B_\beta f\|_{H^p(H^q)}^p = \int_0^1 \left(\int_0^1 \left(\sum_{R \in \mathcal{R}^{N_0}} |a_R|^2 \sum_{Q \in \mathcal{B}_R} |\beta_Q|^2 \mathbb{1}_Q(x, y) \right)^{q/2} dy \right)^{p/q} dx.$$

Recall that $|\beta_{I \times J}| \leq M$ and that by (4.7) $\mathbb{1}_{X_{I \times J}}(x) \mathbb{1}_{Y_{I \times J}}(y) \leq \mathbb{1}_{X_I}(x) \mathbb{1}_{Y_J}(y)$, so we note

$$\|B_\beta f\|_{H^p(H^q)}^p \leq M^p \int_0^1 \left(\int_0^1 \left(\sum_{I \times J \in \mathcal{R}^{N_0}} |a_{I \times J}|^2 \mathbb{1}_{X_I}(x) \mathbb{1}_{Y_J}(y) \right)^{q/2} dy \right)^{p/q} dx. \quad (4.12)$$

If we define $c_J(x) = \sum_{I \in \mathcal{D}^{N_0}} |a_{I \times J}|^2 \mathbb{1}_{X_I}(x)$, (4.12) reads

$$\|B_\beta f\|_{H^p(H^q)}^p \leq M^p \int_0^1 \left(\int_0^1 \left(\sum_{J \in \mathcal{D}^{N_0}} c_J(x) \mathbb{1}_{Y_J}(y) \right)^{q/2} dy \right)^{p/q} dx. \quad (4.13)$$

Lemma 4.6 yields the following identity for the inner integrand of (4.13):

$$\left(\sum_{J \in \mathcal{D}^{N_0}} c_J(x) \mathbb{1}_{Y_J}(y) \right)^{q/2} = \sum_{J \in \mathcal{D}^{N_0}} \tilde{c}_J(x) \mathbb{1}_{Y_J}(y), \quad (4.14)$$

where $\tilde{c}_J(x) = (\sum_{J_1 \supset J} c_{J_1}(x))^{q/2} - (\sum_{J_1 \supsetneq J} c_{J_1}(x))^{q/2} \geq 0$. Integrating (4.14) with respect to y and using that $|Y_J| \leq C_Y |J|$ by (P3), we have

$$\int_0^1 \left(\sum_{J \in \mathcal{D}^{N_0}} c_J(x) \mathbb{1}_{Y_J}(y) \right)^{q/2} dy \leq C_Y \sum_{J \in \mathcal{D}^{N_0}} \tilde{c}_J(x) |J|.$$

Combining the latter estimate with (4.13) yields

$$\|B_\beta f\|_{H^p(H^q)}^p \leq M^p C_Y^{p/q} \int_0^1 \left(\sum_{J \in \mathcal{D}^{N_0}} \tilde{c}_J(x) |J| \right)^{p/q} dx. \quad (4.15)$$

It remains to estimate $\int_0^1 (\sum_{J \in \mathcal{D}^{N_0}} \tilde{c}_J(x) |J|)^{p/q} dx$ from above by a constant multiple of $\|f\|_{H^p(H^q)}^p$. Note that

$$\begin{aligned} \left(\sum_{J_1 \supset J} c_{J_1}(x) \right)^{q/2} &= \left(\sum_{I \in \mathcal{D}^{N_0}} d_{I,J} \mathbb{1}_{X_I}(x) \right)^{q/2}, & \text{where } d_{I,J} &= \sum_{J_1 \supset J} |a_{I \times J_1}|^2, \\ \left(\sum_{J_1 \supsetneq J} c_{J_1}(x) \right)^{q/2} &= \left(\sum_{I \in \mathcal{D}^{N_0}} e_{I,J} \mathbb{1}_{X_I}(x) \right)^{q/2}, & \text{where } e_{I,J} &= \sum_{J_1 \supsetneq J} |a_{I \times J_1}|^2, \end{aligned}$$

and that $\tilde{c}_J(x)$ was defined as the difference between the two quantities, above. By Lemma 4.6, we obtain

$$\begin{aligned} \left(\sum_{I \in \mathcal{D}^{N_0}} d_{I,J} \mathbb{1}_{X_I}(x) \right)^{q/2} &= \sum_{I \in \mathcal{D}^{N_0}} \tilde{d}_{I,J} \mathbb{1}_{X_I}(x), \\ \left(\sum_{I \in \mathcal{D}^{N_0}} e_{I,J} \mathbb{1}_{X_I}(x) \right)^{q/2} &= \sum_{I \in \mathcal{D}^{N_0}} \tilde{e}_{I,J} \mathbb{1}_{X_I}(x), \end{aligned}$$

where

$$\begin{aligned} \tilde{d}_{I,J} &= \left(\sum_{I_1 \supset I} d_{I_1,J} \right)^{q/2} - \left(\sum_{I_1 \supsetneq I} d_{I_1,J} \right)^{q/2} \geq 0, \\ \tilde{e}_{I,J} &= \left(\sum_{I_1 \supset I} e_{I_1,J} \right)^{q/2} - \left(\sum_{I_1 \supsetneq I} e_{I_1,J} \right)^{q/2} \geq 0. \end{aligned}$$

Summing up, in between (4.15) and here, we have shown that

$$\|B_\beta f\|_{H^p(H^q)}^p \leq M^p C_Y^{p/q} \int_0^1 \left(\sum_{I \in \mathcal{D}^{N_0}} f_I \mathbb{1}_{X_I}(x) \right)^{p/q} dx, \quad (4.16)$$

where $f_I = \sum_{J \in \mathcal{D}^{N_0}} |J| (\tilde{d}_{I,J} - \tilde{e}_{I,J})$.

It is important to show that $f_I \geq 0$, for all $I \in \mathcal{D}^{N_0}$. To this end, note the identity

$$\begin{aligned} \tilde{d}_{I,J} - \tilde{e}_{I,J} &= \left(\sum_{\substack{I_1 \supset I \\ J_1 \supset J}} |a_{I_1 \times J_1}|^2 \right)^{q/2} - \left(\sum_{\substack{I_1 \supsetneq I \\ J_1 \supset J}} |a_{I_1 \times J_1}|^2 \right)^{q/2} \\ &\quad - \left(\sum_{\substack{I_1 \supset I \\ J_1 \supsetneq J}} |a_{I_1 \times J_1}|^2 \right)^{q/2} + \left(\sum_{\substack{I_1 \supsetneq I \\ J_1 \supsetneq J}} |a_{I_1 \times J_1}|^2 \right)^{q/2}. \end{aligned}$$

Let $J_0 \in \mathcal{D}_{N_0}$, then grouping together the first with the third term as well as the second with the fourth, and summing the latter identity over $J \supset J_0$ yields

$$\sum_{J \supset J_0} \tilde{d}_{I,J} - \tilde{e}_{I,J} = \left(\sum_{\substack{I_1 \supset I \\ J_1 \supset J_0}} |a_{I_1 \times J_1}|^2 \right)^{q/2} - \left(\sum_{\substack{I_1 \supsetneq I \\ J_1 \supset J_0}} |a_{I_1 \times J_1}|^2 \right)^{q/2} \geq 0.$$

Since we have

$$f_I = \sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \sum_{J \supset J_0} (\tilde{d}_{I,J} - \tilde{e}_{I,J}),$$

we showed that $f_I \geq 0$.

A final application of Lemma 4.6 gives

$$\int_0^1 \left(\sum_{I \in \mathcal{D}^{N_0}} f_I \mathbb{1}_{X_I}(x) \right)^{p/q} dx = \int_0^1 \sum_{I \in \mathcal{D}^{N_0}} \tilde{f}_I \mathbb{1}_{X_I}(x) dx = \sum_{I \in \mathcal{D}^{N_0}} \tilde{f}_I |X_I|,$$

where $\tilde{f}_I = (\sum_{I_1 \supset I} f_{I_1})^{p/q} - (\sum_{I_1 \supsetneq I} f_{I_1})^{p/q} \geq 0$. Using (P3) in the above identity and combining it with (4.16) yields

$$\|B_\beta f\|_{H^p(H^q)}^p \leq C_X M^p C_Y^{p/q} \sum_{I \in \mathcal{D}^{N_0}} \tilde{f}_I |I|.$$

Finally, we remark that

$$\|f\|_{H^p(H^q)}^p = \sum_{I \in \mathcal{D}^{N_0}} \tilde{f}_I |I|.$$

To see this, it suffices to apply Lemma 4.6 as above.

PART 2: THE ESTIMATE FOR A_β . Let $N_0 \in \mathbb{N}$, and define the collections of building blocks \mathcal{B}_{N_0} and \mathcal{R}^{N_0} by

$$\mathcal{B}_{N_0} = \{K_0 \times L_0 \in \mathcal{B}_{I_0 \times J_0} : I_0 \times J_0 \in \mathcal{R}_{N_0}\}$$

and

$$\mathcal{R}^{N_0} = \{K \times L \in \mathcal{B}_{I \times J} : I \times J \in \mathcal{R}^{N_0}\},$$

where \mathcal{R}_{N_0} and \mathcal{R}^{N_0} are defined in (4.11). Taking into account that the bi-parameter Haar system is a 1-unconditional basis of $H^p(H^q)$, it suffices to consider only those f that can be written as follows:

$$f = \sum_{K \times L \in \mathcal{B}^{N_0}} a_{K \times L} h_{K \times L}.$$

We will now estimate $\|A_\beta f\|_{H^p(H^q)}^p$. To this end, note that by the definitions of A_β and the norm in $H^p(H^q)$ we have

$$\|A_\beta f\|_{H^p(H^q)}^p = \int_0^1 \left(\int_0^1 \left(\sum_{R \in \mathcal{R}^{N_0}} \frac{|\langle f, b_R^{(\beta)} \rangle|^2}{\|b_R\|_2^4} \mathbb{1}_R(x, y) \right)^{q/2} dy \right)^{p/q} dx.$$

Since \mathcal{D}_{N_0} is a partition of the unit interval, we obtain that

$$\|A_\beta f\|_{H^p(H^q)}^p = \sum_{I_0 \in \mathcal{D}_{N_0}} \int_{I_0} \left(\sum_{J_0 \in \mathcal{D}_{N_0}} \int_{J_0} \left(\sum_{R \in \mathcal{R}^{N_0}} \frac{|\langle f, b_R^{(\beta)} \rangle|^2}{\|b_R\|_2^4} \mathbb{1}_R(x, y) \right)^{q/2} dy \right)^{p/q} dx.$$

Recall that $|\beta_Q| \leq M$, note that for $I_0, J_0 \in \mathcal{D}_{N_0}$ and $R \in \mathcal{R}^{N_0}$ as in the above sums, $\mathbb{1}_R(x, y) = 1$ exactly when $R \supset I_0 \times J_0$, and apply Lemma 4.4 to obtain

$$\begin{aligned} & \|A_\beta f\|_{H^p(H^q)}^p \\ & \leq M^p C_X^p C_Y^p \sum_{I_0 \in \mathcal{D}_{N_0}} |I_0| \left(\sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \left(\sum_{\substack{R \in \mathcal{R}^{N_0} \\ R \supset I_0 \times J_0}} \left(\sum_{Q \in \mathcal{B}_R} \frac{|a_Q| |Q|}{|R|} \right)^2 \right)^{q/2} \right)^{p/q}. \end{aligned} \quad (4.17)$$

We continue by proving a lower bound for $\|f\|_{H^p(H^q)}^p$. Set

$$w_R = \sum_{Q \in \mathcal{B}_R} |a_Q| h_Q, \quad R \in \mathcal{R}^{N_0},$$

and observe that by (P1) we have

$$\|f\|_{H^p(H^q)}^p = \int_0^1 \left(\int_0^1 \left(\sum_{R \in \mathcal{R}^{N_0}} w_R^2(x, y) \right)^{q/2} dy \right)^{p/q} dx.$$

By (P2) the collections $\{X_{I_0} : I_0 \in \mathcal{D}_{N_0}\}$ and $\{Y_{J_0} : J_0 \in \mathcal{D}_{N_0}\}$ are each pairwise disjoint, thus we obtain

$$\|f\|_{H^p(H^q)}^p \geq \sum_{I_0 \in \mathcal{D}_{N_0}} \int_{X_{I_0}} \left(\sum_{J_0 \in \mathcal{D}_{N_0}} |Y_{J_0}| \int_{Y_{J_0}} \left(\sum_{R \in \mathcal{R}^{N_0}} w_R^2(x, y) \right)^{q/2} \frac{dy}{|Y_{J_0}|} \right)^{p/q} dx.$$

For fixed $I_0, J_0 \in \mathcal{D}_{N_0}$, $x \in X_{I_0}$, $y \in Y_{J_0}$ and $R \in \mathcal{R}^{N_0}$, we have by (4.7) and (P2) that $w_R(x, y) \neq 0$ implies $R \supset I_0 \times J_0$, so we obtain from the latter estimate together with (P3) the following lower estimate for $C_Y^{p/q} \|f\|_{H^p(H^q)}^p$:

$$\sum_{I_0 \in \mathcal{D}_{N_0}} \int_{X_{I_0}} \left(\sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \int_{Y_{J_0}} \left(\sum_{R \supset I_0 \times J_0} w_R^2(x, y) \right)^{q/2} \frac{dy}{|Y_{J_0}|} \right)^{p/q} dx. \quad (4.18)$$

With $I_0, J_0 \in \mathcal{D}_{N_0}$ fixed, we now prepare for the application of Lemma 4.5 to the inner integral of the above estimate. We use the following specification. We put $\Omega = Y_{J_0}$, $d\mu = \frac{dy}{|Y_{J_0}|}$, and $r = q$. In view of (i) of Lemma 4.5 we obtain that

$$\int_{Y_{J_0}} \left(\sum_{R \supset I_0 \times J_0} w_R^2(x, y) \right)^{q/2} \frac{dy}{|Y_{J_0}|} \geq \left(\sum_{R \supset I_0 \times J_0} \left(\int_{Y_{J_0}} |w_R(x, y)| \frac{dy}{|Y_{J_0}|} \right)^2 \right)^{q/2}. \quad (4.19)$$

By (P1) we have $|w_R(x, y)| = \sum_{K \times L \in \mathcal{B}_R} |a_{K \times L}| \mathbb{1}_K(x) \mathbb{1}_L(y)$, hence by (P4) and (P3)

$$\begin{aligned} \int_{Y_{J_0}} |w_R(x, y)| \frac{dy}{|Y_{J_0}|} &= \sum_{K \times L \in \mathcal{B}_R} |a_{K \times L}| \frac{|L \cap Y_{J_0}|}{|Y_{J_0}|} \mathbb{1}_K(x) \\ &\geq C_Y^{-2} \sum_{K \times L \in \mathcal{B}_R} |a_{K \times L}| \frac{|L|}{|J|} \mathbb{1}_K(x) \end{aligned}$$

for all $R \in \mathcal{R}^{N_0}$ with $R = I \times J \supset I_0 \times J_0$. Combining the latter estimate with (4.19) and (4.18) we obtain the following lower estimate for $C_Y^{2p+p/q} \|f\|_{H^p(H^q)}^p$:

$$\sum_{I_0 \in \mathcal{D}_{N_0}} |X_{I_0}| \int_{X_{I_0}} \left(\sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \left(\sum_{R \supset I_0 \times J_0} v_R^2(x) \right)^{q/2} \right)^{p/q} \frac{dx}{|X_{I_0}|}, \quad (4.20)$$

where we put $v_R(x) = \sum_{K \times L \in \mathcal{B}_R} \frac{|a_{K \times L}| |L|}{|J|} \mathbb{1}_K(x)$, if $R = I \times J$. With $I_0 \in \mathcal{D}_{N_0}$ fixed, we now prepare for the application of Lemma 4.5 to obtain a lower bound for the following term:

$$\int_{X_{I_0}} \left(\sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \left(\sum_{R \supset I_0 \times J_0} v_R^2(x) \right)^{q/2} \right)^{p/q} \frac{dx}{|X_{I_0}|}. \quad (4.21)$$

To this end, we use the following specification. We put $\Omega = X_{I_0}$, $d\mu = \frac{dx}{|X_{I_0}|}$, and $r = p$, $s = q$. Invoking (ii) of Lemma 4.5, we find that (4.21) is bounded from below by

$$\left(\sum_{J_0 \in \mathcal{D}_{N_0}} |J_0| \left(\sum_{R \supset I_0 \times J_0} \left(\int_{X_{I_0}} v_R(x) \frac{dx}{|X_{I_0}|} \right)^2 \right)^{q/2} \right)^{p/q}. \quad (4.22)$$

Recall that we defined $v_R(x) = \sum_{K \times L \in \mathcal{B}_R} \frac{|a_{K \times L}| |L|}{|J|} \mathbb{1}_K(x)$, if $R = I \times J$. By (P4) and (P3) we estimate

$$\begin{aligned} \int_{X_{I_0}} v_R(x) \frac{dx}{|X_{I_0}|} &= \sum_{K \times L \in \mathcal{B}_R} \frac{|a_{K \times L}| |L|}{|J|} \frac{|K \cap X_{I_0}|}{|X_{I_0}|} \\ &\geq C_X^{-2} \sum_{Q \in \mathcal{B}_R} \frac{|a_Q| |Q|}{|R|} \end{aligned}$$

for all $R = I \times J \in \mathcal{R}^{\mathbb{N}_0}$ with $R \supset I_0 \times J_0$. Combining the latter estimate with (4.22), (4.21), and (4.20), we obtain the following lower estimate for $C_X^{2p} C_Y^{2p+p/q} \|f\|_{H^p(H^q)}^p$:

$$\sum_{I_0 \in \mathcal{D}_{\mathbb{N}_0}} |X_{I_0}| \left(\sum_{J_0 \in \mathcal{D}_{\mathbb{N}_0}} |J_0| \left(\sum_{R \supset I_0 \times J_0} \left(\sum_{Q \in \mathcal{B}_R} \frac{|a_Q||Q|}{|R|} \right)^2 \right)^{q/2} \right)^{p/q}.$$

Finally, by (P3) the latter estimate yields

$$\begin{aligned} C_X^{2p+1} C_Y^{2p+p/q} \|f\|_{H^p(H^q)}^p &\geq \\ \sum_{I_0 \in \mathcal{D}_{\mathbb{N}_0}} |I_0| \left(\sum_{J_0 \in \mathcal{D}_{\mathbb{N}_0}} |J_0| \left(\sum_{R \supset I_0 \times J_0} \left(\sum_{Q \in \mathcal{B}_R} \frac{|a_Q||Q|}{|R|} \right)^2 \right)^{q/2} \right)^{p/q}. \end{aligned} \quad (4.23)$$

Direct comparison with (4.17) gives

$$\|A_\beta f\|_{H^p(H^q)} \leq M C_X^{3+1/p} C_Y^{3+1/q} \|f\|_{H^p(H^q)}.$$

PART 3: CONCLUSION OF THE PROOF. If additionally, we assume that $m := \inf_Q |\beta_Q| > 0$, Part 2 implies that A_γ is bounded by $m^{-1} C_X^{3+1/p} C_Y^{3+1/q}$. The commutativity of the diagram (4.10) follows from the fact that $\beta_Q \gamma_Q = 1$. \square

4.4. A linear order on \mathcal{R} and Capon's local product condition.

In Section 5, we will iteratively construct collections of dyadic rectangles $\mathcal{B}_R \subset \mathcal{R}$, $R \in \mathcal{R}$ satisfying Capon's local product condition. This will be accomplished by organizing the dyadic rectangles according to the linear order \triangleleft defined in the present section, below. The other purpose of this section is to introduce the auxiliary condition (R1)–(R6) and to show that it implies Capon's local product condition (P1)–(P4).

First, we define the bijective function $\mathcal{O}_{\mathbb{N}_0^2} : \mathbb{N}_0^2 \rightarrow \mathbb{N}_0$ by

$$\mathcal{O}_{\mathbb{N}_0^2}(m, n) = \begin{cases} n^2 + m, & \text{if } m < n, \\ m^2 + m + n, & \text{if } m \geq n. \end{cases}$$

To see that $\mathcal{O}_{\mathbb{N}_0^2}$ is bijective consider that for each $k \in \mathbb{N}$:

- ▷ $\mathcal{O}_{\mathbb{N}_0^2}(0, 0) = 0$,
- ▷ $m \mapsto \mathcal{O}_{\mathbb{N}_0^2}(m, k)$ maps $\{0, \dots, k-1\}$ bijectively onto $\{k^2, \dots, k^2 + k - 1\}$ and preserves the natural order on \mathbb{N}_0 ,
- ▷ $\mathcal{O}_{\mathbb{N}_0^2}(k, 0) = \mathcal{O}_{\mathbb{N}_0^2}(k-1, k) + 1$,
- ▷ $n \mapsto \mathcal{O}_{\mathbb{N}_0^2}(k, n)$ maps $\{0, \dots, k\}$ bijectively onto $\{k^2 + k, \dots, k^2 + 2k\}$ and preserves the natural order on \mathbb{N}_0 ,
- ▷ $\mathcal{O}_{\mathbb{N}_0^2}(0, k+1) = \mathcal{O}_{\mathbb{N}_0^2}(k, k) + 1$.

See Figure 5 for a depiction of $\mathcal{O}_{\mathbb{N}_0^2}$.

Now, let $<_\ell$ denote the lexicographic order on \mathbb{R}^3 . For two dyadic rectangles $I_k \times J_k \in \mathcal{R}$ with $|I_k| = 2^{-m_k}$, $|J_k| = 2^{-n_k}$, $k = 0, 1$, we define $I_0 \times J_0 \triangleleft I_1 \times J_1$ if and only if

$$(\mathcal{O}_{\mathbb{N}_0^2}(m_0, n_0), \inf I_0, \inf J_0) <_\ell (\mathcal{O}_{\mathbb{N}_0^2}(m_1, n_1), \inf I_1, \inf J_1).$$

Associated to the linear ordering \triangleleft is the bijective index function $\mathcal{O}_\triangleleft : \mathcal{R} \rightarrow \mathbb{N}_0$ defined by

$$\mathcal{O}_\triangleleft(R_0) < \mathcal{O}_\triangleleft(R_1) \Leftrightarrow R_0 \triangleleft R_1, \quad R_0, R_1 \in \mathcal{R}.$$

The geometry of a dyadic rectangle is linked to its index by the estimate

$$(2^k - 1)^2 \leq \mathcal{O}_\triangleleft(I \times J) < (2^{k+1} - 1)^2, \quad \text{whenever } \min(|I|, |J|) = 2^{-k}, \quad (4.24)$$




FIGURE 5. This figure depicts the order of the first 16 pairs in \mathbb{N}_0^2 with respect to the map $\mathcal{O}_{\mathbb{N}_0^2}$.

0		4		24									
		3		23									
		6		8		28	32						
		5		7		27	31						
1	2	14		16		18	20	36	40	44	48		
		13		15		17	19	35	39	43	47		
9		10		11		12		34		38		42	46
13		15		17		19		33		37		41	45

FIGURE 6. The first 49 rectangles and their indices $\mathcal{O}_{\triangleleft}$.

and hence,

$$\frac{1}{(1 + \sqrt{i})^2} \leq |I| |J|, \quad i = \mathcal{O}_{\triangleleft}(I \times J). \quad (4.25)$$

The index of a dyadic rectangle and its predecessors are related by

$$\tilde{I} \times J \triangleleft I \times J, \quad \text{for } I \neq [0, 1) \quad \text{and} \quad I \times \tilde{J} \triangleleft I \times J, \quad \text{for } J \neq [0, 1), \quad (4.26)$$

where we recall that for $I \neq [0, 1)$, \tilde{I} is the unique dyadic interval satisfying $\tilde{I} \supset I$ and $|\tilde{I}| = 2|I|$. See Figure 6 for a picture of $\mathcal{O}_{\triangleleft}$.

For a dyadic interval I , we write I^ℓ and I^r for the dyadic intervals which are the left and right halves of I , respectively. In the following definition, we use the notation introduced in (4.5), so that for a collection \mathcal{X}_R (respectively, \mathcal{Y}_R) of dyadic intervals, X_R (respectively, Y_R) denotes its union.

Definition 4.7. Let $\mathcal{A} = \mathcal{R}$ or $\mathcal{A} = \{R \in \mathcal{R} : R \triangleleft R_0\}$ for some $R_0 \in \mathcal{R}$. We say that $\{\mathcal{B}_R : R \in \mathcal{A}\}$ satisfies the *auxiliary condition (R1)–(R6)* if the following properties hold true.

- (R1) For each $R \in \mathcal{A}$, there are non-negative integers $\mu(R)$, $\nu(R)$ and non-empty sets $\mathcal{X}_R \subset \mathcal{D}_{\mu(R)}$ and $\mathcal{Y}_R \subset \mathcal{D}_{\nu(R)}$ such that $\mathcal{B}_R = \{K \times L : K \in \mathcal{X}_R, L \in \mathcal{Y}_R\}$.
- (R2) $\mu([0, 1] \times [0, 1]) = \nu([0, 1] \times [0, 1]) = 0$ and $\mathcal{X}_{[0,1] \times [0,1]} = \mathcal{Y}_{[0,1] \times [0,1]} = \{[0, 1]\}$.
- (R3) For each $I \in \mathcal{D} \setminus \{[0, 1]\}$ with $R = I \times [0, 1] \in \mathcal{A}$

$$X_{I \times [0,1]} = \begin{cases} \bigcup \{K^\ell : K \in \mathcal{D}_{\kappa(R)}, K \subset X_{\tilde{I} \times [0,1]}\} & \text{if } I = \tilde{I}^\ell, \\ \bigcup \{K^r : K \in \mathcal{D}_{\kappa(R)}, K \subset X_{\tilde{I} \times [0,1]}\} & \text{if } I = \tilde{I}^r, \end{cases}$$

where $\kappa(R) = \max\{\mu(S) : S \triangleleft [0, |I|] \times [0, 1]\}$;

- (R4) If $R = I \times J \in \mathcal{A}$ with $|I| < |J|$, then

$$\mu(R) > \max\{\mu(S) : S \triangleleft R\},$$

$X_R = X_{I \times [0,1]}$, and $\mathcal{Y}_R = \mathcal{Y}_{I' \times J}$, where $I' \in \mathcal{D}$ is the unique dyadic interval such that $I' \supset I$ and $|I'| = |J|$.

- (R5) For $J \in \mathcal{D} \setminus \{[0, 1]\}$ with $R = [0, 1] \times J \in \mathcal{A}$

$$Y_{[0,1] \times J} = \begin{cases} \bigcup \{L^\ell : L \in \mathcal{D}_{\lambda(R)}, L \subset Y_{[0,1] \times \tilde{J}}\} & \text{if } J = \tilde{J}^\ell, \\ \bigcup \{L^r : L \in \mathcal{D}_{\lambda(R)}, L \subset Y_{[0,1] \times \tilde{J}}\} & \text{if } J = \tilde{J}^r, \end{cases}$$

where $\lambda(R) = \max\{\nu(S) : S \triangleleft [0, 1] \times [0, |J|]\}$.

- (R6) If $R = I \times J \in \mathcal{A} \setminus \{[0, 1] \times [0, 1]\}$ with $|I| \geq |J|$, then

$$\nu(R) > \max\{\nu(S) : S \triangleleft R\},$$

$Y_R = Y_{[0,1] \times J}$, and $\mathcal{X}_R = \mathcal{X}_{I \times J'}$, where $J' \in \mathcal{D}$ is the unique dyadic interval such that $J' \supset J$ and $|J'| = 2|I|$ if $I \neq [0, 1]$, and $J' = [0, 1]$ if $I = [0, 1]$.

Remark 4.8. Let $\{\mathcal{B}_R : R \in \mathcal{R}\}$ be a collection such that each of the finite sub-collections $\{\mathcal{B}_R : R \triangleleft R_0\}$, $R_0 \in \mathcal{R}$, satisfies the auxiliary condition (R1)–(R6). Then it is easy to see that $\{\mathcal{B}_R : R \in \mathcal{R}\}$ itself satisfies the auxiliary condition (R1)–(R6).

Lemma 4.9. *Let $\{\mathcal{B}_R : R \in \mathcal{R}\}$ satisfy the auxiliary condition (R1)–(R6). Then $\{\mathcal{B}_R : R \in \mathcal{R}\}$ satisfies the local product condition (P1)–(P4) with constants $C_X = C_Y = 1$.*

Proof. The usual linear order \prec on dyadic intervals is given by $I_1 \prec I_0$ if and only if either $|I_1| > |I_0|$ or $|I_1| = |I_0|$ and $\min I_1 < \min I_0$. The proof uses induction with respect to the linear orders \prec and \triangleleft .

VERIFICATION OF (P1). For each $R \in \mathcal{R}$, \mathcal{X}_R consists of pairwise disjoint intervals because \mathcal{X}_R is contained in $\mathcal{D}_{\mu(R)}$. Similarly, $\mathcal{Y}_R \subset \mathcal{D}_{\nu(R)}$ and consists of pairwise disjoint intervals, and therefore the rectangles in \mathcal{B}_R are pairwise disjoint.

Now suppose that $R_0, R_1 \in \mathcal{R}$ are distinct. By relabelling them if necessary, we may suppose that $R_1 \triangleleft R_0$, where $R_0 = I_0 \times J_0 \neq [0, 1] \times [0, 1]$. To establish the disjointness of \mathcal{B}_{R_0} and \mathcal{B}_{R_1} , we must show that either \mathcal{X}_{R_0} and \mathcal{X}_{R_1} are disjoint or \mathcal{Y}_{R_0} and \mathcal{Y}_{R_1} are disjoint. If $|I_0| < |J_0|$, then (R4) implies that $\mu(R_0) > \mu(R_1)$, so that $\mathcal{X}_{R_0} \cap \mathcal{X}_{R_1} \subset \mathcal{D}_{\mu(R_0)} \cap \mathcal{D}_{\mu(R_1)} = \emptyset$. Otherwise $|I_0| \geq |J_0|$, in which case a similar argument based on (R6) shows that $\mathcal{Y}_{R_0} \cap \mathcal{Y}_{R_1} = \emptyset$.

VERIFICATION OF (P2). We begin by observing that (R4) and (R6) imply that the sets X_R, Y_R, X_I , and Y_J defined in (4.5)–(4.6) are given by

$$X_R = X_{I \times [0,1]} = X_I \quad \text{and} \quad Y_R = Y_{[0,1] \times J} = Y_J, \quad R = I \times J \in \mathcal{R}. \quad (4.27)$$

Since the order \prec is linear, and the set \mathcal{D} is countable and has a minimum element $[0, 1)$ with respect to \prec , we may use induction on $I_0 \in \mathcal{D}$ to prove the following two statements:

- (a) $X_{I_0 \times [0, 1)} \cap X_{I_1 \times [0, 1)} = \emptyset$ and $Y_{[0, 1) \times I_0} \cap Y_{[0, 1) \times I_1} = \emptyset$ for each $I_1 \in \mathcal{D}$ with $I_1 \prec I_0$ and $I_0 \cap I_1 = \emptyset$;
- (b) $X_{I_0 \times [0, 1)} \subset X_{I_1 \times [0, 1)}$ and $Y_{[0, 1) \times I_0} \subset Y_{[0, 1) \times I_1}$ for each $I_1 \in \mathcal{D}$ with $I_0 \subset I_1$.

The statements (a) and (b) above together with (4.27) imply (P2). The start of the induction is easy. Indeed, suppose that $I_0 = [0, 1)$. Then no I_1 satisfies $I_1 \prec [0, 1)$, so that (a) is vacuous, while (b) holds trivially because $I_1 = [0, 1)$ is the only dyadic interval which contains $[0, 1)$.

Now let $I_0 \in \mathcal{D} \setminus \{[0, 1)\}$, and assume inductively that (a)–(b) have been established for each $I'_0 \prec I_0$ (that is, (a)–(b) hold whenever I_0 is replaced with I'_0). We shall prove the statements concerning $X_{I_0 \times [0, 1)}$; the proofs for $Y_{[0, 1) \times I_0}$ are similar, requiring only minor adjustments of the notation.

To verify (a), suppose that $I_1 \in \mathcal{D}$ satisfies $I_1 \prec I_0$ and $I_0 \cap I_1 = \emptyset$. Then either $I_1 \cap \tilde{I}_0 = \emptyset$, or $I_1 = (\tilde{I}_0)^\ell$ and $I_0 = (\tilde{I}_0)^r$. (Note that because $I_1 \prec I_0$, we cannot have $I_1 = (\tilde{I}_0)^r$ and $I_0 = (\tilde{I}_0)^\ell$.) In the first case, since $I_1 \prec I_0$ and $\tilde{I}_0 \prec I_0$, the induction hypothesis implies that $X_{\tilde{I}_0 \times [0, 1)} \cap X_{I_1 \times [0, 1)} = \emptyset$, from which the result follows because $X_{I_0 \times [0, 1)} \subset X_{\tilde{I}_0 \times [0, 1)}$ by (R3).

In the second case, we observe that $\tilde{I}_0 = \tilde{I}_1$ and $|I_0| = |I_1|$, so that $\kappa(I_0 \times [0, 1)) = \kappa(I_1 \times [0, 1))$. This implies that $X_{I_0 \times [0, 1)}$ and $X_{I_1 \times [0, 1)}$ are disjoint because $X_{I_0 \times [0, 1)}$ is the disjoint union of the *right* halves of the intervals $K \in \mathcal{D}_{\kappa(I_0 \times [0, 1))}$ with $K \subset X_{\tilde{I}_0 \times [0, 1)}$, while $X_{I_1 \times [0, 1)}$ is the disjoint union of the *left* halves of the same intervals.

Next, to prove (b), suppose that $I_1 \in \mathcal{D}$ with $I_0 \subset I_1$. The inclusion is obvious if $I_0 = I_1$, so we may suppose that $I_0 \subsetneq I_1$. Then we have $\tilde{I}_0 \subset I_1$, so the induction hypothesis implies that $X_{\tilde{I}_0 \times [0, 1)} \subset X_{I_1 \times [0, 1)}$. Hence the statement follows from the fact that $X_{I_0 \times [0, 1)} \subset X_{\tilde{I}_0 \times [0, 1)}$.

VERIFICATION OF (P3). The proofs of (P3) and (P4) both rely on the following two identities:

$$|K \cap X_{I \times [0, 1)}| = \frac{|K \cap X_{\tilde{I} \times [0, 1)}|}{2} \quad \text{and} \quad |L \cap Y_{[0, 1) \times J}| = \frac{|L \cap Y_{[0, 1) \times \tilde{J}}|}{2}, \quad (4.28)$$

valid for $I, J \in \mathcal{D} \setminus \{[0, 1)\}$, $K \in \mathcal{D}^{\kappa(I \times [0, 1))}$, and $L \in \mathcal{D}^{\lambda([0, 1) \times J)}$.

We shall establish the first of these identities; again, the proof of the other requires only notational changes. For $I \in \mathcal{D} \setminus \{[0, 1)\}$ and $K \in \mathcal{D}^{\kappa(I \times [0, 1))}$, set $\mathcal{V}_I(K) = \{K_0 \in \mathcal{D}_{\kappa(I \times [0, 1))} : K_0 \subset K \cap X_{\tilde{I} \times [0, 1)}\}$. We claim that

$$K \cap X_{\tilde{I} \times [0, 1)} = \bigcup \mathcal{V}_I(K) \quad \text{and} \quad K \cap X_{I \times [0, 1)} = \begin{cases} \bigcup \{K_0^\ell : K_0 \in \mathcal{V}_I(K)\} & \text{if } I = \tilde{I}^\ell \\ \bigcup \{K_0^r : K_0 \in \mathcal{V}_I(K)\} & \text{if } I = \tilde{I}^r. \end{cases} \quad (4.29)$$

Indeed, the inclusion $\bigcup \mathcal{V}_I(K) \subset K \cap X_{\tilde{I} \times [0, 1)}$ is clear from the definition of $\mathcal{V}_I(K)$. Conversely, for each $x \in K \cap X_{\tilde{I} \times [0, 1)}$, there is a (necessarily unique) interval $K_0 \in \mathcal{X}_{\tilde{I} \times [0, 1)}$ such that $x \in K_0$. We have $\mu(\tilde{I} \times [0, 1)) \leq \kappa(I \times [0, 1))$ because $\tilde{I} \times [0, 1) \triangleleft [0, |I|) \times [0, 1)$, so we can find $K_1 \in \mathcal{D}_{\kappa(I \times [0, 1))}$ such that $x \in K_1 \subset K_0$. The sets K_1 and K are not disjoint as they both contain x ; combined with the fact that $|K_1| \leq |K|$, this shows that $K_1 \subset K$. Moreover, we have $K_1 \subset K_0 \subset X_{\tilde{I} \times [0, 1)}$, so that $K_1 \in \mathcal{V}_I(K)$, and hence $x \in K_1 \subset \bigcup \mathcal{V}_I(K)$.

Moving on to the second part of (4.29), we obtain the inclusion \supset directly from the definition of $\mathcal{V}_I(K)$ and (R3). Conversely, suppose that $x \in K \cap X_{I \times [0, 1)}$, so

that $x \in K$ and either $x \in K_0^\ell$ or $x \in K_0^r$ (depending on whether $I = (\tilde{I})^\ell$ or $I = (\tilde{I})^r$) for some $K_0 \in \mathcal{D}_{\kappa(I \times [0,1])}$ with $K_0 \subset X_{\tilde{I} \times [0,1]}$. In both cases, we see that $K \cap K_0 \neq \emptyset$ and $|K_0| \leq |K|$, so that $K_0 \subset K$, and hence $K_0 \in \mathcal{V}_I(K)$, from which the inclusion follows.

The first equation in (4.28) is immediate from (4.29) because $\mathcal{V}_I(K)$ consists of disjoint sets and $|K_0^\ell| = |K_0^r| = |K_0|/2$.

We can now easily establish (P3) with $C_X = C_Y = 1$. By (4.27), we must show that

$$|X_{I \times [0,1]}| = |I| \quad \text{and} \quad |Y_{[0,1] \times I}| = |I|, \quad I \in \mathcal{D}. \quad (4.30)$$

We do so by induction on I . The start of the induction, where $I = [0, 1)$, follows immediately from the fact that $X_{[0,1] \times [0,1]} = Y_{[0,1] \times [0,1]} = [0, 1)$ by (R2).

Now let $I \in \mathcal{D} \setminus \{[0, 1)\}$, and assume inductively that the result is true for each $I' \prec I$. Using (4.28) with $K = L = [0, 1)$, we obtain that $|X_{I \times [0,1]}| = |X_{\tilde{I} \times [0,1]}|/2 = |\tilde{I}|/2 = |I|$ because $\tilde{I} \prec I$ and likewise $|Y_{[0,1] \times I}| = |I|$.

VERIFICATION OF (P4). We shall prove that, for each $R_0 = I_0 \times J_0$ and $R = I \times J$ in \mathcal{R} with $R_0 \subset R$,

$$\frac{|K \cap X_{I_0 \times [0,1]}|}{|I_0|} = \frac{|K|}{|I|} \quad \text{and} \quad \frac{|L \cap Y_{[0,1] \times J_0}|}{|J_0|} = \frac{|L|}{|J|}, \quad K \in \mathcal{X}_R, L \in \mathcal{Y}_R. \quad (4.31)$$

By (4.27) and (4.30), this will verify (P4) with $C_X = C_Y = 1$.

The proof of (4.31) is by induction on R_0 . The start of the induction is trivial because the only $R \in \mathcal{R}$ that contains $R_0 = [0, 1) \times [0, 1)$ is R_0 itself.

Now let $R_0 \in \mathcal{R} \setminus \{[0, 1) \times [0, 1)\}$, and assume inductively that (4.31) has been verified for each $R'_0 \triangleleft R_0$. This time, we shall focus on the proof of the second identity in (4.31); the proof of the first identity is similar, but formally slightly easier due to the lack of symmetry between conditions (R4) and (R6): when $|I| = |J|$, we re-use an existing set as \mathcal{X}_R and define a new set \mathcal{Y}_R .

Suppose that $R = I \times J \in \mathcal{R}$ with $R_0 \subset R$, and let $L \in \mathcal{Y}_R$. If $J_0 = J$, then $L \subset Y_{[0,1] \times J_0}$, and the identity is immediate. Hence we may suppose that $J_0 \subsetneq J$. Moreover, we may suppose that $|I| \geq |J|$. Indeed, if not, then by (R4) $\mathcal{Y}_R = \mathcal{Y}_{I' \times J}$, where $I' \in \mathcal{D}$ satisfies $I' \supset I$ and $|I'| = |J|$, so that we may replace I with I' to obtain that $|I| \geq |J|$.

Then we have $|J_0| < |J| = \min\{|I|, |J|\}$, so that $R \triangleleft [0, 1) \times [0, |J_0|)$, and hence $\lambda([0, 1) \times J_0) \geq \nu(R)$; thus $L \in \mathcal{Y}_R \subset \mathcal{D}_{\nu(R)} \subset \mathcal{D}^{\lambda([0,1] \times J_0)}$, so that (4.28) shows that $|L \cap Y_{[0,1] \times J_0}| = |L \cap Y_{[0,1] \times \tilde{J}_0}|/2$. Now $R'_0 = I_0 \times \tilde{J}_0$ satisfies $R'_0 \triangleleft R_0$ and $R'_0 \subset R$, and therefore the induction hypothesis implies that $|L \cap Y_{[0,1] \times \tilde{J}_0}|/|\tilde{J}_0| = |L|/|J|$. Hence the conclusion follows because $|\tilde{J}_0| = 2|J_0|$. \square

Having obtained Theorem 4.3 and Lemma 4.9, we are finally prepared to prove Theorem 3.1.

5. PROOF OF THEOREM 3.1

Here, we prove that the identity operator on $H^p(H^q)$ factors through any operator $T : H^p(H^q) \rightarrow H^p(H^q)$ having large diagonal with respect to the bi-parameter Haar system (see Theorem 3.1). The basic pattern of our argument below is the following: we carefully construct $\{\mathcal{B}_R : R \in \mathcal{R}\}$ satisfying the auxiliary condition (R1)–(R6) (see Section 4). Moreover, these collections are chosen in such a way that we are able to find signs $\varepsilon_Q \in \{\pm 1\}$, $Q \in \bigcup_{R \in \mathcal{R}} \mathcal{B}_R$, for which the block basis $b_R^{(\varepsilon)} = \sum_{Q \in \mathcal{B}_R} \varepsilon_Q h_Q$, $R \in \mathcal{R}$ has the following properties: $|\langle T b_{R_1}^{(\varepsilon)}, b_{R_2}^{(\varepsilon)} \rangle|$ is small in

the precise sense of (5.6a) below whenever $R_1, R_2 \in \mathcal{R}$ are distinct, and

$$|\langle Tb_R^{(\varepsilon)}, b_R^{(\varepsilon)} \rangle| \geq \delta \|b_R^{(\varepsilon)}\|_2^2, \quad R \in \mathcal{R}.$$

Thereafter we apply the two main results of the preceding section, Theorem 4.3 and Lemma 4.9, and finally we construct a factorization of the identity operator through T .

Proof of Theorem 3.1. Let $1 \leq p, q < \infty$ and $\delta > 0$, and let $T : H^p(H^q) \rightarrow H^p(H^q)$ be an operator such that

$$|\langle Th_R, h_R \rangle| \geq \delta |R|, \quad R \in \mathcal{R}. \quad (5.1)$$

We define $\gamma = (\gamma_R : R \in \mathcal{R})$ by

$$\gamma_R = \frac{\overline{\langle Th_R, h_R \rangle}}{|\langle Th_R, h_R \rangle|}, \quad R \in \mathcal{R}.$$

Recall that in (4.2) we defined the Haar multiplier M_γ which satisfies $\|M_\gamma\| = 1$, and $\langle (TM_\gamma)h_R, h_R \rangle \geq \delta |R|$. Thereby, replacing T with TM_γ , it suffices to consider the special case where

$$\langle Th_R, h_R \rangle \geq \delta |R|, \quad R \in \mathcal{R}. \quad (5.2)$$

OVERVIEW. Let $0 < \eta \leq 1$. The main part of the proof consists of choosing collections of dyadic rectangles \mathcal{B}_R , $R \in \mathcal{R}$ and suitable signs $\varepsilon = (\varepsilon_Q)$ such that $b_R^{(\varepsilon)} = \sum_{Q \in \mathcal{B}_R} \varepsilon_Q h_Q$ satisfies the following:

- ▷ The closed linear span of $\{b_R^{(\varepsilon)} : R \in \mathcal{R}\}$ is complemented and isomorphic to $H^p(H^q)$.
- ▷ There is an operator $U : H^p(H^q) \rightarrow H^p(H^q)$ given by

$$U(f) = \sum_{R \in \mathcal{R}} \frac{\langle f, b_R^{(\varepsilon)} \rangle}{\langle Tb_R^{(\varepsilon)}, b_R^{(\varepsilon)} \rangle} b_R^{(\varepsilon)}.$$

- ▷ For every finite linear combination $g = \sum_{R \in \mathcal{R}} \lambda_R b_R^{(\varepsilon)}$ we have

$$\|UTg - g\|_{H^p(H^q)} \leq \frac{\eta}{2} \|g\|_{H^p(H^q)}.$$

PREPARATION. Given $R = I \times J \in \mathcal{R}$ we write

$$Th_R = \alpha_R h_R + r_R, \quad (5.3a)$$

where

$$\alpha_R = \frac{\langle Th_R, h_R \rangle}{|R|} \quad \text{and} \quad r_R = \sum_{S \neq R} \frac{\langle Th_R, h_S \rangle}{|S|} h_S. \quad (5.3b)$$

We note the estimates

$$\delta \leq \alpha_R \leq \|T\| \quad \text{and} \quad \|r_R\|_{H^p(H^q)} \leq 2\|T\| |I|^{1/p} |J|^{1/q}. \quad (5.4)$$

INDUCTIVE CONSTRUCTION OF $b_R^{(\varepsilon)}$. We will now inductively define the block basis $\{b_R^{(\varepsilon)} : R \in \mathcal{R}\}$. For fixed $R \in \mathcal{R}$, the block basis element $b_R^{(\varepsilon)}$ is determined by a collection of dyadic rectangles $\mathcal{B}_R \subset \mathcal{R}$ and a suitable choice of signs $\varepsilon = (\varepsilon_Q)$ and is of the following form:

$$b_R^{(\varepsilon)} = \sum_{Q \in \mathcal{B}_R} \varepsilon_Q h_Q. \quad (5.5)$$

From now on, we systematically use the following rule: whenever $\mathcal{O}_\triangleleft(R) = i$ we set

$$\mathcal{B}_i = \mathcal{B}_R, \quad b_i^{(\varepsilon)} = b_R^{(\varepsilon)}, \quad h_i = h_R.$$

We will construct collections $\{\mathcal{B}_i : i \in \mathbb{N}_0\}$ satisfying the auxiliary condition (R1)–(R6) and choose signs $\varepsilon = (\varepsilon_Q)$ such that

$$\sum_{j=0}^{i-1} |\langle T b_j^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle| + |\langle b_i^{(\varepsilon)}, T^* b_j^{(\varepsilon)} \rangle| \leq \eta \delta 4^{-i-2}, \quad i \in \mathbb{N}, \quad (5.6a)$$

$$|\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle| \geq \delta \|b_i^{(\varepsilon)}\|_2^2, \quad i \in \mathbb{N}_0. \quad (5.6b)$$

The induction begins by putting

$$\mathcal{B}_0 = \{[0, 1) \times [0, 1)\} \quad \text{and} \quad b_0^{(\varepsilon)} = h_{[0,1) \times [0,1)}. \quad (5.7)$$

Consequently, $\mathcal{X}_{[0,1) \times [0,1)} = \mathcal{Y}_{[0,1) \times [0,1)} = \{[0, 1)\}$ and $\mu([0, 1) \times [0, 1)) = 0$, $\nu([0, 1) \times [0, 1)) = 0$. Obviously, $\{\mathcal{B}_0\}$ satisfies (R1)–(R6).

Let $i_0 \in \mathbb{N}$. At this stage we assume that

- ▷ $\{\mathcal{B}_j : 0 \leq j \leq i_0 - 1\}$ satisfies the auxiliary condition (R1)–(R6).
- ▷ the block basis $\{b_j^{(\varepsilon)} : 0 \leq j \leq i_0 - 1\}$ given by (5.5) satisfies (5.6) (for $0 \leq i \leq i_0 - 1$).

Now, we turn to the construction of \mathcal{B}_{i_0} and ε_Q , where $Q \in \mathcal{B}_{i_0}$. In the first step we will find \mathcal{B}_{i_0} in (5.20), and only then we will choose the signs ε_Q , $Q \in \mathcal{B}_{i_0}$ in (5.23). The collection \mathcal{B}_{i_0} and the signs ε_Q , $Q \in \mathcal{B}_{i_0}$ then determine $b_{i_0}^{(\varepsilon)}$.

CONSTRUCTION OF \mathcal{B}_{i_0} . Let $I_0 \times J_0 \in \mathcal{R}$ be such that $\mathcal{O}_{\triangleleft}(I_0 \times J_0) = i_0$. We distinguish between the four cases

$$|I_0| < |J_0|, J_0 = [0, 1), \quad |I_0| < |J_0|, J_0 \neq [0, 1),$$

and

$$|I_0| \geq |J_0|, I_0 = [0, 1), \quad |I_0| \geq |J_0|, I_0 \neq [0, 1).$$


CASE 1: $|I_0| < |J_0|$. Here, we will construct the collection $\mathcal{B}_{I_0 \times J_0}$, for which the index rectangle $I_0 \times J_0$ is “below the diagonal”.

First, we define

$$\nu(I_0 \times J_0) = \nu(I'_0 \times J_0) \quad \text{and} \quad \mathcal{Y}_{I_0 \times J_0} = \mathcal{Y}_{I'_0 \times J_0}, \quad (5.8)$$

where $I'_0 \in \mathcal{D}$ is the unique interval such that $I'_0 \supset I_0$ and $|I'_0| = |J_0|$. We remark that $\mu(I_0 \times J_0)$ will be defined at the end of the proof in (5.21a).

CASE 1.A: $J_0 = [0, 1)$. Here, we know that $I_0 \neq [0, 1)$. Recall that \tilde{I}_0 denotes the dyadic predecessor of I_0 , and note that $\mathcal{B}_{\tilde{I}_0 \times [0,1)}$ has already been defined. The collections indexed by the black rectangles have already been constructed. Here, we determine the collections for the gray rectangles. The white ones will be treated later.



Note that $[0, |I_0|) \times [0, 1) \trianglelefteq I_0 \times [0, 1)$, and define the integer $\kappa(I_0 \times [0, 1))$ by

$$\kappa(I_0 \times [0, 1)) = \max\{\mu(Q) : Q \triangleleft [0, |I_0|) \times [0, 1)\}.$$

Recall that for a dyadic interval K_0 we denote its left half by K_0^ℓ and its right half by K_0^r . Following the basic construction of Gamlen-Gaudet [5], we proceed as follows. The set $X_{\tilde{I}_0 \times [0,1)}$ has already been defined in a previous step of the construction. Now we put

$$X_{I_0 \times [0,1)} = \begin{cases} \bigcup \{K_0^\ell : K_0 \in \mathcal{D}_{\kappa(I_0 \times [0,1))}, K_0 \subset X_{\tilde{I}_0 \times [0,1)}\} & \text{if } I_0 = \tilde{I}_0^\ell, \\ \bigcup \{K_0^r : K_0 \in \mathcal{D}_{\kappa(I_0 \times [0,1))}, K_0 \subset X_{\tilde{I}_0 \times [0,1)}\} & \text{if } I_0 = \tilde{I}_0^r. \end{cases}$$

To finish the construction in Case 1.A, we define the family of high frequency covers of the set $X_{I_0 \times [0,1]} \times [0,1]$ by putting

$$\mathcal{F}_m = \{K \times [0,1] \in \mathcal{R} : K \in \mathcal{D}_m, K \subset X_{I_0 \times [0,1]}\}, \quad (5.9)$$

for all $m > \kappa(I_0 \times [0,1])$, see Figure 7, and observe that

$$\bigcup \mathcal{F}_m = X_{I_0 \times [0,1]} \times [0,1]. \quad (5.10)$$





FIGURE 7. The above figure depicts an instance of \mathcal{F}_m in Case 1.A. $K_0^{(k)}$ is a dyadic interval such that $K_0^{(k)} \times [0,1] \in \mathcal{B}_{\tilde{I}_0 \times [0,1]}$, and K is a dyadic interval such that $K \times [0,1] \in \mathcal{F}_m$.

CASE 1.B: $J_0 \neq [0,1]$. The collections indexed by the black rectangles have already been constructed. Here, we determine the collections for the gray rectangles. The white ones will be treated later.



By our induction hypothesis, $\{\mathcal{B}_{I \times J} : \mathcal{O}_{\triangleleft}(I \times J) \leq i_0 - 1\}$ satisfies (R1)–(R6). Note that the set $X_{I_0 \times [0,1]}$ is already defined. To conclude the construction in Case 1.B, we define the high frequency covers of $X_{I_0 \times [0,1]} \times Y_{I_0 \times J_0}$ by

$$\mathcal{F}_m = \{K \times L_0 \in \mathcal{R} : K \in \mathcal{D}_m, K \subset X_{I_0 \times [0,1]}, L_0 \in \mathcal{Y}_{I_0 \times J_0}\}, \quad (5.11)$$

for $m > \mu(I_0 \times \tilde{J}_0)$.


CASE 2: $|I_0| \geq |J_0|$. In this case, we will construct the collection $\mathcal{B}_{I_0 \times J_0}$, for which the index rectangle $I_0 \times J_0$ is “on or above the diagonal”.

First, we set

$$\mu(I_0 \times J_0) = \mu(I_0 \times J'_0) \quad \text{and} \quad \mathcal{X}_{I_0 \times J_0} = \mathcal{X}_{I_0 \times J'_0}, \quad (5.12)$$

where $J'_0 \in \mathcal{D}$ is the unique dyadic interval such that $J'_0 \supset J_0$ and $|J'_0| = 2|I_0|$ if $I_0 \neq [0,1]$, and $J'_0 = [0,1]$ if $I_0 = [0,1]$. We remark that $\nu(I_0 \times J_0)$ will be defined at the end of the proof in (5.21b).

CASE 2.A: $I_0 = [0,1]$. Note that $J_0 \neq [0,1]$ and $\mathcal{B}_{[0,1] \times \tilde{J}_0}$ has already been constructed. The collections indexed by the black rectangles have already been constructed. Here, we determine the collections for the gray rectangles. The white ones will be treated later.



Note that $[0, 1) \times [0, |J_0|) \trianglelefteq [0, 1) \times J_0$. Define $\lambda([0, 1) \times J_0)$ to be

$$\lambda([0, 1) \times J_0) = \max\{\nu(Q) : Q \triangleleft [0, 1) \times [0, |J_0|)\}. \quad (5.13)$$

Recall that for a dyadic interval L_0 we denote its left (=lower) half by L_0^ℓ and its right (=upper) half by L_0^r . The set $Y_{[0,1) \times \tilde{J}_0}$ has already been defined. Now, put

$$Y_{[0,1) \times J_0} = \begin{cases} \bigcup \{L_0^\ell : L_0 \in \mathcal{D}_{\lambda([0,1) \times J_0)}, L_0 \subset Y_{[0,1) \times \tilde{J}_0}\} & \text{if } J_0 = \tilde{J}_0^\ell, \\ \bigcup \{L_0^r : L_0 \in \mathcal{D}_{\lambda([0,1) \times J_0)}, L_0 \subset Y_{[0,1) \times \tilde{J}_0}\} & \text{if } J_0 = \tilde{J}_0^r. \end{cases}$$

We define the family of high frequency covers of the set $[0, 1) \times Y_{[0,1) \times J_0}$ by

$$\mathcal{F}_m = \{[0, 1) \times L \in \mathcal{R} : L \in \mathcal{D}_m, L \subset Y_{[0,1) \times J_0}\}, \quad (5.14)$$

for all $m > \lambda([0, 1) \times J_0)$, see Figure 8, and observe that

$$\bigcup \mathcal{F}_m = [0, 1) \times Y_{[0,1) \times J_0}. \quad (5.15)$$





FIGURE 8. The above figure depicts an instance of \mathcal{F}_m in Case 2.A. $L_0^{(k)}$ is a dyadic interval such that $[0, 1) \times L_0^{(k)} \in \mathcal{B}_{[0,1) \times \tilde{J}_0}$, and L is a dyadic interval such that $[0, 1) \times L \in \mathcal{F}_m$.

CASE 2.B: $I_0 \neq [0, 1)$. The collections indexed by the black rectangles have already been constructed. Here, we determine the collections for the gray rectangles.



By our induction hypothesis, $\{\mathcal{B}_{I \times J} : \mathcal{O}_{\triangleleft}(I \times J) \leq i_0 - 1\}$ satisfies (R1)–(R6). At this stage of the proof, the set $Y_{[0,1) \times J_0}$ has already been constructed. Now, we define the high frequency covers of $X_{I_0 \times J_0} \times Y_{[0,1) \times J_0}$ by putting

$$\mathcal{F}_m = \{K_0 \times L : K_0 \in \mathcal{X}_{I_0 \times J_0}, L \in \mathcal{D}_m, L \subset Y_{[0,1) \times J_0}\}, \quad (5.16)$$

whenever $m > \nu(\tilde{I}_0 \times J_0)$, see Figure 9.

In each of the above cases (5.9), (5.11), (5.14), and (5.16) we define the following functions. Firstly, let

$$f_m = \sum_{Q \in \mathcal{F}_m} h_Q, \quad (5.17a)$$

and secondly for any choice of signs $\varepsilon_Q \in \{-1, +1\}$, $Q \in \mathcal{F}_m$ put

$$f_m^{(\varepsilon)} = \sum_{Q \in \mathcal{F}_m} \varepsilon_Q h_Q. \quad (5.17b)$$

Now, we specify the value of m . To this end, put

$$k_{i_0} = \max\{\mu(R), \nu(R) : R \in \mathcal{R}, \mathcal{O}_{\triangleleft}(R) \leq i_0 - 1\}, \quad (5.18)$$




FIGURE 9. The above figure depicts an instance of \mathcal{F}_m in Case 2.B. We have $K_1^{(k)} \in \mathcal{X}_{\tilde{I}_0 \times J_0}$, $L_1^{(\ell)} \in \mathcal{Y}_{\tilde{I}_0 \times J_0}$, and the dyadic interval K_0 is in $\mathcal{X}_{I_0 \times J_0}$. \mathcal{F}_m is the collection of all the small gray rectangles. We obtain \mathcal{F}_m by leaving intact the intervals of the x -coordinate ($K_0 \in \mathcal{X}_{I_0 \times J_0}$) and using a high frequency cover – comprised of the intervals L – of the intervals $L_1^{(\ell)} \in \mathcal{Y}_{\tilde{I}_0 \times J_0}$. The intervals $L_1^{(\ell)} \in \mathcal{Y}_{\tilde{I}_0 \times J_0}$ in this Figure are covering the exact same set as the intervals denoted by L in Figure 8, i.e. they cover $Y_{[0,1] \times J_0}$.

and note that each \mathcal{F}_m , $m > k_{i_0}$ can be written as the product of two sets of intervals, i.e.

$$\mathcal{F}_m = \{K \times L : K \in \mathcal{X}_m, L \in \mathcal{Y}_m\}, \quad m > k_{i_0},$$

where the collections \mathcal{X}_m and \mathcal{Y}_m , $m > k_{i_0}$, satisfy the following:

- ▷ \mathcal{X}_m and \mathcal{Y}_m are each a non-empty, finite collection of pairwise disjoint dyadic intervals of equal length, whenever $m > k_{i_0}$;
- ▷ $\mathcal{X}_m \cap \mathcal{X}_n = \emptyset$ or $\mathcal{Y}_m \cap \mathcal{Y}_n = \emptyset$ whenever $m, n > k_{i_0}$ are distinct;
- ▷ the union of the sets in \mathcal{X}_m is independent of $m > k_{i_0}$, and the union of the sets in \mathcal{Y}_m is independent of $m > k_{i_0}$.

Thus, by Lemma 4.1, we have that

- ▷ for each $g \in H^p(H^q)^*$, $\sup_{\gamma \in \Gamma} |\langle M_\gamma f_m, g \rangle| \rightarrow 0$ as $m \rightarrow \infty$;
- ▷ for each $g \in H^p(H^q)$, $\sup_{\gamma \in \Gamma} |\langle M_\gamma g, f_m \rangle| \rightarrow 0$ as $m \rightarrow \infty$;

where we recall that Γ denotes the unit ball of $\ell^\infty(\mathcal{R})$, and that $\gamma = (\gamma_R : R \in \mathcal{R}) \in \Gamma$ defines the operator M_γ (see (4.2)). Hence, we can find an integer $m_{i_0} > k_{i_0}$ such that

$$\sum_{j=0}^{i_0-1} |\langle T b_j^{(\varepsilon)}, f_{m_{i_0}}^{(\varepsilon)} \rangle| + |\langle f_{m_{i_0}}^{(\varepsilon)}, T^* b_j^{(\varepsilon)} \rangle| \leq \eta \delta 4^{-i_0-2}, \quad (5.19)$$

for all choices of signs $\varepsilon_{K \times L}$, $K \times L \in \mathcal{F}_{m_{i_0}}$. Now, we put

$$\mathcal{B}_{I_0 \times J_0} = \mathcal{B}_{i_0} = \mathcal{F}_{m_{i_0}}. \quad (5.20)$$

If $I_0 \times J_0$ is a ‘‘Case 1’’ rectangle, i.e. $|I_0| < |J_0|$, then, by (5.9) and (5.11)

$$\mu(I_0 \times J_0) = m_{i_0} \quad \text{and} \quad \mathcal{X}_{I_0 \times J_0} = \{K \in \mathcal{D}_{m_{i_0}} : K \subset X_{I_0 \times [0,1]}\}, \quad (5.21a)$$

and if $I_0 \times J_0$ is a ‘‘Case 2’’ rectangle, i.e. $|I_0| \geq |J_0|$, then, by (5.14) and (5.16)

$$\nu(I_0 \times J_0) = m_{i_0} \quad \text{and} \quad \mathcal{Y}_{I_0 \times J_0} = \{L \in \mathcal{D}_{m_{i_0}} : L \subset Y_{[0,1] \times J_0}\}. \quad (5.21b)$$

Thereby, we have completed the construction of $\mathcal{B}_{I_0 \times J_0} = \mathcal{B}_{i_0}$.

Reviewing the four cases Case 1.A, Case 1.B, Case 2.A, and Case 2.B of the construction we see that $\{\mathcal{B}_i : i \leq i_0\}$ satisfies (R1)–(R6).

SELECTING THE SIGNS ε . Let $\varepsilon_Q \in \{\pm 1\}$, $Q \in \mathcal{B}_{i_0}$ be fixed. We obtain from (5.3) and (5.17)

$$\langle Tf_{m_{i_0}}^{(\varepsilon)}, f_{m_{i_0}}^{(\varepsilon)} \rangle = \sum_{Q \in \mathcal{B}_{i_0}} \alpha_Q |Q| + \langle f_{m_{i_0}}^{(\varepsilon)}, s_{m_{i_0}}^{(\varepsilon)} \rangle,$$

where

$$s_{m_{i_0}}^{(\varepsilon)} = \sum_{Q \in \mathcal{B}_{i_0}} \varepsilon_Q r_Q.$$

By (5.3) we have $\langle h_Q, r_Q \rangle = 0$, $Q \in \mathcal{B}$, and consequently

$$\langle f_{m_{i_0}}^{(\varepsilon)}, s_{m_{i_0}}^{(\varepsilon)} \rangle = \sum \varepsilon_{Q_0} \varepsilon_{Q_1} \langle h_{Q_0}, r_{Q_1} \rangle, \quad (5.22)$$

where the sum is taken over all $Q_0, Q_1 \in \mathcal{B}_{i_0}$ with $Q_0 \neq Q_1$. Let \mathbb{E}_ε denote the average over all possible choices of signs ε_Q , $Q \in \mathcal{B}_{i_0}$. Taking expectations we obtain from (5.22) that

$$\mathbb{E}_\varepsilon \langle f_{m_{i_0}}^{(\varepsilon)}, s_{m_{i_0}}^{(\varepsilon)} \rangle = 0.$$

This gives us

$$\mathbb{E}_\varepsilon \langle Tf_{m_{i_0}}^{(\varepsilon)}, f_{m_{i_0}}^{(\varepsilon)} \rangle = \sum_{Q \in \mathcal{B}_{i_0}} \alpha_Q |Q|.$$

Hence, in view of (5.4), there exists at least one ε such that

$$|\langle Tf_{m_{i_0}}^{(\varepsilon)}, f_{m_{i_0}}^{(\varepsilon)} \rangle| \geq \sum_{Q \in \mathcal{B}_{i_0}} \alpha_Q |Q| \geq \delta \|f_{m_{i_0}}^{(\varepsilon)}\|_2^2. \quad (5.23)$$

We complete the inductive construction by choosing ε according to (5.23) and define

$$b_{I_0 \times J_0}^{(\varepsilon)} = b_{i_0}^{(\varepsilon)} = f_{m_{i_0}}^{(\varepsilon)}. \quad (5.24)$$

Hence, (5.6b) holds for $i = i_0$, while (5.19) ensures that (5.6a) holds for $i = i_0$.

ESSENTIAL PROPERTIES OF OUR INDUCTIVE CONSTRUCTION. Since each of the finite collections $\{\mathcal{B}_i : i \leq i_0\}$, $i_0 \in \mathbb{N}_0$, satisfies (R1)–(R6), Remark 4.8 asserts that the infinite collection $\{\mathcal{B}_i : i \in \mathbb{N}_0\}$ satisfies (R1)–(R6), and hence, by Lemma 4.9, it satisfies the local product condition (P1)–(P4) with constants $C_X = C_Y = 1$.

For $1 \leq u, v < \infty$ and $I \times J \in \mathcal{R}$, Lemma 4.1(i)–(ii) together with (4.7) and (P3) gives us the following mixed-norm estimates for $b_{I \times J}^{(\varepsilon)}$:

$$\|b_{I \times J}^{(\varepsilon)}\|_{H^u(H^v)} = |I|^{1/u} |J|^{1/v} = \|h_{I \times J}\|_{H^u(H^v)}, \quad (5.25a)$$

$$\|b_{I \times J}^{(\varepsilon)}\|_{H^u(H^v)^*} = |I|^{1-1/u} |J|^{1-1/v} = \|h_{I \times J}\|_{H^u(H^v)^*}. \quad (5.25b)$$

The estimates (5.6a) and (5.6b) show that the block basis $\{b_i^{(\varepsilon)}\}$ almost-diagonalizes T in the following precise sense:

$$|\langle Tb_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle| \geq \delta \|b_i^{(\varepsilon)}\|_2^2, \quad i \in \mathbb{N}_0, \quad (5.26)$$

$$\sum_{j=0}^{i-1} |\langle Tb_j^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle| + |\langle Tb_i^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle| \leq \eta \delta 4^{-i-2}, \quad i \in \mathbb{N}. \quad (5.27)$$

PUTTING IT TOGETHER. The basic model of argument presented below can be traced to the seminal paper of Alspach, Enflo, and Odell [1]. Since $\{\mathcal{B}_{I \times J}\}$ satisfies the local product condition (P1)–(P4) with constants $C_X = C_Y = 1$, we obtain from Theorem 4.3 the following. First, let $Y = \overline{\text{span}}\{b_i^{(\varepsilon)} : i \in \mathbb{N}_0\} \subset H^p(H^q)$ and let

$B_\varepsilon : H^p(H^q) \rightarrow Y$ denote the unique linear extension of $B_\varepsilon h_i = b_i^{(\varepsilon)}$, $i \in \mathbb{N}_0$, then by Theorem 4.3

$$\begin{array}{ccc} H^p(H^q) & \xrightarrow{I_{H^p(H^q)}} & H^p(H^q) \\ B_\varepsilon \downarrow & & \uparrow A_\varepsilon|_Y \\ Y & \xrightarrow{I_Y} & Y \end{array} \quad \|B_\varepsilon\| = \|A_\varepsilon\| = 1, \quad (5.28)$$

where we recall that $A_\varepsilon : H^p(H^q) \rightarrow H^p(H^q)$ denotes the operator given by

$$A_\varepsilon f = \sum_{i=0}^{\infty} \frac{\langle f, b_i^{(\varepsilon)} \rangle}{\|b_i\|_2^2} h_i, \quad f \in H^p(H^q).$$

Secondly, we put

$$\gamma_i = \frac{\|b_i\|_2^2}{\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle}, \quad i \in \mathbb{N}_0.$$

Recall that M_γ was defined in (4.2) as the linear extension of $M_\gamma h_i = \gamma_i h_i$, $i \in \mathbb{N}_0$. The operator norm of M_γ is $\sup_{i \in \mathbb{N}_0} |\gamma_i| \leq \frac{1}{\delta}$ by (5.26). Define $U : H^p(H^q) \rightarrow Y$ by $U = B_\varepsilon M_\gamma A_\varepsilon$ and note that

$$U(f) = \sum_{i=0}^{\infty} \frac{\langle f, b_i^{(\varepsilon)} \rangle}{\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle} b_i^{(\varepsilon)}, \quad f \in H^p(H^q). \quad (5.29)$$

The above estimates for the norms of the operators A_ε , B_ε , and M_γ yield

$$\|U : H^p(H^q) \rightarrow Y\| \leq \|M_\gamma\| \|B_\varepsilon\| \|A_\varepsilon\| \leq \frac{1}{\delta}. \quad (5.30)$$

Thirdly, observe that for all $g = \sum_{i=0}^{\infty} \lambda_i b_i^{(\varepsilon)} \in Y$, we have the identity

$$UTg - g = \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} \lambda_j \frac{\langle T b_j^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle}{\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle} b_i^{(\varepsilon)} + \lambda_i \frac{\langle T b_i^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle}{\langle T b_j^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle} b_j^{(\varepsilon)}. \quad (5.31)$$

Using that $\|b_j^{(\varepsilon)}\|_{H^p(H^q)} \leq 1$, $j \in \mathbb{N}_0$, we obtain

$$\|UTg - g\|_{H^p(H^q)} \leq \sum_{i=1}^{\infty} \sum_{j=0}^{i-1} |\lambda_j| \frac{|\langle T b_j^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle|}{|\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle|} + |\lambda_i| \frac{|\langle T b_i^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle|}{|\langle T b_j^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle|}. \quad (5.32)$$

Now, we will make the following two observations: The first is that (5.25b) implies $\|b_j^{(\varepsilon)}\|_{H^p(H^q)^*} \leq 1$, and thus by (5.25a) and (4.25), we obtain

$$\|g\|_{H^p(H^q)} \geq \|g\|_{H^p(H^q)} \|b_j^{(\varepsilon)}\|_{H^p(H^q)^*} \geq |\langle g, b_j^{(\varepsilon)} \rangle| = |\lambda_j| \|b_j^{(\varepsilon)}\|_2^2 \geq \frac{1}{(1+\sqrt{j})^2} |\lambda_j|,$$

for all $j \in \mathbb{N}_0$. The second observation is that $|\langle T b_j^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle| \geq \frac{\delta}{(1+\sqrt{j})^2}$, $j \in \mathbb{N}_0$, which is a consequence of (5.26), (5.25a), and (4.25). These two observations yield the following estimate:

$$\frac{|\lambda_j|}{|\langle T b_i^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle|} \leq \frac{1}{\delta} \|g\|_{H^p(H^q)} (1+\sqrt{i})^2 (1+\sqrt{j})^2, \quad j \neq i.$$

Inserting this estimate into (5.32) and applying (5.27) yields

$$\begin{aligned} \|UTg - g\|_{H^p(H^q)} &\leq \frac{1}{\delta} \|g\|_{H^p(H^q)} \sum_{i=1}^{\infty} (1+\sqrt{i})^4 \sum_{j=0}^{i-1} |\langle T b_j^{(\varepsilon)}, b_i^{(\varepsilon)} \rangle| + |\langle T b_i^{(\varepsilon)}, b_j^{(\varepsilon)} \rangle| \\ &\leq \frac{\eta}{2} \|g\|_{H^p(H^q)}. \end{aligned}$$

To see the latter estimate note that $\sum_{i=1}^{\infty} (1 + \sqrt{i})^4 4^{-i} \leq 4 \sum_{i=1}^{\infty} (1 + i)^2 4^{-i} = 4 \frac{53}{27} \leq 8$.

Finally, let $J : Y \rightarrow H^p(H^q)$ denote the inclusion operator given by $Jy = y$. Since we assumed that $0 < \eta \leq 1$, the operator UTJ is invertible, and its inverse has norm at most $(1 - \frac{\eta}{2})^{-1} \leq 1 + \eta$. Now we define the operator $V : H^p(H^q) \rightarrow Y$ by $(UTJ)^{-1}U$ and observe that

$$\begin{array}{ccc}
 Y & \xrightarrow{I_Y} & Y \\
 \downarrow J & \searrow UTJ & \nearrow (UTJ)^{-1} \\
 & Y & \\
 \downarrow & \swarrow U & \uparrow V \\
 H^p(H^q) & \xrightarrow{T} & H^p(H^q)
 \end{array}
 \quad \|J\| \|V\| \leq (1 + \eta)/\delta. \quad (5.33)$$

Merging the commutative diagram (5.28) with (5.33) yields

$$\begin{array}{ccc}
 H^p(H^q) & \xrightarrow{I_{H^p(H^q)}} & H^p(H^q) \\
 \downarrow B_\varepsilon & & \uparrow A_\varepsilon|Y \\
 Y & \xrightarrow{I_Y} & Y \\
 \downarrow J & \searrow UTJ & \nearrow (UTJ)^{-1} \\
 & Y & \\
 \downarrow & \swarrow U & \uparrow V \\
 H^p(H^q) & \xrightarrow{T} & H^p(H^q)
 \end{array}$$

where $\|B_\varepsilon\| = \|A_\varepsilon\| = 1$ and $\|J\| \|V\| \leq (1 + \eta)/\delta$, which concludes the proof of Theorem 3.1. \square

REFERENCES

- [1] D. Alspach, P. Enflo, and E. Odell. On the structure of separable \mathcal{L}_p spaces ($1 < p < \infty$). *Studia Math.*, 60(1):79–90, 1977.
- [2] A. D. Andrew. Perturbations of Schauder bases in the spaces $C(K)$ and L^p , $p > 1$. *Studia Math.*, 65(3):287–298, 1979.
- [3] M. Capon. Primarité de $L^p(L^r)$, $1 < p, r < \infty$. *Israel J. Math.*, 42(1-2):87–98, 1982.
- [4] S. R. Caradus, W. E. Pfaffenberger, and B. Yood. *Calkin algebras and algebras of operators on Banach spaces*. Marcel Dekker, Inc., New York, 1974. Lecture Notes in Pure and Applied Mathematics, Vol. 9.
- [5] J. L. B. Gamlen and R. J. Gaudet. On subsequences of the Haar system in $L_p [0, 1]$ ($1 \leq p \leq \infty$). *Israel J. Math.*, 15:404–413, 1973.
- [6] D. J. H. Garling. *Inequalities: a journey into linear analysis*. Cambridge University Press, Cambridge, 2007.
- [7] W. T. Gowers. A solution to Banach’s hyperplane problem. *Bull. London Math. Soc.*, 26(6):523–530, 1994.
- [8] W. T. Gowers and B. Maurey. Banach spaces with small spaces of operators. *Math. Ann.*, 307(4):543–568, 1997.
- [9] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge University Press, 1952. 2nd ed.
- [10] W. B. Johnson, B. Maurey, G. Schechtman, and L. Tzafriri. Symmetric structures in Banach spaces. *Mem. Amer. Math. Soc.*, 19(217):v+298, 1979.
- [11] R. Lechner. Factorization in mixed norm Hardy and BMO spaces. *Studia Math.*, to appear. Preprint available on ArXiv.
- [12] R. Lechner. Factorization in SL^∞ . *Israel J. Math.*, to appear. Preprint available on ArXiv.

- [13] R. Lechner and P. F. X. Müller. Localization and projections on bi-parameter BMO. *Q. J. Math.*, 66(4):1069–1101, 2015.
- [14] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces. I*. Springer-Verlag, Berlin-New York, 1977. Sequence spaces, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 92.
- [15] B. Maurey. Système de Haar. *Séminaire Maurey-Schwartz 1974–1975: Espaces L^p , applications radonifiantes et géométrie des espaces de Banach, Exp. Nos. I et II*, 1975.
- [16] B. Maurey. Isomorphismes entre espaces H_1 . *Acta Math.*, 145(1-2):79–120, 1980.
- [17] P. F. X. Müller. Orthogonal projections on martingale H^1 spaces of two parameters. *Illinois J. Math.*, 38(4):554–573, 1994.
- [18] P. F. X. Müller. *Isomorphisms between H^1 spaces*, volume 66 of *Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) [Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series)]*. Birkhäuser Verlag, Basel, 2005.
- [19] A. Pełczyński. Projections in certain Banach spaces. *Studia Math.*, 19:209–228, 1960.
- [20] G. Pisier. *Martingales in Banach Spaces*. Cambridge University Press, 2016.

NIELS JAKOB LAUSTSEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE,
LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UNITED KINGDOM
Email address: `n.laustsen@lancaster.ac.uk`

RICHARD LECHNER, INSTITUTE OF ANALYSIS, JOHANNES KEPLER UNIVERSITY LINZ, AL-
TENBERGER STRASSE 69, A-4040 LINZ, AUSTRIA
Email address: `Richard.Lechner@jku.at`

PAUL F.X. MÜLLER, INSTITUTE OF ANALYSIS, JOHANNES KEPLER UNIVERSITY LINZ, AL-
TENBERGER STRASSE 69, A-4040 LINZ, AUSTRIA
Email address: `Paul.Mueller@jku.at`