

Third-generation cephalosporin resistance conferred by a chromosomally encoded *bla*_{CMY-23} gene in the *Escherichia coli* ST131 reference strain EC958

Minh-Duy Phan¹, Kate M. Peters¹, Sohinee Sarkar¹, Brian M. Forde¹, Alvin W. Lo¹, Mitchell Stanton-Cook¹, Leah W. Roberts¹, Mathew Upton², Scott A. Beatson¹ and Mark A. Schembri^{1*}

¹Australian Infectious Diseases Research Centre, School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia; ²Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth PL4 8AA, UK

*Corresponding author. Tel: +617-33653306; Fax: +617-33654699; E-mail: m.schembri@uq.edu.au

Received 5 January 2015; returned 30 January 2015; revised 12 February 2015; accepted 23 February 2015

Objectives: *Escherichia coli* ST131 is a globally disseminated MDR clone originally identified due to its association with the *bla*_{CTX-M-15} gene encoding an ESBL. It is thus assumed that *bla*_{CTX-M-15} is the major determinant for resistance to β -lactam antibiotics in this clone. The complete sequence of EC958, a reference strain for *E. coli* ST131, revealed that it contains a chromosomally located *bla*_{CMY-23} gene with an upstream *ISEcp1* element as well as several additional plasmid-encoded β -lactamase genes. Here, we examined the genetic context of the *bla*_{CMY-23} element in EC958 and other *E. coli* ST131 strains and investigated the contribution of *bla*_{CMY-23} to EC958 resistance to a range of β -lactam antibiotics.

Methods: The genetic context of *bla*_{CMY-23} and its associated mobile elements was determined by PCR and sequencing. Antibiotic susceptibility testing was performed using Etests. The activity of the *bla*_{CMY-23} promoter was assessed using *lacZ* reporter assays. Mutations were generated using λ -Red-recombination.

Results: The genetic structure of the *ISEcp1*-*IS5*-*bla*_{CMY-23} mobile element was determined and localized within the *betU* gene on the chromosome of EC958 and five other *E. coli* ST131 strains. The transcription of *bla*_{CMY-23}, driven by a previously defined promoter within *ISEcp1*, was significantly higher than other β -lactamase genes and could be induced by cefotaxime. Deletion of the *bla*_{CMY-23} gene resulted in enhanced susceptibility to cefoxitin, cefotaxime and ceftazidime.

Conclusions: This is the first known report to demonstrate the chromosomal location of *bla*_{CMY-23} in *E. coli* ST131. In EC958, CMY-23 plays a major role in resistance to third-generation cephalosporins and cephamycins.

Keywords: *E. coli* ST131, antibiotic resistance, AmpC β -lactamases

Introduction

Escherichia coli ST131 is a globally disseminated MDR clone originally identified due to its close association with the spread of the *bla*_{CTX-M-15} ESBL gene.^{1–3} Two studies have recently reported the global epidemiology of *E. coli* ST131 using WGS, and revealed the *bla*_{CTX-M-15} allele is highly prevalent within a fluoroquinolone-resistant FimH30 (H30) ST131 sublineage (referred to as clade C2⁴ or H30-Rx⁵). We have characterized and completely sequenced the reference ST131 strain EC958.⁶ *E. coli* EC958 is a urinary tract infection (UTI)-derived fluoroquinolone-resistant, *fimH30* *E. coli* ST131 strain.⁷ The strain is a representative member of the UK epidemic strain A, one of the major pathogenic lineages causing UTI across the UK,³ and belongs to the recently defined phylogenetic C2 lineage of ST131.⁴ EC958 contains a large multidrug resistance plasmid (pEC958; HG941719) that harbours a copy of *bla*_{CTX-M-15} gene. The

*bla*_{CTX-M-15} gene has always been seen with an upstream *ISEcp1* insertion element.⁸ *ISEcp1* contains a well-described outward-facing promoter that drives the transcription of the downstream *bla*_{CTX-M-15} gene,⁹ and this transposition unit has been identified on multiple plasmids characterized from ST131 strains.¹⁰ Several different insertions of IS26 in the region upstream of *bla*_{CTX-M-15} have been reported, including the insertion at 24 bp upstream of the right inverted repeat (IRR) of *ISEcp1*, which results in reduced resistance to cephalosporins.¹⁰ This particular insertion was first identified in UK epidemic strain A isolates and was also found on plasmid pEC958 from the *E. coli* ST131 strain EC958.

In addition to the production of ESBLs, resistance to third-generation cephalosporins can also be mediated by chromosomal and plasmid-encoded AmpC (or class C) β -lactamases.¹¹ The plasmid-encoded *bla*_{CMY-1} gene and its variants are related to chromosomal AmpC enzymes from *Aeromonas* spp., while

*bla*_{CMY-2} and its variants are related to AmpC β -lactamases from *Citrobacter freundii*.¹¹ *bla*_{CMY-2} gene is the most frequently reported plasmid-mediated AmpC β -lactamase worldwide; it confers resistance to cephamycins (cefoxitin and cefotetan) and oxyimino-cephalosporins (cefotaxime and ceftazidime) and is poorly inhibited by β -lactamase inhibitors, including clavulanate, sulbactam and tazobactam.¹² Several studies have reported the identification of *bla*_{CMY-2} in *E. coli* ST131.^{13,14} In addition, a variant of *bla*_{CMY-2}, named *bla*_{CMY-23}, has been identified in several *E. coli* ST131 isolates belonging to the UK epidemic *E. coli* strain A.¹⁵ The *bla*_{CMY-2} and *bla*_{CMY-23} genes differ by a single nucleotide, which results in a Glu239Gly amino acid substitution in CMY-23.

A *bla*_{CMY-23} gene accompanied by an upstream *ISEcp1* insertion element was found inserted in the chromosome of EC958. This led us to hypothesize that CMY-23, instead of CTX-M-15, is the β -lactamase responsible for cephalosporin resistance in EC958. Here, we investigated the contribution of *bla*_{CMY-23} to EC958 resistance to a range of β -lactam antibiotics, and demonstrated that it is the primary mediator of resistance to third-generation cephalosporins and cephamycins.

Materials and methods

Bacterial strains and growth conditions

The strains used in this study are listed in Table S1 (available as Supplementary data at JAC Online). All *E. coli* strains were routinely cultured at 37°C on solid or in liquid Lysogeny Broth (LB) medium supplemented with an appropriate antibiotic: chloramphenicol (30 mg/L), gentamicin (20 mg/L) or ampicillin (100 mg/L).

Antimicrobial susceptibility testing

The MICs of β -lactam antibiotics were determined by Etest (bioMérieux Australia) on Mueller–Hinton agar at 37°C using EUCAST breakpoints (http://www.eucast.org/clinical_breakpoints/).

Molecular methods

Genomic DNA preparation and DNA sequencing were performed as described previously.¹⁶ Oligonucleotides used in this study are listed in Table S2. Plasmids pQF50::ISEcp1 and pQF50::IS5+ISEcp1 were created by PCR amplification of the fragments indicated in Table S1 and cloned into pQF50 via BamHI–HindIII digestion. The resulting plasmids were electroporated into *E. coli* TOP10. Mutants containing deletions in the *bla*_{CMY-23} gene and IS5 were constructed by λ -Red-mediated recombination as previously described.⁷ β -Galactosidase assays were performed as previously described.¹⁶

Results and discussion

The *bla*_{CMY-23} gene is located on the chromosome of *E. coli* EC958

E. coli EC958 contains a chromosomally encoded *bla*_{CMY-23} gene (EC958_4781). Closer inspection of its genomic context suggests an insertion event of a 4277 bp mobile element (nucleotides 4937433–4941709, including *ISEcp1*, IS5 and *bla*_{CMY-23}) into the *betU* gene (encodes a betaine uptake system) within the GI-*leuX* genomic island, creating 5 bp direct repeats (DRs) of TATAT (Figure 1a). This insertion was not found in the *E. coli* ST131 strain JJ1886 (CP006784), which harbours two tandem copies of the

GI-*leuX* island (Figure S1). *ISEcp1* is known to mediate the transposition of its downstream genes using an imperfect IRR (IRR').¹⁷ Indeed, we identified an IRR' with the sequence 5'-gtGTAGctcCccGa-3' (nucleotides matching the *ISEcp1* IRR are capitalized) immediately before the right DR. The IS5 element is located 152 bp before the end of *ISEcp1* (Figure 1b and c) and flanked by a 4 bp DR of TTAA. This location does not interrupt the *ISEcp1* transposase gene and leaves the promoter region at the end of *ISEcp1* intact. It is not clear whether this IS5 insertion occurred prior to or after the movement of *ISEcp1*-*bla*_{CMY-23} to the chromosome.

Prevalence of *ISEcp1*-*bla*_{CMY-23}

The *bla*_{CMY-23} gene (DQ438952) was first reported from isolates within the UK epidemic strain A, which was later shown to be part of the *E. coli* ST131 lineage.¹⁵ A second *bla*_{CMY-23} sequence has been reported from *Salmonella enterica* serovar Senftenberg (DQ463751). However, neither of these entries includes information about the genomic context nor location of the *bla*_{CMY-23} gene.

Analysis of a global collection of previously sequenced ST131 strains⁴ identified five clade C2 strains (S30EC, accession number ERR161246; S39EC, accession number ERR161248; S43EC, accession number ERR161249; S47EC, accession number ERR161250; and S53EC, accession number ERR161251) that possess the *bla*_{CMY-23} gene. Further PCR and sequencing of S30EC, S39EC and S43EC confirmed that they contain an identical *ISEcp1*-IS5-*bla*_{CMY-23} sequence inserted at the same location on the chromosome as in EC958 (Figure 1b). Of note, EC958 and these five strains were all isolated from the same hospital in the UK (from 2004 to 2007) where the first *bla*_{CMY-23} positive strain was reported.¹⁵

We also identified one strain (S79EC, accession number ERR161305) from the ST131 clade A phylogeny that contained the *bla*_{CMY-2} gene within a transposition unit identical to that found in IncI1 plasmids pSTM709 (HG428759) and pCVM29188_101 (CP001121). This unit lacks IS5, has a longer region downstream of *ISEcp1* (2253 bp compared with 1422 bp) and based on its flanking genes may be located on a plasmid in S79EC (Figure 1b).

The *bla*_{CMY-23} gene is the major contributor to EC958 resistance to third-generation cephalosporins

EC958 harbours several genes that could contribute to its resistance to β -lactam antibiotics, including *bla*_{TEM-1}, *bla*_{OXA-1}, *bla*_{CTX-M-15}, *bla*_{CMY-23} and the chromosomal *ampC* gene. To specifically demonstrate the contribution of *bla*_{CMY-23} to EC958 resistance, we mutated the *bla*_{CMY-23} gene using λ -Red-mediated homologous recombination. The resultant strain, designated EC958 Δ *bla*_{CMY-23}, was then tested in parallel with WT EC958 against a panel of β -lactam antibiotics. Despite the presence of pEC958 (which harbours *bla*_{TEM-1}, *bla*_{OXA-1} and *bla*_{CTX-M-15}), EC958 Δ *bla*_{CMY-23} displayed significantly enhanced susceptibility to cefoxitin, cefotaxime and ceftazidime (8–16-fold reduction in MIC; Table 1) compared with WT EC958. Taken together, our data confirm that the *bla*_{CMY-23} gene is the main determinant that mediates resistance to second- and third-generation cephalosporins in EC958.

*bla*_{CMY-23} expression is induced by cefotaxime, but not affected by IS5

We used qRT–PCR to evaluate the transcription of the five β -lactamase genes in EC958. Our results showed that *bla*_{CMY-23}

Table 1. Antimicrobial susceptibility (expressed as MIC in mg/L) of EC958 WT and mutant strains

Antibiotic	EC958	EC958Δ <i>bla</i> _{CMY-23}	EC958ΔIS5
Ampicillin	≥256 (R)	≥256 (R)	≥256 (R)
Amoxicillin/clavulanic acid	24 (R)	12 (R)	16 (R)
Piperacillin	≥256 (R)	≥256 (R)	≥256 (R)
Piperacillin/tazobactam	16 (I)	12 (I)	16 (I)
Cefoxitin	48 (NA)	4 (NA)	48 (NA)
Cefotaxime	12 (R)	1.5 (I)	12 (R)
Ceftazidime	16 (R)	1 (S)	16 (R)
Cefpirome	3 (NA)	1 (NA)	3 (NA)
Aztreonam	2 (I)	1 (S)	2 (I)

R, resistant; I, intermediate; S, susceptible; NA, breakpoints not available.

activity was detected in the vector control strain grown under the same conditions.

IS5 has been shown to alter transcription by enhancing the activity of a promoter downstream of its site of insertion.^{19,20} Thus, we investigated whether the IS5 element located within *ISEcp1* affects expression of the *bla*_{CMY-23} gene using two complementary approaches. First, we cloned the *ISEcp1* promoter region together with the upstream IS5 in front of the promoterless *lacZ* gene in plasmid pQF50 to generate plasmid pQF50::IS5+*ISEcp1* and showed that there was no difference in the β-galactosidase activity between TOP10(pQF50::IS5+*ISEcp1*) and TOP10(pQF50::*ISEcp1*) (Figure 1d). Second, we deleted the IS5 element using λ-Red-mediated homologous recombination. The resultant strain, designated EC958ΔIS5, did not show any difference in antibiotic resistance profile from that of WT EC958 (Table 1). These results demonstrate that the *bla*_{CMY-23} promoter provided by *ISEcp1* is functional and can be induced by cefotaxime.

Conclusions

*bla*_{CTX-M-15} is considered to be the main determinant for resistance to third-generation cephalosporins in *E. coli* ST131. Here, we showed that in EC958, which contains a plasmid-located *bla*_{CTX-M-15} gene, cephalosporin resistance is conferred by a chromosomal *bla*_{CMY-23} gene that encodes an AmpC type β-lactamase. We also showed that the *bla*_{CMY-23} in EC958 is strongly transcribed from a promoter located within *ISEcp1* that can be induced by cefotaxime.

Funding

This work was supported by grants from the Australian National Health and Medical Research Council to M. A. S. and S. A. B. (APP1012076 and APP1067455). M. A. S. is supported by an Australian Research Council (ARC) Future Fellowship (FT100100662).

Transparency declarations

None to declare.

Supplementary data

Tables S1 and S2 and Figures S1 and S2 are available as Supplementary data at JAC Online (<http://jac.oxfordjournals.org/>).

References

- Nicolas-Chanoine MH, Blanco J, Leflon-Guibout V et al. Intercontinental emergence of *Escherichia coli* clone O25:H4-ST131 producing CTX-M-15. *J Antimicrob Chemother* 2008; **61**: 273–81.
- Coque TM, Novais A, Carattoli A et al. Dissemination of clonally related *Escherichia coli* strains expressing extended-spectrum β-lactamase CTX-M-15. *Emerg Infect Dis* 2008; **14**: 195–200.
- Lau SH, Kaufmann ME, Livermore DM et al. UK epidemic *Escherichia coli* strains A-E, with CTX-M-15 β-lactamase, all belong to the international O25:H4-ST131 clone. *J Antimicrob Chemother* 2008; **62**: 1241–4.
- Petty NK, Ben Zakour NL, Stanton-Cook M et al. Global dissemination of a multidrug resistant *Escherichia coli* clone. *Proc Natl Acad Sci USA* 2014; **111**: 5694–9.
- Price LB, Johnson JR, Aziz M et al. The epidemic of extended-spectrum-β-lactamase-producing *Escherichia coli* ST131 is driven by a single highly pathogenic subclone, H30-Rx. *MBio* 2013; **4**: e00377–13.
- Forde BM, Ben Zakour NL, Stanton-Cook M et al. The complete genome sequence of *Escherichia coli* EC958: a high quality reference sequence for the globally disseminated multidrug resistant *E. coli* O25b:H4-ST131 clone. *PLoS One* 2014; **9**: e104400.
- Totsika M, Beatson SA, Sarkar S et al. Insights into a multidrug resistant *Escherichia coli* pathogen of the globally disseminated ST131 lineage: genome analysis and virulence mechanisms. *PLoS One* 2011; **6**: e26578.
- Partridge SR, Zong Z, Iredell JR. Recombination in IS26 and Tn2 in the evolution of multiresistance regions carrying *bla*_{CTX-M-15} on conjugative IncF plasmids from *Escherichia coli*. *Antimicrob Agents Chemother* 2011; **55**: 4971–8.
- Poirel L, Decousser JW, Nordmann P. Insertion sequence *ISEcp1B* is involved in expression and mobilization of a *bla*_{CTX-M} β-lactamase gene. *Antimicrob Agents Chemother* 2003; **47**: 2938–45.
- Dhanji H, Patel R, Wall R et al. Variation in the genetic environments of *bla*_{CTX-M-15} in *Escherichia coli* from the faeces of travellers returning to the United Kingdom. *J Antimicrob Chemother* 2011; **66**: 1005–12.
- Jacoby GA. AmpC β-lactamases. *Clin Microbiol Rev* 2009; **22**: 161–82.
- Bauernfeind A, Stemplinger I, Jungwirth R et al. Characterization of the plasmidic β-lactamase CMY-2, which is responsible for cephamycin resistance. *Antimicrob Agents Chemother* 1996; **40**: 221–4.
- Nicolas-Chanoine MH, Bertrand X, Madec JY. *Escherichia coli* ST131, an intriguing clonal group. *Clin Microbiol Rev* 2014; **27**: 543–74.
- Sidjabat HE, Seah KY, Coleman L et al. Expansive spread of Inc11 plasmids carrying *bla*_{CMY-2} amongst *Escherichia coli*. *Int J Antimicrob Agents* 2014; **44**: 203–8.
- Woodford N, Reddy S, Fagan EJ et al. Wide geographic spread of diverse acquired AmpC β-lactamases among *Escherichia coli* and *Klebsiella* spp. in the UK and Ireland. *J Antimicrob Chemother* 2007; **59**: 102–5.
- Allsopp LP, Beloin C, Gomes Moriel D et al. Functional heterogeneity of the UpaH autotransporter protein from uropathogenic *Escherichia coli*. *J Bacteriol* 2012; **194**: 5769–82.
- Poirel L, Lartigue MF, Decousser JW et al. *ISEcp1B*-mediated transposition of *bla*_{CTX-M} in *Escherichia coli*. *Antimicrob Agents Chemother* 2005; **49**: 447–50.
- Hossain A, Reisbig MD, Hanson ND. Plasmid-encoded functions compensate for the biological cost of AmpC overexpression in a clinical isolate of *Salmonella typhimurium*. *J Antimicrob Chemother* 2004; **53**: 964–70.
- Zhang Z, Saier MH Jr. A novel mechanism of transposon-mediated gene activation. *PLoS Genet* 2009; **5**: e1000689.
- Schnetz K, Rak B. IS5: a mobile enhancer of transcription in *Escherichia coli*. *Proc Natl Acad Sci USA* 1992; **89**: 1244–8.