

1 **Treatment of Dragonfruit (*Hylocereus polyrhizus*) with Salicylic Acid and**
2 **Methyl Jasmonate improves Postharvest Physico-chemical Properties and**
3 **Antioxidant Activity during Cold Storage**

4
5 MAYSOUN A. MUSTAFA¹, ASGAR ALI^{1*}, GRAHAM SEYMOUR² AND GREGORY
6 TUCKER³

7
8 ¹Centre of Excellence for Postharvest Biotechnology (CEPB), School of Biosciences,
9 The University of Nottingham Malaysia Campus, 43500 Semenyih,
10 Selangor, Malaysia

11 ²Division of Plant and Crop Sciences; and ³Division of Nutritional Sciences, School of
12 Biosciences, The University of Nottingham, Sutton Bonington Campus, Leicestershire,
13 United Kingdom

14
15 ***E-mail address: asgar.ali@nottingham.edu.my**

16

17

18 As the market for tropical fruit constantly expands, cold storage is increasingly used for
19 transporting fruits over long distances. This is an economic postharvest tool, yet challenges
20 tropical fruits by exposure to chilling injury. An assessment of the effect of abiotic stresses,
21 induced by cold storage, on dragonfruit was conducted. Dragonfruit was treated with salicylic
22 acid (SA) and methyl jasmonate (MJ) and subjected to cold storage for three weeks. Fruits
23 were treated with either SA or MJ, administered at four different concentrations, along with
24 an untreated control, and stored at 6 °C. Changes in biochemical quality parameters, along
25 with bioactive content and antioxidant activity were assessed during storage. Application of
26 SA was found to reduce the metabolic activity of the fruit, as determined by soluble solids
27 content and titratable acidity. Meanwhile, MJ significantly enhanced the betacyanin content
28 and antioxidant activity. We demonstrate that cold storage can be applied for dragonfruit, by
29 combining the treatment with the application of hormones, especially MJ which can enhance
30 the antioxidant activity of dragonfruit under cold storage.

31

32

33 **Keywords:** abiotic stress, metabolic activity, ripening, nutraceutical value, stress hormones

34

35 **Running title:** Dragonfruit responses to stress hormones and chilling

36 **1. Introduction**

37 Cold storage is an economic postharvest tool for extending shelf life of perishable plant
38 products. However, it is a source of postharvest stress on tropical fruits, which may result in
39 physiological changes in the plant organ (Wang and Frei, 2011). Chilling injury of fruits is
40 closely associated with degradation of the membrane integrity, resulting in oxidation of
41 phenolic compounds. This negatively affects the physical and biochemical qualities of fruits,
42 manifesting through browning and shrivelling of fruit surface, as well as depletion of
43 phenolic compounds.

44 Phenolic compounds and antioxidants are of particular importance, as they play a role
45 in acclimation to cold stress (Gonzalez-Aguilar et al., 2004). An increase in accumulation of
46 these compounds during chilling tolerance has been reported, which serves to regulate the
47 oxidative stress on the biological system (Chen et al., 2006). This demonstrates an interesting
48 phenomenon where abiotic stresses, such as chilling, can trigger intrinsic defence
49 mechanisms of the plant and ultimately enhance the bioactive content of the fruit. Exploiting
50 these natural resistance mechanisms for extending the shelf life of products is increasingly
51 being explored as an alternative to traditional postharvest tools (Ashgari and Hassanlooe,
52 2015). The authors have reported this on other tropical fruits such as mangosteen and
53 carambola (Mustafa et al., 2016), and this research will explore this approach on the tropical
54 fruit, dragonfruit.

55 Dragonfruit (*Hylocerus* sp.) is an exotic fruit of the family Cactaceae commonly
56 found in the tropical regions of South America and Southeast Asia. It has been increasing in
57 popularity due to its attractive purple coloration and distinct nutraceutical health properties
58 and anti-inflammatory characteristics (Macias-Ceja et al., 2016; Dembitsky et al., 2011). The
59 characteristic skin colour is attributed to the presence of betacyanins and the flesh is also rich
60 in betacyanins and phenolic compounds, and is characterized by potent antioxidant activity

61 (Wu et al., 2006). As a tropical fruit, dragonfruit is best stored at 14 - 20 °C, and is
62 susceptible to chilling injury at 6 °C (Hoa et al., 2006; Nerd et al., 1999). Moreover, the fruit
63 is preferably harvested at maturity, which minimises the incidence of chilling injury during
64 cold storage, allowing the fruit to maintain desirable quality during marketing (Wanitchang et
65 al., 2010; Nerd et al. 1999). Nonetheless, relatively scarce information is available on the
66 changes in the physico-chemical properties and antioxidant activity of the fruit during cold
67 storage.

68 The stress hormones salicylic acid (SA) and methyl jasmonate (MJ) are also involved
69 in improving natural stress tolerance in crops by enhancing secondary metabolism (Gonzalez-
70 Aguilar et al., 2010; Heredia and Cisneros-Zevallos, 2009). SA is involved throughout plant
71 growth and development and is directly associated with accumulation of protective
72 compounds during systemic acquired resistance (Chen et al., 2006). MJ enhances the
73 defense-signaling pathway, operating through the octadecanoid pathway and eliciting
74 production of secondary metabolites such as phenolic acids (Heredia and Cisneros-Zevallos,
75 2009; Kim et al., 2006), some of which may act as antioxidants. These secondary metabolites
76 can react with reactive oxygen species (ROS) triggered during stress, and quench ROS
77 activity directly or indirectly. The nutraceutical properties of the fruit can be closely
78 associated to changes in the physico-chemical qualities of the fruit. Thus, the importance of
79 plant-based secondary metabolites as a strong contributor to aesthetic and nutritional
80 properties of fruits and vegetables (Rautiainen et al., 2012; Marnewick et al., 2011).

81 Nutraceutical properties of fruits and vegetables are of significant interest to
82 consumers, dually as protective compounds for maintaining the fruit quality and for their
83 nutritional benefits. Moreover, Thus, this study aims to explore the effect of SA and MJ in
84 combination with storage at 6 °C could on shelf life and the bioactive content of dragonfruit.

85

86 2. Materials and Methods

87 2.1 Plant material

88 Disease free and uniform sized dragonfruit were purchased from a commercial farm in the
89 state of Negeri Sembilan, Malaysia. Freshly harvested fruit that were free from disease and of
90 uniform size and shape were selected, and then washed with 0.05% sodium hypochlorite
91 followed by distilled water and air-dried at ambient temperature.

92

93 2.2 Treatments of fruit

94 The fruits were subjected to one of two sets of treatments. The first being the MJ treatment,
95 with fruits subjected to MJ vapour at concentrations of 0.01 mM, 0.1 mM, 0.2 mM and 0.5
96 mM MJ, along with an untreated control. The fruits were incubated in 45 L airtight storage
97 boxes for 16 h at room temperature, and then allowed to ventilate for 2 h. The second set was
98 the SA treatment, which involved dipping the fruits in SA solution at concentrations of 0.1
99 mM, 1 mM, 2 mM and 5 mM SA, along with a control dipped in water. After treatment,
100 fruits were packed in cardboard boxes and stored for 21 d at 6 °C, 60-70 % relative humidity,
101 and were assessed at 7 day intervals.

102

103 2.3 Biochemical quality parameters

104 Fruit pulp was mixed with distilled water at a ration of 1:10 and homogenized in a blender.
105 Soluble solids content (SSC) was determined using a Palette Digital Refractometer (Model:
106 PR-32 α , Atago Co., Ltd. Japan) that was standardised with distilled water. The refractometer
107 was calibrated against sucrose.

108 The fruit homogenate was titrated against 0.1 N NaOH, using 0.1 % phenolphthalein
109 as an indicator, to determine titratable acidity (Ranganna, 1977). The results were expressed
110 as percentage of citric acid (% citric acid).

111
$$\%TA = \frac{\text{titre value} \times \text{vol. made up} \times \text{Eq wt of citric acid (64)} \times 100}{\text{sample wt} \times \text{vol sample used} \times 1000}$$

112

113 2.4 Betacyanin content

114 The betacyanin content and soluble phenolics content of dragonfruit was measured to assess
115 their contribution to the bioactive content of the fruit. Betacyanin content was determined
116 using the spectrophotometric method (Nerd et al., 1999) where fruit pulp was homogenised
117 with 80 % methanol (v/v) and extract was filtered with Whatman no. 1 paper. The extract was
118 mixed with 0.1 M acetate buffer and absorbance measured at 538 nm.

119

120 2.5 Soluble phenolics content

121 Soluble phenolics content was assessed using Folin Ciocalteu spectrophotometric method
122 (Singleton and Rossi, 1965). Fruit pulp was subjected to extraction with 60 % methanol (v/v),
123 at a ration of solute to solvent of 1:10. The extract was reacted with undiluted Folin Ciocalteu
124 reagent (Sigma-Aldrich, USA) at 35 °C for 2 h. Absorbance readings were then measured at
125 765 nm, and calibrated against a standard curve constructed using freshly prepared gallic acid
126 solution (0 - 1000 µM).

127

128 2.6 Antioxidant activity

129 Total antioxidant activity was determined using the fruit pulp extracted with 60 % methanol
130 (Section 2.5). Two methods were adopted for measuring antioxidant activity: Ferric Reducing
131 Antioxidant Power (FRAP) assay (Benzie and Strain, 1996) and 2,2-diphenyl-1-
132 picrylhydrazyl (DPPH) adapted from the method of Brand-Williams and colleagues
133 (Molyneux, 2004).

134 Fruit extract was reacted with freshly prepared FRAP reagent and the absorbance was
135 measured at 593 nm after 4 min of incubation at 37 °C, with the FRAP reagent reacted with

136 60 % methanol as a blank. Results were expressed as mM kg⁻¹ Trolox equivalents on a fresh
137 weight basis based on a standard curve prepared beforehand using Trolox (0 – 1000 µM).

138 For DPPH analysis, fruit extract was mixed with 0.1 M Tris-HCl and 0.3 mM DPPH
139 dissolved in methanol and allowed to react for 20 min at room temperature under dark
140 conditions. Absorbance readings were recorded at 517 nm with a methanol only blank for
141 baseline correction and methanol in DPPH as a control. Results were expressed as mM kg⁻¹
142 Trolox equivalents on a fresh weight basis.

143

144 2.7 Statistical analysis

145 The experiment followed a completely randomized design (CRD) with four replicates per
146 treatment, **consisting of three fruits per replicate**. Analysis of Variance (ANOVA) was used
147 to measure the treatment effect using SAS® 9.1 (SAS Institute Inc., USA), and means were
148 separated using Duncan's Multiple Range Test (DMRT) test at ($P < 0.05$).

149

150 3. Results

151 3.1 Biochemical quality of fruits

152 SSC and TA were assessed as indicators of the metabolic activity and ripening stage of the
153 fruit. A significant decline ($P < 0.05$) in SSC (Figure 1a) was observed during the storage
154 period. Treatment with 0.5 mM MJ maintained the highest SSC values by the final day of
155 storage. Similarly, TA (Figure 2a) declined during the storage period but at a slower rate, and
156 significantly lower ($P < 0.05$) values were observed for the fruits treated with 0.1 mM MJ. By
157 the final day of storage, all the treatments were at par ($P > 0.05$) with the exception of the
158 0.01 mM MJ treatment, which exhibited the lowest TA values.

159 The decline in SSC (Figure 1b) was consistent with the observed ripening pattern for
160 dragonfruit (Figure 1a), however, treatment with 0.1 mM and 1 mM SA maintained

161 significantly higher ($P < 0.05$) SSC levels throughout the storage period. An initial decline in
162 TA (Figure 4b) ($P < 0.05$) was observed for all treatments including the control, during the
163 storage period. However, TA values of fruits treated with 1 mM SA were significantly higher
164 ($P < 0.05$) on the final day of storage.

165

166 3.2 Bioactive content of fruits

167 A gradual increase ($P < 0.05$) in betacyanin content was observed during the storage period
168 (Figure 3a). Fruits treated with 0.5 mM MJ exhibited a peak in betacyanin content by the 14th
169 day of storage, however, the lower doses of MJ (0.01 and 0.1 mM) exhibited the highest
170 levels of betacyanin at the end of the storage period. Meanwhile, no significant effect ($P >$
171 0.05) of treatment was observed for the soluble phenolics content of the fruits (Figure 4a).
172 Although a decline was observed throughout the storage period, the treatments were at par (P
173 > 0.05).

174 Similarly, an increase ($P < 0.05$) in betacyanin content was observed for the SA treatment
175 (Figure 3b). Moreover, the control maintained higher betacyanin levels during the first two
176 weeks of storage. Lower betacyanin levels were observed for fruits treated with the higher
177 doses of SA (2 and 5 mM) during the first 2 weeks of storage, however, betacyanin levels
178 peaked for these two treatments by the final day of storage. Moreover, soluble phenolic
179 content gradually declined during storage (Figure 4b), with lower values reported for the
180 fruits treated with 2 mM SA.

181

182 3.3 Antioxidant activity of fruits

183 A significant effect ($P < 0.05$) of MJ treatment was observed for FRAP antioxidant activity
184 (Figure 5a) throughout the storage period. The antioxidant activity peaked on the 14th day of
185 storage, with the highest peak observed for the higher concentrations of MJ. A 4.2 fold

186 increase in the initial antioxidant levels were observed for fruits treated with 0.5 mM MJ. By
187 the final day of storage, all the treatments were at par ($P > 0.05$) except for 0.5 mM MJ which
188 exhibited significantly lower levels. DPPH radical scavenging activity (Figure 6a) increased
189 linearly throughout the storage period. The highest level of activity was observed for the
190 control, with the MJ treated fruits displaying significantly lower ($P < 0.05$) activity.

191 The control (Figure 5b) maintained the highest antioxidant activity during the initial
192 storage period. However, towards the end of the storage period, treatment with SA resulted in
193 significant ($P < 0.05$) increases in the antioxidant activity. The increase in FRAP was
194 positively correlated to the dose of SA applied. This was different to the trend observed for
195 DPPH radical scavenging activity (Figure 6b). The highest activity was reported for the
196 untreated control throughout the storage period, with lower levels of activity observed as the
197 SA dose increased. By the final day of storage, the radical scavenging activity was par ($P >$
198 0.05) for all the treatments.

199

200 **4. Discussion**

201 The fruits used in this experiment were harvested at maturity, thus at a more advanced
202 ripening stage, which is a common commercial practice for dragonfruit. Gradual decline in
203 the chemical composition of dragonfruit were observed in the experiments. The decline in
204 SSC is attributed to the hydrolysis of insoluble polysaccharides into sugars (Vyas et al.,
205 2015). Organic acids can also be utilized as energy reserves during respiration, accounting for
206 the decline in TA (Valero and Serrano, 2010). SSC and TA both serve as a measure of the
207 changes in the energy reserves that fuel metabolic activities of the plant (Wills et al., 2007).
208 Thus, the gradual decline of SSC and TA can be regarded as an indicator of the metabolic
209 status of the fruit, representing progressive ripening of the fruit. High respiration of the fruit

210 is linked to a relatively short shelf life; thus, the fruits were assessed for only three weeks of
211 storage.

212 MJ promotes both ripening and defence mechanisms of plants through the regulation
213 of gene expression. A positive feedback on the biosynthesis pathway of jasmonic acid has
214 also been reported upon exposure to MJ (Ziosi et al., 2008). However, the results of this
215 experiment indicate that MJ did not influence ripening of dragonfruit. On the other hand, SA
216 treatment did influence ripening, as the higher doses increased the metabolic activity of the
217 fruit, while lower doses decreased the metabolic activity. Reduced metabolism rate and
218 delayed fruit ripening upon exposure to SA has been previously reported in various fruits
219 (Ashgari and Aghdam, 2010). SA has been successfully demonstrated to enhance chilling
220 tolerance in various fruits, such as sponge gourd (Han et al., 2017), plum (Luo et al., 2011)
221 and pomegranate (Sayyari et al., 2011). This protective activity of SA has been primarily
222 associated with enhanced antioxidant activity, increased heat shock proteins and enhanced
223 membrane integrity (Han et al., 2017). However, the protective activity of SA during cold
224 storage has also been attributed to decreasing levels of total phenolic content and antioxidant
225 enzymes in fruits during cold storage (Han et al., 2017; Luo et al., 2011; Sayyari et al., 2011).

226 Previous studies have accounted an increase in antioxidant activity of various fruits
227 upon subjection to MJ (Szymanowska et al., 2015; Wang et al., 2007). Enhanced antioxidant
228 activity, as reported in grapes (Flores et al., 2015) and pomegranates (Sayyari et al., 2011),
229 was associated with enhanced phenolic content and anthocyanin content. While hormonal
230 treatment did not induce a significant effect ($P > 0.05$) on the soluble phenolics content of the
231 fruit, betacyanin content and antioxidant activity was promoted in fruits exposed to MJ.
232 Betacyanin, being the principal bioactive compound in dragonfruit (Muhammad et al., 2014),
233 could have increased during ripening as a defense response towards the induced stresses
234 (Zhang et al., 2003). Fan et al., (2016) reported enhanced antioxidant activity in eggplant,

235 which was related to delayed senescence of the fruit. Higher doses of MJ are reported to have
236 a more pronounced impact on antioxidant activity (Kim et al., 2006), and this was similarly
237 observed for FRAP activity of the dragonfruit in this study.

238 While MJ enhanced betacyanin content and antioxidant activity, SA treatment at all
239 doses did not enhance betacyanin content or soluble phenolic content. However, SA
240 treatment did result in an increase in antioxidant activity during the final days of storage.
241 Clearly, another factor is at play in extending the storage period of the fruits treated with
242 lower doses of SA. Studies have related application of SA to increased activity of antioxidant
243 enzymes, such as catalase and superoxide dismutase (Dokhanieh and Aghdam, 2016; Asghari
244 and Aghdam, 2010), which could enhance the antioxidant capacity of the fruit. Nie et al.,
245 (2015) reported increased activity of catalase gene expression during cold stress of
246 dragonfruit explants, demonstrating a clear role of this enzyme in cold tolerance. This could
247 be explored further by examining the response of this enzyme in the mature fruit to SA and
248 MJ in combination with the cold stress. Moreover, Asghari and Hasanlooe (2015) reported
249 the importance of positive crosstalk between SA and MJ in enhancing antioxidant activity
250 and natural resistance in strawberry. The interaction of these two stress hormones might also
251 prove beneficial for treating dragonfruit during cold storage.

252

253 **5. Conclusion**

254 An increasing interest in nutritional composition of fruits has been witnessed among
255 consumers lately, however the aesthetic appearance of fruits remains an important parameter
256 for consumer perception of fruit quality. Progressive changes in the physical parameters
257 occur during ripening, accompanied with changes in the chemical and bioactive content
258 affecting both the nutritional and aesthetic properties of the fruit. However, these changes do
259 not essentially follow the same pattern and are largely influenced by external factors such as

260 cold storage and stress hormones. The effect of the stress hormones SA and MJ on
261 dragonfruit in combination with cold storage have not been previously explored, and these
262 results illustrate the ongoing changes in the fruit metabolic activity and redox homeostasis
263 during cold storage. While lower doses of SA exhibited a role in delaying ripening, it did not
264 enhance the bioactivity of the fruit. Meanwhile, MJ did not have a significant effect on
265 ripening, yet resulted in enhanced betacyanin and antioxidant activity. Future studies could
266 explore the effect of the interaction of these two hormones on dragonfruit.

267

268 **Acknowledgement**

269 This work was supported by the University of Nottingham Doctoral Training Grant (DTG)

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285 **References**

- 286 Ashgari, M., Aghdam, M.S., 2010. Impact of salicylic acid on post-harvest physiology of horticultural
287 crops. Trends Food Sci. Technol. 21, 502 – 509.
- 288 Ashgari, M., Hasanlooe, A.R., 2015. Interaction effects of salicylic acid and methyl jasmonate on
289 total antioxidant content, catalase and peroxidase enzymes activity in “Sabrosa” strawberry fruit during
290 storage. Sci. Hort. 197, 490 – 495.
- 291 Benzie, I.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of
292 “antioxidant power”: the FRAP assay. Anal. Biochem. 239, 70–6.
- 293 Chen, J.-Y., Wen, P.-F., Kong, W.-F., Pan, Q.-H., Zhan, J.-C., Li, J.-M., Wan, S.-B., Huang, W.-D.,
294 2006. Effect of salicylic acid on phenylpropanoids and phenylalanine ammonia-lyase in harvested
295 grape berries. Postharvest Biol. Tec. 40, 64–72.
- 296 Dembitsky, V.M., Poovarodom, S., Leontowicz, H., Leontowicz, M., Veerasilp, S., Trakhtenberg, S.,
297 Gorinstein, S., 2011. The multiple nutrition properties of some exotic fruits: Biological activity and
298 active metabolites. Food Res. Int. 44, 1671–1701.
- 299 Dokhanieh, A.Y., Aghdam, M.S., 2016. Postharvest browning alleviation of *Agaricus bisporus* using
300 salicylic acid treatment. Sci. Hort. 207, 146 – 151.
- 301 Fan, L., Shi, J., Zuo, J., Gao, L., Lv, J., Wang, Q., 2016. Methyl jasmonate delays postharvest
302 ripening and senescence in the non-climacteric eggplant (*Solanum melongena* L.) fruit. Postharvest
303 Biol. Tec. 120, 76 – 83.
- 304 Flores, G., Blanch, G.P., del Castillo, M.L.R., 2015. Postharvest treatment with (-) and (+) methyl
305 jasmonate stimulates anthocyanin accumulation in grapes. LWT Food Sci. Technol. 62, 807 – 812.
- 306 Gonzalez-Aguilar, G.A., Villa-Rodriguez, J.A., Ayala-Zavala, J.F., Yahia, E.M., 2010. Improvement
307 of the antioxidant status of tropical fruits as a secondary response to some postharvest treatments.
308 Trends Food Sci. Tech. 21, 475–482.
- 309 González-Aguilar, G.A., Tiznado-Hernandez, M.E., Zavaleta-Gatica, R., Martinez-Tellez, M.A.,
310 2004. Methyl jasmonate treatments reduce chilling injury and activate the defense response of guava
311 fruits. Biochem. Biophys. Res. Commun. 313, 694–701.

- 312 Han, C., Zuo, J.-H., Wang, Q., Dong, H.-Z., Gao, L.-P., 2017. Salicylic acid alleviates postharvest
313 chilling injury of sponge gourd (*Luffa cylindrica*). J. Integr. Agr. 16, 735 – 741.
- 314 Heredia, J.B., Cisneros-Zevallos, L., 2009. The effect of exogenous ethylene and methyl jasmonate on
315 pal activity, phenolic profiles and antioxidant capacity of carrots (*Daucus carota*) under different
316 wounding intensities. Postharvest Biol. Tec. 51, 242–249.
- 317 Hoa, T.T., Clark, C.J., Waddell, B.C., Woolf, A.B., 2006. Postharvest quality of Dragon fruit
318 (*Hylocereus undatus*) following disinfesting hot air treatments. Postharvest Biol. Tec. 41, 62–69.
- 319 Kim, H.J., Chen, F., Wang, X., Rajapakse, N.C., 2006. Effect of methyl jasmonate on secondary
320 metabolites of sweet basil (*Ocimum basilicum* L.). J. Agric. Food Chem. 54, 2327–32.
- 321 Luo, Z., Chen, C., Xie, J., 2011. Effect of salicylic acid treatment on alleviating postharvest chilling
322 injury of “Qingnai” plum fruit. Postharvest Biol. Tec. 62, 115–120.
- 323 Macias-Ceja, D.C., Cosin-Roger, J., Ortiz-Masia, D., Salvador, P., Hernandez, C., Calatayud, S.,
324 2016. The flesh ethanolic extract of *Hylocereus polyrhizus* exerts anti-inflammatory effects and
325 prevents murine colitis. Clin. Nutr. 35, 1333 – 1339.
- 326 Marnewick, J.L., Rautenbach, F., Venter, I., Neethling, H., Blackhurst, D.M., Wolmarans, P., 2011.
327 Effects of rooibos (*Aspalathus linearis*) on oxidative stress and biochemical parameters in adults at
328 risk for cardiovascular disease. J. Ethnopharmacol. 133, 46 - 52.
- 329 Molyneux, P., 2004. The use of the stable free radical diphenylpicryl- hydrazyl (DPPH) for
330 estimating antioxidant activity. Songklanakarin J. Sci. Technol. 26, 211 – 219.
- 331 Muhammad, K., Mohd Zahari, N.I., Gannasin, S.P., Mohd Adzahan, N., Bakar, J., 2014. High
332 methoxyl pectin from dragon fruit (*Hylocereus polyrhizus*) peel. Food Hydrocolloid. 42, 289–297.
- 333 Mustafa, M.A., Ali, A., Seymour, G., Tucker, G., 2016. Enhancing the antioxidant content of
334 carambola (*Averrhoa carambola*) during cold storage and methyl jasmonate treatments. Postharvest
335 Biol. Tec. 118, 79 - 86.
- 336 Nerd, A., Gutman, F., Mizrahi, Y., 1999. Ripening and postharvest behaviour of fruits of two
337 *Hylocereus* species (Cactaceae). Postharvest Biol. Tec. 17, 39–45.

- 338 Nie, Q., Gao, G.-L., Fan, Q.-J., Qiao, G., Wen, X.-P., Liu, T., Peng, Z.J., Cai, Y.Q., 2015. Isolation
339 and characterisation of a catalase gene “HuCAT3” from pitaya (*Hylocereus undatus*) and its
340 expression under abiotic stress. *Gene* 563, 63 – 71.
- 341 Ranganna, S., 1977. *Handbook of Analysis and Quality Control for Fruit and Vegetable Products*,
342 Tata McGraw Hill: New Delhi.
- 343 Rautiainen, S., Levitan, E.B., Orsini, N., Akesson, A., Morgenstern, R., Mittleman, M.A., Wolk, A.,
344 2012. Total antioxidant capacity from diet and risk of myocardial infarction: A prospective cohort of
345 women. *Am. J. Med.* 125, 874 - 980.
- 346 Sayyari, M., Babalar, M., Kalantari, S., Martinez-Romero, D., Gullen, F., Serrano, M., Valero, D.,
347 2011. Vapour treatments with methyl salicylate or methyl jasmonate alleviated chilling injury and
348 enhanced antioxidant potential during postharvest storage of pomegranates. *Food Chem.* 124, 964 –
349 970.
- 350 Singleton, V.L., Rossi, J.A., 1965. Colorimetry of Total Phenolics with Phosphomolybdic-
351 Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* 16, 144–158.
- 352 Szymanowska, U., Zlotek, U., Karas, M., Baraniak, B., 2015. Anti-inflammatory and antioxidative
353 activity of anthocyanins from purple basil leaves induced by selected abiotic elicitors. *Food Chem.*
354 172, 71–7.
- 355 Valero, D., Serrano, M., 2010. *Postharvest Biology and Technology for Preserving Fruit Quality*,
356 Boca Raton: CRC Press-Taylor & Francis.
- 357 Vyas, P.B., Ramana Rao, T.V., Thakkar, V.R., 2015. Combined effects of chemical and physical
358 elicitors on postharvest quality of custard apple (*Annona squamosa* L., cv. Balanagar). *Sci. Hort.* 187,
359 50 - 57.
- 360 Wang, S., Bowman, L., Ding, M., 2007. Methyl jasmonate enhances antioxidant activity and
361 flavonoid content in blackberries (*Rubus* sp.) and promotes antiproliferation of human cancer cells.
362 *Food Chem.* 107, 1261–1269.
- 363 Wang, Y., Frei, M., 2011. Stressed food – The impact of abiotic environmental stresses on crop
364 quality. *Agric. Ecosyst. Environ.* 141, 271–286.
- 365 Wanitchang, J., Terdwongworakul, A., Wanitchang, P., Noypitak, S., 2010. Maturity sorting index of
366 dragon fruit: *Hylocereus polyrhizus*. *J. Food Eng.* 100, 409–416

- 367 Wills, R.B.H., McGlasson, B., Graham, D., Joyce, D., 2007. Postharvest- An introduction to the
368 physiology and handling of fruit, vegetables and ornamentals. Fifth edit., CAB International.
- 369 Wu, L.-C., Hsu, H.-W., Chen, Y.-C., Chiu, C.-C., Lin, Y.-Y., Ho, J.-A., 2006. Antioxidant and
370 antiproliferative activities of red pitaya. *Food Chem.* 95, 319 – 327.
- 371 Zhang, Y., Chen, K., Zhang, S., Ferguson, I., 2003. The role of salicylic acid in postharvest ripening
372 of kiwifruit. *Postharvest Biol. Tec.* 28, 67–74.
- 373 Ziosi, V., Bonghi, C., Begoli, A.M., Trainotti, L., Biondi, S., Sutthiwal, S., Kondo, S., Costa, G.,
374 Torrigiani, P., 2008. Jasmonate-induced transcriptional changes suggest a negative interference with
375 the ripening syndrome in peach fruit. *J. Exp. Bot.* 59, 563–73.
- 376
- 377
378
379
380
381
382