

# ATMOSPHERIC TIDAL OSCILLATIONS

## PART 2. DIURNAL VARIATION OF PRESSURE OVER INDIA\*

R. ANANTHAKRISHNAN, S. S. ARALIKATTI and J. A. MALIEKAL

*Indian Institute of Tropical Meteorology, Pune 411005, India.*

### ABSTRACT

Hourly surface pressure data (annual and monthly) over a network of Indian stations have been harmonically analysed. The spatial and temporal variation of the amplitude, phase and percentage contribution towards variance of the time series in respect of the first four harmonics of the barometric oscillation are discussed.

### INTRODUCTION

THE discovery of atmospheric tidal oscillations in the seventeenth century and the historical developments leading to the present thermo-tidal theory of these oscillations were dealt within Part 1 of this paper. Here we propose to give an account of the salient observational features of the barometric oscillation over India. The diurnal variation of many meteorological elements including pressure at Indian stations have been discussed in a series of publications towards the end of the last century by Blanford<sup>1</sup> and Eliot<sup>2</sup>. A global study of the semi-diurnal pressure oscillation utilising the data of over 200 stations was made by Simpson<sup>3</sup> while working in the India Meteorological Department at Simla. Alvi and Jagannathan<sup>4</sup> studied the barometric oscillation at 38 Indian stations. We have recently examined the diurnal variation of pressure at 62 Indian stations using the hourly pressure data for periods ranging from 5 to 12 years at individual stations. Figure 1 gives the location of the stations.

### HARMONIC ANALYSIS OF HOURLY PRESSURE

The annual and monthly mean values of the hourly pressure data at the stations were subjected to harmonic analysis to determine the amplitude, phase and the percentage contri-

bution to the variance of the time series in respect of the first four harmonics denoted (following the usual convention) by  $S_1(p)$ ,  $S_2(p)$ ,  $S_3(p)$  and  $S_4(p)$ . As example, the results of the analysis for six selected stations—Trivandrum, Kodaikanal, Madras, Bombay, Nagpur and Jodhpur—are shown in figures 2(a), (b), (c). In general  $S_1(p)$  and  $S_2(p)$ , whose amplitudes are comparable, account for most of the variance of the pressure time series. The residual variance is practically fully accounted for by  $S_3(p)$  and  $S_4(p)$  whose amplitudes are an order of magnitude less than those of the first two harmonics. For each station, the figure gives the monthly variation of the amplitude, variance and phase. The term phase is here used to denote the time at which the first maximum of the oscillation occurs, expressed in local mean solar time (LMT) reckoned from zero at local midnight. The annual values of the parameters are indicated by thick dots for  $S_1(p)$  and  $S_3(p)$  and by open circles for  $S_2(p)$  and  $S_4(p)$ .

### ANNUAL VARIATIONS OF THE HARMONIC COMPONENTS

*24-hourly oscillation,  $S_1(p)$ :* The amplitude and phase of  $S_1(p)$  show large spatial variations. The amplitudes range from 0.5 to 1.0 mb at coastal stations and from 1.0 to 1.5 mb at inland stations. Large amplitudes are found over north and central India. A diagram showing isopleths of amplitude over the globe for annual  $S_1(p)$  by Haurwitz and Cowley<sup>5</sup> indicates the maximum amplitude over India as 1.04 mb which is an underestimate. The present study reveals that the

\* Part 1 of this article was appeared in September 20, 1984 issue of *Curr. Sci.*






Figure 2(a). Monthly progression of the amplitudes, phases and variance contribution of the harmonic components: Trivandrum and Kodaikanal.






Figure 2(b). Monthly progression of the amplitudes, phases and variance contribution of the harmonic components: Madras and Bombay.




Figure 2(c). Monthly progression of the amplitudes, phases and variance contribution of the harmonic components: Nagpur and Jodhpur.

phase is found to be between 0250 and 0450 hr at individual stations. Its contribution to variance is of the order of 0.2% or less.

Figures 3 and 4 give a pictorial representation of  $S_1(p)/S_2(p)$  and  $S_3(p)/S_4(p)$  at selected stations. The harmonic components are depicted as vectors originating from the stations, the length of the vector being proportional to the amplitude. The direction of the vector gives the phase of the oscillation as per the harmonic dials shown on the diagrams.

**SEASONAL VARIATIONS OF THE HARMONIC COMPONENTS**

*The 24-hourly oscillation,  $S_1(p)$ :* The diurnal oscillation attains maximum amplitudes in the hot weather months March to May, the amplitudes ranging from 1.0 to 2.0 mb, with high

values at the north and central Indian stations. The minimum amplitudes from 0.2 to 1.0 mb occur mostly in the southwest monsoon months of July—August. Very low amplitudes are noticed at the west coast stations in July. The ratio of the minimum to the maximum amplitude of  $S_1(p)$  at individual stations varies from 0.2 to 0.7 with a mean value of about 0.45. The phase of the oscillation is earliest in the monsoon months (mean value for July 6.84 hr LMT) and latest in the winter months (mean value for February 7.89 hr LMT). The variance accounted for by  $S_1(p)$  ranges from a maximum of 40% to 70% in the hot weather months and a minimum of 5% to 30% in the monsoon/winter months.

*The 12-hourly oscillation,  $S_2(p)$ :* The semi-diurnal oscillation shows maximum amplitude of 1.0 to 1.7 mb in February—March and minimum




Figure 3. Vectorial representation of  $S_1(p)$  and  $S_2(p)$  at selected stations.




Figure 4. Vectorial representation of  $S_3(p)$  and  $S_4(p)$  at selected stations.


amplitude of 0.6 to 1.2 mb in June—July. The amplitudes decrease with increasing latitude generally following  $\cos^3 \phi$ . The ratio of the minimum to the maximum amplitude at individual stations varies from 0.6 to 0.9 with a mean value of about 0.73. Thus the seasonal variation of the amplitude of  $S_2(p)$  is less than that of  $S_1(p)$ . The phase of  $S_2(p)$  is earliest in November (mean value for all stations 9.78 hr LMT) and latest in July (mean value 10.43 hr LMT). It may be noted that the seasonal variations of the phases of  $S_2(p)$  and  $S_1(p)$  show roughly opposite behaviour. The percentage variance accounted for by  $S_2(p)$  has a seasonal variation opposite to that of  $S_1(p)$ ; the minimum values are found in the months April—May—June and the maximum values generally in July—August. It may be noted that since  $S_1(p)$  and  $S_2(p)$  together make up practically the entire variance of the time series, the variance curves

corresponding to these oscillations in figure 2 appear nearly as mirror images of each other about the horizontal line corresponding to 50% variance.

The monthly progression of the ratio of the amplitudes  $S_1(p)/S_2(p)$  for 16 stations is shown in figure 5. The low values of the ratio at stations along the west coast during the southwest monsoon months may be noted. At most of the interior stations the ratio exceeds unity in the hot weather months. The thick dots in this diagram indicate the value of the ratio in respect of the annual data.

*The 8-hourly oscillation,  $S_3(p)$ :* The third harmonic shows interesting seasonal variations in amplitude and phase. The amplitude has two maxima, a primary maximum in December—January and a secondary maximum in June—





**Figure 5.** Monthly variation of the ratio of the amplitudes of the diurnal and semi-diurnal components of the barometric oscillation at selected coastal and interior stations.

July. Pronounced minima are found in April—May and in August—September. There is a reversal of phase of the oscillation from about 0205 hr during the epoch of the primary maximum in winter to 0605 hr during the secondary maximum in summer. During the winter months  $S_3(p)$  accounts for about 0.5 to 1.5% of the variance at the southern stations and 2 to 6% of the variance at the north Indian stations.

*The 6-hourly oscillation,  $S_4(p)$ :* The amplitude of this oscillation is about half that of  $S_3(p)$ .

Similar to the third harmonic, the fourth harmonic is also most pronounced in the winter months with amplitudes increasing from south to north. However, it does not show a secondary maximum nor does it show a reversal of phase from winter to summer. These features can be appreciated from figure 2. The phase of the oscillation varies between 2.7 hr and 3.7 hr LMT. It accounts for about 0.2% of the variance at the southern stations and 0.5 to 1% of the variance at the northern stations during the winter months.







Figure 6. Monthly variation of the amplitudes and phases of  $S_1(p)$  and  $S_2(p)$  in five-degree zonal belts.

Figure 7. Monthly variation of the amplitudes and phases of  $S_3(p)$  and  $S_4(p)$  in five-degree zonal belts.

## ZONALLY AVERAGED PRESENTATION OF THE HARMONIC COMPONENTS

A graphical presentation of the monthly progression of the amplitudes and phases of the four harmonic components of the atmospheric tidal pressure oscillation over India averaged for stations in five degree zonal belts ( $05^{\circ}$ – $10^{\circ}$ ), . . . ( $30^{\circ}$ – $35^{\circ}$ ) N is given in figures 6(a), (b) and 7(a), (b). The curves A, B, . . . F in these diagrams refer to the six zonal belts. The broken curve G represents the average for all the 62 stations. The maximum/minimum values of the amplitudes and phases are indicated on the respective curves which summarise the major features discussed in the preceding sections.

## ACKNOWLEDGEMENTS

One of us (RA) is grateful to the Indian Space Research Organisation (ISRO) for a research grant.

20 June 1984

1. Blanford, H. F., *Indian Met. Mem.*, 1876–1895, Vols. 1, 5, 9.
2. Eliot, J., *Indian Met. Mem.*, 1895–1899, Vols. 5, 9, 10.
3. Simpson, G. C., *Q. J. R. Met. Soc.*, 1918, 44, 1.
4. Alvi, S. M. A. and Jagannathan, P., *Indian Met. Mem.*, 1972, 32, Part I.
5. Haurwitz, B. and Cowley, A. D., *Pure Appl. Geophys.*, 1973, 102, 193.

---

## NEWS

---

### VEB Carl Zeiss JENA

CURRENT SCIENCE wishes to greet JENA<sup>er</sup> GLAS during its centenary year 1984.

Leipzig, the famous metropolis of international trade, once again demonstrated the world's progress in science and technology and thus indicates possibilities for dynamic economic growth. The display of products of VEB Carl Zeiss JENA in Leipzig is an example of continuous advancement even under the untoward conditions of the world's economic situation. More than 75% of the products range has been replaced by new products in the last five years, which is in keeping with the present pace of world-wide innovation. The 1984 exhibits include more than 30 new or improved items. Highly efficient microelectronic systems have further increased the speed of instrument processes and added to new optical and mechanical solutions, upgraded the quality, precision and performance capabilities of the products on display.

Following are some of the products of Kombinat VEB Carl Zeiss JENA at Spring 1984 Leipzig Fair:

(1) DZT 90 × 120 RGS Computer-Assisted Stereoplotting System, (2) ABBE 01-200 C Vertical Metroscope, (3) AE-80 Digital Meter Unit, (4) IKF-

100 Incremental Displacement Pickup, (5) IDL-01 Length Measuring System with AE-100 Meter Unit, (6) IGR-M2 Incremental Rotary Pulse Transmitter with analogue output, (7) Computer Input Adapter for Incremental Displacement Pickups, (8) 3D Telemetric Probe, (9) MS-4 Size Control Device /SG-112 Controller, (10) Laser Coagulator, (11) OAP-310/M Eye Examination Setup, (12) SPECORD-M80 and SPECORD-M85, (13) SPEKOL-11 Spectrophotometer, (14) Mercury Hydride Generation System for the AAS-3 Atomic Absorption Spectrophotometer, (15) TZ-6 Texture Attachment to the HZG-4 Universal X-ray Diffractometer, (16) JENAVERT Incident Light Microscopes, (17) GSM Stereomicroscope (18) 1m Reflector Telescope, (19) COSMORAMA Large-Size Planetarium (20) Improved 50 mm f/1.4 PRAKTICAR-MC Lens (21) SF-11/1535 Optical Glass, (22) Component blanks for multifocal ophthalmic lenses (23) Distilling Plant for Corrosive Substances.

Further details may be had from: Dr. H. C. Wolfgang Biermann, Director General, Kombinat VEB Carl Zeiss JENA-DDR, DDR -6900 Jena, Carl-Zeiss-Strasse 1.