



**Michigan
Technological
University**

Michigan Technological University
Digital Commons @ Michigan Tech

Michigan Tech Patents

Vice President for Research Office

3-9-2010

Isolated cellulose synthase promoter regions

Chandrashekhar P. Joshi

Michigan Technological University, cpjoshi@mtu.edu

Follow this and additional works at: <https://digitalcommons.mtu.edu/patents>


 Part of the [Biological Engineering Commons](#)

Recommended Citation

Joshi, Chandrashekhar P., "Isolated cellulose synthase promoter regions" (2010). *Michigan Tech Patents*. 111.

<https://digitalcommons.mtu.edu/patents/111>

Follow this and additional works at: <https://digitalcommons.mtu.edu/patents>

 Part of the [Biological Engineering Commons](#)



(12) **United States Patent**
Joshi

(10) **Patent No.:** **US 7,674,951 B1**
(45) **Date of Patent:** ***Mar. 9, 2010**

(54) **ISOLATED CELLULOSE SYNTHASE PROMOTER REGIONS**

WO WO 2007/019245 2/2007

(75) Inventor: **Chandrashekar P. Joshi**, Houghton, MI (US)

(73) Assignee: **Michigan Technological University**, Houghton, MI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 275 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **11/760,877**

(22) Filed: **Jun. 11, 2007**

Related U.S. Application Data

(60) Continuation-in-part of application No. 11/437,368, filed on May 19, 2006, now Pat. No. 7,232,941, which is a division of application No. 09/980,043, filed as application No. PCT/US00/13637 on May 18, 2000, now Pat. No. 7,049,481.

(60) Provisional application No. 60/135,280, filed on May 21, 1999.

(51) **Int. Cl.**
C12N 15/09 (2006.01)
C12N 15/82 (2006.01)
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **800/287; 800/290; 800/298; 800/295; 800/278; 435/468**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,268,274	A	12/1993	Ben-Bassat et al.	
5,633,439	A	5/1997	Walter et al.	
5,646,023	A	7/1997	Secor et al.	
5,654,190	A	8/1997	Matsunaga et al.	
6,013,860	A	1/2000	Himmel et al.	
6,271,443	B1	8/2001	Stalker et al.	
6,303,847	B1	10/2001	Kawaoka et al.	
6,867,352	B2	3/2005	Allen	
7,049,481	B1	5/2006	Chiang et al.	
7,232,941	B2*	6/2007	Chiang et al.	800/287
7,288,409	B1	10/2007	Chiang et al.	
2005/0108791	A1	5/2005	Edgerton	
2008/0222752	A1	9/2008	Joshi	

FOREIGN PATENT DOCUMENTS

CA	2012311	9/1990	
EP	875575	11/1998	
WO	WO 98/00549	* 1/1998	
WO	WO 98/18949	* 5/1998	
WO	WO 00/22092	4/2000	
WO	WO 00/71670	11/2000	

OTHER PUBLICATIONS

U.S. Appl. No. 11/437,368, filed May 19, 2006.*
U.S. Appl. No. 09/980,043, filed Apr. 5, 2002.*
Accession No. T10797—Sequence Search Result (1996) 1-2.
Accession No. AF072131 [gi: 4115904], pp. 1-2, Sequence Search Result, Jan. 8, 1999.
Accession No. AF072131 [gi: 3243277], pp. 1, Sequence Search Result, Jun. 23, 1998.
Accession No. AF062485 [gi: 3135610], pp. 1-3, Sequence Search Result, Aug. 8, 1998.
Altmorbe et al., *Mol. Plant-Microbe Interac.* 2:301, (1989).
Animal Cell Culture: A Practical Approach, R.I. Freshney, ed., "Immobilised Cells and Enzymes," (1986), IRL Press.
Arioli et al., *Science*, 279: 717-720, (1998).
Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).
Benton and Davis, *Science* 196:180, (1977).
Bird et al., "Manipulation of Plant Gene expression by Antisense RNA," *Biotechnology & Genetic Engineering Reviews* (1991) 9:207-227.
Brown et al., *Trends Pl Sci.*, 1: 149-156 (1996).
Bugos et al., *Biotechniques* 19:734-737, (1995).
Database EMBL Online, Jun. 23, 1998, Wu, L., Joshi, CP, Chiang, VL, "Populus tremuloides secondary xylem cellulose synthase (ce1A) mRNA, complete cds." EBI Database accession No. af072131, abstract.
Database SWALL Online, May 1, 1997, "Cellulose synthase, CELA1", accession No. P93155.
Database SWALL Online, Nov. 1, 1998, Secondary xylem cellulose synthase (ce1A) mRNA, complete cds. EBI Database accession No. 081368, abstract.
Delmer and Amor, *Pl Cell*, 7: 987-1000, (1995).
DNA Cloning: A Practical Approach, vols. I and II (D.N. Glover ed. 1985).
Esau, K., *Anatomy of Seed Plants*, New York: John Wiley and Sons. (1960).
Esau, K., *Plant Anatomy*, 2nd ed. New York: Wiley (1953).
Esau, K., *Vascular Differentiation in Plants*, New York: Hold, Rinehart & Winston (1965).
Esau, K., et al., "Observations on Cytokinesis", *Planta* (Berl.) 67, 168-181 (1965).
Esau, K., "Anatomy of Plant Virus Infections", *Annu. Rev. Phytopathol.* 5:45-76 (1967).
Fukuda, *Ann Rev Pl Physiol Pl Mol Biol*, 47: 299-325, (1996).
Fullner and Nester, *J. Bacteriol.* 178: 1498, (1996).
Fullner et al., *Science* 273: 1107, (1996).
Ge and Chiang, 1996, *Pl Physiol*, 112: 861, (1996).

(Continued)

Primary Examiner—Medina A Ibrahim
(74) *Attorney, Agent, or Firm*—Michael Best & Friedrich LLP

(57) **ABSTRACT**

The invention relates to methods of inducing expression of coding sequences including cellulose synthase coding sequences in transgenic plants using promoters of cellulose synthase genes from *Populus* plant species and transgenic plants produced by the methods.

OTHER PUBLICATIONS

- Grunstein and Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961, (1975).
- Haigler and Blanton, PNAS, 93: 12082-12085, (1996).
- Higuchi, "Using PCR to Engineer DNA", in *PCR Technology: Principles and Applications for DNA Amplification*, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70. (1989).
- Higuchi, *Biochemistry and Molecular Biology of Wood*, Springer Verlag (1997).
- Hoffman and Stoffel, *Biol Chem*, Hoppe-Seyler 374: 166, (1993).
- Hu et al., *Nature Biotechnology*, In Press, (1999).
- Hu et al., PNAS, 95: 5407-5412, (1998).
- Hutchinson, C., et al., "Mutagenesis at a Specific Position in a DNA Sequence", *J. Biol. Chem.*, 253:6551 (1978).
- Hutchinson et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:710, (1986).
- Joshi et al., *PMB*, 35: 993-1001, (1997).
- Joshi, 1987, *NAR*, 15: 6643-6653, (1987).
- Joshi, 1987, *NAR*, 15: 9627-9640, (1987).
- Lazar et al., "Transforming Growth Factor α : Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities," *Molec. & Cell. Biol.* (1988) 8:3:1247-1252.
- Lu, Shan-Fa, et al., Molecular cloning and characterization of three cellulose synthases associated with xylem development in *Eucalyptus grandis*, Poster presented at American Society of Plant Biologists annual meeting at Denver, CO, Aug. 2-7, 2002.
- Matton, et al., "Identification of cis-acting elements involved in the Regulation of the Pathogenesis-related Gene STH-2 in Potato" *Plant Molecular Biology*, 102:279-291, (1993).
- Nakai, et al., Control of expression by the cellulose synthase (bcsA) promoter region from *Acetobacter xylinum* BPR 2002 *Gene*, 213(1-2):93-100, (1998).
- Needleman & Wunsch, *J. Mol. Biol.*, 48:443-453, (1970).
- Nucleic Acid Hybridization*, B.D. Hames & S.J. Higgins eds. (1985).
- Oligonucleotide Synthesis* (M.J. Gait ed. 1984).
- Oliphant et al., *Gene* 44:177, (1986).
- Pear, et al., "Higher Plants Contain Homologs of the Bacterial *celA* Genes Encoding The Catalytic Subunit of Cellulose Synthase" *PNAS*, 93:12637-12642, (1996).
- Perbal, B., *A Practical Guide to Molecular Cloning* (1984).
- Salvucci, et al., "Identification of the Uridine-Binding Domain of Sucrose-Phosphate Synthase" *Plant Physiology*, 102:529-536, (1993).
- Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, (1989).
- Saxena, et al., "Cloning and Sequencing of the Cellulose Synthase Catalytic Subunit Gene of the *Acetobacter Xylinum*" *Plant Molecular Biology*, 15:673-683, (1990).
- Smith et al., "Antisense RNA Inhibition of Polygalacturonase Gene Expression in Transgenic Tomatoes," *Nature* (1988) 334:724-726.
- Timmell, *Compression Wood in Gymnosperms*, Springer Verlag, (1986).
- Transcription and Translation*, B.D. Hames & S.J. Higgins, eds. (1984).
- Whetten et al., *Ann Rev Pl Physiol in Mol Biol*, 49: 585-609, (1998).
- Wu and Wu, *J. Biol. Chem.* 263:14621-14624, (1988).
- Wu et al., *J. Biol. Chem.* 267:963-967, (1992).
- Wu et al., *Pl Physiol*, 117: 1125, (1998).
- Ye and Varner, *Plant Physiology* 103(3):805-813 (Nov. 1993).
- Ye and Varner, *Methods in Enzymology* 218:671-681 (1993).
- Zhou Y., et al., "Molecular cloning of cellulose synthase genes from loblolly pine", Poster presented at American Society of Plant Biologists annual meeting at Denver, CO, Aug. 2-7, 2002.
- Zoller and Smith, *DNA* 3:479-488, (1984).
- Djerbi, S. et al., "Identification and expression analysis of genes encoding putative cellulose synthases (CesA) in the hybrid aspen, *Populus tremula* (L.) x *P. tremuloides* (Michx.)," *Cellulose* (2004) 11:301-312.
- Joshi, C.P. et al., "Genomics of cellulose biosynthesis in poplars," *New Phytologist* (2004) 164:53-61.
- Joshi, J.P. et al., "Xylem-specific and tension stress-responsive expression of cellulose synthase genes from aspen trees," *Appl. Biochem. Biotech.* (2003) 105-108:17-25.
- Joshi, J.P., "Molecular biology of cellulose biosynthesis in plants," *Recent Res. Devel. Plant Mol. Biol.* (2003) 1:19-38.
- Kalluri U.C. et al., "Differential expression patterns of two cellulose synthase genes are associated with primary and secondary cell wall development in aspen trees," *Planta*. (2004) 220:47-55.
- Kalluri, U.C. et al., "Isolation and characterization of a new, full-length cellulose synthase cDNA, PtrCesA5 from developing xylem of aspen trees," *J. Exp. Botany* (2003) 54(390):2187-2188.
- Li, L. et al., "Combinatorial modification of multiple lignin traits in trees through multigene cotransformation," *Proc. Natl. Acad. Sci. USA* (2003) 100:4939-4944.
- Liang, X. et al., "Molecular cloning of ten distinct hypervariable regions from the cellulose synthase gene superfamily in aspen trees," *Tree Physiology* (2004) 24:543-550.
- Samuga, A. et al., Accession No. AY095297, "A new cellulose synthase gene (PtrCesA2) from Aspen xylem is orthologous to Arabidopsis AtCesA7 (irx3) gene associated with secondary cell wall synthesis," *Gene* (2002) 296(1-2):37-44.
- Tsai, C.J. et al., "Agrobacterium-mediated transformation of quaking aspen (*Populus tremuloides*) and regeneration of transgenic plants," *Plant Cell Reports* (1994) 14:94-97.
- Tsai, C.J. et al., "Suppression of O-methyltransferase gene by homologous sense transgene in quaking aspen causes red-brown wood phenotypes," *Plant Physiology* (1998) 117(5):101-112.
- Wu, L. et al., Accession No. AF072131, "A xylem-specific cellulose synthase gene from Aspen (*Populus tremuloides*) is responsive to mechanical stress," *Plant J.* (2000) 22(6):495-502.
- International Search Report for Application No. PCT/US2000/71670 dated Oct. 18, 2000 (7 pages).
- Written Opinion for Application No. PCT/US2000/71670 dated Oct. 29, 2001 (7 pages).

* cited by examiner

Fig. 1: DNA and predicted protein sequence of PtCela cDNA

```

1  GTCGACCCACGCGTCCCGTCTTGAAAGAATATGAAAGTTGTAAAGAGCTGGTAAAGTGGTAA  60
61  TAAGCAAGATGATGGAATCTGGGGCTCCTATATGCCATACCTGTGGTGAACAGGTGGGGC  120
    M M E S G A P I C H T C G E Q V G H
121  ATGATGCAAATGGGGAGCTATTTGTGGCTTGCCATGAGTGTAGCTATCCCATGTGCAAGT  180
    D A N G E L F V A C H E C S Y P M C K S
181  CTTGTTTCGAGTTTGAAATCAATGAGGGCCGGAAAGTTTGCTTGCGGTGTGGCTCGCCAT  240
    C F E F E I N E G R K V C L R C G S P Y
241  ATGATGAGAACTTGCTGGATGATGTAGAAAAGAAGGGTCTGGCAATCAATCCACAATGG  300
    D E N L L D D V E K K G S G N Q S T M A
301  CATCTCACCTCAACGATTCTCAGGATGTCGGAATCCATGCTAGACATATCAGTAGTGTGT  360
    S H L N D S Q D V G I H A R H I S S V S
361  CCACTGTGGATAGTGAATGAATGATGAATATGGGAATCCAATTTGGAAGAATCGGGTGA  420
    T V D S E M N D E Y G N P I W K N R V K
421  AGAGCTGTAAGGATAAAGAGAACAAGAAGAAAAAGAGAAGTCCTAAGGCTGAAACTGAAC  480
    S C K D K E N K K K R S P K A E T E P
481  CAGCTCAAGTTCCTACAGAACAGCAGATGGAAGAGAAACCGTCTGCAGAGGCTTCGGAGC  540
    A Q V P T E Q Q M E E K P S A E A S E P
541  CGCTTTCAATGTTTATCCAATTCCACGCAACAAGCTCACACCATACAGAGCAGTGATCA  600
    L S I V Y P I P R N K L T P Y R A V I I
601  TTATGCGACTGGTCAATCTGGGCCCTCTTCTTCCACTTCAGAATAACAAATCCTGTCGATA  660
    M R L V I L G L F F H F R I T N P V D S
661  GTGCCCTTGGCCCTGTGGCTTACTTCTGTCAATATGTGAGATCTGGTTTGCATTTCTTGGG  720
    A F G L W L T S V I C E I W F A F S W V
721  TGTGGATCAGTTCCTCAAGTGGAAATCCTGTCAATAGAGAAACGTATATCGAAAGGCTGT  780
    L D Q F P K W N P V N R E T Y I E R L S
781  CGGCAAGGTATGAAAGAGAGGGTGAGCCTTCTCAGCTTGCTGGTGTGGATTTTTCGTGA  840
    A R Y E R E G E P S Q L A G V D F F V S
841  GTACTGTTGATCCGCTGAAGGAACCGCCATTGATCACTGCCAATACAGTCCCTTCCATCC  900
    T V D P L K E P P L I T A N T V L S I L
901  TTGCTGTGGACTATCCCGTCGATAAAGTCTCCTGCTACGTGTCTGATGATGGTGCAGCTA  960
    A V D Y P V D K V S C Y V S D D G A A M
961  TGCTTTCATTTGAATCTCTGTAGAAACAGCTGAGTTTGCAAGGAAGTGGGTCCGTTCT  1020
    L S F E S L V E T A E F A R K W V P F C
1021  GCAAAAAATTCTCAATTGAACCAAGAGCACCGGAGTTTACTTCTCACAGAAAATTGATT  1080
    K K F S I E P R A P E F Y F S Q K I D Y
1081  ACTTGAAAGACAAGGTTCAACCTTCTTTCGTGAAAGAACGTAGAGCAATGAAAAGGATT  1140
    L K D K V Q P S F V K E R R A M K R D Y
1141  ATGAAGAGTACAAAGTCCGAGTTAATGCCCTGGTAGCAAAGGCTCAGAAAACACCTGAAG  1200
    E E Y K V R V N A L V A K A Q K T P E E
    
```

Fig. 1 continued

1201 AAGGATGGACTATGCAAGATGGAACACCTTGGCCTGGGAATAACACACGTGATCACCTG 1260
G W T M Q D G T P W P G N N T R D H P G

1261 GGCATGATTCAGGTCTTCTTGGGAAATACTGGGAGCTCGTGACATTGAAGGAAATGAAC 1320
H D S G L P W E I L G A R D I E G N E L

1321 TACCTCGTCTAGTATATGTCTCCAGGGAGAAGAGACCTGGCTACCAGCACCACAAAAAGG 1380
P R L V Y V S R E K R P G Y Q H H K K A

1381 CTGGTGCAGAAAATGCTCTGGTGAGAGTGTCTGCAGTACTCACAAATGCTCCCTACATCC 1440
G A E N A L V R V S A V L T N A P Y I L

1441 TCAATGTTGATTTGTGATCACTATGTAAACAATAGCAAGGCTGTTCGAGAGGCAATGTGCA 1500
N V D C D H Y V N N S K A V R E A M C I

1501 TCCTGATGGACCCACAAGTAGGTGAGATGTATGCTATGTGCAGTTCCCTCAGAGGTTTG 1560
L M D P Q V G R D V C Y V Q F P Q R F D

1561 ATGGCATAGATAAGAGTGTGCTACGCCAATCGTAACGTAGTTTTCTTTGATGTTAACA 1620
G I D K S D R Y A N R N V V F F D V N M

1621 TGAAAGGTTGGATGGCATTCAAGGACCAGTATACGTAGGAACFGGTTGTGTTTTCAACA 1680
K G L D G I Q G P V Y V G T G C V F N R

1681 GGCAAGCACTTTACGGCTACGGCCCTCCTTCTATGCCAGCTTACGCAAGAGAAAGGATT 1740
Q A L Y G Y G P P S M P S L R K R K D S

1741 CTTCATCCTGCTTCTCATGTGCTGCCCTCAAAGAAGAAGCCTGCTCAAGATCCAGCTG 1800
S S C F S C C C P S K K K P A Q D P A E

1801 AGGTATACAGAGATGCAAAAAGAGAGGATCTCAATGCTGCCATATTTAATCTTACAGAGA 1860
V Y R D A K R E D L N A A I F N L T E I

1861 TTGATAATTATGACGAGCATGAAAGGTCAATGCTGATCTCCCAGTTGAGCTTTGAGAAAA 1920
D N Y D E H E R S M L I S Q L S F E K T

1921 CTTTGGCTTATCTTCTGTCTTCATTGAGTCTACACTAATGGAGAATGGAGGAGTACCCG 1980
F G L S S V F I E S T L M E N G G V P E

1981 AGTCTGCCAACTCACCACCATTCATCAAGGAAGCGATTCAAGTCAATCGGCTGTGGCTATG 2040
S A N S P P F I K E A I Q V I G C G Y E

2041 AAGAGAAGACTGAATGGGAAAACAGATTGGTTGGATATATGGGTCAGTCACTGAGGATA 2100
E K T E W G K Q I G W I Y G S V T E D I

2101 TCTTAAGTGGCTTCAAGATGCACTGCCGAGGATGGAGATCAATTTACTGCATGCCCGTAA 2160
L S G F K M H C R G W R S I Y C M P V R

2161 GGCTGCATTCAAAGGATCTGCACCCATCAACCTGTCTGATAGATTGCACCAGGTCCTCC 2220
P A F K G S A P I N L S D R L H Q V L R

2221 GATGGGCTCTTGGTTCTGTGGAAATTTTCTTTAGCAGACTGTCCCTCTGGTACGGGT 2280
W A L G S V E I F F S R H C P L W Y G F

2281 TTGGAGGAGGCCGCTTAAATGGCTCCAAAGGCTTGGCTATATAAACACCATTTGTGTACC 2340
G G G R L K W L Q R L A Y I N T I V Y P

2341 CATTACATCCCTCCCTCTCATTCCTTATGCAAAATCCTGCAGTTTGTCTGCTCACCG 2400
F T S L P L I A Y C T I P A V C L L T G

2401 GAAAATTCATCATACCAACGCTCTCAAACCTGGCAAGCATGCTGTTTCTTGGCCTCTTTA 2460
K F I I P T L S N L A S M L F L G L F I

2461 TCTCCATCATTGTAAC TGCGGTGCTTGAGCTAAGATGGAGCGGTGTCAGCATTGAAGATT 2520

Fig. 1 continued

S I I V T A V L E L R W S G V S I E D L

2521 TATGGCGTAATGAACAATTCGGGTGATCGGAGGTGTTTCAGCCCATCTCTTTGCGGTCT 2580
W R N E Q F W V I G G V S A H L F A V F

2581 TCCAGGGATTCTFAAAAATGTTGGCTGGCATCGATACGAACTTCACTGTCACAGCAAAG 2640
Q G F L K M L A G I D T N F T V T A K A

2641 CAGCCGAAGATGCAGAATTGGGGAGCTATATATGGTCAAGTGGACAACACTTTTGATT 2700
A E D A E F G E L Y M V K W T T L L I P

2701 CTCCAACCACACTTCTCATTATCAATATGTCGGGTGCTGGATTCTCTGATGCACTCA 2760
P T T L L I I N M S G C A G F S D A L N

2761 ACAAAGGATATGAAGCATGGGGCCTCTCTTTGGCAAGGTGTTCTTTGCTTTCTGGGTGA 2820
K G Y E A W G P L F G K V F F A F W V I

2821 TTCTTCATCTCTATCCATTCTTAAAGGTCTAATGGGTCGCCAAAACCTAACACCAACCA 2880
L H L Y P F L K G L M G R Q N L T P T I

2881 TTGTTGTTCTCTGGTCAGTCTGTTGGCCTCTGTCCTCTCTCGTTTGGGTCAAGATCA 2940
V V L W S V L L A S V F S L V W V K I N

2941 ATCCATTCGTTAACAAAGTTGATAACACCTTGGTTGCGGAGACCTGCATTTCCATTGATT 3000
P F V N K V D N T L V A E T C I S I D C

3001 GCTGAGCTACCTCCAATAAGTCTCTCCAGTATTTTGGGGTTACAAAACCTTTGGGAATT 3060
*

3061 GGAATATGATCCTCGTTGTAGTTTCCCTCAAGAAAGCACATATCGCTGTCAGTATTTAAA 3120

3121 TGAAC TGCAAGATGATTGTTCTCTATGAAGTTTGAACAGTTTGAATGATATTATGTTA 3180

3181 AAATACAGGTTTTGATTGTGTTGAAAAAAAAAAGAAAAAAAAAAAAAAAAAAAA 3232




FIG. 2




FIG. 3

Fig. 4: DNA sequence of PtCelAP, the 5' flanking region of PtCela coding sequence

```

1 GAATTCGCCCTTTTGAATTCAGGAGACGATAGTTCCGGTTCGTTGAATGGCTTTGTTC 60
61 CTTCTGGTCTAGCAATTTGCAAAGAAGTTACAAAACAAATGCATATTATGTAAATTTAA 120
121 CAAGAGATGGGTTCTATGGTCACTTATTTATGCCCATCATTTGTTCTGGGGTTACTCTTT 180
181 ATAGTCTGATTCGAAGTTGCAAACGCGTTTCTGGTATTGCAATTATGTAGCCATAAAC 240
241 TGTTAATCCTGTAGCTATTAGCGGACCAACAACCAGATATACGGGATCAGCGTCGTA AAA 300
301 GAGATCTCCATTCTACGTTTCTTTCTAATTTTTCCGTTTTCAGTGAGAGAATTACCCTGAT 360
361 ACATTGACATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT 420
421 GGATCCTGCCAGTTTTTCAGTTCCACATGGCATCTCAGCCCAAGATCATGTGTTTATACGCC 480
481 TAATGACTTGTATTGAAAGTTTGGTAAAGTTGAAGATGTGCTCTGCCCAACAGAAACCTTC 540
541 CTTAAATTTCCAGCAAATCTTTCAAACCTTGGCCTTACACCCCGAAAATAGACGTGCTTCT 600
601 ACTTGGGTCTTTGGAAACCATGCACCAACCGCCATACCCCAACCAACCCACCCCTCAAC 660
661 CTTCTCTCGCCATTACAAAATGTCAGTACCACCCCTCTGAAAGACACCAACACACCCCTA 720
721 GCTTTGGTTAGGGTATTTGATATAAAAAACAAGGCCAAAACAAAAGATTGGAAGGAAGCAG 780
781 AGGAAGACCCCTCTTGAAGAATGAAAGTTGTAAGAGCTGGTAAAGTGGTAATAAGCAAG 840
841 ATGATGGAATCTGGGGCTCCTATATGCCATACCTGTGGTGAACAGGTGGGGCATGATGCA 900
M M E S G A P I C H T C G E Q V G H D A
901 AATGGGGAGCTATTTGTGGCTTGCCATGAGTGTAGCTATCCCATGTGCAAGTCTTGTTTC 960
N G E L F V A C H E C S Y P M C K S C F
961 GAGTTTGAAATCAAAGAGGGCCGGAAAGTTTGCTTCCGGTGTGGCTCGAG 1010
E F E I K E G R K V C L R C G S

```

FIG. 5




FIG. 6




FIG. 7

Arabidopsis thaliana cellulose synthase mRNA SEQ ID NO: 4

```

1  ggggcccggg ttaatcgccg gttctcacia caggaatgag tttgtcctca ttaatgccga
61  tgagaatgcc cgaataagat cagtccaaga gctgagtgga cagacatgtc aaatctgcag
121 agatgagatc gaattgactg ttgatggaga accgtttggt gcatgtaacg aatgtgcatt
181 cctgtgtgtg agaccttget atgagtacga aagacgagaa ggcaatcaag cttgtccaca
241 gtgcaaaaacc cgtttcaaac gtcttaaagg aagtccaaga gttgaagggt atgaagagga
301 agatgacatt gatgatttag acaatgagtt tgagta tggga aataatggga ttggatttga
361 tcaggtttct gaaggatgtg caatctctcg tcgcaactcc ggtttccacc aatctgattt
421 ggattcagct ccacctgget ctacagattcc attgctgact tacggcgagc aggacgttga
481 gatttctctc gatagacatg ctcttattgt tcctccttca cttggtgggt atggcaatag
541 agttcctcct gttctctctt ctgaccggca cgtggctgca catogaagcc tgatggacc
601 tcagaaagat cttgcccgtt atggttatgg aagtgtcgct tggaaagatc ggtggagga
661 atggaagaga aagcagaatg agaaacttca ggttgttagg catgaaggag atcctgattt
721 tgaagatggt gatgatgctg attttccaat gatgtagag ggaaggcagc cattgtctat
781 gaagatacca atcaaatcga gcaagataaa tccttaccgg atgttaattg tgcatcgtct
841 tgtgattctt ggtctctctc ttcactaccg tattctcac cccgtcaaaq atgcatatgc
901 tttgtggctt atttctgtta tatgtgagat atggtttgcct gtttcatggg ttcttgatca
961 gttccctaaa tggtaaccta tcgagcgaga aacgtacttg gaccgactct cattaagata
1021 tgagaaagaa gggaaaccgt cgggactatc ccctgtggat gtatttgtta gtacagtgga
1081 tccattgaaa gagcctccgc ttattactgc aaatactgtc ttgtctatc ttgctgttga
1141 ttatcctgtc gataagggtg cttggttacc atctgatgat ggtgctgcta tgctactttt
1201 cgaagctctt tctgagaccg ctgaattcgc aaggaaatgg gttcctttct gcaagaaata
1261 ttgtattgag cctcgtgctc ccgaatggtg tttctgccat aaaatggact acttgaagaa
1321 taaagttcat cccgcatttg ttagggagcg gcgagccatg aagagagatt atgaagaatt
1381 caaagtaaaq atcaatgctt tagtagcaac agcacagaaa gtgcctgagg atggttggac
1441 tatgcaagac ggtacacctt ggcccggtaa tagtgtgcca gatcctcctg gcatgattca
1501 ggtcttctct ggaagtgcag gtgttcgtga tgtcgaaaac aacgagttgc ctcgattagt
1561 ttacgtttct cgtgagaaga gaccgggatt tgatcaccat aagaaggctg gagctatgaa
1621 ttccctgata caggtctctg ggttctctatc aaatgctcct tacctcttga atgtcgattg
1681 tgatcactac atcaacaata gcaagctctc tagagaagca atgtgttca tgatggatcc
1741 tgatcagga aagaaaatct gttatgttca gttccctcaa aggttcgatg aggtgatag
1801 gcacgatcga tactcaaatc gcaatgttgt gttctttgat atcaatatga aaggtttgga
1861 tgggtcaciaa gggcctatat acgtcggtac aggttgtgtt ttcaggaggc aagcgttta
1921 cggatttgat gcaccgaaga agaagaaggg cccacgtaag acatgcaat tctggccaaa
1981 atggtgtctc ctatgttttg gttcaagaaa gaatcgtaaa gcaaaagcag tgggtcggga
2041 taagaagaag aagaataggg aagcgtcaaa gcagatccac gcattagaaa atatogaaga
2101 gggcccggtt cataaagttc ttaacgtaga acagtcaccc gaggcaatgc aaatgaagtt
2161 gcagaagaaa tatgggcagt ctctgttatt tgttgcctct gcgcctctgg agaattggtg
2221 gatggctaga aacgcaagcc cggcttgtct gcttaagaa gccatccaag tcattagtcg
2281 cggatagtaa gataaaactg aatggggaaa agagattggg tggatctatg gttctgttac
2341 cgaagatatt cttacgggtt ctaagatgca ttctcatggt tggagacatg tttattgtac
2401 accaaagtta gggccttcca aaggatcagc tccaatcaat ctttcggatc gttccatca
2461 agttctctga tgggcgcttg ggtcggttga gattttcttg agtaggcatt gtctatttg
2521 gtatggttat ggagggtggg tgaatggctc tgagcggttg tcctacatta actctgtggt
2581 ttaccctggg acctctctac cgtcatcctg ttactgttct ctccctgcca tctgtctct
2641 cactggaaaa ttcatcgttc ccgagattag caactatgct agtatcctct tcatggcct
2701 cttctcgtcg attgcaataa cgggtattct cgagatgcaa tggggcaaaq ttgggatcga
2761 tgattggtgg agaaaacgaac agttttgggt catggagggt gtttctgcgc atctgtttgc
2821 tctctccaa ggtctctca aggtctcttg ttgtgtgcac actaaactca cagtcacatc
2881 aaaagcagct gatgatggag agttctctga cctttacctg tccaatgga cttcactct
2941 catccctcca atgactctac tcatcataaa cgtcattgga gtcatagtcg gagtctctga
3001 tgccatcagc aatggatacg actcgtgggg accgcttttc ggaagactgt tctttgact
3061 ttgggtcatc attcatcttt acccgttctc taaaggtttg cttgggaaac aagatagaat
3121 gccaacatt attgtcgtct ggtccatcct cctggcctcg attcttacac tctttgggt
3181 cgggtttaat ccgtttgtgg cgaaggcggg tcttattctc gagatctgtg gtttagactg
3241 cttgtgatcc gattgaccgg tggatgggtt ggtgaaaaag gtttaattcc cacggatcaa
3301 agagaggtaa gagagatatt gttttacctc taaaagactc cttcattgtg ttcattagat
3361 gatgaaaaat gaaaagaaa agaagattta atlttgttac gagaattggt atttttgcaa
3421 gaatgtggtg tagatagcgg ccgc

```

FIG. 8

Arabidopsis thaliana cellulose synthase SEQ ID NO: 5

RPRLIAGSHNRNEFVLINADENARIRSVQELSGQTCQICRDEIE
LTVDGEPFVACNECAFVCRPCYEYERREGNQACPQCKTRFKRLKGS PRVEGDEEEDD
IDDL DNEF EYGNNGIGFDQVSEGMSISRNSGFPQSDLDSAPPGSQI PLLTYGDEDVE
ISSDRHALIVPPSLGGHGNRVHPVSLSDPTVAHRRLMVPQKDLAVYGYGSVAWKDRM
EEWKRKQNEKLQVVRHEGDPDFEDGDDADFPMMDEGRQPLSMKIPKSSKINPYRMLI
VLRLVILGLFFHYRILHPVKDAYALWLISVICEIWFVAVSWVLDQFPKWYPIERETYLD
RLSLRYEKEGKPSGLSPVDVVFVSTVDPLKEPPLITANTVLSILAVDYPVDKVACYVSD
DGAAMLTFEALSETAEFARKWVPFCKKYCIEPRAPEWYFCHKMDYLKKNVHPAFVRER
RAMKRDIYEEFKVKINALVATAQKVPEDGWTMQDGTWPWPGNSVRDHPGMIQVFLGSDGV
RDVENNELPRLVYVSREKRPFGDHHKKAGAMNSLIRVSGVLSNAPYLLNVDCDHYINN
SKALREAMCFMMDPQSGKKICYVQFPQRFDGDIDRHDYRNSNRNVVFFDINMKGLDGLQG
PIYVGTGCVFRRQALYGFDA PKKKK GPRKTCNCWPKWCLLCFGSRKNRKAKTVAADKK
KKNREASKQIHALENIIEGRGHKVLNVEQSTEAMQMKLQKKYGQSPV FVASARLENGG
MARNASPACLLKEAIQVISRGYEDKTEWGKEIGWIYGSVTEDILTGSKMHSWGWRHVY
CTPKLAAFKGSAPINLSDRHLHQVLRWALGSVEIFLSRHCPWIWYGYGGGLKWLERLSYI
NSVVYPWTS LPLIVYCSLPAICLLTGKFIVPEISNYASILFMALFSSIAITGILEMQW
GKVGIDDWWRNEQFWVIGGVSAHLFALFQGLLKVLAVGVD TNFTVTSKAADDGEFSDLY
LFWKTSLLIPPMTLLIINVIGVIVGVSDAISNGYDSWGPLFGRLFFALWV I IHLYPFL
KGLLGKQDRMPTIIVVWSILLASILTLLWVRVNPVFAKGGP ILEICGLDCL

gaattogccc ttttgattc aggagacgat agtttccgggt tcgttgaatg gctttgttca
 cttotgggtot agcaatttgc aaaagaagtt acaaaacaaa tgcatattat gtaaatttaa
caagagatgg gttctatggt cacttattta tgccatcat ttgttctggg gttactcttt
 atagtctgat tcgaagttgc aaactgccgt ttctggtatt gcaattatgt agccataaac
 tgtaatoct gtagctatta gcggaaccaac aaccagatat acgggatcag cgtcgtaaaa
 gagatctoca ttctacgttt ctttctaatt tttccgtttc agtgagagaa ttaccctgat
 acattgacat gatgattgat gattatggga accattccga tgtagacac gagaccatct
 ggatcctgcc agttttcagt tcacatggca tctcagcca agatcatgtg tttatacgcc
 taatgacttg tattgaaagt ttggtaagtt gaagatgtgc ttgccaac agaaaccttc
cttaaatttc cagcaaatct ttcaaacttg gccttacacc ccgaaaatag acgtgcttct
 acttgggttc ttggaacca tgcaccaacc gccataccc accaaccac caccctcaac
 cttctcttcg ccattacaaa aatgtcagta ccaccctctg aaagacacca acacacctta
gctttggtta gggtatttga tataaaaaca aggccaaaac aaaagattgg aaggaagcag
 aggaagacc tcttgaaaga attgaagttg taaagagctg gtaaagtgt aataagcaag

FIGURE 9

1

ISOLATED CELLULOSE SYNTHASE PROMOTER REGIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is continuation-in-part of U.S. application Ser. No. 11/437,368 filed May 19, 2006, now U.S. Pat. No. 7,232,941, which is a divisional of U.S. application Ser. No. 09/980,043 filed Apr. 5, 2002, now U.S. Pat. No. 7,049,481, which is a §371 of International Application No. PCT/US00/13637 filed May 18, 2000, which claims the benefit of priority to U.S. Provisional Application No. 60/135,280 filed May 21, 1999, each of which are incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

Not Applicable

FIELD OF THE INVENTION

This invention relates to polynucleotide molecules encoding cellulose synthase, promoters of cellulose synthase and cellulose synthase polypeptides, methods for genetically altering cellulose and lignin biosynthesis, and methods for improving strength properties of juvenile wood and fiber in trees. The invention further relates to methods for identifying regulatory elements in a cellulose synthase promoter and transcription factors that bind to such regulatory elements, and to methods for augmenting expression of polynucleotides operably linked to a cellulose synthase promoter.

BACKGROUND OF THE INVENTION

Lignin and cellulose are the two major building blocks of plant cell walls that provide mechanical strength and rigidity. In plants, and especially in trees, these two organic materials exist in a dynamic equilibrium conferring mechanical strength, water transporting ability and protection from biotic and abiotic environmental stresses. Normally, oven-dry wood contains 30 to 50% cellulose, 20 to 30% lignin and 20 to 30% hemicellulose (Higuchi, 1997).

Proportions of lignin and cellulose are known to change with variation in the natural environment. For example, during the development of compression wood in conifers, the percentage of lignin increases from 30 to 40%, and cellulose content proportionally decreases from 40 to 30% (Timmell, 1986). Conversely, in angiosperm tension wood the percentage of cellulose increases from 30 to 40%, while lignin content decreases from 30 to 20% (Timmell, 1986).

It was recently discovered that the genetic down-regulation of a key tissue-specific enzyme from the lignin biosynthesis pathway, 4CL, results in reduction of lignin content by up to 45% in transgenic aspen trees (Hu et al., 1999). This down-regulation is also associated with a 15% increase in the cellulose content. If the converse were true, i.e., that increasing cellulose content by genetic up-regulation of cellulose biosynthesis results in reduction of lignin content, then the pulp yield could be increased. This would allow tremendous savings in chemical and energy costs during pulping because, for example, lignin must be degraded and removed during the pulping process.

Cellulose is a linear glucan consisting of β -D-1,4-linked glucose residues. It is formed by a cellulose synthase enzyme which catalyzes assembly of UDP-glucose units in plasma membrane complexes known as "particle rosettes" (Delmer

2

and Amor, 1995). Cellulose synthase is thought to be anchored to the membrane by eight transmembrane binding domains to form the basis of the cellulose biosynthesis machinery in the plant cell wall (Pear et al., 1996).

5 In higher plants, the glucan chains in cellulose microfibrils of primary and secondary cell walls are different in their degree of polymerization (Brown et al., 1996). For example, secondary cell walls are known to contain cellulose having a high degree of polymerization, while in primary cell walls the degree of polymerization is lower. In another example, 10 woody cell walls suffering from tension stress produce tension wood on the upper side of a bent angiosperm tree in response to the stress. In these cells, there are elevated quantities of cellulose which have very high crystallinity. The 15 formation of highly crystalline cellulose is important to obtain a higher tensile strength of the wood fiber. Woody cell walls located at the under side of the same stem experience a compression stress, but do not produce highly crystalline cellulose. Such variation in the degree of polymerization in 20 cell walls during development is believed to be due to different types of cellulose synthases for organizing glucose units into different paracrystalline arrays (Haigler and Blanton, 1996). Therefore, it would be advantageous to determine the molecular basis for the synthesis of highly crystalline cellulose so that higher yields of wood pulp having superior 25 strength properties can be obtained from transgenic trees. Production of highly crystalline cellulose in transgenic trees would also markedly improve the mechanical strength properties of juvenile wood formed in normal trees. This would be a great benefit to the industry because juvenile wood is generally undesirable for solid wood applications because it has inferior mechanical properties.

Since the deposition of cellulose and lignin in trees is regulated in a compensatory fashion, genetic augmentation of 35 cellulose biosynthesis might have a repressive effect on lignin deposition. Since the degree of polymerization and crystallinity may depend upon the type of cellulose synthase incorporated in the cellulose biosynthesis machinery, the expression of heterologous cellulose synthase or a UDP-glucose binding region thereof (e.g., sweetgum protein expression in loblolly pine), could increase the quality of cellulose in transgenic plants. Over-expression of a heterologous cellulose 40 synthase may also increase cellulose quantity in transgenic plants. Thus, genetic engineering of cellulose biosynthesis can provide a strategy to augment cellulose quality and quantity, while reducing lignin content in transgenic plants.

A better understanding of the biochemical processes that lead to wood formation would enable the pulp and paper industries to more effectively use genetic engineering as a 50 tool to meet the increasing demands for wood from a decreasing production area. With this objective, many xylem-specific genes, including most lignin biosynthesis genes, have been isolated from developing xylem tissues of various plants including tree species (Ye and Varner, 1993; Fukuda, 1996; Whetten et al., 1998). Genes regulating cellulose biosynthesis in crop plants (Pear et al., 1996 and Arioli et al., 1998), versus 55 in trees, have also been isolated. However, isolation of tree genes which are directly involved in cellulose biosynthesis has remained a great challenge.

60 For more than 30 years, no gene encoding higher plant cellulose synthase (Ce1A) was identified. Recently, Pear et al. (1996) isolated the first putative higher plant Ce1A cDNA, GhCe1A (GenBank No. GHU58283), by searching for UDP-glucose binding sequences in a cDNA library prepared from cotton fibers having active secondary wall cellulose synthesis. GhCe1A was considered to encode a cellulose synthase catalytic subunit because it is highly expressed in cotton

fibers, actively synthesizes secondary wall cellulose, contains eight transmembrane domains, binds UDP-glucose, and contains two other domains unique to plants.

Recently, Arioli et al. (1998) cloned a CelA homolog, RSW1 (radial swelling) (GenBank No. AF027172), from *Arabidopsis* by chromosome walking to a defective locus of a temperature sensitive cellulose-deficient mutant. Complementation of the RSW1 mutant with a wild type full-length genomic RSW1 clone restored the normal phenotype. This complementation provided the first genetic proof that a plant CelA gene encodes a catalytic subunit of cellulose synthase and functions in the biosynthesis of cellulose microfibrils. The full-length *Arabidopsis* RSW1 represents the only known, currently available cellulose synthase cDNA available for further elucidating cellulose biosynthesis in transgenic systems (Wu et al., 1998).

The discovery of the RSW1 gene substantiated the belief that the assembly of a cellulose synthase into the plasma membrane is required for functional cellulose biosynthetic machinery and for manufacturing crystalline cellulose microfibrils in plant cell walls. Most significantly, a single CelA gene, e.g. RSW1, is sufficient for the biosynthesis of cellulose microfibrils in plants, e.g. *Arabidopsis*. Thus, RSW1 is a prime target for engineering augmented cellulose formation in transgenic plants.

Since many of society's fiber, chemical and energy demands are met through the industrial-scale production of cellulose from wood, genetic engineering of the cellulose biosynthesis machinery in trees could produce higher pulp yields. This would allow greater returns on investment by pulp and paper industries. Therefore, it would be advantageous to isolate and characterize genes from trees that are involved in cellulose biosynthesis in order to improve the properties of wood.

SUMMARY OF THE INVENTION

The present invention relates to polynucleotides comprising a nucleotide sequence that encodes a cellulose synthase, regulatory sequences, including a stress-inducible promoter, of the cellulose synthase, a cellulose synthase protein or a functional domain thereof and methods for augmenting cellulose biosynthesis in plants.

Thus, in one aspect, the invention provides a polynucleotide comprising a sequence that encodes a cellulose synthase, or a polynucleotide fragment thereof, the fragment encoding a functional domain of cellulose synthase, such as a UDP-glucose binding domain. The invention also provides a cellulose synthase or a functional domain or fragment thereof, including a UDP-glucose binding domain and at least one of eight transmembrane domains. The invention further provides a cellulose synthase promoter, or a functional fragment thereof, which fragment contains one or more mechanical stress response elements (MSRE).

In another aspect, the present invention is directed to a method of improving the quality of wood by altering the quantity of cellulose in plant cells, and optionally decreasing the content of lignin in the cell. The invention also relates to a method of altering the growth or the cellulose content of a plant by expressing an exogenous polynucleotide encoding a cellulose synthase or a UDP-glucose binding domain thereof in the plant. The invention further provides a method for causing a stress-induced gene expression in a plant cell by expressing a polynucleotide of choice using a stress-inducible cellulose synthase promoter.

In yet another aspect, the invention relates to a method for determining a mechanical stress responsive element (MSRE)

in a cellulose synthase promoters and a method for identifying transcription factors that binds to the MSRE.

In a further aspect, the invention provides a method for altering (increasing or decreasing) i.e., regulating, the expression of a cellulose synthase in a plant by expressing an exogenous polynucleotide encoding a transcription factor having the property of binding a positive MSRE of a cellulose synthase promoter or by expressing an antisense polynucleotide encoding a transcription factor having the property of binding a negative MSRE to block the expression of the transcription factor.

Other aspects of the invention will be appreciated by a consideration of the detailed description of the invention drawings and appended claims.

DESCRIPTION OF THE DRAWINGS

FIG. 1 represents a nucleic acid sequence encoding a cellulose synthase from *Populus tremuloides* [SEQ ID NO: 1] and the protein sequence thereof [SEQ ID NO: 2].

FIG. 2 a-c (collectively referred to as FIG. 2) represent a Southern blot analysis of aspen genomic DNA probed with a fragment of the aspen cDNA represented in FIG. 1 under low (panel a) and high stringency conditions (panel b), and a northern blot analysis of the total aspen RNA from stem internodes using the same probe (panel c).

FIG. 3 a-d (collectively referred to as FIG. 3) represent in situ localization of the cellulose synthase gene transcripts as shown in the transverse sections from second (panel a), fourth (panel b), sixth (panel c) and fifth (panel d) internode.

FIG. 4 represents a nucleic acid sequence of the 5' region of aspen cellulose synthase gene including the promoter region and the 5' portion of the coding sequence [SEQ ID NO: 3] and the peptide sequence deduced from the coding sequence [SEQ ID NO: 6].

FIG. 5 a-f (collectively referred to as FIG. 5) represents a histochemical analysis (panels a-d and f) and fluorescence microscopy (panel e) of transgenic tobacco for GUS gene expression driven by a cellulose synthase promoter of the invention.

FIG. 6 a-d (collectively referred to as FIG. 6) represents a histochemical analysis of GUS gene expression driven by aspen cellulose synthase promoter of the invention; tangential and longitudinal sections were harvested before bending (panel a), and 4 (panel b), 20 (panel c) and 40 (panel d) hours after bending and stained for GUS expression.

FIG. 7 represents a cDNA encoding cellulose synthase isolated from *Arabidopsis* [SEQ ID NO:4].

FIG. 8 represents an *Arabidopsis* cellulose synthase [SEQ ID NO:5] encoded by the cDNA represented in FIG. 7.

FIG. 9 represents a nucleic acid sequence from position 1 to position 840 of SEQ ID NO: 3, which includes the promoter region that is 5' of the aspen cellulose synthase coding sequence, with the position of primers used to clone sub fragments indicated in bold and underlined, and the direction of the primers indicated by arrows.

DETAILED DESCRIPTION OF THE INVENTION

All patents, patent applications and references cited in this specification are hereby incorporated herein by reference in their entirety. In case of any inconsistency, the present disclosure governs.

Definitions

The terms used in this specification generally have their ordinary meanings in the art, within the context of the inven-

tion, and in the specific context where each term is used. Certain terms are discussed below, or elsewhere in the specification, to provide additional guidance to the person of skill in the art in describing the compositions and methods of the invention and how to make and use them. It will be appreciated that the same thing can be said in more than one way. Consequently, alternative language and synonyms may be used for any one or more of the terms discussed herein, nor is any special significance to be placed upon whether or not a term is elaborated or discussed herein. Synonyms for certain terms are provided. A recital of one or more synonyms does not exclude the use of other synonyms. The use of examples anywhere in this specification, including examples of any terms discussed herein, is illustrative only, and in no way limits the scope and meaning of the invention or of any exemplified term. Likewise, the invention is not limited to the preferred embodiments.

The term "plant" includes whole plants and portions of plants, including plant organs (e.g. roots, stems, leaves, etc.).

The term "angiosperm" refers to plants which produce seeds encased in an ovary. A specific example of an angiosperm is *Liquidambar styraciflua* (L.) [sweetgum].

The term "gymnosperm" refers to plants which produce naked seeds, that is, seeds which are not encased in an ovary. Specific examples of a gymnosperm include *Pinus taeda* (L.) [loblolly pine].

The term "polynucleotide" or "nucleic acid molecule" is intended to include double or single stranded genomic and cDNA, RNA, any synthetic and genetically manipulated polynucleotide, and both sense and anti-sense strands together or individually (although only sense or anti-sense stand may be represented herein). This includes single- and double-stranded molecules, i.e., DNA-DNA, DNA-RNA and RNA-RNA hybrids, as well as "protein nucleic acids" (PNA) formed by conjugating bases to an amino acid backbone. This also includes nucleic acids containing modified bases, for example thio-uracil, thio-guanine and fluoro-uracil.

An "isolated" nucleic acid molecule or polynucleotide refers to a component that is removed from its original environment (for example, its natural environment if it is naturally occurring). An isolated nucleic acid or polypeptide may contain less than about 50%, preferably less than about 75%, and most preferably less than about 90%, of the cellular components with which it was originally associated. A polynucleotide amplified using PCR so that it is sufficiently and easily distinguishable (on a gel, for example) from the rest of the cellular components is considered "isolated". The polynucleotides and polypeptides of the invention may be "substantially pure," i.e., having the highest degree of purity that can be achieved using purification techniques known in the art.

The term "hybridization" refers to a process in which a strand of nucleic acid joins with a complementary strand through base pairing. Polynucleotides are "hybridizable" to each other when at least one strand of one polynucleotide can anneal to a strand of another polynucleotide under defined stringency conditions. Hybridization requires that the two polynucleotides contain substantially complementary sequences; depending on the stringency of hybridization, however, mismatches may be tolerated. Typically, hybridization of two sequences at high stringency (such as, for example, in an aqueous solution of 0.5×SSC at 65° C.) requires that the sequences exhibit some high degree of complementarity over their entire sequence. Conditions of intermediate stringency (such as, for example, an aqueous solution of 2×SSC at 65° C.) and low stringency (such as, for example, an aqueous solution of 2×SSC at 55° C.), require correspondingly less overall complementarity between the

hybridizing sequences. (1×SSC is 0.15 M NaCl, 0.015 M Na citrate.) As used herein, the above solutions and temperatures refer to the probe-washing stage of the hybridization procedure. The term "a polynucleotide that hybridizes under stringent (low, intermediate) conditions" is intended to encompass both single and double-stranded polynucleotides although only one strand will hybridize to the complementary strand of another polynucleotide.

A "sequence-conservative variant" is a polynucleotide that contains a change of one or more nucleotides in a given codon position, as compared with another polynucleotide, but the change does not result in any alteration in the amino acid encoded at that position.

A "function-conservative variant" is a polypeptide (or a polynucleotide encoding the polypeptide) having a given amino acid residue that has been changed without altering the overall conformation and function of the polypeptide, including, but not limited to, replacement of an amino acid with one having similar physico-chemical properties (such as, for example, acidic, basic, hydrophobic, and the like). Amino acids having similar physico-chemical properties are well known in the art. For example, arginine, histidine and lysine are hydrophilic-basic amino acids and may be interchangeable. Similarly, isoleucine, a hydrophobic amino acid, may be replaced with leucine, methionine or valine. Sequence- and function-conservative variants are discussed in greater detail below with respect to degeneracy of the genetic code.

A "functional domain" or a "functional fragment" refers to any region or portion of a protein or polypeptide or polynucleotide which is a region or portion of a larger protein or polynucleotide, the region or portion having the specific activity or specific function attributable to the larger protein or polynucleotide, e.g., a functional domain of cellulose synthase is the UDP-glucose binding domain.

The term "% identity" refers to the percentage of the nucleotides/amino acids of one polynucleotide/polypeptide that are identical to the nucleotides/amino acids of another sequence of polynucleotide/polypeptide as identified by program GAP from Genetics Computer Group Wisconsin (GCG) package (version 9.0) (Madison, Wis.). GAP uses the algorithm of Needleman and Wunsch (J. Mol. Biol. 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. When parameters required to run the above algorithm are not specified, the default values offered by the program are contemplated. The following parameters are used by the GCG program GAP as default values (for polynucleotides): gap creation penalty: 50; gap extension penalty: 3; scoring matrix: nwsgapdna.cpm (local data file).

The "% similarity" or "% homology" between two polypeptide sequences is a function of the number of similar positions shared by two sequences on the basis of the scoring matrix used divided by the number of positions compared and then multiplied by 100. This comparison is made when two sequences are aligned (by introducing gaps if needed) to determine maximum homology. PowerBlast program, implemented by the National Center for Biotechnology Information, can be used to compute optimal, gapped alignments. GAP program from Genetics Computer Group Wisconsin package (version 9.0) (Madison, Wis.) can also be used. GAP uses the algorithm of Needleman and Wunsch (J Mol Biol 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. When parameters required to run the above algorithm are not specified, the default values offered by the program are contemplated. The following parameters are used by the GCG program GAP as default

values (for polypeptides): gap creation penalty: 12; gap extension penalty: 4; scoring matrix: Blosum62.cpm (local data file).

The term "oligonucleotide" refers to a nucleic acid, generally of at least 10, preferably at least 15, and more preferably at least 20 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, or an mRNA molecule encoding a gene, mRNA, cDNA, or other nucleic acid of interest. Oligonucleotides can be labeled, e.g., with ³²P-nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. In one embodiment, a labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid. In another embodiment, oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of CelA, or to detect the presence of nucleic acids encoding CelA. In a further embodiment, an oligonucleotide of the invention can form a triple helix with a CelA DNA molecule. In still another embodiment, a library of oligonucleotides arranged on a solid support, such as a silicon wafer or chip, can be used to detect various polymorphisms of interest. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer. Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

The term "coding sequence" refers to that portion of the gene that contains the information for encoding a polypeptide. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences.

A "promoter" is a polynucleotide containing elements (e.g., a TATA box) which are capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Examples of promoters that can be used in the present invention include PtCelAP, 4CL-1 and 35S.

The term "constitutive promoter" refers to a promoter which typically, does not require positive regulatory proteins to activate expression of an associated coding sequence, i.e., a constitutive promoter maintains some basal level of expression. A constitutive promoter is commonly used in creation of an expression cassette. An example of a constitutive promoter are 35S CaMV (Cauliflower Mosaic Virus), available from Clontech, Palo Alto, Calif.

The term "inducible promoter" refers to the promoter which requires a positive regulation to activate expression of an associated coding sequence. An example of such a promoter is a stress-inducible cellulose synthase promoter from aspen described herein.

A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced and translated into the protein encoded by the coding sequence.

A "vector" is a recombinant nucleic acid construct, such as plasmid, phage genome, virus genome, cosmid, or artificial chromosome to which a polynucleotide of the invention may be attached. In a specific embodiment, the vector may bring about the replication of the attached segment, e.g., in the case of a cloning vector.

The term "expression cassette" refers to a polynucleotide which contains both a promoter and a protein coding sequence such that expression of a given protein is achieved upon insertion of the expression cassette into a cell.

A cell has been "transfected" by exogenous or heterologous polynucleotide when such polynucleotide has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous polynucleotide when the transfected polynucleotide effects a phenotypic change. Preferably, the transforming polynucleotide should be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

"Exogenous" refers to biological material, such as a polynucleotide or protein, that has been isolated from a cell and is then introduced into the same or a different cell. For example, a polynucleotide encoding a cellulose synthase of the invention can be cloned from xylem cells of a particular species of tree, inserted into a plasmid and reintroduced into xylem cells of the same or different species. The species thus contains an exogenous cellulose synthase polynucleotide.

"Heterologous polynucleotide" refers to an exogenous polynucleotide not naturally occurring in the cell into which it is introduced.

"Homologous polynucleotide" refers to an exogenous polynucleotide that naturally exists in the cells into which it is introduced.

The present invention relates to isolation and characterization of polynucleotides encoding cellulose synthases from plants, especially trees, including full length or naturally occurring forms of cellulose synthases, functional domains, promoters and regulatory elements. Therefore, in accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D. N. Glover ed. 1985); *Oligonucleotide Synthesis* (M. J. Gait ed. 1984); *Nucleic Acid Hybridization* [B. D. Hames & S. J. Higgins eds. (1985)]; *Transcription And Translation* [B. D. Hames & S. J. Higgins, eds. (1984)]; *Animal Cell Culture* [R. I. Freshney, ed. (1986)]; *Immobilized Cells And Enzymes* [IRL Press, (1986)]; B. Perbal, *A Practical Guide To Molecular Cloning* (1984); F. M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

The present invention relates to a novel, full-length cellulose synthase gene (CelA), a novel stress inducible promoter of cellulose synthases (CelAP), and cellulose synthase proteins from trees, including UDP-glucose catalytic domains thereof. The invention enables the development of transgenic tree varieties having increased cellulose content, decreased lignin content and, therefore, improved wood fiber characteristics. Production of increased cellulose quantity and quality in multiple varieties of commercially relevant, transgenic forest tree species in operational production scenarios are further contemplated. The invention further provides a new experimental system for study of CelA gene expression and function in trees.

Polynucleotides Encoding Cellulose Synthase and Fragments Thereof

The present invention relates to polynucleotides which comprise the nucleotide sequence that encodes cellulose synthase of the invention or a functional fragment thereof. In a preferred embodiment, the polynucleotide comprises the sequence encoding a tree cellulose synthase and most preferably, the sequence encoding a cellulose synthase from aspen. In one embodiment, a polynucleotide of the invention includes the entire cellulose synthase coding region, e.g., nucleotides 69 to 3,005 of SEQ ID NO: 1. In another aspect of the invention, the polynucleotide encoding an *Arabidopsis* cellulose synthase is provided (see SEQ ID NO:4 and the translated protein of SEQ ID NO:5).

Also within the scope of the invention are fragments of the polynucleotides encoding cellulose synthase of the invention, which fragments encode at least one transmembrane domain and/or a UDP-glucose binding domain. For example, a polynucleotide comprising the nucleotides encoding a UDP-glucose binding domain of aspen cellulose synthase (e.g., nucleotides 660 to 2250 of SEQ ID NO:1) or corresponding nucleotides of SEQ ID NO:4 are within the scope of the invention. The nucleotides encoding the UDP-glucose binding domain can be determined by, for example, alignment of protein sequences as described below.

The invention further relates to sequence conservative variants of the coding portion of SEQ ID NOS: 1 and 4.

Polynucleotides that hybridize under conditions of low, medium, and high stringency to SEQ ID NOS: 1 and 4, and their respective coding portions are also within the scope of the invention. Preferably, the polynucleotide that hybridizes to any of SEQ ID NOS: 1 and 4, or their respective coding portions, is about the same length as that sequence, for example, not more than about 10 to about 20 nucleotides longer or shorter. In another embodiment of the invention, the hybridizable polynucleotide is at least 1500 nucleotides long, preferably at least 2500 nucleotides long and most preferably at least 3000 nucleotides long. In yet another embodiment, the hybridizable polynucleotide comprises the UDP-glucose binding domain as found in SEQ ID NO: 1 or 4, or at least the conserved region QVLRW [SEQ ID NO: 7]. Most preferably, the hybridizable polynucleotide has a UDP-glucose binding activity.

The polynucleotides that occur originally in nature may be isolated from the organisms that contain them using methods described herein or well known in the art. The non-naturally occurring polynucleotides may be prepared using various manipulations known in the field of recombinant DNA. For example, the cloned CelA polynucleotide can be modified according to methods described by Sambrook et al., 1989. The sequence can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated in vitro. In the production of the modified polynucleotides, for example, care should be taken to ensure that the modified polynucleotide remains within the appropriate translational reading frame (if to be expressed) or uninterrupted by translational stop signals. As a further example, a CelA-encoding nucleic acid sequence can be mutated in vitro or in vivo, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy preexisting ones, to facilitate further in vitro modification. Preferably, such mutations enhance the functional activity of the mutated CelA polynucleotide. Any technique for mutagenesis known in the art can be used, including but not limited to, in vitro site-directed mutagenesis (Hutchinson, C., et al., 1978, J. Biol.

Chem. 253:6551; Zoller and Smith, 1984, DNA 3:479-488; Oliphant et al., 1986, Gene 44:177; Hutchinson et al., 1986, Proc. Natl. Acad. Sci. U.S.A. 83:710), use of TAB linkers (Pharmacia), etc. PCR techniques are preferred for site directed mutagenesis (see Higuchi, 1989, "Using PCR to Engineer DNA", in *PCR Technology: Principles and Applications for DNA Amplification*, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70).

The polynucleotides of the present invention may be introduced into various vectors adapted for plant or non-plant replication. These are well known in the art, thus, choice, construction and use of such vectors is well within the skill of a person skilled in the art. Possible vectors include, but are not limited to, plasmids or modified viruses of plants, but the vector system must be compatible with the host cell used. An example of a suitable vector is Ti plasmid. The insertion into a cloning vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary cohesive termini. However, if the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. An expression cassette containing cellulose synthase or recombinant molecules thereof can be introduced into host cells via silicon carbide whiskers, transformed protoplasts, transformation, e.g., *Agrobacterium* vectors (discussed below), electroporation, infection, etc., so that many copies of the gene sequence are generated. Preferably, the cloned gene is contained on a shuttle vector plasmid, which provides for expansion in a cloning cell, e.g., *E. coli*, and facile purification for subsequent insertion into an appropriate expression cell line, if such is desired. For example, a shuttle vector, which is a vector that can replicate in more than one type of organism, can be prepared for replication in both *E. coli* and *Saccharomyces cerevisiae* by linking sequences from an *E. coli* plasmid with sequences from the yeast 2 μ plasmid.

Transgenic plants containing the polynucleotides described herein are also within the scope of the invention. Methods for introducing exogenous polynucleotides into plant cells and regenerating transgenic plants are well known. Some are provided below.

In one embodiment, to introduce a plasmid containing a CelA coding sequence or promoter of the invention into a plant, a 1:1 mixture of plasmid DNA containing a selectable marker expression cassette and plasmid DNA containing a cellulose synthase expression cassette is precipitated with gold to form microprojectiles. The microprojectiles are rinsed in absolute ethanol and aliquots are dried onto a suitable macrocarrier such as the macrocarrier available from BioRad in Hercules, Calif. Prior to bombardment, embryogenic tissue is preferably desiccated under a sterile laminar-flow hood. The desiccated tissue is transferred to semi-solid proliferation medium. The prepared microprojectiles are accelerated from the macrocarrier into the desiccated target cells using a suitable apparatus such as a BioRad PDS-1000/HE particle gun. In a preferred method, each plate is bombarded once, rotated 180 degrees, and bombarded a second time. Preferred bombardment parameters are 1350 psi rupture disc pressure, 6 mm distance from the rupture disc to macrocarrier (gap distance), 1 cm macrocarrier travel distance, and 10 cm distance from macrocarrier stopping screen to culture plate (macrocarrier travel distance). Tissue is then transferred to semi-solid

proliferation medium containing a selection agent, such as hygromycin B, for two days after bombardment.

Cellulose Synthase Protein and Fragment Thereof

A cellulose synthase of the invention is a plant protein that contains a catalytic subunit which has UDP-glucose binding activity for the synthesis of glucan from glucose, and eight transmembrane domains for localizing the cellulose synthase to the cell membrane. The cellulose synthase of the invention has eight transmembrane binding domains; two at the amino terminal and six at the carboxyl terminal. The UDP-glucose binding domain is located between transmembrane domains two and three. Examples of this protein structure are seen in the aspen cellulose synthase as well as in those of RSW1 and GhCelA. The location of the transmembrane domain may be identified as described below and as exemplified in the Example. Preferably, the cellulose synthase of the invention has an amino acid sequence of a tree cellulose synthase.

In one embodiment, the cellulose synthase protein of the invention is isolated from aspen. Aspen cellulose synthase contains about 978 amino acids and has a molecular weight of about 110 KDa and a pI of about 6.58. In one embodiment, the aspen cellulose synthase has the amino acid sequence of SEQ ID NO:2 as represented in FIG. 1. In another aspect, the invention relates to cellulose synthase of SEQ ID NO: 5.

The invention further relates to fragments of plant cellulose synthases, such as fragments containing at least one transmembrane region and/or a UDP-glucose binding domain. The transmembrane regions may be identified as described in the Example by using the method of Hoffman and Stoffel (1993).

The cellulose synthase fragment containing the UDP-glucose binding domain is functional without the presence of the rest of the protein. This separable activity is as shown in the Example. This result was surprising and unexpected because previously identified UDP-glucose binding domains were not known to be functional when isolated from other portions of the protein. Thus, a fragment of any cellulose synthase (such as PtCelA, RSW1, GhCelA and SEQ ID NO:5) that contains a UDP-glucose binding domain and is independently functional is within the scope of the invention. The function of the UDP-glucose binding domain may be determined using the assay described in the Example. The UDP-glucose binding domain of the invention is located between the second and third transmembrane region of the cellulose synthase and has conserved amino acid sequences for UDP-glucose binding, such as the sequence QVLRW and conserved D residues. The UDP-glucose binding domain and the conserved regions therein may be located in a cellulose synthase using the guidance of the present specification and the general knowledge in the art, for example Brown, 1996. In one embodiment, the UDP-glucose binding domain and the conserved regions therein may be identified by comparing the amino acid sequence of cellulose synthase of interest with the amino acid sequence of aspen cellulose synthase using the algorithms described in the specification or generally known in the art. For example, the UDP-glucose binding domain of SEQ ID NO:2 is in the position amino acids 220 to 749. The conserved QVLRW sequence is located at positions 715-719 of SEQ ID NO:2.

Polypeptides having at least 75%, preferably at least 85% and most preferably at least 95% similarity to the amino acid sequence of SEQ ID NO: 2, amino acids 220-749 of SEQ ID NO:2, SEQ ID NO:5 or its UDP-glucose binding domain using Power Blast or GAP algorithm described above. In a preferred embodiment, these polypeptides are of about the same length as the polypeptide of SEQ ID NO: 2 or amino

acids 220-749 of SEQ ID NO:2. For example, the polypeptide may be from about 2-3 to about 5-7 and to about 10-15 amino acids longer or shorter. In another embodiment, the polypeptides described in this paragraph are not originally found (i.e., naturally occurring) in *Arabidopsis* or cotton. These polypeptides may be prepared by, for example, altering the nucleic acid sequence of a cloned polynucleotide encoding the protein of SEQ ID NO:2 or SEQ ID NO:5 using the methods well known in the art.

Function conservative variants of cellulose synthase are also within the scope of the invention and can be prepared by altering the sequence of a cloned polynucleotide encoding cellulose synthase or fragments thereof. Conventional methods used in the art can be used to make substitutions, additions or deletions in one or more amino acids, to provide functionally equivalent molecules. For example, a function conservative variant that has substitutions, deletions and/or additions in the amino and/or carboxyl terminus of the protein, outside of the UDP-glucose binding domain is within the scope of the invention. Preferably, variants are made that have enhanced or increased functional activity relative to native cellulose synthase. Methods of directed evolution can be used for this purpose.

The invention also includes function conservative variants which include altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a conservative amino acid substitution. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point. Particularly preferred substitutions are: (i) Lys for Arg and vice versa such that a positive charge may be maintained; (ii) Glu for Asp and vice versa such that a negative charge may be maintained; (iii) Ser for Thr such that a free —OH can be maintained; and (iv) Gln for Asn such that a free CONH₂ can be maintained. Amino acid substitutions may also be introduced to substitute an amino acid with a particularly preferable property. For example, a Cys may be introduced a potential site for disulfide bridges with another Cys. A His may be introduced as a particularly "catalytic" site (i.e., His can act as an acid or base and is the most common amino acid in biochemical catalysis). Pro may be introduced because of its particularly planar structure, which induces b-turns in the protein's structure.

The cellulose synthase of the invention can be isolated by expressing a cloned polynucleotide encoding the cellulose synthase as well as using direct protein purification techniques. These methods will be apparent to those of skill in the art.

Polynucleotides Containing Cellulose Synthase Promoter

The present invention further relates to a cellulose synthase promoter. The promoter is a stress-inducible promoter and may be used to synthesize greater quantities of high crystal-

line cellulose in plant, and preferably in trees. This permits an increase in the proportion of cellulose in transgenic plants, greater strength of juvenile wood and fiber, and acceleration of overall growth rate.

In one embodiment, the promoter of the invention is from aspen and is represented in FIG. 4. The promoter sequence is located within the region of nucleotides 1-840 of SEQ ID NO:3. The inventors have identified sub-regions within this region sufficient for promoter function by performing deletion analysis. Thus, functional fragments of SEQ ID NO:3 are within the scope of the invention.

Polynucleotides that hybridize under conditions of low, medium, and high stringency to SEQ ID NO:3, and its non-coding portion are also within the scope of the invention. The hybridizable polynucleotide may be about the same length as the sequence to which it hybridizes, for example, not more than about 10 to about 20 nucleotides longer or shorter. In another embodiment, the hybridizable polynucleotide is at least about 200 nucleotides long, at least about 400 nucleotides long or at least 500 nucleotides long. In yet another embodiment, the hybridizable polynucleotide comprises at least one MSRE element identified according to the method described below.

The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, for example, other plants, or to synthesize polynucleotides having promoter activity. Methods such as PCR, hybridization, synthetic gene construction and the like can be used to identify or generate such sequences based on their sequence homology to the SEQ ID NO:3. Accordingly, sequences isolated or constructed based on their sequence identity to the whole of or any portion of the regions of SEQ ID NO:3 are encompassed by the present invention.

In a PCR approach, primers can be designed to amplify corresponding DNA sequences from genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide sequences present in a population of cloned DNA fragments (i.e., libraries) from a chosen organism. The hybridization probes may be labeled with a detectable group such as ^{32}P , or any other detectable marker. Probes for hybridization can be made by labeling synthetic oligonucleotides based on particular regions of SEQ ID NO:3. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art.

A region of SEQ ID NO:3, for example, a region falling between positions 334 to 838, 334 to 522, 334 to 422, 423 to 838 or 423 to 522 of SEQ ID NO:3, may be used as a probe capable of specifically hybridizing to corresponding sequences. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique among the sequences to be screened and are suitably at least about 10 nucleotides in length, at least about 20 nucleotides, at least about 25 nucleotides, at least about 30 nucleotides, at least about 35 nucleotides, at least about 40 nucleotides, at least about 50 nucleotides or at least about 60 nucleotides in length. Such sequences may alternatively be used to amplify corresponding sequences from a chosen plant by PCR. This technique may be used to isolate sequences from a desired plant or as a diagnostic assay to determine the presence of sequences in a plant. Hybridization techniques may include hybridization screening of DNA libraries plated as either plaques or colonies.

Hybridization of such sequences may be carried out under stringent conditions. By "stringent conditions" or "stringent hybridization conditions" is intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing).

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulfate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl and 0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1.0 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60° C. to 65° C.

Specificity is also the function of post-hybridization washes, with factors including the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation $T_m = 81.5^\circ\text{C.} + 16.6(\log M) + 0.41(\% \text{GC}) - 0.61(\% \text{form.}) - 500/L$, where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form. is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted for sequences of the desired identity to hybridize. For example, if sequences with 90% identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4° C. lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6° C., 7° C., 8° C., 9° C., or 11° C. lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11° C. to 20° C. lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution), the SSC concentration may be increased so that a higher temperature can be used.

Thus, isolated sequences that have promoter activity and which hybridize under stringent conditions to a region falling between positions 324 to 838, 324 to 522, 324 to 423, 423 to

838 or 423 to 522 of SEQ ID NO:3, or to fragments thereof, are encompassed by the present invention. Sequences may be suitably at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to sequences from positions 334 to 838, positions 334 to 522, positions 334 to 423, positions 423 to 838 or positions 423 to 522 of SEQ ID NO:3. The promoter regions of the invention may be used to isolate substantially identical sequences from any plant species, including, but not limited to, any plant species described herein.

Suitably the promoter sequences comprise any sequence of at least 30 consecutive nucleotides from position 334 to 522 of SEQ ID NO:3 having promoter activity. Suitably, the promoter sequences may comprise at least 40 consecutive nucleotides, at least 45 consecutive nucleotides, at least 50 consecutive nucleotides, at least 55 consecutive nucleotides, at least 60 consecutive nucleotides, at least 65 consecutive nucleotides, at least 70 consecutive nucleotides, at least 75 consecutive nucleotides, or at least 80 consecutive nucleotides from position 334 to 522 of SEQ ID NO:3 having promoter activity. Suitably the promoter sequences may show at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to the consecutive nucleotide sequences falling between positions 334 and 522 of SEQ ID NO:3.

Suitably, the promoter sequences may be operably linked to a coding sequence, and the coding sequence may be suitably expressed in a cell, including a plant cell. The coding sequence may suitably encode a polypeptide, including, but not limited to, reporter polypeptides, such as fluorescent or luminescent polypeptides and antibiotics, structural polypeptides, such as cell wall or cell membrane polypeptides, enzymes, such as enzymes involved in the metabolism of cellulose, lignin, starch, sucrose, nitrogen and secondary metabolites, and immuno-polypeptides, such as antigens and antibodies.

A cellulose synthase promoter of the invention typically provides tissue-specific gene regulation in xylem, but also permits up-regulation of gene expression in other tissues as well, e.g., phloem under tension stress. Furthermore, expression of cellulose synthase is localized to an area of the plant under stress.

This stress-inducible phenomenon is regulated by positive and negative mechanical stress response elements (MSREs). These MSREs upregulate (positive) or downregulate (negative) the expression of a cellulose synthase polynucleotide under stress conditions through binding of transcription factors. MSRE-regulated expression of cellulose synthase permits synthesis of cellulose with high crystallinity.

The MSREs of cellulose synthase can be modified or employed otherwise in methods to regulate expression of a polynucleotide, including a cellulose synthase, operatively linked to a promoter containing an MSRE in response to mechanical stress (e.g., tension or compression) to a transgenic plant.

Negative MSREs of a cellulose synthase promoter can be modified, removed or blocked to improve expression of a cellulose synthase, and thereby increase cellulose production and improve wood quality. Alternatively, positive MSREs can be removed or blocked to decrease expression of a cellulose synthase, which decreases cellulose production and increases lignin deposition. This is useful for increasing the fuel value of wood because lignin has a higher BTU value than cellulose. Moreover, a modified cellulose synthase promoter can be operatively linked to a polynucleotide of interest to control its expression upon mechanical stress to a plant harboring it.

The location of MSRE elements in the SEQ ID NO:3 may be identified, for example, using promoter deletion analysis, DNase Foot Print Analysis, and Southwestern screening of an expression library for an MSRE. In one embodiment, cellulose synthase promoter that has one or more portions deleted, and is operatively linked to a reporter sequence, is introduced into a plant or a plant cell. A positive MSRE is detected by observing no relative change or increase in the amount of reporter in a transgenic plant or tissue, e.g., phloem after inducing a stress to the plant, and a negative MSRE is detected by observing increases in the amount of reporter in the plant in the absence of any stress to the plant. A positive element is detected when by removing it, GUS expression goes down and by adding it kept at the same level or more. The negative element does not support, or suppress, expression of GUS and by removing it, normal or enhanced GUS expression is observed as compared to when negative element is present.

Manipulation of a MSRE binding sites and/or providing transcription factors that bind thereto, provides a mechanism to continuously produce high crystalline cellulose in woody plant cell walls of transgenic plants. For example, one having ordinary skill in the art can delete or block negative MSRE elements, or provide cDNA encoding protein(s) that bind the positive MSREs, to enable constitutive expression of a cellulose synthase without the requirement of a mechanical stress. The increased cellulose synthase, and therefore, increased cellulose content, can improve the strength properties of juvenile wood and fiber. It is also contemplated that the positive MSREs can be deleted or blocked, or cDNA in an antisense direction, which in the sense direction encodes a protein that binds a positive MSRE, can be provided, to reduce cellulose synthase activity and decrease cellulose production.

Method of Isolating Polynucleotides Encoding Cellulose Synthase

The invention further relates to identifying and isolating polynucleotides encoding cellulose synthase in plants, e.g., trees, (in addition to those polynucleotides provided in the Example and represented in FIG. 1 and FIG. 7). These polynucleotides may be used to manipulate expression of cellulose synthase with an objective to improve the cellulose content and properties of wood.

The method comprises identifying a nucleic acid fragment containing a sequence encoding cellulose synthase or a portion thereof by using a fragment of SEQ ID NOS: 1 or 4 as a probe or a primer. Once identified, the nucleic acid fragment containing a sequence encoding cellulose synthase or a portion thereof is isolated.

Polynucleotides encoding cellulose synthases of the invention, whether genomic DNA, cDNA, or fragments thereof, can be isolated from many sources, particularly from cDNA or genomic libraries from plants, preferably trees (e.g. aspen, sweetgum, loblolly pine, eucalyptus, and other angiosperms and gymnosperms). Molecular biology methods for obtaining polynucleotides encoding a cellulose synthase are well known in the art, as described above (see, e.g., Sambrook et al., 1989, supra).

Accordingly, cells from any species of plant can potentially serve as a nucleic acid source for the molecular cloning of a polynucleotide encoding a cellulose synthase of the invention. The DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a DNA "library"), and preferably is obtained from a cDNA library prepared from tissues with high level expression of a cellulose synthase (e.g., xylem tissue, since cells in this tissue evidence very high levels of expression of CelA), by chemical synthesis, by

cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from a desired cell (see, for example, Sambrook et al., 1989, supra; Glover, D. M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will not contain intron sequences. Whatever the source, a polynucleotide should be molecularly cloned into a suitable vector for its propagation.

In another embodiment for the molecular cloning of a polynucleotide encoding a cellulose synthase of the invention from genomic DNA, DNA fragments are generated from a genome of interest, such as from a plant, or more particularly a tree genome, part of which will correspond to a desired polynucleotide. The DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, as for example, by sonication. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing a desired CelA sequence may be accomplished in a number of ways. For example, if an amount of a portion of a CelA sequence or its specific RNA, or a fragment thereof, is available and can be purified and labeled, the generated DNA fragments may be screened by nucleic acid hybridization to a labeled probe (Benton and Davis, 1977, Science 196:180; Grunstein and Hogness, 1975, Proc. Natl. Acad. Sci. U.S.A. 72:3961). For example, a set of oligonucleotides corresponding to the partial amino acid sequence information obtained for a CelA protein from trees can be prepared and used as probes for DNA encoding cellulose synthase, or as primers for cDNA or mRNA (e.g., in combination with a poly-T primer for RT-PCR). Preferably, a fragment is selected that is highly unique to a cellulose synthase of the invention, such as the UDP-glucose binding regions. Those DNA fragments with substantial homology to the probe will hybridize. As noted above, the greater the degree of homology, the more stringent hybridization conditions can be used. In a specific embodiment, stringency hybridization conditions can be used to identify homologous CelA sequences from trees or other plants.

Thus, in one embodiment, a labeled cellulose synthase cDNA from, e.g., *Populus tremuloides* (PtCelA), can be used to probe a library of genes or DNA fragments from various species of plants, especially angiosperm and gymnosperm, to determine whether any bind to a CelA of the invention. Once genes or fragments are identified, they can be amplified using standard PCR techniques, cloned into a vector, e.g., pBlue-script vector (StrataGene of LaJolla, Calif.), and transformed into a bacteria, e.g., DH5 α *E. coli* strain (Gibco BRL of Gaithersburg, Md.). Bacterial colonies are typically tested to determine whether any contains a cellulose synthase-encoding nucleic acid. Once a positive clone is identified through binding, it is sequenced from an end, preferably the 3' end.

cDNA libraries can be constructed in various hosts, such as lambda ZAPII, available from Stratagene, LaJolla, Calif., using poly(A) + RNA isolated from aspen xylem, according to the methods described by Bugos et al. (Biotechniques 19:734-737, 1995). The above mentioned probes are used to assay the aspen cDNA library to locate cDNA which codes for enzymes involved in production of cellulose synthases.

Once a cellulose synthase sequence is located, it is then cloned and sequenced according to known methods in the art.

Further selection can be carried out on the basis of the properties of the gene, e.g., if the gene encodes a protein product having the isoelectric, electrophoretic, hydrophathy plot, amino acid composition, or partial amino acid sequence of a cellulose synthase protein of the invention, as described herein. Thus, the presence of the gene may be detected by assays based on the physical, chemical, or immunological properties of its expressed product. For example, cDNA clones or DNA clones which hybrid-select the proper mRNAs can be used to produce a protein that has similar properties known for cellulose synthases of the invention. Such properties may include, for example, similar or identical electrophoretic migration patterns, isoelectric focusing or non-equilibrium pH gel electrophoresis behavior, proteolytic digestion maps, hydrophathy plots, or functional properties (such as isolated, functional UDP-glucose binding domains).

A cellulose synthase polynucleotide of the invention can also be identified by mRNA selection, i.e., by nucleic acid hybridization followed by in vitro translation. In this procedure, nucleotide fragments are used to isolate complementary mRNAs by hybridization. Such DNA fragments may represent available, purified CelA DNA, or may be synthetic oligonucleotides designed from the partial amino acid sequence information. Functional assays (e.g., UDP-glucose activity) of the in vitro translation products of the products of the isolated mRNAs identifies the mRNA and, therefore, the complementary DNA fragments, that contain the desired sequences.

A radiolabeled CelA cDNA can be synthesized using a selected mRNA as a template. The radiolabeled mRNA or cDNA may then be used as a probe to identify homologous CelA DNA fragments from amongst other genomic DNA fragments.

It will be appreciated that other polynucleotides, in addition to a CelA of the invention can be operatively linked to a CelA promoter to control expression of the polynucleotide upon application of a mechanical stress.

Expression of CelA Polypeptides

The nucleotide sequence coding for CelA, or a functional fragment, derivative or analog thereof, including chimeric proteins, can be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. Preferably, an expression vector includes an origin of replication. The elements are collectively termed herein a "promoter." Thus, a nucleic acid encoding CelA of the invention can be operatively associated with a promoter in an expression vector of the invention. Both cDNA and genomic sequences can be cloned and expressed under control of such regulatory sequences. The necessary transcriptional and translational signals can be provided on a recombinant expression vector, or they may be supplied by the native gene encoding CelA and/or its flanking regions.

In addition to a CelAP, expression of cellulose synthase can be controlled by any promoter/enhancer element known in the art, but these regulatory elements must be functional in the host selected for expression. Promoters which may be used to control CelA polynucleotide expression include, constitutive, development-specific and tissue-specific. Examples of these promoters include 35S Cauliflower Mosaic Virus, terminal flower and 4CL-1. Thus, there are various ways to alter the growth of a plant using different promoters, depending on the needs of the practitioner.

The nucleotide sequence may be inserted in a sense or antisense direction depending on the needs of the practitioner. For example, if augmentation of cellulose biosynthesis is desired then polynucleotides encoding, e.g., cellulose synthase, can be inserted into the expression vector in the sense direction to increase cellulose synthase production and thus cellulose biosynthesis. Alternatively, if it is desired that cellulose biosynthesis is reduced or lignin content is increased, then polynucleotides encoding, e.g., cellulose synthase, can be inserted in the antisense direction so that upon transcription the antisense mRNA hybridizes to other complementary transcripts in the sense orientation to prevent translation. In other embodiments, the polynucleotide encodes a UDP-glucose binding domain and is used in a similar manner as described.

A recombinant CelA protein of the invention, or functional fragment, derivative, chimeric construct, or analog thereof, may be expressed chromosomally, after integration of the coding sequence by recombination. In this regard, any of a number of amplification systems for plants may be used to achieve high levels of stable gene expression, as discussed above. Any of the methods previously described for the insertion of DNA fragments into a cloning vector may be used to construct expression vectors containing a gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and synthetic techniques and in vivo recombination (genetic recombination).

Expression vectors containing a nucleic acid encoding a CelA of the invention can be identified by four general approaches: (a) PCR amplification of the desired plasmid DNA or specific mRNA, (b) nucleic acid hybridization, (c) presence or absence of selection marker gene functions, (d) analyses with appropriate restriction endonucleases, and (e) expression of inserted sequences. In the first approach, the nucleic acids can be amplified by PCR to provide for detection of the amplified product. In the second approach, the presence of a foreign gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted marker gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "selection marker" gene functions (e.g., β -glucuronidase activity, resistance to antibiotics, transformation phenotype, etc.) caused by the insertion of foreign genes in the vector. In another example, if the nucleic acid encoding CelA is inserted within the "selection marker" gene sequence of the vector, recombinants containing the CelA insert can be identified by the absence of the CelA gene function. In the fourth approach, recombinant expression vectors are identified by digestion with appropriate restriction enzymes. In the fifth approach, recombinant expression vectors can be identified by assaying for the activity, biochemical, or immunological characteristics of the gene product expressed by the recombinant, provided that the expressed protein assumes a functionally active conformation.

After a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to those vectors or their derivatives described above.

Vectors are introduced into the desired host cells by methods known in the art, e.g., *Agrobacterium*-mediated transformation (described in greater detail below), transfection, elec-

trporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; Hartmut et al., Canadian Patent Application No. 2,012,311, filed Mar. 15, 1990).

The cell into which the recombinant vector comprising the nucleic acid encoding CelA is cultured in an appropriate cell culture medium under conditions that provide for expression of CelA by the cell. In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (such as glycosylation, cleavage, e.g., of a signal sequence) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed.

Agrobacterium-Mediated Transformation and Inducing Somatic Embryos

The culture media used in the invention, and for transforming *Agrobacterium*, contain an effective amount of each of the medium components (e.g. basal medium, growth regulator, carbon source) described above. As used in describing the present invention, an "effective amount" of a given medium component is the amount necessary to cause a recited effect. For example, an effective amount of a growth hormone in the primary callus growth medium is the amount of the growth hormone that induces callus formation when combined with other medium components. Other compounds known to be useful for tissue culture media, such as vitamins and gelling agents, may also be used as optional components of the culture media of the invention.

Transformation of cells from plants, e.g., trees, and the subsequent production of transgenic plants using *Agrobacterium*-mediated transformation procedures known in the art, and further described herein, is one example of a method for introducing a foreign gene into trees. Transgenic plants may be produced by various methods, such as by the following steps: (i) culturing *Agrobacterium* in low-pH induction medium at low temperature and preconditioning, i.e., coculturing bacteria with wounded tobacco leaf extract in order to induce a high level of expression of the *Agrobacterium* vir genes whose products are involved in the T-DNA transfer; (ii) coculturing a desired plant tissue explants, including zygotic and/or somatic embryo tissues derived from cultured explants, with the incited *Agrobacterium*; (iii) selecting transformed callus tissue on a medium containing antibiotics; and (v) and converting the embryos into plantlets.

Any non-tumorigenic *A. tumefaciens* strain harboring a disabled Ti plasmid may be used in the method of the invention. Any *Agrobacterium* system may be used. For example, Ti plasmid/binary vector system or a cointegrative vector system with one Ti plasmid may be used. Also, any marker gene or polynucleotide conferring the ability to select transformed cells, callus, embryos or plants and any other gene, such as, for example, a gene conferring resistance to a disease, or one improving cellulose content, may also be used. Any promoter desired can be used, such as, for example, a PtCelA of the invention, and those promoters as described above. A person of ordinary skill in the art can determine which markers and genes are used depending on particular needs.

For purposes of the present invention, "transformed" or "transgenic" means that at least one marker gene or poly-

nucleotide conferring selectable marker properties is introduced into the DNA of a plant cell, callus, embryo or plant. Additionally, any gene may also be introduced.

To increase the infectivity of the bacteria, *Agrobacterium* is cultured in low-pH induction medium, i.e., any bacterium culture media with a pH value adjusted to from 4.5 to 6.0, most preferably about 5.2, and at low temperature such as for example about 19-30° C., preferably about 21-26° C. The conditions of low-pH and low temperature are among the well-defined factors for inducing virulence activity in *Agrobacterium* (e.g., Altmore et al., Mol. Plant-Microbe. Interac. 2: 301, 1989; Fullner et al., Science 273: 1107, 1996; Fullner and Nester, J. Bacteriol. 178: 1498, 1996).

The bacteria is preconditioned by coculturing with wounded tobacco leaf extract (prepared according to methods known generally known in the art) to induce a high level of expression of the *Agrobacterium* vir genes. Prior to inoculation of plant somatic embryos, *Agrobacterium* cells can be treated with a tobacco extract prepared from wounded leaf tissues of tobacco plants grown in vitro. To achieve optimal stimulation of the expression of *Agrobacterium* vir genes by wound-induced metabolites and other cellular factors, tobacco leaves can be wounded and pre-cultured overnight. Culturing of bacteria in low pH medium and at low temperature can be used to further enhance the bacteria vir gene expression and infectivity. Preconditioning with tobacco extract and the vir genes involved in the T-DNA transfer process are generally known in the art.

Agrobacterium treated as described above is then cocultured with a plant tissue explant, such as for example zygotic and/or somatic embryo tissue. Non-zygotic (i.e., somatic) or zygotic tissues can be used. Any plant tissue may be used as a source of explants. For example, cotyledons from seeds, young leaf tissue, root tissues, parts of stems including nodal explants, and tissues from primary somatic embryos such as the root axis may be used. Generally, young tissues are a preferred source of explants.

The invention also relates to methods of altering the growth of a plant by expressing the polynucleotide of the invention, which as a result alters the growth of the plant. The polynucleotide used in the method may be a homologous polynucleotide or a heterologous polynucleotide and are described in detail above. For example, both full-length and UDP-glucose binding region containing fragments may be expressed. Additionally, depending on the aim of the method, the polynucleotide may be introduced into the plant in the sense or in the antisense orientation. Any suitable promoter may be used to provide expression. The promoter or a functional fragment thereof is operatively linked to the polynucleotide. The promoter may be a constitutive promoter, a tissue-specific promoter or a development-specific plant promoter. Examples of suitable promoters are Cauliflower Mosaic Virus 35S, 4CL, cellulose synthase promoter, PtCelAP and terminal flower promoter.

The invention further relates to a method of altering the cellulose content in a plant by expressing the polynucleotide of the invention as described above. The method may be used to increased the ratio of cellulose to lignin in the plant that have an exogenous polynucleotide of the invention introduced therein.

The invention further relates to a method for altering expression of a cellulose synthase in a plant cell by introducing into the cell a vector comprising a polynucleotide of the invention and expressing the polynucleotide. The polynucleotides and promoters described above may be used.

A method for causing stress-induced gene expression in a plant cell is also within the scope of the invention. The method

comprises (i) introducing into the plant or a plant cell an expression cassette comprising a cellulose synthase promoter or a functional fragment thereof or providing a plant or a plant cell that comprises the expression cassette (The promoter of the cassette is operatively linked to a coding sequence of choice.); and (ii) applying mechanical stress to the plant to induce expression of the desired coding sequence.

A method for determining a positive mechanical stress responsive element (MSRE) in a cellulose synthase promoter is also within the scope of the invention and comprises (i) making serial deletions in the cellulose synthase promoter, such as for example, SEQ ID NO:3; (ii) introducing the deletion linked to a polynucleotide encoding a reporter sequence into a plant cell, and (iii) detecting a decrease in the amount of reporter in the plant after inducing a stress to the plant. Similarly, a method for determining a negative MSRE in a cellulose synthase promoter is provided. It comprises (i) making serial deletions in the cellulose synthase promoter, such as for example, SEQ ID NO:3; (ii) introducing the deletion linked to a polynucleotide encoding a reporter sequence into a plant cell, and (iii) detecting an increase in the amount of reporter in the plant after inducing a stress to the plant.

The following methods are also within the scope of the invention: a method for expressing cellulose synthase in a tissue-specific manner comprising transforming a plant with a tissue specific promoter operatively linked to a polynucleotide encoding a cellulose synthase; a method for inducing expression of a cellulose synthase in a plant comprising introducing into a plant a cDNA encoding a protein that binds to a positive MSRE of a cellulose synthase promoter, thereby resulting in increased expression of cellulose in the plant, wherein the binding to the positive MSRE results in expression of a cellulose synthase; a method for reducing expression of a cellulose synthase comprising introducing into a plant a cDNA in an antisense orientation, wherein the cDNA in a sense orientation encodes a protein that binds to a positive MSRE and results in expression of a cellulose synthase; a method for increasing cellulose biosynthesis in a plant comprising introducing into a plant a cDNA encoding a protein that binds to a positive MSRE of a cellulose synthase promoter, whereby binding of the protein to the positive MSRE results in expression of a cellulose synthase, and a method for reducing cellulose biosynthesis in a plant comprising introducing into a plant a cDNA in an antisense orientation, wherein the cDNA in a sense orientation encodes a protein that binds to a positive MSRE of a cellulose synthase promoter.

EXAMPLE

Molecular Cloning of Cellulose Synthase

This Example describes the first tree cellulose synthase cDNA (PtCelA, GenBank No. AF072131) cloned from developing secondary xylem of aspen trees using RSW1 cDNA.

Prior to the present invention, only partial clones of cellulose synthases from crop species and cotton GhCelA have been discovered, which have significant homology to each other. The present inventors have discovered and cloned a new full-length cellulose synthase cDNA, AraxCelA (GenBank No. AF062485) (FIG. 7, [SEQ ID NO: 4]), from an *Arabidopsis* primary library. AraxCelA is a new member of cellulose synthase and shows 63-85% identity and 72-90% similarity in amino acid sequence with other *Arabidopsis* CelA members.

Another cellulose synthase was cloned in aspen using a ³²P-labeled 1651-bp long EcoRI fragment of *Arabidopsis* CelsA cDNA, which encodes a centrally located UDP-glucose binding domain, was used as a probe to screen about 500,000 pfu of a developing xylem cDNA library from aspen (*Populus tremuloides*) (Ge and Chiang, 1996). Four positive clones were obtained after three rounds of plaque purification. Sequencing the 3' ends of these four cDNAs showed that they were identical clones. The longest cDNA clone was fully sequenced and determined to be a full-length cDNA having a 3232 bp nucleotide sequence (FIG. 1) [SEQ ID NO: 1], which encodes a protein of 978 amino acids [SEQ ID NO: 2].

Characterization of a Cellulose Synthase from Aspen

The first AUG codon of PtCelsA was in the optimum context for initiation of transcription on the basis of optimal context sequence described by Joshi (1987a) and Joshi et al. (1997). A putative polyadenylation signal (AATACA) was found 16 bp upstream of a polyadenylated tail of 28 bp, which is similar to the proposed plant structure (Joshi, 1987b). The 5' untranslated leader was determined to have 68 bp and the 3' untranslated trailer was 227 bp. Both of these regions have a typical length observed in many plant genes (Joshi, 1987a and Joshi, 1987b). This cDNA clone exhibited 90% amino acid sequence similarity with cellulose synthase from cotton (Gh-CelsA) and 71% with cellulose synthase from *Arabidopsis* (RSW1), suggesting that this particular tree homolog also encodes a cellulose synthase.

The full length cDNA was designated PtCelsA, and encodes a 110,278 Da polypeptide having an isoelectric point (pI) of 6.58 and 8 charged molecules. The hydropathy curve indicated that this particular cellulose synthase has eight transmembrane binding domains; two at the amino terminal and six at the carboxyl terminal, using the method of Hoffman and Stoffel (1993). This protein structure is analogous to those of RSW1 and GhCelsA. All of the conserved domains for UDP-glucose binding, such as QVLRW and conserved D residues, are also present in a cellulose synthase of the invention, e.g., PtCelsA (Brown et al., 1996). Thus, based on sequence and molecular analyses, it was concluded that PtCelsA encodes a catalytic subunit which, like RSW1 in *Arabidopsis*, is essential for the cellulose biosynthesis machinery in aspen.

In situ localization of PtCelsA mRNA transcripts along the developmental gradient defined by stem primary and secondary growth demonstrated that cellulose synthase expression is confined exclusively to developing xylem cells undergoing secondary wall thickening. This cell-type-specific nature of PtCelsA gene expression was also consistent with xylem-specific activity of cellulose synthase promoter (PtCelsAP) based on heterologous promoter-β-glucuronidase (GUS) fusion analysis. Overall, the results provide several lines of evidence that cellulose synthase is the gene primarily responsible for cellulose biosynthesis during secondary wall formation in woody xylem of trees, such as aspen. Previous results by the inventors (Hu et al., 1999) showed that cellulose and lignin are deposited in a compensatory fashion in wood. The discovery of a cellulose synthase in trees, such as aspen, permits the up-regulation of the protein to elevate cellulose production. Surprisingly, expression of CelsA in trees suppressed lignin biosynthesis to further improve wood properties of trees.

Preparation of Transgenic Plants

The UDP-glucose binding sequence was subcloned into pBI121, which was used to prepare transgenic tobacco plants

(Hu et al., 1998). The expression of a heterologous UDP-glucose binding sequence resulted in a remarkable growth-accelerating effect. This was surprising because current knowledge of the function of plant cellulose synthases teaches that a UDP-glucose sequence must remain intact with other functional domains in CelsA, e.g., the transmembrane domains, in order for cellulose synthase to initiate cellulose biosynthesis. The remarkable growth and tremendous increase in plant biomass observed in transgenic tobacco was due likely to an augmented deposition of cellulose, indicating that the UDP-glucose domain alone is sufficient for genetic augmentation of cellulose biosynthesis in plants.

Genome Organization and Expression of a Novel Cellulose Synthase

To confirm that the cDNA clone of FIG. 1 [SEQ ID No: 1] was a cellulose synthase, genomic Southern blot analysis was performed under both high and low stringency conditions using the cDNA. Genomic DNA (25 μg per lane) from aspen was digested with PstI (lane P), HindIII (lane H) and EcoRI (lane E), and probed using a 1 kb ³²P-labeled fragment from the 5' end of a cellulose synthase of FIG. 1. The Southern blot suggested the presence of a small family of cellulose synthase genes in aspen genome (FIG. 2, panels a and b). Repeated screening of the aspen xylem cDNA library with various plant CelsA gene-related probes always resulted in the isolation of the same cellulose synthase cDNA clone. This suggested that the cellulose synthase cDNA cloned (FIG. 1) [SEQ ID NO: 1], represents the primary and most abundant cellulose synthase-encoding gene in developing xylem of trees, such as aspen, where active cellulose deposition takes place. It also indicates that manipulation of cellulose synthase gene expression can have a profound influence on cellulose biosynthesis in trees.

In Situ Hybridization

Northern blot analysis of total RNA (40 μg per lane) from the 1st and 2nd internodes (lane 1), 3rd and 4th internodes (lane 2), 5th and 6th internodes (lane 3), and 9th and 10th internodes (lane 4) of aspen seedling stems (FIG. 2, panel c) using the labeled probe (as described above) revealed the near absence of cellulose synthase transcripts in tissues undergoing primary growth (internodes 1 to 4), and that the presence of cellulose synthase transcripts occurs during the secondary growth of stem tissues (internodes 5 to 11). However, weak northern signals in primary growth may only suggest that cellulose synthase gene expression is specific to xylem, of which there is little in primary growth tissue.

Xylogenesis in higher plants offers a unique model that involves sequential execution of cambium cell division, commitment to xylem cell differentiation, and culmination in xylem cell death (Fukuda, 1996). Although primary and secondary xylem cells originate from different types of cambia, namely procambium and inter/intrafascicular cambium, both exhibit conspicuous secondary wall development with massive cellulose and lignin deposition (Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons). To further investigate spatial and temporal cellulose synthase gene expression patterns at the cellular level, in situ hybridization was used to localize cellulose synthase mRNA along the developmental gradient defined by stem primary and secondary growth.

Localization of cellulose synthase gene transcripts (RNA) in stem at various growth stages was also observed. FIG. 3 shows transverse sections from 2nd (panel a), 4th (panel b),

and 6th (panel c) internodes hybridized with digoxigenin (DIG)-labeled cellulose synthase antisense or sense (control, from the 5th internode, panel d) RNA probes, as described. Positive RNA-RNA hybridization signals were stained. In FIG. 3, arrows indicate the cellular localization of PtCelA transcripts. No hybrid signal was detectable in the control section.

PtCelA transcripts were detected in young aspen stem sections by in situ hybridization with transcripts of highly variable 5' region of PtCelA cDNA (a 771 bp long fragment generated from PstI and SacI). This region was first subcloned in the plasmid vector, pGEM, -3Zf (+) (Promega) for the production of digoxigenin (DIG)-labeled transcripts using T7 (for antisense transcripts) and SP6 (for sense transcripts) RNA polymerase (DIG system: Boehringer Mannheim). Probes were subjected to mild alkaline hydrolysis by incubation in 100 mM NaHCO₃, pH 10.2 at 60° C., which produced approximately 200 bp fragments.

Aspen young stems were prepared for sectioning by fixation in 4% (w/v) paraformaldehyde in 100 mM phosphate buffer (pH 7.0) at 4° C. overnight, dehydrated through an ethanol series on ice, and embedded in Paraplast medium (Sigma). Ten µm sections (bars=100 µm in FIG. 3) were mounted on Superfrost/plus (Fisher) slides at 42° C. overnight, dewaxed and then rehydrated through a descending ethanol series. The sections were incubated with proteinase K (10 µg/ml in 100 mM Tris-HCl, 50 mM EDTA, pH 7.5) for 30 min and were post-fixed with FAA. The sections were acetylated with 0.33% (v/v) acetic anhydride in 0.1 M triethanolamine-HCl (pH 8.0) prior to hybridization. The sections were then incubated in a hybridization mixture (approximately 2 µg/ml DIG-labeled probes, 50% (v/v) formamide, 2×SSPE, 10% (w/v) dextran sulfate, 125 µg/ml tRNA, pH 7.5) at 45° C. for 12-16 hrs. Nonhybridized single-stranded RNA probe was removed by treatment with 20 µg/ml RNase A in TE buffer with 500 mM NaCl. The sections were washed at 50° C. Hybridized DIG-labelled probe was detected on sections using anti-digoxigenin antiserum at a 1:1500 dilution, as described in the manufacturer's instruction (DIG system: Boehringer Mannheim). Sections were examined by Eclipse 400 light microscope (Nikon) and photographed.

During the primary growth stage (FIG. 3, panels a and b), strong expression of cellulose synthase was found localized exclusively to primary xylem (PX) cells. At this stage, young internodes are elongating, resulting in thickening of primary xylem cells through formation of secondary walls (Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons). The concurrence of shoot elongation with high expression of cellulose synthase strongly suggests the association of cellulose synthase protein with secondary cell wall cellulose synthesis. Later stages of primary growth (FIG. 3, panel b) are characterized by the appearance of an orderly alignment of primary xylem cells. Active cellulose biosynthesis accompanies cell elongation-induced wall thickening, as indicated by the strong expression of cellulose synthase in these primary xylem cells.

At the beginning of secondary growth in older internodes, it was observed that expression of cellulose synthase is also exclusively localized to xylem cells (FIG. 3, panel c). Instead of elongation in internodes distal to the meristematic activity, growth at this stage is mainly radial due to thickening in secondary cell walls of secondary xylem. At the same time, expression of PtCelA gene becomes localized to the secondary developing xylem cells (SX in FIG. 3, panel c), which is again consistent with the idea that PtCelA encodes a secondary cell wall cellulose synthase. At this stage, secondary xylem cells cover the elongated and differentiated primary

xylem cells in which PtCelA gene expression is no longer detectable (FIG. 3, panel c). These results demonstrate that expression of PtCelA gene is xylem-specific and the cellulose synthase of FIG. 1 [SEQ ID NO: 1] encodes a cellulose synthase associated with cellulose biosynthesis in secondary walls of xylem cells. To further confirm xylem-specific expression of cellulose synthase, a cellulose synthase gene promoter sequence was cloned and characterized for regulatory activities.

Characterization of Expression Regulated by Cellulose Synthase Promoter

A 5' 1,200 bp cDNA fragment of a cellulose synthase of FIG. 1 [SEQ ID NO: 1] was used as a probe to screen an aspen genomic library for 5' regulatory sequences of a novel cellulose synthase gene, PtCelA. The library was constructed by cloning aspen genomic DNA fragments, generated from an Sau3AI partial-digest and sucrose gradient-selected, into the BamHI site of a Lambda DASH II vector (Stratagene, La Jolla, Calif.). Five positive clones were obtained from about 150,000 pfu and Lambda DNA was purified. One clone having about a 20 kb DNA insert size was selected for restriction mapping and partial sequencing. This resulted in the identification of a 5' flanking region of PtCelA gene of approximately 1 kb. This genomic fragment, designated PtCelAP (FIG. 4) [SEQ ID NO: 3], contained about 800 bp of promoter sequence, 68 bp of 5' end untranslated region and 160 bp of coding sequence. To investigate regulation of tissue-specific cellulose synthase expression at the cellular level, promoter activity was analyzed in transgenic tobacco plants by histochemical staining of a GUS protein. A PtCelAP-GUS fusion binary vector was constructed in pBI121 with the 35S promoter replaced with PtCelAP [SEQ ID NO: 3] and introduced into tobacco (*Nicotiana tabacum*) as per Hu et al. (1998).

Eleven independent transgenic lines harboring a CelAP-GUS fusion were generated. FIG. 5 shows a histochemical analysis of GUS expression driven by a cellulose synthase promoter of the invention in transgenic tobacco plants. Transverse sections from the 3rd (panel a), 5th (panel b), 7th (panel c), and 8th (panels d and f) internodes were stained for GUS activity, and fluorescence microscopy (panel e, 8th internode from panel d) was used to visualize expression under UV radiation (bar=100 µm in a, b, c, d, e; bar=1.5 mm in f).

GUS staining was detected exclusively in xylem tissue of stems, roots and petioles. In stems, strong GUS activity was found localized to xylem cells undergoing primary (FIG. 5, panel a) and secondary growth (FIG. 5 panels b-d and f). GUS expression was confined to xylem cells in the primary growth stage and became more localized in developing secondary xylem cells during secondary growth. An entire section from the 8th internode stained for GUS activity (FIG. 5, panel f). These results are consistent with the in vivo expression patterns of cellulose synthase in aspen stems. Lignin autofluorescence was visualized after UV radiation. Phloem fibers, which are also active in cellulose and lignin biosynthesis (FIG. 5, panels d and e), did not show GUS activity, suggesting that cellulose synthase gene expression is not associated with cellulose biosynthesis in cell types other than xylem. Examination of GUS activity in roots, stems, leaves, anthers and fruit also showed GUS expression in xylem tissue of all these organs suggesting that cellulose synthases of the invention are xylem-specific cellulose and expressed in all plant organs.

Characterization of promoter activity and cellular expression of a cellulose synthase of the invention from one particu-

lar source (aspen) indicated that expression produces a protein that encodes a secondary cell wall-specific cellulose synthase and is specifically compartmentalized in developing xylem cells. Characterization of the cellulose synthase gene promoter sequence not only confirms cell type-specific expression of cellulose synthase, but also provides a method for over-expressing cellulose synthase in a tissue-specific manner to augment cellulose production in xylem.

Expression of Cellulose Synthase Under Tension Stress

As described earlier, a cellulose synthase promoter of the invention is involved in a novel gene regulatory phenomenon of cellulose synthase. To further characterize a cellulose synthase of the invention, GUS expression driven by an aspen cellulose synthase promoter (PtCelAP) was observed in transgenic tobacco plants without or under tension stress. The stress was induced by bending and affixing the plants to maintain the bent position (e.g., tying) over a 40 hour period. Tangential and longitudinal sections were taken before bending, and 4 hrs, 20 hrs and 40 hours after bending (panels a-d in FIG. 6, respectively) and stained for GUS expression. Arrows in FIG. 6 indicate the bend sites.

The cellulose synthase promoter-GUS fusion binary constructs showed exclusive xylem-specific expression of GUS without any tension stress (FIG. 6, panel a). However, under tension stress conditions endured by angiosperms in nature, the transgenic tobacco plants induced xylem and phloem-specific expression on the upper side of the stem within the first four hours of stress (FIG. 6, panel b).

This observation was surprising because during tension wood development fibers produce highly crystalline cellulose in order to provide essential mechanical strength to a bending stem. The present observation was the first showing of transcriptional up-regulation of a cellulose synthase, mediated through a cellulose synthase promoter that is directly responsible for development of highly crystalline cellulose in trees. Furthermore, after 20 hrs of tension stress, both xylem and phloem exhibited GUS expression, but only on the upper side of the stem that was under tensile stress, i.e., GUS expression on the lower side was inhibited (FIG. 6, panel c). With extended stress (up to 40 hrs), GUS expression was restricted to only one small region on the upper side of the stem where maximum tension stress was present (FIG. 6, panel d). Based on the observation of GUS signal in woody cells upon tension stress and the absence of GUS under compression or no stress, it was concluded that a cellulose synthase promoter of the invention has mechanical stress responsive elements (MSREs) that turn cellulose synthase genes on and off depending on the presence and type of stress to the stem.

The results indicate that positive MSREs exist in a cellulose synthase promoter of the invention to bind transcription factors in response to tension stress for regulating the expression of cellulose synthase and increasing biosynthesis of higher crystalline cellulose. This is evident based on the expression of GUS in xylem and phloem tissue at the upper side of the stem subjected to tension stress, but not when tissue on the lower side was subjected to compression or no stress. Furthermore, the tissue at the lower side of the stem, which was subjected to compression stress, showed no GUS expression, i.e., expression was turned off. This indicated the presence of negative MSREs, which bind transcription factors to turn off expression of cellulose synthase at the lower side of the stem. Negative MSREs likely suppress development of highly crystalline cellulose in normal wood.

These results provide a mechanism for genetically engineering synthesis of highly crystalline cellulose in juvenile wood for enhancing strength properties, and for synthesizing a higher percentage of cellulose in reaction wood. The positive MSREs and their cognate transcription factors are important in the synthesis of highly crystalline cellulose of high tensile strength, as are the negative MSREs and inhibition of cognate transcription factors thereto. The present invention thus provides a starting point for cloning cDNAs for the transcription factors that bind to positive and negative MSREs according to methods known in the art. Constitutive expression of cDNAs for positive MSRE transcription factors allows the continuous production of highly crystalline cellulose in transgenic trees, while expression of antisense cDNAs for negative MSRE transcription factors inhibits those transcription factors so that cellulose synthase cannot turn off. This combination will assure continuous production of highly crystalline cellulose in trees.

Genetic Engineering of Cellulose Synthase in Transgenic Plants

As discussed above, the nucleotide sequence of a cellulose synthase of the invention, e.g., PtCelA cDNA from aspen, shows significant homology with other polynucleotides encoding cellulose synthase proteins that have been suggested as authentic cellulose synthase clones. To further characterize the activity of a cellulose synthase, four constructs were prepared in a pBI121 plasmid.

1) A constitutive plant promoter Cauliflower mosaic Virus 35S was operatively linked to PtCelA (35SP-PtCelA-s) and overexpressed in transgenic plants. This causes excess production of cellulose, resulting in a reduction in lignin content. Tobacco and aspen have been transformed with this construct.

2) Cauliflower mosaic Virus 35S was operatively linked to antisense RNA from PtCelA (35S-PtCelA-a) and constitutively expressed to reduce production of cellulose and increase lignin content in transgenic plants. This negative control construct may not result in healthy plants since cellulose is essential for plant growth and development. Aspen plants have been transformed with this construct.

3) Aspen 4CL-1 promoter (Hu et al., 1998) was operatively linked to PtCelA (Pt4CLP-PtCelA) (the 35S promoter of pBI121 was removed in this construct) and expressed in a tissue-specific manner in developing secondary xylem of transgenic aspen. This expression augments the native cellulose production and reduces lignin content of angiosperm tissues. Tobacco and aspen have transformed with this construct.

4) The cytoplasmic domain of PtCelA which contains three conserved regions thought to be involved in UDP-glucose binding during cellulose biosynthesis, was linked to a 35S promoter to produce binary constructs (35S-PtCelA UDP-glucose). Expression by this promoter permits constitutive expression of a UDP glucose binding domain of PtCelA in transgenic plants. Tobacco and aspen have been transformed with this construct.

35S-GUS constructs (pBI121, ClonTech, CA) were used as controls for each experiment with the constructs. Transgenic tobacco plants were transformed with the constructs. The following table shows the general growth measurements of the T0 tobacco plants. Plants carrying a PtCelA construct grew much faster than control plants carrying a pBI121 (control) construct. In comparing developmental 4CL and constitutive 35S promoter control of PtCelA expression, the 35S was more effective, permitting faster growth of transgenic

tobacco plants. The fastest growth was seen in transgenic plants carrying a 35S promoter driven UDP-G domain from PtCelA.

It is noted that TO generation plants can have carry over effects from their tissue culture treatments. Therefore, seeds were collected for testing this growth phenomenon in T1 generations. The transgenic tobacco plants were analyzed for presence of the transferred genes and all tested positive for the respective gene constructs.

TABLE 1

Transgenic tobacco plant measurements after transfer in soil for about 1.5 months (N = 2)					
Construct	Height	Diameter	Internode length	No. of leaves	Longest leaf
35S-GUS	17	0.5	1	11	17
35S-PtCelA	77	1.0	6	13	37
35S-UDPG	83	1.0	6	13	37
4CLP-PtCelA	41	0.8	5	10	29

Note: All values were measured in centimeters, excluding number of leaves.

It will be appreciated by persons of ordinary skill in the art that the examples and preferred embodiments herein are illustrative, and that the invention may be practiced in a variety of embodiments which share the same inventive concept.

Deletion Analysis and Activity of PtrCesA1 Promoter

Sub-regions within the promoter region of PtrCesA1 corresponding to nucleotides at positions 1 to 840 of SEQ ID NO: 3 were generated and evaluated for promoter activity. The sequence of the region showing the positions of the primers used in the deletion analysis is set forth in FIG. 9. For each deletion, a forward primer was designed at intervals of approximately 100 bp (shown by forward arrows in FIG. 9) and a HindIII site was added at the 5' end of each forward primer. A reverse primer was designed based on the sequence near the 3' end of the available promoter sequence (shown by reverse arrow in FIG. 9) and included an XbaI site at the 3' end. The sequence of each forward primer and the reverse primer along the PtrCesA1 promoter region (bases 1 to 840 of SEQ ID NO:3) is shown as underlined in FIG. 9.

Using these sets of primer pairs, a total of eight deletion fragments were made in separate PCR reactions. Each of the eight PCR products was subcloned into a pCR2.1 vector. After digesting with HindIII and XbaI restriction enzymes, promoter deletion fragments were ligated to pBI121 that had been digested with the same pair of enzymes. Restriction digestion of pBI121 with HindIII and XbaI removed the 35S promoter leaving the GUS and NOS terminators intact. Thus, the resulting constructs had eight different PtrCesA1 promoter fragments driving the expression of the GUS coding sequence. PCR reactions using the same primer pairs were performed to verify that the size of the inserts of each of these eight constructs was correct. Further verification was performed by digestion of the constructs with HindIII and XbaI. DNA sequencing of each of the eight amplified PCR products was also performed, and confirmed that the sequences were each correct. The deletion fragments were designated 18-838, 120-838, 258-838, 334-838, 423-838, 523-838, 620-838 and 717-838, based on the position from the start of the nucleotides recited in SEQ ID NO:3.

The 18-838 fragment comprised 821 base pairs corresponding from position 18 to position 838 of SEQ ID NO. 3. The 120-838 fragment comprised 719 base pairs correspond-

ing from position 120 to position 838 of SEQ ID NO. 3. The 258-838 fragment comprised 581 base pairs corresponding from position 258 to position 838 of SEQ ID NO. 3. The 334-838 fragment comprised 505 base pairs corresponding from position 334 to position 838 of SEQ ID NO. 3. The 423-838 fragment comprised 416 bases from position 423 to position 838 of SEQ ID NO. 3. The 523-838 fragment comprised 316 base pairs corresponding from position 523 to position 838 of SEQ ID NO. 3. The 620-838 fragment comprised 219 base pairs corresponding from position 620 to position 838 of SEQ ID NO. 3. The 717-838 fragment comprised 122 base pairs corresponding from position 717 to position 838 of SEQ ID NO. 3.

Each of the eight constructs was mobilized into *Agrobacterium*, and the presence of the correct construct in *Agrobacterium* was confirmed by PCR amplification of DNA from the transformed *Agrobacterium*. The eight expression vectors were subsequently transformed into tobacco, and at least two independent lines per construct were transferred to the greenhouse. Genomic PCR testing of these plants showed the expected promoter fragments integrated into the genome. Our previous studies have indicated that tobacco transformants exhibit transgene expression patterns similar to aspen (Wu et al., 2000).

GUS staining of transgenic tobacco stems from each of these eight types of transgenic plants was performed as described earlier (Wu et al., 2000). The results of these experiments are shown in Table 2.

TABLE 2

Fragment of SEQ ID NO:3	Intensity of GUS Staining
18-838	Strong
120-838	Strong
258-838	Strong
334-838	Strong
423-838	Some
523-838	Very weak
620-838	None
717-838	None

GUS staining was unaffected in PtrCesA1 promoter fragments comprising nucleotides 18-838, 120-838, 258-838 and 334-838 of SEQ ID NO:3, but in promoter fragments comprising nucleotides 423-838 of SEQ ID NO:3 staining was reduced. Staining was further reduced in the promoter fragment comprising nucleotides 523-838 of SEQ ID NO:3 with very weak staining noted. No GUS staining was noted in promoter fragments comprising nucleotides 620-838 and 717-838 of SEQ ID NO:3. Thus, sequences comprising nucleotides corresponding to positions 334 to 422, 423 to 522, 334 to 522, 334 to 838 and 423 to 838 of SEQ ID NO. 3 each show some promoter activity. The sequences showing promoter activity may be useful in driving the expression of any coding sequence to which they are operably linked.

Sequences comprising nucleotides from 334 to 422 and 423 to 522 of SEQ ID NO: 3 may therefore drive transcription of polynucleotides in a plant xylem, including the xylem-specific expression of PtrCesA1. Sequences comprising nucleotides from 334 to 422 and 423 to 522 of SEQ ID NO: 3 may be responsible for tension-stress responsive expression of PtrCesA1 coding sequence.

BIBLIOGRAPHY

- Hu et al., 1999, Nature Biotechnology, In Press
 Whetten et al., 1998, Ann Rev P1 Physiol P1 Mol Biol, 49: 585-609
 Arioli et al., 1998, Science, 279: 717-720
 Wu et al., 1998, P1 Physiol, 117: 1125
 Hu et al., 1998, PNAS, 95: 5407-5412
 Joshi et al., 1997, PMB, 35: 993-1001
 Fukuda, 1996, Ann Rev P1 Physiol P1 Mol Biol, 47: 299-325
 Pear et al., 1996, PNAS, 93: 12637-12642
 Haigler and Blanton, 1996, PNAS, 93: 12082-12085
 Ge and Chiang, 1996, P1 Physiol, 112: 861

- Brown et al., 1996, Trends P1 Sci., 1: 149-156
 Delmer and Amor, 1995, P1 Cell, 7: 987-1000
 Hoffman and Stoffel, 1993, Biol Chem, Hoppe-Seyler 374: 166
 Joshi, 1987, NAR, 15: 6643-6653
 Joshi, 1987, NAR, 15: 9627-9640
 Timmell, 1986, Compression Wood in Gymnosperms, Springer Verlag
 Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons
 Higuchi, 1997, Biochemistry and Molecular Biology of Wood, Springer Verlag

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 3232

<212> TYPE: DNA

<213> ORGANISM: Populus tremuloides

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (69)..(3002)

<400> SEQUENCE: 1

```

gtcgaccac gcgtccgtct tgaagaata tgaagttgta aagagctggt aaagtggtaa      60
taagcaag atg atg gaa tct ggg gct cct ata tgc cat acc tgt ggt gaa      110
      Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu
      1          5          10
cag gtg ggg cat gat gca aat ggg gag cta ttt gtg gct tgc cat gag      158
Gln Val Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu
      15          20          25          30
tgt agc tat ccc atg tgc aag tct tgt ttc gag ttt gaa atc aat gag      206
Cys Ser Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Asn Glu
      35          40          45
ggc cgg aaa gtt tgc ttg cgg tgt ggc tcg cca tat gat gag aac ttg      254
Gly Arg Lys Val Cys Leu Arg Cys Gly Ser Pro Tyr Asp Glu Asn Leu
      50          55          60
ctg gat gat gta gaa aag aag ggg tct ggc aat caa tcc aca atg gca      302
Leu Asp Asp Val Glu Lys Lys Gly Ser Gly Asn Gln Ser Thr Met Ala
      65          70          75
tct cac ctc aac gat tct cag gat gtc gga atc cat gct aga cat atc      350
Ser His Leu Asn Asp Ser Gln Asp Val Gly Ile His Ala Arg His Ile
      80          85          90
agt agt gtg tcc act gtg gat agt gaa atg aat gat gaa tat ggg aat      398
Ser Ser Val Ser Thr Val Asp Ser Glu Met Asn Asp Glu Tyr Gly Asn
      95          100          105          110
cca att tgg aag aat cgg gtg aag agc tgt aag gat aaa gag aac aag      446
Pro Ile Trp Lys Asn Arg Val Lys Ser Cys Lys Asp Lys Glu Asn Lys
      115          120          125
aag aaa aag aga agt cct aag gct gaa act gaa cca gct caa gtt cct      494
Lys Lys Lys Arg Ser Pro Lys Ala Glu Thr Glu Pro Ala Gln Val Pro
      130          135          140
aca gaa cag cag atg gaa gag aaa ccg tct gca gag gct tcg gag ccg      542
Thr Glu Gln Gln Met Glu Glu Lys Pro Ser Ala Glu Ala Ser Glu Pro
      145          150          155
ctt tca att gtt tat cca att cca cgc aac aag ctc aca cca tac aga      590
Leu Ser Ile Val Tyr Pro Ile Pro Arg Asn Lys Leu Thr Pro Tyr Arg
      160          165          170
gca gtg atc att atg cga ctg gtc att ctg ggc ctc ttc ttc cac ttc      638

```

-continued

Ala Val Ile Ile Met Arg Leu Val Ile Leu Gly Leu Phe Phe His Phe 175 180 185 190	
aga ata aca aat cct gtc gat agt gcc ttt ggc ctg tgg ctt act tct Arg Ile Thr Asn Pro Val Asp Ser Ala Phe Gly Leu Trp Leu Thr Ser 195 200 205	686
gtc ata tgt gag atc tgg ttt gca ttt tct tgg gtg ttg gat cag ttc Val Ile Cys Glu Ile Trp Phe Ala Phe Ser Trp Val Leu Asp Gln Phe 210 215 220	734
ccc aag tgg aat cct gtc aat aga gaa acg tat atc gaa agg ctg tcg Pro Lys Trp Asn Pro Val Asn Arg Glu Thr Tyr Ile Glu Arg Leu Ser 225 230 235	782
gca agg tat gaa aga gag ggt gag cct tct cag ctt gct ggt gtg gat Ala Arg Tyr Glu Arg Glu Gly Glu Pro Ser Gln Leu Ala Gly Val Asp 240 245 250	830
ttt ttc gtg agt act gtt gat ccg ctg aag gaa ccg cca ttg atc act Phe Phe Val Ser Thr Val Asp Pro Leu Lys Glu Pro Pro Leu Ile Thr 255 260 265 270	878
gcc aat aca gtc ctt tcc atc ctt gct gtg gac tat ccc gtc gat aaa Ala Asn Thr Val Leu Ser Ile Leu Ala Val Asp Tyr Pro Val Asp Lys 275 280 285	926
gtc tcc tgc tac gtg tct gat gat ggt gca gct atg ctt tca ttt gaa Val Ser Cys Tyr Val Ser Asp Asp Gly Ala Ala Met Leu Ser Phe Glu 290 295 300	974
tct ctt gta gaa aca gct gag ttt gca agg aag tgg gtt ccg ttc tgc Ser Leu Val Glu Thr Ala Glu Phe Ala Arg Lys Trp Val Pro Phe Cys 305 310 315	1022
aaa aaa ttc tca att gaa cca aga gca ccg gag ttt tac ttc tca cag Lys Lys Phe Ser Ile Glu Pro Arg Ala Pro Glu Phe Tyr Phe Ser Gln 320 325 330	1070
aaa att gat tac ttg aaa gac aag gtt caa cct tct ttc gtg aaa gaa Lys Ile Asp Tyr Leu Lys Asp Lys Val Gln Pro Ser Phe Val Lys Glu 335 340 345 350	1118
cgt aga gca atg aaa agg gat tat gaa gag tac aaa gtc cga gtt aat Arg Arg Ala Met Lys Arg Asp Tyr Glu Glu Tyr Lys Val Arg Val Asn 355 360 365	1166
gcc ctg gta gca aag gct cag aaa aca cct gaa gaa gga tgg act atg Ala Leu Val Ala Lys Ala Gln Lys Thr Pro Glu Glu Gly Trp Thr Met 370 375 380	1214
caa gat gga aca cct tgg cct ggg aat aac aca cgt gat cac cct ggg Gln Asp Gly Thr Pro Trp Pro Gly Asn Asn Thr Arg Asp His Pro Gly 385 390 395	1262
cat gat tca ggt ctt cct tgg gaa ata ctg gga gct cgt gac att gaa His Asp Ser Gly Leu Pro Trp Glu Ile Leu Gly Ala Arg Asp Ile Glu 400 405 410	1310
gga aat gaa cta cct cgt cta gta tat gtc tcc agg gag aag aga cct Gly Asn Glu Leu Pro Arg Leu Val Tyr Val Ser Arg Glu Lys Arg Pro 415 420 425 430	1358
ggc tac cag cac cac aaa aag gct ggt gca gaa aat gct ctg gtg aga Gly Tyr Gln His His Lys Lys Ala Gly Ala Glu Asn Ala Leu Val Arg 435 440 445	1406
gtg tct gca gta ctc aca aat gct ccc tac atc ctc aat gtt gat tgt Val Ser Ala Val Leu Thr Asn Ala Pro Tyr Ile Leu Asn Val Asp Cys 450 455 460	1454
gat cac tat gta aac aat agc aag gct gtt cga gag gca atg tgc atc Asp His Tyr Val Asn Asn Ser Lys Ala Val Arg Glu Ala Met Cys Ile 465 470 475	1502
ctg atg gac cca caa gta ggt cga gat gta tgc tat gtg cag ttc cct Leu Met Asp Pro Gln Val Gly Arg Asp Val Cys Tyr Val Gln Phe Pro 480 485 490	1550

-continued

cag agg ttt gat ggc ata gat aag agt gat cgc tac gcc aat cgt aac	1598
Gln Arg Phe Asp Gly Ile Asp Lys Ser Asp Arg Tyr Ala Asn Arg Asn	
495 500 505 510	
gta gtt ttc ttt gat gtt aac atg aaa ggg ttg gat ggc att caa gga	1646
Val Val Phe Phe Asp Val Asn Met Lys Gly Leu Asp Gly Ile Gln Gly	
515 520 525	
cca gta tac gta gga act ggt tgt gtt ttc aac agg caa gca ctt tac	1694
Pro Val Tyr Val Gly Thr Gly Cys Val Phe Asn Arg Gln Ala Leu Tyr	
530 535 540	
ggc tac ggg cct cct tct atg ccc agc tta cgc aag aga aag gat tct	1742
Gly Tyr Gly Pro Pro Ser Met Pro Ser Leu Arg Lys Arg Lys Asp Ser	
545 550 555	
tca tcc tgc ttc tca tgt tgc tgc ccc tca aag aag aag cct gct caa	1790
Ser Ser Cys Phe Ser Cys Cys Pro Ser Lys Lys Lys Pro Ala Gln	
560 565 570	
gat cca gct gag gta tac aga gat gca aaa aga gag gat ctc aat gct	1838
Asp Pro Ala Glu Val Tyr Arg Asp Ala Lys Arg Glu Asp Leu Asn Ala	
575 580 585 590	
gcc ata ttt aat ctt aca gag att gat aat tat gac gag cat gaa agg	1886
Ala Ile Phe Asn Leu Thr Glu Ile Asp Asn Tyr Asp Glu His Glu Arg	
595 600 605	
tca atg ctg atc tcc cag ttg agc ttt gag aaa act ttt ggc tta tct	1934
Ser Met Leu Ile Ser Gln Leu Ser Phe Glu Lys Thr Phe Gly Leu Ser	
610 615 620	
tct gtc ttc att gag tct aca cta atg gag aat gga gga gta ccc gag	1982
Ser Val Phe Ile Glu Ser Thr Leu Met Glu Asn Gly Gly Val Pro Glu	
625 630 635	
tct gcc aac tca cca cca ttc atc aag gaa gcg att caa gtc atc ggc	2030
Ser Ala Asn Ser Pro Pro Phe Ile Lys Glu Ala Ile Gln Val Ile Gly	
640 645 650	
tgt ggc tat gaa gag aag act gaa tgg gga aaa cag att ggt tgg ata	2078
Cys Gly Tyr Glu Glu Lys Thr Glu Trp Gly Lys Gln Ile Gly Trp Ile	
655 660 665 670	
tat ggg tca gtc act gag gat atc tta agt ggc ttc aag atg cac tgc	2126
Tyr Gly Ser Val Thr Glu Asp Ile Leu Ser Gly Phe Lys Met His Cys	
675 680 685	
cga gga tgg aga tca att tac tgc atg ccc gta agg cct gca ttc aaa	2174
Arg Gly Trp Arg Ser Ile Tyr Cys Met Pro Val Arg Pro Ala Phe Lys	
690 695 700	
gga tct gca ccc atc aac ctg tct gat aga ttg cac cag gtc ctc cga	2222
Gly Ser Ala Pro Ile Asn Leu Ser Asp Arg Leu His Gln Val Leu Arg	
705 710 715	
tgg gct ctt ggt tct gtg gaa att ttc ttt agc aga cac tgt ccc ctc	2270
Trp Ala Leu Gly Ser Val Glu Ile Phe Phe Ser Arg His Cys Pro Leu	
720 725 730	
tgg tac ggg ttt gga gga ggc cgt ctt aaa tgg ctc caa agg ctt gcg	2318
Trp Tyr Gly Phe Gly Gly Arg Leu Lys Trp Leu Gln Arg Leu Ala	
735 740 745 750	
tat ata aac acc att gtg tac cca ttt aca tcc ctc cct ctc att gcc	2366
Tyr Ile Asn Thr Ile Val Tyr Pro Phe Thr Ser Leu Pro Leu Ile Ala	
755 760 765	
tat tgc aca att cct gca gtt tgt ctg ctc acc gga aaa ttc atc ata	2414
Tyr Cys Thr Ile Pro Ala Val Cys Leu Leu Thr Gly Lys Phe Ile Ile	
770 775 780	
cca acg ctc tca aac ctg gca agc atg ctg ttt ctt ggc ctc ttt atc	2462
Pro Thr Leu Ser Asn Leu Ala Ser Met Leu Phe Leu Gly Leu Phe Ile	
785 790 795	
tcc atc att gta act gcg gtg ctt gag cta aga tgg agc ggt gtc agc	2510
Ser Ile Ile Val Thr Ala Val Leu Glu Leu Arg Trp Ser Gly Val Ser	
800 805 810	

-continued

```

att gaa gat tta tgg cgt aat gaa caa ttc tgg gtg atc gga ggt gtt 2558
Ile Glu Asp Leu Trp Arg Asn Glu Gln Phe Trp Val Ile Gly Gly Val
815                820                825                830

tca gcc cat ctc ttt gcg gtc ttc cag gga ttc tta aaa atg ttg gct 2606
Ser Ala His Leu Phe Ala Val Phe Gln Gly Phe Leu Lys Met Leu Ala
                835                840                845

ggc atc gat acg aac ttc act gtc aca gca aaa gca gcc gaa gat gca 2654
Gly Ile Asp Thr Asn Phe Thr Val Thr Ala Lys Ala Ala Glu Asp Ala
                850                855                860

gaa ttt ggg gag cta tat atg gtc aag tgg aca aca ctt ttg att cct 2702
Glu Phe Gly Glu Leu Tyr Met Val Lys Trp Thr Thr Leu Leu Ile Pro
                865                870                875

cca acc aca ctt ctc att atc aat atg tcg ggt tgt gct gga ttc tct 2750
Pro Thr Thr Leu Leu Ile Ile Asn Met Ser Gly Cys Ala Gly Phe Ser
                880                885                890

gat gca ctc aac aaa gga tat gaa gca tgg ggg cct ctc ttt ggc aag 2798
Asp Ala Leu Asn Lys Gly Tyr Glu Ala Trp Gly Pro Leu Phe Gly Lys
895                900                905                910

gtg ttc ttt gct ttc tgg gtg att ctt cat ctc tat cca ttc ctt aaa 2846
Val Phe Phe Ala Phe Trp Val Ile Leu His Leu Tyr Pro Phe Leu Lys
                915                920                925

ggg cta atg ggt cgc caa aac cta aca cca acc att gtt gtt ctc tgg 2894
Gly Leu Met Gly Arg Gln Asn Leu Thr Pro Thr Ile Val Val Leu Trp
                930                935                940

tca gtg ctg ttg gcc tct gtc ttc tct ctc gtt tgg gtc aag atc aat 2942
Ser Val Leu Leu Ala Ser Val Phe Ser Leu Val Trp Val Lys Ile Asn
                945                950                955

cca ttc gtt aac aaa gtt gat aac acc ttg gtt gcg gag acc tgc att 2990
Pro Phe Val Asn Lys Val Asp Asn Thr Leu Val Ala Glu Thr Cys Ile
                960                965                970

tcc att gat tgc tgagctacct ccaataagtc tctcccagta ttttggggtt 3042
Ser Ile Asp Cys
975

acaaaacctt tgggaattgg aatgatgcc tcggtgtagt ttccctcaag aaagcacata 3102

tcgctgtcag tatttaaatg aactgcaaga tgattgttct ctatgaagtt ttgaacagtt 3162

tgaaatgata ttatgttaaa atacaggttt tgattgttt gaaaaaaaaa aagaaaaaaaa 3222

aaaaaaaaaa 3232

<210> SEQ ID NO 2
<211> LENGTH: 978
<212> TYPE: PRT
<213> ORGANISM: Populus tremuloides

<400> SEQUENCE: 2

Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val
 1                5                10                15

Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser
20                25                30

Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Asn Glu Gly Arg
35                40                45

Lys Val Cys Leu Arg Cys Gly Ser Pro Tyr Asp Glu Asn Leu Leu Asp
50                55                60

Asp Val Glu Lys Lys Gly Ser Gly Asn Gln Ser Thr Met Ala Ser His
65                70                75                80

Leu Asn Asp Ser Gln Asp Val Gly Ile His Ala Arg His Ile Ser Ser
85                90                95

```

-continued

Val	Ser	Thr	Val	Asp	Ser	Glu	Met	Asn	Asp	Glu	Tyr	Gly	Asn	Pro	Ile	100	105	110	
Trp	Lys	Asn	Arg	Val	Lys	Ser	Cys	Lys	Asp	Lys	Glu	Asn	Lys	Lys	Lys	115	120	125	
Lys	Arg	Ser	Pro	Lys	Ala	Glu	Thr	Glu	Pro	Ala	Gln	Val	Pro	Thr	Glu	130	135	140	
Gln	Gln	Met	Glu	Glu	Lys	Pro	Ser	Ala	Glu	Ala	Ser	Glu	Pro	Leu	Ser	145	150	155	160
Ile	Val	Tyr	Pro	Ile	Pro	Arg	Asn	Lys	Leu	Thr	Pro	Tyr	Arg	Ala	Val	165	170	175	
Ile	Ile	Met	Arg	Leu	Val	Ile	Leu	Gly	Leu	Phe	Phe	His	Phe	Arg	Ile	180	185	190	
Thr	Asn	Pro	Val	Asp	Ser	Ala	Phe	Gly	Leu	Trp	Leu	Thr	Ser	Val	Ile	195	200	205	
Cys	Glu	Ile	Trp	Phe	Ala	Phe	Ser	Trp	Val	Leu	Asp	Gln	Phe	Pro	Lys	210	215	220	
Trp	Asn	Pro	Val	Asn	Arg	Glu	Thr	Tyr	Ile	Glu	Arg	Leu	Ser	Ala	Arg	225	230	235	240
Tyr	Glu	Arg	Glu	Gly	Glu	Pro	Ser	Gln	Leu	Ala	Gly	Val	Asp	Phe	Phe	245	250	255	
Val	Ser	Thr	Val	Asp	Pro	Leu	Lys	Glu	Pro	Pro	Leu	Ile	Thr	Ala	Asn	260	265	270	
Thr	Val	Leu	Ser	Ile	Leu	Ala	Val	Asp	Tyr	Pro	Val	Asp	Lys	Val	Ser	275	280	285	
Cys	Tyr	Val	Ser	Asp	Asp	Gly	Ala	Ala	Met	Leu	Ser	Phe	Glu	Ser	Leu	290	295	300	
Val	Glu	Thr	Ala	Glu	Phe	Ala	Arg	Lys	Trp	Val	Pro	Phe	Cys	Lys	Lys	305	310	315	320
Phe	Ser	Ile	Glu	Pro	Arg	Ala	Pro	Glu	Phe	Tyr	Phe	Ser	Gln	Lys	Ile	325	330	335	
Asp	Tyr	Leu	Lys	Asp	Lys	Val	Gln	Pro	Ser	Phe	Val	Lys	Glu	Arg	Arg	340	345	350	
Ala	Met	Lys	Arg	Asp	Tyr	Glu	Glu	Tyr	Lys	Val	Arg	Val	Asn	Ala	Leu	355	360	365	
Val	Ala	Lys	Ala	Gln	Lys	Thr	Pro	Glu	Glu	Gly	Trp	Thr	Met	Gln	Asp	370	375	380	
Gly	Thr	Pro	Trp	Pro	Gly	Asn	Asn	Thr	Arg	Asp	His	Pro	Gly	His	Asp	385	390	395	400
Ser	Gly	Leu	Pro	Trp	Glu	Ile	Leu	Gly	Ala	Arg	Asp	Ile	Glu	Gly	Asn	405	410	415	
Glu	Leu	Pro	Arg	Leu	Val	Tyr	Val	Ser	Arg	Glu	Lys	Arg	Pro	Gly	Tyr	420	425	430	
Gln	His	His	Lys	Lys	Ala	Gly	Ala	Glu	Asn	Ala	Leu	Val	Arg	Val	Ser	435	440	445	
Ala	Val	Leu	Thr	Asn	Ala	Pro	Tyr	Ile	Leu	Asn	Val	Asp	Cys	Asp	His	450	455	460	
Tyr	Val	Asn	Asn	Ser	Lys	Ala	Val	Arg	Glu	Ala	Met	Cys	Ile	Leu	Met	465	470	475	480
Asp	Pro	Gln	Val	Gly	Arg	Asp	Val	Cys	Tyr	Val	Gln	Phe	Pro	Gln	Arg	485	490	495	
Phe	Asp	Gly	Ile	Asp	Lys	Ser	Asp	Arg	Tyr	Ala	Asn	Arg	Asn	Val	Val	500	505	510	
Phe	Phe	Asp	Val	Asn	Met	Lys	Gly	Leu	Asp	Gly	Ile	Gln	Gly	Pro	Val				

-continued

515					520					525					
Tyr	Val	Gly	Thr	Gly	Cys	Val	Phe	Asn	Arg	Gln	Ala	Leu	Tyr	Gly	Tyr
530						535					540				
Gly	Pro	Pro	Ser	Met	Pro	Ser	Leu	Arg	Lys	Arg	Lys	Asp	Ser	Ser	Ser
545				550						555					560
Cys	Phe	Ser	Cys	Cys	Cys	Pro	Ser	Lys	Lys	Lys	Pro	Ala	Gln	Asp	Pro
			565						570					575	
Ala	Glu	Val	Tyr	Arg	Asp	Ala	Lys	Arg	Glu	Asp	Leu	Asn	Ala	Ala	Ile
			580					585					590		
Phe	Asn	Leu	Thr	Glu	Ile	Asp	Asn	Tyr	Asp	Glu	His	Glu	Arg	Ser	Met
		595				600						605			
Leu	Ile	Ser	Gln	Leu	Ser	Phe	Glu	Lys	Thr	Phe	Gly	Leu	Ser	Ser	Val
610						615					620				
Phe	Ile	Glu	Ser	Thr	Leu	Met	Glu	Asn	Gly	Gly	Val	Pro	Glu	Ser	Ala
625					630					635					640
Asn	Ser	Pro	Pro	Phe	Ile	Lys	Glu	Ala	Ile	Gln	Val	Ile	Gly	Cys	Gly
				645					650					655	
Tyr	Glu	Glu	Lys	Thr	Glu	Trp	Gly	Lys	Gln	Ile	Gly	Trp	Ile	Tyr	Gly
			660					665					670		
Ser	Val	Thr	Glu	Asp	Ile	Leu	Ser	Gly	Phe	Lys	Met	His	Cys	Arg	Gly
		675				680						685			
Trp	Arg	Ser	Ile	Tyr	Cys	Met	Pro	Val	Arg	Pro	Ala	Phe	Lys	Gly	Ser
690						695					700				
Ala	Pro	Ile	Asn	Leu	Ser	Asp	Arg	Leu	His	Gln	Val	Leu	Arg	Trp	Ala
705				710						715					720
Leu	Gly	Ser	Val	Glu	Ile	Phe	Phe	Ser	Arg	His	Cys	Pro	Leu	Trp	Tyr
				725					730					735	
Gly	Phe	Gly	Gly	Gly	Arg	Leu	Lys	Trp	Leu	Gln	Arg	Leu	Ala	Tyr	Ile
			740					745					750		
Asn	Thr	Ile	Val	Tyr	Pro	Phe	Thr	Ser	Leu	Pro	Leu	Ile	Ala	Tyr	Cys
		755					760					765			
Thr	Ile	Pro	Ala	Val	Cys	Leu	Leu	Thr	Gly	Lys	Phe	Ile	Ile	Pro	Thr
	770					775					780				
Leu	Ser	Asn	Leu	Ala	Ser	Met	Leu	Phe	Leu	Gly	Leu	Phe	Ile	Ser	Ile
785				790						795					800
Ile	Val	Thr	Ala	Val	Leu	Glu	Leu	Arg	Trp	Ser	Gly	Val	Ser	Ile	Glu
				805					810					815	
Asp	Leu	Trp	Arg	Asn	Glu	Gln	Phe	Trp	Val	Ile	Gly	Gly	Val	Ser	Ala
			820					825					830		
His	Leu	Phe	Ala	Val	Phe	Gln	Gly	Phe	Leu	Lys	Met	Leu	Ala	Gly	Ile
		835						840					845		
Asp	Thr	Asn	Phe	Thr	Val	Thr	Ala	Lys	Ala	Ala	Glu	Asp	Ala	Glu	Phe
	850					855					860				
Gly	Glu	Leu	Tyr	Met	Val	Lys	Trp	Thr	Thr	Leu	Leu	Ile	Pro	Pro	Thr
865					870					875					880
Thr	Leu	Leu	Ile	Ile	Asn	Met	Ser	Gly	Cys	Ala	Gly	Phe	Ser	Asp	Ala
				885					890					895	
Leu	Asn	Lys	Gly	Tyr	Glu	Ala	Trp	Gly	Pro	Leu	Phe	Gly	Lys	Val	Phe
			900					905					910		
Phe	Ala	Phe	Trp	Val	Ile	Leu	His	Leu	Tyr	Pro	Phe	Leu	Lys	Gly	Leu
		915					920						925		
Met	Gly	Arg	Gln	Asn	Leu	Thr	Pro	Thr	Ile	Val	Val	Leu	Trp	Ser	Val
930						935						940			

-continued

Leu Leu Ala Ser Val Phe Ser Leu Val Trp Val Lys Ile Asn Pro Phe
945 950 955 960

Val Asn Lys Val Asp Asn Thr Leu Val Ala Glu Thr Cys Ile Ser Ile
965 970 975

Asp Cys

<210> SEQ ID NO 3
<211> LENGTH: 1010
<212> TYPE: DNA
<213> ORGANISM: Populus tremuloides
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (841)..(1008)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: 5' flanking region of PtCelA coding sequence

<400> SEQUENCE: 3

```

gaattcgccc ttttgaattc aggagacgat agtttccggt tcggtgaatg gctttgttca    60
cttctggtct agcaatttgc aaaagaagtt acaaaacaaa tgcattatg gtaaattdaa    120
caagagatgg gttctatggt cacttattta tgcccatcat ttgttctggg gttactcttt    180
atagtctgat tcgaagtgc aaactgccgt ttctggtatt gcaattatgt agccataaac    240
tgtaaatcct gtatgtatta geggaccaac aaccagatat acgggatcag cgtcgtaaaa    300
gagatctcca ttctacgttt ctttctaatt tttccgtttc agtgagagaa ttaccctgat    360
acattgacat gatgattgat gattatggga accattccga tgtagacac gagaccatct    420
ggatcctgcc agttttcagt tcacatggca tctcagccca agatcatgtg tttatacgcc    480
taatgacttg tattgaaagt ttggaagtt gaagatgtgc tctgccaac agaaaccttc    540
cttaaatttc cagcaaatct tcaaaacttg gccttacacc ccgaaaatag acgtgcttct    600
acttgggttc ttgaaacca tgcaccaacc gccatacccc accaacccac caccctcaac    660
cttctcttgc ccattacaaa aatgtcagta ccaccctctg aaagacacca acacacccta    720
gctttggtta gggtatttga tataaaaaaca aggcctcaaac aaaagattgg aaggaagcag    780
aggaagacc tcttgaaga attgaagttg taaagagctg gtaaagtggg aataagcaag    840
atg atg gaa tct ggg gct cct ata tgc cat acc tgt ggt gaa cag gtg    888
Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val
  1         5         10        15
ggg cat gat gca aat ggg gag cta ttt gtg gct tgc cat gag tgt agc    936
Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser
  20        25        30
tat ccc atg tgc aag tct tgt ttc gag ttt gaa atc aaa gag ggc cgg    984
Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Lys Glu Gly Arg
  35        40        45
aaa gtt tgc ttg cgg tgt ggc tcg ag    1010
Lys Val Cys Leu Arg Cys Gly Ser
  50        55

```

<210> SEQ ID NO 4
<211> LENGTH: 3444
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 4

```

gctggcgcgg ttaatcgccg gttctcacia caggaatgag tttgtcctca ttaatgccga    60
tgagaatgcc cgaataagat cagtccaaga gctgagtggg cagacatgct aatctgcag    120

```

- cont inued

agatgagatc gaattgactg ttgatggaga accgtttgty gcatgtaacg aatgtgcatt	180
ccctgtgtgt agacctgct atgagtacga aagacgagaa ggcaatcaag cttgtccaca	240
gtgcaaaacc cgtttcaaac gtcttaaagg aagtccaaga gttgaaggty atgaagagga	300
agatgacatt gatgatthag acaatgagtt tgagtatgga aataatggga ttggatttga	360
tcaggtttct gaaggatgt caatctctcg tcgcaactcc ggtttcccac aatctgattt	420
ggattcagct ccacctggct ctacagattcc attgctgact tacggcgacg aggacgttga	480
gatttcttct gatagacatg ctcttattgt tcctccttca cttggtyggtc atggcaatag	540
agttcatcct gtttctcttt ctgaccgac cgtggtgca catcgaaggc tgatggtacc	600
tcagaagat cttcggttt atggttatgg aagtgtcgtc tggaaagatc ggatggagga	660
atggaagaga aagcagaatg agaaacttca gttgttagg catgaaggag atcctgattt	720
tgaagatggt gatgatgctg attttccaat gatggatgag ggaaggcagc cattgtctat	780
gaagatacca atcaaatcga gcaagataaa tccttaccgg atgttaattg tgctacgtct	840
tgtgattctt ggtctcttct ttcactaccg tattcttcac cccgtcaaag atgcatatgc	900
tttgtggctt atttctgtta tatgtgagat atggtttgct gtttcatggg ttcttgatca	960
gttcctaaa tggtaacctc tcgagcgaga aacgtacttg gaccgactct cattaagata	1020
tgagaagaa gggaaaccgt cgggactatc ccctgtggat gtatttgta gtacagtgga	1080
tcattgaaa gagcctccgc ttattactgc aaactactgc ttgtctattc ttgctgttga	1140
ttatctgtc gataaggttg cttggtacgt atctgatgat ggtgctgcta tgcttacttt	1200
cgaagctctt tctgagaccg ctgaattcgc aaggaaatgg gttccttct gcaagaata	1260
ttgtattgag cctcgtgctc ccgaatggta tttctgcat aaaatggact acttgaagaa	1320
taaagttcat cccgcatttg ttagggagcg gcgagccatg aagagagatt atgaagaatt	1380
caaagtaaag atcaatgctt tagtagcaac agcacagaaa gtgcctgagg atggttggac	1440
tatgcaagac ggtacacctt ggcccggtaa tagtgtgca gatcatcctg gcatgattca	1500
ggtcttctt ggaagtacg gtgttcgtga tgtcgaaaac aacgagttgc ctgattagt	1560
ttacgtttct cgtgagaaga gaccgggatt tgatcacctc aagaaggctg gagctatgaa	1620
ttcctgata cgagtctctg ggttctatc aaatgctcct taccttctga atgtcgattg	1680
tgatcactac atcaacaata gcaagctct tagagaagca atgtgtttca tgatggatcc	1740
tcagtcagga aagaaaatct gttatgttca gttccctcaa aggttcgatg ggattgatag	1800
gcacgatcga tactcaaatc gcaatgttgt gttctttgat atcaatatga aaggtttga	1860
tgggtacaaa gggctatat acgtcggtag aggttggtt ttcaggaggc aagcgcttta	1920
cggatttgat gcaccgaaga agaagaaggg cccacgtaag acatgcaatt gctggccaaa	1980
atggtgtctc ctatgttttg gttcaagaaa gaatcgtaaa gcaaaagacag tggctgcgga	2040
taagaagaag aagaataggg aagcgtcaaa gcagatccac gcattagaaa atatcgaaga	2100
ggcccgcggt cataaagttc ttaacgtaga acagtcaacc gaggcaatgc aatgaagtt	2160
gcagaagaaa tatgggcagt ctctgtatt tgttgcctc gcgctctgg agaatggtgg	2220
gatggctaga aacgcaagcc cggcttctc gcttaagaaa gccatccaag tcattagtcg	2280
cggatgatga gataaaactg aatggggaaa agagattggg tggatctatg gttctgttac	2340
cgaagatatt cttacgggtt ctaagatgca ttctcatggt tggagacatg tttattgtac	2400
accaaagtta gcgctttca aaggatcagc tccaatcaat ctttcggatc gtctccatca	2460
agttctctga tggcgcttg ggtcggttga gatttcttg agtaggcatt gtcctatttg	2520

-continued

```

gtatggttat ggaggtgggt tgaatggct tgagegggtg tectacatta actctgtggt 2580
ttaccctggg acctctctac cgtcatcgt ttactgttct ctccctgcc tctgtcttct 2640
cactggaaaa ttcacgttc cggagattag caactatgcg agtatcctct tcatggcgct 2700
cttctcgtcg attgcaataa cgggtattct cgagatgcaa tggggcaaag ttgggatcga 2760
tgattggtgg agaaacgaac agttttgggt cattggaggt gtttctgcgc atctgtttgc 2820
tctcttccaa ggtctcctca aggttcttgc tgggtgcgac actaacttca cagtcaatc 2880
aaaagcagct gatgatggag agttctctga cctttacctc tcaaatgga ctteacttct 2940
catccctcca atgactctac tcatcataaa cgtcattgga gtcatagtcg gagtctctga 3000
tgccatcagc aatggatagc actcgtgggg accgcttttc ggaagactgt tctttgcaact 3060
ttgggtcatc attcatcttt acccgttcct taaaggtttg cttgggaaac aagatagaat 3120
gcccaaccatt attgtcgtct ggtccatcct cctggcctcg attcttacac ttctttgggt 3180
ccgggttaat ccgtttggg cgaaggcgg tctattctc gagatctgtg gtttagactg 3240
cttgtgattc gattgaccgg tggatgggtt ggtgaaaaag gtttaattcc cacggatcaa 3300
agagaggtaa gagagatatt gttttacctc taaaagactc cttcattgtg ttcattagat 3360
gatgaaaaat gaaaagaaaa agaagattta atttgttac gagaattggt atttttgcaa 3420
gaatgtgttg tagatagcgg ccgc 3444

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1080
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: cellulose synthase

```

```

<400> SEQUENCE: 5

```

```

Arg Pro Arg Leu Ile Ala Gly Ser His Asn Arg Asn Glu Phe Val Leu
  1           5           10
Ile Asn Ala Asp Glu Asn Ala Arg Ile Arg Ser Val Gln Glu Leu Ser
  20          25          30
Gly Gln Thr Cys Gln Ile Cys Arg Asp Glu Ile Glu Leu Thr Val Asp
  35          40          45
Gly Glu Pro Phe Val Ala Cys Asn Glu Cys Ala Phe Pro Val Cys Arg
  50          55          60
Pro Cys Tyr Glu Tyr Glu Arg Arg Glu Gly Asn Gln Ala Cys Pro Gln
  65          70          75          80
Cys Lys Thr Arg Phe Lys Arg Leu Lys Gly Ser Pro Arg Val Glu Gly
  85          90          95
Asp Glu Glu Glu Asp Asp Ile Asp Asp Leu Asp Asn Glu Phe Glu Tyr
 100         105         110
Gly Asn Asn Gly Ile Gly Phe Asp Gln Val Ser Glu Gly Met Ser Ile
 115         120         125
Ser Arg Arg Asn Ser Gly Phe Pro Gln Ser Asp Leu Asp Ser Ala Pro
 130         135         140
Pro Gly Ser Gln Ile Pro Leu Leu Thr Tyr Gly Asp Glu Asp Val Glu
 145         150         155         160
Ile Ser Ser Asp Arg His Ala Leu Ile Val Pro Pro Ser Leu Gly Gly
 165         170         175
His Gly Asn Arg Val His Pro Val Ser Leu Ser Asp Pro Thr Val Ala
 180         185         190

```

-continued

Ala His Arg Arg Leu Met Val Pro Gln Lys Asp Leu Ala Val Tyr Gly
195 200 205

Tyr Gly Ser Val Ala Trp Lys Asp Arg Met Glu Glu Trp Lys Arg Lys
210 215 220

Gln Asn Glu Lys Leu Gln Val Val Arg His Glu Gly Asp Pro Asp Phe
225 230 235 240

Glu Asp Gly Asp Asp Ala Asp Phe Pro Met Met Asp Glu Gly Arg Gln
245 250 255

Pro Leu Ser Met Lys Ile Pro Ile Lys Ser Ser Lys Ile Asn Pro Tyr
260 265 270

Arg Met Leu Ile Val Leu Arg Leu Val Ile Leu Gly Leu Phe Phe His
275 280 285

Tyr Arg Ile Leu His Pro Val Lys Asp Ala Tyr Ala Leu Trp Leu Ile
290 295 300

Ser Val Ile Cys Glu Ile Trp Phe Ala Val Ser Trp Val Leu Asp Gln
305 310 315 320

Phe Pro Lys Trp Tyr Pro Ile Glu Arg Glu Thr Tyr Leu Asp Arg Leu
325 330 335

Ser Leu Arg Tyr Glu Lys Glu Gly Lys Pro Ser Gly Leu Ser Pro Val
340 345 350

Asp Val Phe Val Ser Thr Val Asp Pro Leu Lys Glu Pro Pro Leu Ile
355 360 365

Thr Ala Asn Thr Val Leu Ser Ile Leu Ala Val Asp Tyr Pro Val Asp
370 375 380

Lys Val Ala Cys Tyr Val Ser Asp Asp Gly Ala Ala Met Leu Thr Phe
385 390 395 400

Glu Ala Leu Ser Glu Thr Ala Glu Phe Ala Arg Lys Trp Val Pro Phe
405 410 415

Cys Lys Lys Tyr Cys Ile Glu Pro Arg Ala Pro Glu Trp Tyr Phe Cys
420 425 430

His Lys Met Asp Tyr Leu Lys Asn Lys Val His Pro Ala Phe Val Arg
435 440 445

Glu Arg Arg Ala Met Lys Arg Asp Tyr Glu Glu Phe Lys Val Lys Ile
450 455 460

Asn Ala Leu Val Ala Thr Ala Gln Lys Val Pro Glu Asp Gly Trp Thr
465 470 475 480

Met Gln Asp Gly Thr Pro Trp Pro Gly Asn Ser Val Arg Asp His Pro
485 490 495

Gly Met Ile Gln Val Phe Leu Gly Ser Asp Gly Val Arg Asp Val Glu
500 505 510

Asn Asn Glu Leu Pro Arg Leu Val Tyr Val Ser Arg Glu Lys Arg Pro
515 520 525

Gly Phe Asp His His Lys Lys Ala Gly Ala Met Asn Ser Leu Ile Arg
530 535 540

Val Ser Gly Val Leu Ser Asn Ala Pro Tyr Leu Leu Asn Val Asp Cys
545 550 555 560

Asp His Tyr Ile Asn Ser Lys Ala Leu Arg Glu Ala Met Cys Phe
565 570 575

Met Met Asp Pro Gln Ser Gly Lys Lys Ile Cys Tyr Val Gln Phe Pro
580 585 590

Gln Arg Phe Asp Gly Ile Asp Arg His Asp Arg Tyr Ser Asn Arg Asn
595 600 605

-continued

Val	Val	Phe	Phe	Asp	Ile	Asn	Met	Lys	Gly	Leu	Asp	Gly	Leu	Gln	Gly
610						615					620				
Pro	Ile	Tyr	Val	Thr	Gly	Cys	Val	Phe	Arg	Arg	Gln	Ala	Leu	Tyr	Gly
625					630					635					640
Phe	Asp	Ala	Pro	Lys	Lys	Lys	Lys	Gly	Pro	Arg	Lys	Thr	Cys	Asn	Cys
				645					650					655	
Trp	Pro	Lys	Trp	Cys	Leu	Leu	Cys	Phe	Gly	Ser	Arg	Lys	Asn	Arg	Lys
			660					665					670		
Ala	Lys	Thr	Val	Ala	Ala	Asp	Lys	Lys	Lys	Lys	Asn	Arg	Glu	Ala	Ser
		675					680					685			
Lys	Gln	Ile	His	Ala	Leu	Glu	Asn	Ile	Glu	Glu	Gly	Arg	Gly	His	Lys
	690					695					700				
Val	Leu	Asn	Val	Glu	Gln	Ser	Thr	Glu	Ala	Met	Gln	Met	Lys	Leu	Gln
705					710					715					720
Lys	Lys	Tyr	Gly	Gln	Ser	Pro	Val	Phe	Val	Ala	Ser	Ala	Arg	Leu	Glu
				725					730					735	
Asn	Gly	Gly	Met	Ala	Arg	Asn	Ala	Ser	Pro	Ala	Cys	Leu	Leu	Lys	Glu
			740					745					750		
Ala	Ile	Gln	Val	Ile	Ser	Arg	Gly	Tyr	Glu	Asp	Lys	Thr	Glu	Trp	Gly
		755					760					765			
Lys	Glu	Ile	Gly	Trp	Ile	Tyr	Gly	Ser	Val	Thr	Glu	Asp	Ile	Leu	Thr
	770					775					780				
Gly	Ser	Lys	Met	His	Ser	His	Gly	Trp	Arg	His	Val	Tyr	Cys	Thr	Pro
785					790					795					800
Lys	Leu	Ala	Ala	Phe	Lys	Gly	Ser	Ala	Pro	Ile	Asn	Leu	Ser	Asp	Arg
				805					810					815	
Leu	His	Gln	Val	Leu	Arg	Trp	Ala	Leu	Gly	Ser	Val	Glu	Ile	Phe	Leu
			820					825					830		
Ser	Arg	His	Cys	Pro	Ile	Trp	Tyr	Gly	Tyr	Gly	Gly	Gly	Leu	Lys	Trp
		835					840					845			
Leu	Glu	Arg	Leu	Ser	Tyr	Ile	Asn	Ser	Val	Val	Tyr	Pro	Trp	Thr	Ser
	850					855					860				
Leu	Pro	Leu	Ile	Val	Tyr	Cys	Ser	Leu	Pro	Ala	Ile	Cys	Leu	Leu	Thr
865					870					875					880
Gly	Lys	Phe	Ile	Val	Pro	Glu	Ile	Ser	Asn	Tyr	Ala	Ser	Ile	Leu	Phe
				885					890					895	
Met	Ala	Leu	Phe	Ser	Ser	Ile	Ala	Ile	Thr	Gly	Ile	Leu	Glu	Met	Gln
			900					905					910		
Trp	Gly	Lys	Val	Gly	Ile	Asp	Asp	Trp	Trp	Arg	Asn	Glu	Gln	Phe	Trp
		915					920					925			
Val	Ile	Gly	Gly	Val	Ser	Ala	His	Leu	Phe	Ala	Leu	Phe	Gln	Gly	Leu
	930					935					940				
Leu	Lys	Val	Leu	Ala	Gly	Val	Asp	Thr	Asn	Phe	Thr	Val	Thr	Ser	Lys
945					950					955					960
Ala	Ala	Asp	Asp	Gly	Glu	Phe	Ser	Asp	Leu	Tyr	Leu	Phe	Lys	Trp	Thr
				965					970					975	
Ser	Leu	Leu	Ile	Pro	Pro	Met	Thr	Leu	Leu	Ile	Ile	Asn	Val	Ile	Gly
			980					985					990		
Val	Ile	Val	Gly	Val	Ser	Asp	Ala	Ile	Ser	Asn	Gly	Tyr	Asp	Ser	Trp
	995						1000					1005			
Gly	Pro	Leu	Phe	Gly	Arg	Leu	Phe	Phe	Ala	Leu	Trp	Val	Ile	Ile	His
	1010					1015					1020				
Leu	Tyr	Pro	Phe	Leu	Lys	Gly	Leu	Leu	Gly	Lys	Gln	Asp	Arg	Met	Pro

-continued

```

1025          1030          1035          1040
Thr Ile Ile Val Val Trp Ser Ile Leu Leu Ala Ser Ile Leu Thr Leu
          1045          1050          1055
Leu Trp Val Arg Val Asn Pro Phe Val Ala Lys Gly Gly Pro Ile Leu
          1060          1065          1070
Glu Ile Cys Gly Leu Asp Cys Leu
          1075          1080

```

```

<210> SEQ ID NO 6
<211> LENGTH: 56
<212> TYPE: PRT
<213> ORGANISM: Populus tremuloides

```

```

<400> SEQUENCE: 6

```

```

Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val
  1          5          10          15
Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser
  20          25          30
Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Lys Glu Gly Arg
  35          40          45
Lys Val Cys Leu Arg Cys Gly Ser
  50          55

```

```

<210> SEQ ID NO 7
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: concensus sequence

```

```

<400> SEQUENCE: 7

```

```

Gln Val Leu Arg Trp
  1          5

```

What is claimed is:

1. A method of expressing a polypeptide in a plant cell comprising transforming a plant cell with a recombinant construct comprising a sub-fragment of SEQ ID NO:3 having at least 40 consecutive nucleotides from positions 334 to 522 of SEQ ID NO:3 operably linked to a coding sequence, wherein expression of the coding sequence results in expression of the polypeptide.

2. The method of claim 1, wherein the polypeptide comprises a cellulose synthase or a UDP-glucose binding domain.

3. The method of claim 1, wherein the sub-fragment comprises nucleotides from positions 334 to 522 of SEQ ID NO:3.

4. The method of claim 1, wherein the sub-fragment comprises nucleotides from positions 334 to 422 of SEQ ID NO:3.

5. The method of claim 1, wherein the sub-fragment comprises nucleotides from positions 423 to 522 of SEQ ID NO:3.

6. A transgenic plant cell produced by the method of claim 1.

7. The plant cell of claim 6, wherein the plant cell is a tree cell.

8. A method of inducing expression of a coding sequence in a plant cell comprising the coding sequence operably linked to a sub-fragment of SEQ ID NO:3 having at least 40 consecutive nucleotides from positions 334 to 522 of SEQ ID NO:3, comprising exposing the cell to mechanical stress thereby inducing expression of the coding sequence.

40 9. The method of claim 8, wherein the coding sequence encodes a cellulose synthase, and wherein expression of the coding sequence alters a characteristic of a plant selected from growth, cellulose content, lignin content, juvenile wood strength, fiber strength, or a combination thereof, compared to a control plant.

10. A plant produced by the method of claim 8.

11. The plant of claim 10, wherein the plant is a tree.

12. The method of claim 1, wherein the plant cell is a tree cell.

50 13. The method of claim 1, wherein the coding sequence encodes a cellulose synthase, and wherein expression of the coding sequence alters a characteristic of a plant selected from growth, cellulose content, lignin content, juvenile wood strength, fiber strength, or a combination thereof, compared to a control plant.

55 14. The method of claim 8, wherein the coding sequence encodes a cellulose synthase or a UDP-glucose binding domain.

60 15. The method of claim 8, wherein the sub-fragment comprises nucleotides from positions 334 to 522 of SEQ ID NO:3.

16. The method of claim 8, wherein the sub-fragment comprises nucleotides from positions 334 to 422 of SEQ ID NO:3.

17. The method of claim 8, wherein the sub-fragment comprises nucleotides from positions 423 to 522 of SEQ ID NO:3.

* * * * *