



**Michigan
Technological
University**

Michigan Technological University
Digital Commons @ Michigan Tech

Michigan Tech Patents

Vice President for Research Office

6-19-2007

Cellulose synthase promoter and method for modifying cellulose and lignin biosynthesis in plants

Vincent Lee C. Chiang
Michigan Technological University

Chandrashekhar P. Joshi
Michigan Technological University, cpjoshi@mtu.edu

Luguang Wu

Daniel T. Carraway


Follow this and additional works at: <https://digitalcommons.mtu.edu/patents>

 Part of the [Engineering Commons](#)

Recommended Citation

Chiang, Vincent Lee C.; Joshi, Chandrashekhar P.; Wu, Luguang; and Carraway, Daniel T., "Cellulose synthase promoter and method for modifying cellulose and lignin biosynthesis in plants" (2007). *Michigan Tech Patents*. 10.
<https://digitalcommons.mtu.edu/patents/10>

Follow this and additional works at: <https://digitalcommons.mtu.edu/patents>

 Part of the [Engineering Commons](#)



US007232941B2

(12) **United States Patent**
Chiang et al.

(10) **Patent No.:** **US 7,232,941 B2**
(45) **Date of Patent:** **Jun. 19, 2007**

(54) **CELLULOSE SYNTHASE PROMOTER AND METHOD FOR MODIFYING CELLULOSE AND LIGNIN BIOSYNTHESIS IN PLANTS**

(75) Inventors: **Vincent L. Chiang**, Hancock, MI (US); **Luguang Wu**, Kenmore (AU); **Chandrashekhar P. Joshi**, Houghton, MI (US); **Daniel T. Carraway**, Bainbridge, GA (US)

(73) Assignee: **Board of Control of Michigan Technological University**, Houghton, MI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/437,368**

(22) Filed: **May 19, 2006**

(65) **Prior Publication Data**

US 2006/0206964 A1 Sep. 14, 2006

Related U.S. Application Data

(62) Division of application No. 09/980,043, filed as application No. PCT/US00/13637 on May 18, 2000, now Pat. No. 7,049,481.

(60) Provisional application No. 60/135,280, filed on May 21, 1999.

(51) **Int. Cl.**

C12N 15/09 (2006.01)
C12N 15/82 (2006.01)
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **800/287**; 800/278; 800/298; 800/295; 800/290; 536/24.1; 435/320.1; 435/468

(58) **Field of Classification Search** 800/287, 800/278, 290, 298; 536/24.1; 435/468, 435/320.1

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,268,274 A 12/1993 Ben-Bassat et al.
5,633,439 A 5/1997 Walter et al.
5,646,023 A 7/1997 Secor et al.
5,654,190 A 8/1997 Matsunaga et al.
6,013,860 A 1/2000 Himmel et al.
6,271,443 B1 8/2001 Stalker et al.
6,303,847 B1 10/2001 Kawaoka et al.

FOREIGN PATENT DOCUMENTS

CA 2012311 9/1990
EP 875575 11/1998
WO WO 98/00549 1/1998
WO WO 98/18949 5/1998
WO WO 00/22092 4/2000

OTHER PUBLICATIONS

Accession No. T10797—Sequence Search Result (1996) 1-2.

Accession No. AF072131 [gi: 4115904], pp. 1-2, Sequence Search Result, Jan. 8, 1999.

Accession No. AF072131 [gi: 3243277], pp. 1, Sequence Search Result, Jun. 23, 1998.

Altmorbe et al., *Mol. Plant-Microbe Interac.* 2: 301, (1989).

Animal Cell Culture, R.I. Freshney, ed. (1986), "Immobilised Cells and Enzymes" IRL Press.

Arioli et al., *Science*, 279: 717-720, (1998).

Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

Benton and Davis, *Science* 196:180, (1977).

Bird et al., "Manipulation of Plant Gene expression by Antisense RNA," *Biotechnology & Genetic Engineering Reviews* (1991) 9:207-227.

Brown et al., *Trends Pl Sci.*, 1: 149-156 (1996).

Bugos et al., *Biotechniques* 19:734-737, (1995).

Database EMBL Online, Jun. 23, 1998, Wu, L., Joshi, CP, Chiang, VL, "Populus tremuloides secondary xylem cellulose synthase (ce1A) mRNA, complete cds." EBI Database accession No. af072131, abstract.

Database Swall Online, May 1, 1997, "Cellulose synthase, CELAI", accession No. P93155.

Database Swall Online, Nov. 1, 1998, Secondary xylem cellulose synthase (ce1A) mRNA, complete cds. EBI Database accession No. 081368, abstract.

Delmer and Amor, *Pl Cell*, 7: 987-1000, (1995).

DNA Cloning: A Practical Approach, vols. I and II (D.N. Glover ed. 1985).

Esau, K., *Anatomy of Seed Plants*, New York: John Wiley and Sons. (1960).

Esau, K., *Plant Anatomy*, 2nd ed. New York: Wiley (1953).

Esau, K., *Vascular Differentiation in Plants*, New York: Hold, Rinehart & Winston (1965).

Esau, K., et al., "Observations on Cytokinesis", *Planta (Berl.)* 67, 168-181 (1965).

Esau, K., "Anatomy of Plant Virus Infections", *Annu. Rev. Phytopathol.* 5:45-76 (1967).

Fukuda, *Ann Rev Pl Physiol Pl Mol Biol*, 47: 299-325, (1996).

Fullner and Nester, *J. Bacteriol.* 178: 1498, (1996).

Fullner et al., *Science* 273: 1107, (1996).

Ge and Chiang, 1996, *Pl Physiol*, 112: 861, (1996).

Grunstein and Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961, (1975).

(Continued)

Primary Examiner—Medina A. Ibrahim

(74) *Attorney, Agent, or Firm*—Michael, Best & Friedrich LLP

(57) **ABSTRACT**

This invention relates to an isolated cellulose synthase promoter, methods for genetically altering cellulose and lignin biosynthesis, and to methods for improving strength properties of juvenile wood and fiber in trees. The invention further relates to methods for identifying regulatory elements in a cellulose synthase promoter and to methods for augmenting expression of polynucleotides operably linked to a cellulose synthase promoter.

OTHER PUBLICATIONS

- Haigler and Blanton, PNAS, 93: 12082-12085, (1996).
- Higuchi, "Using PCR to Engineer DNA", in PCR Technology: Principles and Applications for DNA Amplification, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70. (1989).
- Higuchi, Biochemistry and Molecular Biology of Wood, Springer Verlag (1997).
- Hoffman and Stoffel, Biol Chem, Hoppe-Seyler 374: 166, (1993).
- Hu et al., Nature Biotechnology, In Press, (1999).
- Hu et al., PNAS, 95: 5407-5412, (1998).
- Hutchinson, C., et al., "Mutagenesis at a Specific Position in a DNA Sequence", J. Biol. Chem., 253:6551 (1978).
- Hutchinson et al., Proc. Natl. Acad. Sci. U.S.A. 83:710, (1986).
- Joshi et al., PMB, 35: 993-1001, (1997).
- Joshi, 1987, NAR, 15: 6643-6653, (1987).
- Joshi, 1987, NAR, 15: 9627-9640, (1987).
- Lazar et al., "Transforming Growth Factor α Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities," Molec. & Cell. Biol. (1988) 8:3:1247-1252.
- Lu, Shan-Fa, et al., Molecular cloning and characterization of three cellulose synthases associated with xylem development in *Eucalyptus grandis*, Poster presented at American Society of Plant Biologists annual meeting at Denver, CO, Aug. 2-7, 2002.
- Matton, et al., "Identification of cis-acting elements involved in the Regulation of the Pathogenesis-related Gene STH-2 in Potato" Plant Molecular Biology, 102:279-291, (1993).
- Nakai, et al., Control of expression by the cellulose synthase (bcsA) promoter region from *Acetobacter xylinum* BPR 2002 Gene, 213(1-2):93-100, (1998).
- Needleman & Wunsch, J. Mol. Biol., 48:443-453, (1970).
- Nucleic Acid Hybridization, B.D. Hames & S.J. Higgins eds. (1985).
- Oligonucleotide Synthesis (M.J. Gait ed. 1984).
- Oliphant et al., Gene 44:177, (1986).
- Pear, et al., "Higher Plants Contain Homologs of the Bacterial celsA Genes Encoding The Catalytic Subunit of Cellulose Synthase" PNAS, 93:12637-12642, (1996).
- Perbal, B., A Practical Guide To Molecular Cloning (1984).
- Salvucci, et al., "Identification of the Uridine-Binding Domain of Sucrose-Phosphate Synthase" Plant Physiology, 102:529-536, (1993).
- Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, (1989).
- Saxena, et al., "Cloning and Sequencing of the Cellulose Synthase Catalytic Subunit Gene of the *Acetobacter xylinum*" Plant Molecular Biology, 15:673-683, (1990).
- Smith et al., "Antisense RNA Inhibition of Polygalacturonase Gene Expression in Transgenic Tomatoes," Nature (1988) 334:724-726.
- Timmell, Compression Wood in Gymnosperms, Springer Verlag, (1986).
- Transcription And Translation, B.D. Hames & S.J. Higgins, eds. (1984).
- Whetten et al., Ann Rev Pl Physiol Pl Mol Biol, 49: 585-609, (1998).
- Wu and Wu, J. Biol. Chem. 263:14621-14624, (1988).
- Wu et al., J. Biol. Chem. 267:963-967, (1992).
- Wu et al., Pl Physiol, 117: 1125, (1998).
- Ye and Varner, Plant Physiology 103(3):805-813 (Nov. 1993).
- Ye and Varner, Methods in Enzymology 218:671-681 (1993).
- Zhou Y., et al., "Molecular cloning of cellulose synthase genes from loblolly opine", Poster presented at American Society of Plant Biologists annual meeting at Denver, CO, Aug. 2-7, 2002.
- Zoller and Smith, DNA 3:479-488, (1984).

Fig. 1: DNA and predicted protein sequence of PtCela cDNA

1 GTCGACCCACGCGTCCGTCTTGAAGAATATGAAGTTGTAAAGAGCTGGTAAAGTGGTAA 60
61 TAAGCAAGATGATGGAATCTGGGGCTCTATATGCCATACCTGTGGTGAACAGGTGGGGC 120
M M E S G A P I C H T C G E Q V G H
121 ATGATGCAAATGGGGAGCTATTTGTGGCTTGCCATGAGTGTAGCTATCCCATGTGCAAGT 180
D A N G E L F V A C H E C S Y P M C K S
181 CTTGTTTCGAGTTTGAATCAATGAGGGCCGAAAGTTTGCCTGCGGTGTGGCTCGCCAT 240
C F E F E I N E G R K V C L R C G S P Y
241 ATGATGAGAACTTGCTGGATGATGTAGAAAAGAAGGGGTCTGGCAATCAATCCACAATGG 300
D E N L L D D V E K K G S G N Q S T M A
301 CATCTCACCTCAACGATTCTCAGGATGTCGGAATCCATGCTAGACATATCAGTAGTGTGT 360
S H L N D S Q D V G I H A R H I S S V S
361 CCACTGTGGATAGTGAATGAATGATGAATATGGGAATCCAATTTGGAAGAATCGGGTGA 420
T V D S E M N D E Y G N P I W K N R V K
421 AGAGCTGTAAGGATAAAGAGAACAAGAAGAAAAGAGAAGTCTAAGGCTGAAACTGAAC 480
S C K D K E N K K K K R S P K A E T E P
481 CAGCTCAAGTTCTACAGAACAGCAGATGGAAGAGAAAACCGTCTGCAGAGGCTTCGGAGC 540
A Q V P T E Q Q M E E K P S A E A S E P
541 CGCTTTC AAT TGT TTTATCCAATCCACGCAACAAGCTCACACCATACAGAGCAGTGATCA 600
L S I V Y P I P R N K L T P Y R A V I I
601 TTATGCGACTGGTCATTCTGGGCCCTTCTTCCACTTCAGAATAACAAATCCTGTGCATA 660
M R L V I L G L F F H F R I T N P V D S
661 GTGCCTTTGGCCTGTGGCTTACTTCTGTGCATATGTGAGATCTGGTTTGCATTTTCTTGGG 720
A F G L W L T S V I C E I W F A F S W V
721 TGTTGGATCAGTTCCCAAGTGAATCCTGTCAATAGAGAAACGTATATCGAAAGGCTGT 780
L D Q F P K W N P V N R E T Y I E R L S
781 CGGCAAGGTATGAAAGAGAGGGTGAGCCTTCTCAGCTTGCTGGTGTGGATTTTTTCGTGA 840
A R Y E R E G E P S Q L A G V D F F V S
841 G T A C T G T T G A T C C G C T G A A G G A A C C G C A T T G A T C A C T G C C A A T A C A G T C C T T T C C A T C C 900
T V D P L K E P P L I T A N T V L S I L
901 T T G C T G T G G A C T A T C C C G T C G A T A A A G T C T C C T G C T A C G T G T C T G A T G A T G G T G C A G C T A 960
A V D Y P V D K V S C Y V S D D G A A M
961 T G C T T T C A T T T G A A T C T C T T G T A G A A C A G C T G A G T T T G C A A G G A A G T G G G T T C C G T T C T 1020
L S F E S L V E T A E F A R K W V P F C
1021 G C A A A A A A T T C T C A A T T G A A C C A A G A G C A C C G G A G T T T T A C T T C T C A C A G A A A A T T G A T T 1080
K K F S I E P R A P E F Y F S Q K I D Y
1081 A C T T G A A G A C A A G G T T C A A C C T T C T T T C G T G A A G A A C G T A G A G C A A T G A A A A G G G A T T 1140
L K D K V Q P S F V K E R R A M K R D Y
1141 A T G A A G A G T A C A A A G T C C G A G T T A A T G C C C T G G T A G C A A A G G C T C A G A A A C A C C T G A A G 1200
E E Y K V R V N A L V A K A Q K T P E E

Fig. 1 (continued)

1201 AAGGATGGACTATGCAAGATGGAACACCTTGGCCTGGGAATAACACACGATCACCCCTG 1260
G W T M Q D G T P W P G N N T R D H P G

1261 GGCATGATTCAGGTCTTCCCTGGGAAATAC TGGGAGCTCGTGACATTGAAGGAAATGAAC 1320
H D S G L P W E I L G A R D I E G N E L

1321 TACCTCGTCTAGTATATGTCTCCAGGGAGAGACCTGGCTACCAGCACCACAAAAGG 1380
P R L V Y V S R E K R P G Y Q H H K K A

1381 CTGGTGCAGAAAATGCTCTGGTGAGAGTGTCTGCAGTACTCACAATGCTCCCTACATCC 1440
G A E N A L V R V S A V L T N A P Y I L

1441 TCAATGTTGATGTGTGATCACTATGTA AACAA TAGCAAGGCTGTTTCGAGAGGCAATGTGCA 1500
N V D C D H Y V N N S K A V R E A M C I

1501 TCCTGATGGACCCACAAGTAGGTCGAGATGTATGCTATGTGCAGTTCCCTCAGAGGTTTG 1560
L M D P Q V G R D V C Y V Q F P Q R P D

1561 ATGGCATAGATAAGAGTGTATCGCTACGCCAATCGTAACGTAGTTTTCTTTGATGTTAACA 1620
G I D K S D R Y A N R N V V F F D V N M

1621 TGAAAGGGTTGGATGGCATTCAAGGACCAGTATACGTAGGAAGCTGGTTGTGTTTCAACA 1680
K G L D G I Q G P V Y V G T G C V F N R

1681 GGCAAGCACTTTACGGCTACGGGCTCCTTCTATGCCAGCTTACGCAAGAGAAAAGGATT 1740
Q A L Y G Y G P P S M P S L R K R K D S

1741 CTTTCATCCTGCTTCTCATGTTGCTGCCCTCAAAGAAGAAGCCTGCTCAGATCCAGCTG 1800
S S C F S C C C P S K K K P A Q D P A E

1801 AGGTATACAGAGATGCAAAAAGAGAGGATCTCAATGCTGCCATATTTAATCTTACAGAGA 1860
V Y R D A K R E D L N A A I F N L T E I

1861 TTGATAATTATGACGAGCATGAAAGGTCATGCTGATCTCCAGTTGAGCTTTGAGAAAA 1920
D N Y D E H E R S M L I S Q L S F E K T

1921 CTTTTGGCTTATCTTCTGTCTTCATTGAGTCTACACTAATGGAGAATGGAGGAGTACCCG 1980
F G L S S V F I E S T L M E N G G V P E

1981 AGTCTGCCAACTCACCACCATTCATCAAGGAAGCGATTCAAGTCATCGGCTGTGGCTATG 2040
S A N S P P F I K E A I Q V I G C G Y E

2041 AAGAGAAGACTGAATGGGAAAACAGATTGGTTGGATATATGGGTCAGTCACTGAGGATA 2100
E K T E W G K Q I G W I Y G S V T E D I

2101 TCTTAAGTGGCTTCAAGATGCACTGCCGAGGATGGAGATCAATTTACTGCATGCCCGTAA 2160
L S G F K M H C R G W R S I Y C M P V R

2161 GGCCTGCATTCAAAGGATCTGCACCCATCAACCTGTCTGATAGATTGCACCAGGTCCTCC 2220
P A F K G S A P I N L S D R L H Q V L R

2221 GATGGGCTCTTGGTCTGTGGAAATTTCTTTAGCAGACACTGTCCCCTCTGGTACGGGT 2280
W A L G S V E I F F S R H C P L W Y G F

2281 TTGGAGGAGCCGCTTAAATGGCTCCAAGGCTTGGGTATATAAACACCATTTGTGTACC 2340
G G G R L K W L Q R L A Y I N T I V Y P

2341 CATTACATCCCTCCCTCTCATTGCCATATGCACAATTCCTGCAGTTTGTCTGCTCACCG 2400
F T S L P L I A Y C T I P A V C L L T G

2401 GAAATTCATCATACCAACCGCTCTCAAACCTGGCAAGCATGCTGTTTCTTGGCCTCTTTA 2460
K F I I P T L S N L A S M L F L G L F I

2461 TCTCCAATCATGTAAC TGGGCTTGAAGTGGAGCGGTGTCAGCATTGAAGATT 2520

Fig. 1 (continued)

S I I V T A V L E L R W S G V S I E D L

2521 TATGGCGTAATGAACAATTCGGGTGATCGGAGGTGTTTCAGCCCATCTCTTTGCGGTCT 2580
W R N E Q F W V I G G V S A H L F A V P

2581 TCCAGGGATTCCTAAAAATGTTGGCTGGCATCGATACGAACTTCACTGTCACAGCAAAG 2640
Q G F L K M L A G I D T N F T V T A K A

2641 CAGCCGAAGATGCAGAATTTGGGGAGCTATATATGGTCAAGTGGACAACACTTTTGATTTC 2700
A E D A E F G E L Y M V K W T T L L I P

2701 CTCCAACCACACTTCTCATTATCAATATGTCGGGTTGTGCTGGATTCTCTGATGCACTCA 2760
F T T L L I I N M S G C A G P S D A L N

2761 ACAAAGGATATGAAGCATGGGGCTCTCTTTGGCAAGGTGTTCTTTGCTTTCTGGGTGA 2820
K G Y E A W G P L F G K V F F A F W V I

2821 TTCTTCATCTCTATCCATTCTTAAAGGTCTAATGGGTCGCCAAAACCTAACACCAACCA 2880
L H L Y P F L K G L M G R Q N L T P T I

2881 TTGTGTTCTCTCGGTCACTGCTGTGGCCTCTGTCTTCTCTCTCGTTGGGTCAAGATCA 2940
V V L W S V L L A S V F S L V W V K I N

2941 ATCCATTGTTAACAAGTTGATAACACCTTGGTTGCGGAGACCTGCATTTCCATTGATT 3000
P F V N K V D N T L V A E T C I S I D C

3001 GCTGAGCTACCTCCAATAAGTCTCTCCAGTATTTTGGGGTTACAAAACCTTTGGGAATT 3060

3061 GGAATATGATCCTCGTTGTAGTTCCCTCAAGAAAGCACATATCGCTGTCAGTATTTAAA 3120

3121 TGAAGTCAAGATGATTGTTCTCTATGAAGTTTGAACAGTTTGAAATGATATTATGTTA 3180

3181 AAATACAGGTTTTGATTGTGTTGAAAAAAAAAAGAAAAAAAAAAAAAAAAA 3232




Figure 2. Southern blot analysis of aspen genomic DNA at low (a) and high (b) stringency. DNA (25 μ g per lane) was digested with Pst I (lane P), HindIII (H), and EcoRI (E), respectively, and probed with a 32 P-labeled 1-kb 5' end *PtCela* cDNA fragment. (c) Northern blot analysis of total RNA (40 μ g per lane) from aspen stem internodes, probed with 32 P-labeled *PtCela* cDNA. 1-2, 1st and 2nd; 3-4, 3rd and 4th; 5-6, 5th and 6th; 9-10, 9th and 10th internodes.

FIG. 2




Figure 3. *In situ* localization of *PtCelA* gene transcripts. Transverse sections (10 μ m) from 2nd (a), 4th (b) and 6th (c) aspen stem internodes were hybridized with DIG-labeled *PtCelA* antisense RNA probe or with a DIG-labeled *PtCelA* sense RNA probe as the control (d, sections from the 5th internode) according to Seidman with modifications. Positive RNA-RNA hybridization signals are stained purple blue. Arrows indicate the cellular localization of the *PtCelA* transcripts. No hybridization signal is detectable in control section. PX, primary xylem cells; SX, secondary xylem cells. Bars = 100 μ m.

FIG. 3

Fig. 4: DNA sequence of PtCelAP, the 5' flanking region of PtCela coding sequence

```

1  GAATTCGCCCTTTTGAATTCAGGAGACGATAGTTTCCGGTTCGTTGAATGGCTTTGTTCÄ 60
61  CTCTGGTCTAGCAATTTGCAAAAGAAGTTACAAAACAAATGCATATTATGTAAATTTAA 120
121 CAAGAGATGGGTTCTATGGTCACTTATTTATGCCCATCATTTGTTCTGGGGTTACTCTTT 180
181 ATAGTCTGATTCGAAGTTGCAAACGCCGTTTCTGGTATTGCAATTATGTAGCCATAAAC 240
241 TGTTAATCCTGTAGCTATTAGCGGACCAACAACCAGATATACGGGATCAGCGTCGTAÄÄÄ 300
301 GAGATCTCCATTCTACGTTTCTTTCTAATTTTCCGTTTCAGTGAGAGAATTACCCTGAT 360
361 ACATTGACATGATGATTGATGATTATGGGAACCATCCGATGTTAGACACGAGACCATCT 420
421 GGATCCTGCCAGTTTTCAGTTCACATGGCATCTCAGCCCAAGATCATGTGTTTATACGCC 480
481 TAATGACTTGTATTGAAAGTTTGGTAAGTTGAAGATGTGCTCTGCCCAACAGAAACCTTC 540
541 CTAAATTTCCAGCAAATCTTTCAAACCTGGCCCTTACACCCCGAAAATAGACGTGCTTCT 600
601 ACTTGGGTTCTTGGAAACCATGCACCAACCGCCATACCCACCAACCCACCACCCTCAAC 660
661 CTCTCTTCGCCATTACAAAATGTCAAGTACCACCCTCTGAAAGACACCAACACACCCTA 720
721 GCTTTGGTTAGGGTATTTGATATAAAAACAAGGCCAAAACAAAAGATTGGAAGGAAGCAG 780
781 AGGAAGACCTCTTGAAAGAATTGAAGTTGTAAAGAGCTGGTAAAGTGGTAATAAGCAAG 840
841 ATGATGGAATCTGGGGCTCCTATATGCCATACCTGTGGTGAACAGGTGGGGCATGATGCA 900
    M M E S G A P I C H T C G E Q V G H D A
901 AATGGGAGCTATTTGTGGCTTGCCATGAGTGTAGCTATCCCATGTGCAAGTCTTGTTTC 960
    N G E L F V A C H E C S Y P M C K S C F
961 GAGTTTGAAATCAAAGAGGGCCGAAAGTTTGCTTGCGGTGTGGCTCGAG 1010
    E F E I K E G R K V C L R C G S

```

FIG. 5




Figure 5. Histochemical analysis of transgenic tobacco for *GUS* gene expression driven by aspen *PtCelA* gene promoter. Stem transverse sections from the 3rd (a), 5th (b), 7th (c), and 8th (d and f) internodes were stained for *GUS* activity. Fluorescence microscopy (e) showing the identical section as in (d). Lignin autofluorescence was visualized after UV irradiation. An entire section from the 8th internode stained for *GUS* activity is shown in (f). Bars = 100 μ m in a to e; bar = 1.5 mm in f.

FIG. 6




Figure 6. GUS gene expression driven by aspen PtCelA gene promoter in transgenic tobacco plants under tension stress. Tension stress was induced by bending the transgenic plants. Tangential and longitudinal sections were harvested before bending (a) and 4 (b), 20 (c) and 40 (d) hours after bending and stained for GUS expression. Arrows indicate the bend sites.

FIG. 7

Arabidopsis thaliana cellulose synthase mRNA SEQ ID NO: 4

```

1  ggggcccggg ttaatcgccg gttctcacia caggaatgag tttgtcctca ttaatgccga
61  tgagaatgcc cgaataagat cagtccaaga gctgagtgga cagacatgct aaatctgcag
121  agatgagatc gaattgactg ttgatggaga accgtttggt gcatgtaacg aatgtgcatt
181  ccctgtgtgt agaccttgct atgagtacga aagacgagaa ggcaatcaag cttgtccaca
241  gtgcaaaacc cgtttcaaac gtcttaaagg aagtccaaga gttgaagggt atgaagagga
301  agatgacatt gatgatttag acaatgagtt tgagtatgga aataatggga ttggatttga
361  tcaggtttct gaaggatgtg caatctctcg tcgcaactcc ggtttcccac aatctgattt
421  ggattcagct ccacctggct ctcagattcc attgctgact tacggcgacg aggacgttga
481  gatttcttct gatagacatg ctcttattgt tcctccttca cttggtggtc atggcaatag
541  agttcatcct gtttctcttt ctgacctgac cgtggctgca catcgaaggc tgaaggtagc
601  tcagaaagat cttgcggttt atggttatgg aagtgtcgct tggaaagatc ggatggagga
661  atggaagaga aagcagaatg agaaacttca ggttggttag catgaaggag atcctgattt
721  tgaagatggt gatgatgctg attttccaat gatggatgag ggaaggcagc cattgtctat
781  gaagatacca atcaaatcga gcaagataaa tccttaccgg atgtaattg  tgctacgtct
841  tgtgattctt ggtctcttct ttcactaccg tattcttcac cccgtcaaag atgcatatgc
901  tttgtggctt atttctgtta tatgtgagat atggtttgct gtttcatggg ttcttgatca
961  gttccctaaa tgytacccta tcgagcgaga aacgtacttg gaccgactct cattaagata
1021  tgaaaaagaa gggaaaccgt cgggactatc ccctgtggat gtatttgtaa gtacagtgga
1081  tccattgaaa gagcctccgc ttattactgc aaatactgtc ttgtctattc ttgctgttga
1141  ttatcctgtc gataaggttg cttgttacgt atctgatgat ggtgctgcta tgcttacttt
1201  cgaagctctt tctgagaccg ctgaattcgc aaggaaatgg gttcctttct gcaagaaata
1261  ttgtattgag cctcgtgctc ccgaatggta tttctgccat aaaatggact acttgaagaa
1321  taaagtctat cccgcatttg ttagggagcg gcgagccatg aagagagatt atgagaatt
1381  caaagtaaag atcaatgctt tagtagcaac agcacagaaa gtgcctgagg atggtggac
1441  tatgcaagac ggtacacctt ggccccgtaa tagtgtgcca gatcatcctg gcatgattca
1501  ggtcttctt ggaagtgacg gtgttcgtga tgtcgaaac aacgagttgc ctcgattagt
1561  ttacgtttct cgtgagaaga gaccgggatt tgatcaccat aagaaggctg gagctatgaa
1621  ttccctgata cgagtctctg gggttctatc aaatgctctt taccttctga atgtcgattg
1681  tgatcactac atcaacaata gcaaagctct tagagaagca atgtgttca tgatggatcc
1741  tcagtcaagga aagaaaatct gttatgttca gttccctcaa aggttcgatg ggattgatag
1801  gcacgatcga tactcaaatc gcaatgttgt gttctttgat atcaatatga aaggtttggg
1861  tgggctacaa gggcctatat acgtcggtac aggttggttt ttcaggaggc aagcgttta
1921  cggatttgat gcaccgaaga agaagaaggg cccacgtaag acatgcaat gctggccaaa
1981  atggtctctc ctatgttttg gttcaagaaa gaatcgtaaa gcaaagacag ttgctgcca
2041  taagaagaag aagaataggg aagcgtcaaa gcagatccac gcattagaaa atatcgaaga
2101  gggccgcggt cataaagttc ttaacgtaga acagtcaacc gaggcaatgc aaatgaagtt
2161  gcagaagaaa tatgggcagt ctctgtatct tgttgcatct gcgctctgg agaatggtgg
2221  gatggctaga aacgcaagcc cggcttctct gcttaaagaa gccatccaag tcattagtcg
2281  cggatagaaa gataaaactg aatggggaaa agagattggg tggatctatg gttctgttac
2341  cgaagatatt ctacgggtt ctaagatgca ttctcatggt tggagacatg tttattgtac
2401  accaaagtta gcggctttca aaggatcagc tccaatcaat ctttcggatc gtctccatca
2461  agttcttcga tgggcgcttg ggtcggttga gattttcttg agtaggcatt gtcctatttg
2521  gtatggttat ggaggtgggt tgaaatggct tgagcggttg tctacatta actctgtggt
2581  ttaccctggt acctctctac cgctatcgt ttactgttct ctcctgcca tctgtcttct
2641  cactggaaaa ttcatcgttc ccgagattag caactatgct agtatcctct tcatggcgtc
2701  cttctcgtcg attgcaataa cgggtattct cgagatgcaa tggggcaag ttgggatcga
2761  tgattggtgg agaaacgaac agttttgggt cattggaggt gtttctgcgc atctgtttgc
2821  tctcttccaa ggtctcctca aggttcttgc tgggtgctgac actaacttca cagtcacatc
2881  aaaagcagct gatgatggag agttctctga cctttacctc ttcaaatgga cttcactctc
2941  catccctcca atgactctac tcatcataaa cgtcattgga gtcatagtcg gagtctctga
3001  tgccatcagc aatggatagc actcgtgggg accgcttttc ggaagactgt tctttgcact
3061  ttgggtcatc attcatcttt acccgttctc taaaggtttg cttgggaaac aagatagaat
3121  gccaacctat attgtcgtct ggtccactct cctggcctcg attcttacac ttctttgggt
3181  ccgggttaat ccgtttggtg cgaaggcgg tcctattctc gagatctgtg gtttagactg
3241  cttgtgattc gattgaccgg tggatgggtt ggtgaaaaag gtttaattcc cacggatcaa
3301  agagaggtaa gagagatatt gttttacctc taaaagactc cttcattgtg ttcattagat
3361  gatgaaaaat gaaaagaaaa agaagattta attttgttac gagaattggt atttttgcaa
3421  gaatgtggtg tagatagcgg ccgc

```

FIG. 8

Arabidopsis thaliana cellulose synthase **SEQ ID NO: 5**

RPRLIAGSHNRNEFVLINADENARIRSVQELSGQTCQICRDEIE
LTVDGEPFVACNECAFVCRPCYEYERREGNQACPQCKTRFKRLKGS PRVEGDEEEDD
IDDLDFEFYGNNGIGFDQVSEGMSISRNSGFPQSDLDSAPPGSQIPLITYGDEDVE
ISSDRHALIVPPSLGGHGNRVHPVSLSDPTVAHRRLMVPQKDLAVYGYGSVAWKDRM
EEWKRKQNEKLQVVRHEGDPDFEDGDDADFPMMDEGRQPLSMKIPIKSSKINPYRMLI
VLRLVILGLFFHYRILHPVKDAYALWLISVICEIWFVSWVLDQFPKWYPIERETYLD
RLSLRYEKEGKPSGLSPVDVVFVSTVDPLKEPPLITANTVLSILAVDYPVDKVACYVSD
DGAAMLTFEALSETAEFARKWVPFCKKYCIEPRAPEWYFCHKMDYLKNKVHFAFVRER
RAMKRDYEEFKVKINALVATAQKVPEDGWTM QDGTWPWPGNSVRDHGMIQVFLGSDGV
RDVENNELPRLVYVSREKRPGFDHKKAGAMNSLIRVSGVLSNAPYLLNVDCDHYINN
SKALREAMCFMMDPQSGKKICYVQFPQRFDGIDRHDRYSNRNVVFFDINMKGLDGLQG
PIYVGTGCVFRRQALYGF DAPKKKKGPRKTCNCWPKWCLLCFGSRKNRKAKTVAADKK
KKNREASKQIHALENI EEGRGHKVLNVEQSTEAMQMKLQKKYGQSPVFVASARLENGG
MARNASPACLLKEAIQVISRGYEDKTEWGKEIGWIYGSVTE DILTGSKMHSHGWRHVY
CTPKLAAPKGSAPINLSDRHLHQVLRWALGSVEIFLSRHCP IWYGYGGGLKWLERLSYI
NSVVYPWTSPLIVYCSLPAICLLTGKFIVPEISNYASILFMALFSSIAITGILEMQW
KVGIDDWWRNEQFWVIGGVS AHLFALFQGLLKVLAGVD TNFTVTSKAADDGEFS DLY
LFWTSLLLIPMTLLIINVIGVIVGVSDAISNGYDSWG PLFGRLFFALWV I IHLYPFL
KGLLGKQDRMPTIIVVWSILLASILTLLWVRVNP FVAKGGPILEICGLDCL

1

**CELLULOSE SYNTHASE PROMOTER AND
METHOD FOR MODIFYING CELLULOSE
AND LIGNIN BIOSYNTHESIS IN PLANTS**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a divisional of U.S. application Ser. No. 09/980,043 filed Apr. 5, 2002, now U.S. Pat. No. 7,049,481, which is a 371 of International Application No. PCT/US00/13637 filed May 18, 2000, which claims priority to U.S. Provisional Application No. 60/135,280 filed May 21, 1999.

STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH OR DEVELOPMENT

Not Applicable

FIELD OF THE INVENTION

This invention relates to polynucleotide molecules encoding cellulose synthase, promoters of cellulose synthase and cellulose synthase polypeptides, methods for genetically altering cellulose and lignin biosynthesis, and methods for improving strength properties of juvenile wood and fiber in trees. The invention further relates to methods for identifying regulatory elements in a cellulose synthase promoter and transcription factors that bind to such regulatory elements, and to methods for augmenting expression of polynucleotides operably linked to a cellulose synthase promoter.

BACKGROUND OF THE INVENTION

Lignin and cellulose are the two major building blocks of plant cell walls that provide mechanical strength and rigidity. In plants, and especially in trees, these two organic materials exist in a dynamic equilibrium conferring mechanical strength, water transporting ability and protection from biotic and abiotic environmental stresses. Normally, oven-dry wood contains 30 to 50% cellulose, 20 to 30% lignin and 20 to 30% hemicellulose (Higuchi, 1997).

Proportions of lignin and cellulose are known to change with variation in the natural environment. For example, during the development of compression wood in conifers, the percentage of lignin increases from 30 to 40%, and cellulose content proportionally decreases from 40 to 30% (Timmell, 1986). Conversely, in angiosperm tension wood the percentage of cellulose increases from 30 to 40%, while lignin content decreases from 30 to 20% (Timmell, 1986).

It was recently discovered that the genetic down-regulation of a key tissue-specific enzyme from the lignin biosynthesis pathway, 4CL, results in reduction of lignin content by up to 45% in transgenic aspen trees (Hu et al., 1999). This down-regulation is also associated with a 15% increase in the cellulose content. If the converse were true, i.e., that increasing cellulose content by genetic up-regulation of cellulose biosynthesis results in reduction of lignin content, then the pulp yield could be increased. This would allow tremendous savings in chemical and energy costs during pulping because, for example, lignin must be degraded and removed during the pulping process.

Cellulose is a linear glucan consisting of β -D-1,4-linked glucose residues. It is formed by a cellulose synthase enzyme which catalyzes assembly of UDP-glucose units in plasma membrane complexes known as "particle rosettes" (Delmer and Amor, 1995). Cellulose synthase is thought to

2

be anchored to the membrane by eight transmembrane binding domains to form the basis of the cellulose biosynthesis machinery in the plant cell wall (Pear et al., 1996).

In higher plants, the glucan chains in cellulose microfibrils of primary and secondary cell walls are different in their degree of polymerization (Brown et al., 1996). For example, secondary cell walls are known to contain cellulose having a high degree of polymerization, while in primary cell walls the degree of polymerization is lower. In another example, woody cell walls suffering from tension stress produce tension wood on the upper side of a bent angiosperm tree in response to the stress. In these cells, there are elevated quantities of cellulose which have very high crystallinity. The formation of highly crystalline cellulose is important to obtain a higher tensile strength of the wood fiber. Woody cell walls located at the under side of the same stem experience a compression stress, but do not produce highly crystalline cellulose. Such variation in the degree of polymerization in cell walls during development is believed to be due to different types of cellulose synthases for organizing glucose units into different paracrystalline arrays (Haigler and Blanton, 1996). Therefore, it would be advantageous to determine the molecular basis for the synthesis of highly crystalline cellulose so that higher yields of wood pulp having superior strength properties can be obtained from transgenic trees. Production of highly crystalline cellulose in transgenic trees would also markedly improve the mechanical strength properties of juvenile wood formed in normal trees. This would be a great benefit to the industry because juvenile wood is generally undesirable for solid wood applications because it has inferior mechanical properties.

Since the deposition of cellulose and lignin in trees is regulated in a compensatory fashion, genetic augmentation of cellulose biosynthesis might have a repressive effect on lignin deposition. Since the degree of polymerization and crystallinity may depend upon the type of cellulose synthase incorporated in the cellulose biosynthesis machinery, the expression of heterologous cellulose synthase or a UDP-glucose binding region thereof (e.g., sweetgum protein expression in loblolly pine), could increase the quality of cellulose in transgenic plants. Over-expression of a heterologous cellulose synthase may also increase cellulose quantity in transgenic plants. Thus, genetic engineering of cellulose biosynthesis can provide a strategy to augment cellulose quality and quantity, while reducing lignin content in transgenic plants.

A better understanding of the biochemical processes that lead to wood formation would enable the pulp and paper industries to more effectively use genetic engineering as a tool to meet the increasing demands for wood from a decreasing production area. With this objective, many xylem-specific genes, including most lignin biosynthesis genes, have been isolated from developing xylem tissues of various plants including tree species (Ye and Vamer, 1993; Fukuda, 1996; Whetten et al., 1998). Genes regulating cellulose biosynthesis in crop plants (Pear et al., 1996 and Arioli et al., 1998), versus in trees, have also been isolated. However, isolation of tree genes which are directly involved in cellulose biosynthesis has remained a great challenge.

For more than 30 years, no gene encoding higher plant cellulose synthase (CelA) was identified. Recently, Pear et al. (1996) isolated the first putative higher plant CelA cDNA, GhCelA (GenBank No. GHU58283), by searching for UDP-glucose binding sequences in a cDNA library prepared from cotton fibers having active secondary wall cellulose synthesis. GhCelA was considered to encode a

cellulose synthase catalytic subunit because it is highly expressed in cotton fibers, actively synthesizes secondary wall cellulose, contains eight transmembrane domains, binds UDP-glucose, and contains two other domains unique to plants.

Recently, Arioli et al. (1998) cloned a C_{el}A homolog, RSW1 (radial swelling) (GenBank No. AF027172), from *Arabidopsis* by chromosome walking to a defective locus of a temperature sensitive cellulose-deficient mutant. Complementation of the RSW1 mutant with a wild type full-length genomic RSW1 clone restored the normal phenotype. This complementation provided the first genetic proof that a plant C_{el}A gene encodes a catalytic subunit of cellulose synthase and functions in the biosynthesis of cellulose microfibrils. The full-length *Arabidopsis* RSW1 represents the only known, currently available cellulose synthase cDNA available for further elucidating cellulose biosynthesis in transgenic systems (Wu et al., 1998).

The discovery of the RSW1 gene substantiated the belief that the assembly of a cellulose synthase into the plasma membrane is required for functional cellulose biosynthetic machinery and for manufacturing crystalline cellulose microfibrils in plant cell walls. Most significantly, a single C_{el}A gene, e.g. RSW1, is sufficient for the biosynthesis of cellulose microfibrils in plants, e.g. *Arabidopsis*. Thus, RSW1 is a prime target for engineering augmented cellulose formation in transgenic plants.

Since many of society's fiber, chemical and energy demands are met through the industrial-scale production of cellulose from wood, genetic engineering of the cellulose biosynthesis machinery in trees could produce higher pulp yields. This would allow greater returns on investment by pulp and paper industries. Therefore, it would be advantageous to isolate and characterize genes from trees that are involved in cellulose biosynthesis in order to improve the properties of wood.

SUMMARY OF THE INVENTION

The present invention relates to polynucleotides comprising a nucleotide sequence that encodes a cellulose synthase, regulatory sequences, including a stress-inducible promoter, of the cellulose synthase, a cellulose synthase protein or a functional domain thereof and methods for augmenting cellulose biosynthesis in plants.

Thus, in one aspect, the invention provides a polynucleotide comprising a sequence that encodes a cellulose synthase, or a polynucleotide fragment thereof, the fragment encoding a functional domain of cellulose synthase, such as a UDP-glucose binding domain. The invention also provides a cellulose synthase or a functional domain or fragment thereof, including a UDP-glucose binding domain and at least one of eight transmembrane domains. The invention further provides a cellulose synthase promoter, or a functional fragment thereof, which fragment contains one or more mechanical stress response elements (MSRE).

In another aspect, the present invention is directed to a method of improving the quality of wood by altering the quantity of cellulose in plant cells, and optionally decreasing the content of lignin in the cell. The invention also relates to a method of altering the growth or the cellulose content of a plant by expressing an exogenous polynucleotide encoding a cellulose synthase or a UDP-glucose binding domain thereof in the plant. The invention further provides a method for causing a stress-induced gene expression in a plant cell by expressing a polynucleotide of choice using a stress-inducible cellulose synthase promoter.

In yet another aspect, the invention relates to a method for determining a mechanical stress responsive element (MSRE) in a cellulose synthase promoters and a method for identifying transcription factors that binds to the MSRE.

In a further aspect, the invention provides a method for altering (increasing or decreasing) i.e., regulating, the expression of a cellulose synthase in a plant by expressing an exogenous polynucleotide encoding a transcription factor having the property of binding a positive MSRE of a cellulose synthase promoter or by expressing an antisense polynucleotide encoding a transcription factor having the property of binding a negative MSRE to block the expression of the transcription factor.

Other aspects of the invention will be appreciated by a consideration of the detailed description of the invention drawings and appended claims.

DESCRIPTION OF THE DRAWINGS

FIG. 1 represents a nucleic acid sequence encoding a cellulose synthase from *Populus tremuloides* [SEQ ID NO: 1] and the protein sequence thereof [SEQ ID NO: 2].

FIG. 2 a-c (collectively referred to as FIG. 2) represent a Southern blot analysis of aspen genomic DNA probed with a fragment of the aspen cDNA represented in FIG. 1 under low (panel a) and high stringency conditions (panel b), and a northern blot analysis of the total aspen RNA from stem internodes using the same probe (panel c).

FIG. 3 a-d (collectively referred to as FIG. 3) represent in situ localization of the cellulose synthase gene transcripts as shown in the transverse sections from second (panel a), fourth (panel b), sixth (panel c) and fifth (panel d) internode.

FIG. 4 represents a nucleic acid sequence of the 5' region of aspen cellulose synthase gene including the promoter region and the 5' portion of the coding sequence [SEQ ID NO: 3] and the peptide sequence deduced from the coding sequence [SEQ ID NO: 6].

FIG. 5 a-f (collectively referred to as FIG. 5) represents a histochemical analysis (panels a-d and f) and fluorescence microscopy (panel e) of transgenic tobacco for GUS gene expression driven by a cellulose synthase promoter of the invention.

FIG. 6 a-d (collectively referred to as FIG. 6) represents a histochemical analysis of GUS gene expression driven by aspen cellulose synthase promoter of the invention; tangential and longitudinal sections were harvested before bending (panel a), and 4 (panel b), 20 (panel c) and 40 (panel d) hours after bending and stained for GUS expression.

FIG. 7 represents a cDNA encoding cellulose synthase isolated from *Arabidopsis* [SEQ ID NO:4].

FIG. 8 represents an *Arabidopsis* cellulose synthase [SEQ ID NO:5] encoded by the cDNA represented in FIG. 7.

DETAILED DESCRIPTION OF THE INVENTION

All patents, patent applications and references cited in this specification are hereby incorporated herein by reference in their entirety. In case of any inconsistency, the present disclosure governs.

Definitions

The terms used in this specification generally have their ordinary meanings in the art, within the context of the invention, and in the specific context where each term is used. Certain terms are discussed below, or elsewhere in the specification, to provide additional guidance to the person of

skill in the art in describing the compositions and methods of the invention and how to make and use them. It will be appreciated that the same thing can be said in more than one way. Consequently, alternative language and synonyms may be used for any one or more of the terms discussed herein, nor is any special significance to be placed upon whether or not a term is elaborated or discussed herein. Synonyms for certain terms are provided. A recital of one or more synonyms does not exclude the use of other synonyms. The use of examples anywhere in this specification, including examples of any terms discussed herein, is illustrative only, and in no way limits the scope and meaning of the invention or of any exemplified term. Likewise, the invention is not limited to the preferred embodiments.

The term "plant" includes whole plants and portions of plants, including plant organs (e.g. roots, stems, leaves, etc.).

The term "angiosperm" refers to plants which produce seeds encased in an ovary. A specific example of an angiosperm is *Liquidambar styraciflua* (L.) [sweetgum].

The term "gymnosperm" refers to plants which produce naked seeds, that is, seeds which are not encased in an ovary. Specific examples of a gymnosperm include *Pinus taeda* (L.) [loblolly pine].

The term "polynucleotide" or "nucleic acid molecule" is intended to include double or single stranded genomic and cDNA, RNA, any synthetic and genetically manipulated polynucleotide, and both sense and anti-sense strands together or individually (although only sense or anti-sense stand may be represented herein). This includes single- and double-stranded molecules, i.e., DNA-DNA, DNA-RNA and RNA-RNA hybrids, as well as "protein nucleic acids" (PNA) formed by conjugating bases to an amino acid backbone. This also includes nucleic acids containing modified bases, for example thio-uracil, thio-guanine and fluorouracil.

An "isolated" nucleic acid molecule or polynucleotide refers to a component that is removed from its original environment (for example, its natural environment if it is naturally occurring). An isolated nucleic acid or polypeptide may contain less than about 50%, preferably less than about 75%, and most preferably less than about 90%, of the cellular components with which it was originally associated. A polynucleotide amplified using PCR so that it is sufficiently and easily distinguishable (on a gel, for example) from the rest of the cellular components is considered "isolated". The polynucleotides and polypeptides of the invention may be "substantially pure," i.e., having the highest degree of purity that can be achieved using purification techniques known in the art.

The term "hybridization" refers to a process in which a strand of nucleic acid joins with a complementary strand through base pairing. Polynucleotides are "hybridizable" to each other when at least one strand of one polynucleotide can anneal to a strand of another polynucleotide under defined stringency conditions. Hybridization requires that the two polynucleotides contain substantially complementary sequences; depending on the stringency of hybridization, however, mismatches may be tolerated. Typically, hybridization of two sequences at high stringency (such as, for example, in an aqueous solution of 0.5×SSC at 65°C) requires that the sequences exhibit some high degree of complementarity over their entire sequence. Conditions of intermediate stringency (such as, for example, an aqueous solution of 2×SSC at 65°C) and low stringency (such as, for example, an aqueous solution of 2×SSC at 55°C), require correspondingly less overall complementarity between the hybridizing sequences. (1×SSC is 0.15 M NaCl, 0.015 M Na

citrate.) As used herein, the above solutions and temperatures refer to the probe-washing stage of the hybridization procedure. The term "a polynucleotide that hybridizes under stringent (low, intermediate) conditions" is intended to encompass both single and double-stranded polynucleotides although only one strand will hybridize to the complementary strand of another polynucleotide.

A "sequence-conservative variant" is a polynucleotide that contains a change of one or more nucleotides in a given codon position, as compared with another polynucleotide, but the change does not result in any alteration in the amino acid encoded at that position.

A "function-conservative variant" is a polypeptide (or a polynucleotide encoding the polypeptide) having a given amino acid residue that has been changed without altering the overall conformation and function of the polypeptide, including, but not limited to, replacement of an amino acid with one having similar physico-chemical properties (such as, for example, acidic, basic, hydrophobic, and the like). Amino acids having similar physico-chemical properties are well known in the art. For example, arginine, histidine and lysine are hydrophilic-basic amino acids and may be interchangeable. Similarly, isoleucine, a hydrophobic amino acid, may be replaced with leucine, methionine or valine. Sequence- and function-conservative variants are discussed in greater detail below with respect to degeneracy of the genetic code.

A "functional domain" or a "functional fragment" refers to any region or portion of a protein or polypeptide or polynucleotide which is a region or portion of a larger protein or polynucleotide, the region or portion having the specific activity or specific function attributable to the larger protein or polynucleotide, e.g., a functional domain of cellulose synthase is the UDP-glucose binding domain.

The term "% identity" refers to the percentage of the nucleotides/amino acids of one polynucleotide/polypeptide that are identical to the nucleotides/amino acids of another sequence of polynucleotide/polypeptide as identified by program GAP from Genetics Computer Group Wisconsin (GCG) package (version 9.0) (Madison, Wis.). GAP uses the algorithm of Needleman and Wunsch (J. Mol. Biol. 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. When parameters required to run the above algorithm are not specified, the default values offered by the program are contemplated. The following parameters are used by the GCG program GAP as default values (for polynucleotides): gap creation penalty:50; gap extension penalty:3; scoring matrix: nwsgapdna.cpm (local data file).

The "% similarity" or "% homology" between two polypeptide sequences is a function of the number of similar positions shared by two sequences on the basis of the scoring matrix used divided by the number of positions compared and then multiplied by 100. This comparison is made when two sequences are aligned (by introducing gaps if needed) to determine maximum homology. PowerBlast program, implemented by the National Center for Biotechnology Information, can be used to compute optimal, gapped alignments. GAP program from Genetics Computer Group Wisconsin package (version 9.0) (Madison, Wis.) can also be used. GAP uses the algorithm of Needleman and Wunsch (J Mol Biol 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. When parameters required to run the above algorithm are not specified, the default values offered by the program are contemplated. The

following parameters are used by the GCG program GAP as default values (for polypeptides): gap creation penalty:12; gap extension penalty:4; scoring matrix:Blosum62.cpm (local data file).

The term "oligonucleotide" refers to a nucleic acid, generally of at least 10, preferably at least 15, and more preferably at least 20 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, or an mRNA molecule encoding a gene, mRNA, cDNA, or other nucleic acid of interest. Oligonucleotides can be labeled, e.g., with ³²P-nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. In one embodiment, a labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid. In another embodiment, oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of CelA, or to detect the presence of nucleic acids encoding CelA. In a further embodiment, an oligonucleotide of the invention can form a triple helix with a CelA DNA molecule. In still another embodiment, a library of oligonucleotides arranged on a solid support, such as a silicon wafer or chip, can be used to detect various polymorphisms of interest. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer. Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

The term "coding sequence" refers to that portion of the gene that contains the information for encoding a polypeptide. The boundaries of the coding sequence are determined by a start codon at the 5N (amino) terminus and a translation stop codon at the 3N (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences.

A "promoter" is a polynucleotide containing elements (e.g., a TATA box) which are capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3N direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3N terminus by the transcription initiation site and extends upstream (5N direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Examples of promoters that can be used in the present invention include PtCelAP, 4CL-1 and 35S.

The term "constitutive promoter" refers to a promoter which typically, does not require positive regulatory proteins to activate expression of an associated coding sequence, i.e., a constitutive promoter maintains some basal level of expression. A constitutive promoter is commonly used in creation of an expression cassette. An example of a constitutive promoter are 35S CaMV (Cauliflower Mosaic Virus), available from Clonetech, Palo Alto, Calif.

The term "inducible promoter" refers to the promoter which requires a positive regulation to activate expression of an associated coding sequence. An example of such a promoter is a stress-inducible cellulose synthase promoter from aspen described herein.

A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into

mRNA, which is then trans-RNA spliced and translated into the protein encoded by the coding sequence.

A "vector" is a recombinant nucleic acid construct, such as plasmid, phage genome, virus genome, cosmid, or artificial chromosome to which a polynucleotide of the invention may be attached. In a specific embodiment, the vector may bring about the replication of the attached segment, e.g., in the case of a cloning vector.

The term "expression cassette" refers to a polynucleotide which contains both a promoter and a protein coding sequence such that expression of a given protein is achieved upon insertion of the expression cassette into a cell.

A cell has been "transfected" by exogenous or heterologous polynucleotide when such polynucleotide has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous polynucleotide when the transfected polynucleotide effects a phenotypic change. Preferably, the transforming polynucleotide should be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

"Exogenous" refers to biological material, such as a polynucleotide or protein, that has been isolated from a cell and is then introduced into the same or a different cell. For example, a polynucleotide encoding a cellulose synthase of the invention can be cloned from xylem cells of a particular species of tree, inserted into a plasmid and reintroduced into xylem cells of the same or different species. The species thus contains an exogenous cellulose synthase polynucleotide.

"Heterologous polynucleotide" refers to an exogenous polynucleotide not naturally occurring in the cell into which it is introduced.

"Homologous polynucleotide" refers to an exogenous polynucleotide that naturally exists in the cells into which it is introduced.

The present invention relates to isolation and characterization of polynucleotides encoding cellulose synthases from plants, especially trees, including full length or naturally occurring forms of cellulose synthases, functional domains, promoters and regulatory elements. Therefore, in accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D. N. Glover ed. 1985); *Oligonucleotide Synthesis* (M. J. Gait ed. 1984); *Nucleic Acid Hybridization* [B. D. Hames & S. J. Higgins eds. (1985)]; *Transcription And Translation* [B. D. Hames & S. J. Higgins, eds. (1984)]; *Animal Cell Culture* [R. I. Freshney, ed. (1986)]; *Immobilized Cells And Enzymes* [IRL Press, (1986)]; B. Perbal, *A Practical Guide To Molecular Cloning* (1984); F. M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

The present invention relates to a novel, full-length cellulose synthase gene (CelA), a novel stress inducible promoter of cellulose synthases (CelAP), and cellulose synthase proteins from trees, including UDP-glucose catalytic domains thereof. The invention enables the development of transgenic tree varieties having increased cellulose content, decreased lignin content and, therefore, improved wood fiber characteristics. Production of increased cellulose quantity and quality in multiple varieties of commercially relevant, transgenic forest tree species in operational production scenarios are further contemplated. The invention

further provides a new experimental system for study of CelA gene expression and function in trees.

Polynucleotides Encoding Cellulose Synthase and Fragments Thereof.

The present invention relates to polynucleotides which comprise the nucleotide sequence that encodes cellulose synthase of the invention or a functional fragment thereof. In a preferred embodiment, the polynucleotide comprises the sequence encoding a tree cellulose synthase and most preferably, the sequence encoding a cellulose synthase from aspen. In one embodiment, a polynucleotide of the invention includes the entire cellulose synthase coding region, e.g., nucleotides 69 to 3,005 of SEQ ID NO: 1. In another aspect of the invention, the polynucleotide encoding an *Arabidopsis* cellulose synthase is provided (see SEQ ID NO:4 and the translated protein of SEQ ID NO:5).

Also within the scope of the invention are fragments of the polynucleotides encoding cellulose synthase of the invention, which fragments encode at least one transmembrane domain and/or a UDP-glucose binding domain. For example, a polynucleotide comprising the nucleotides encoding a UDP-glucose binding domain of aspen cellulose synthase (e.g., nucleotides 660 to 2250 of SEQ ID NO:1) or corresponding nucleotides of SEQ ID NO:4 are within the scope of the invention. The nucleotides encoding the UDP-glucose binding domain can be determined by, for example, alignment of protein sequences as described below.

The invention further relates to sequence conservative variants of the coding portion of SEQ ID NOS: 1 and 4.

Polynucleotides that hybridize under conditions of low, medium, and high stringency to SEQ ID NOS: 1 and 4, and their respective coding portions are also within the scope of the invention. Preferably, the polynucleotide that hybridizes to any of SEQ ID NOS: 1 and 4, or their respective coding portions, is about the same length as that sequence, for example, not more than about 10 to about 20 nucleotides longer or shorter. In another embodiment of the invention, the hybridizable polynucleotide is at least 1500 nucleotides long, preferably at least 2500 nucleotides long and most preferably at least 3000 nucleotides long. In yet another embodiment, the hybridizable polynucleotide comprises the UDP-glucose binding domain as found in SEQ ID NO: 1 or 4, or at least the conserved region QVLRW [SEQ ID NO: 7]. Most preferably, the hybridizable polynucleotide has a UDP-glucose binding activity.

The polynucleotides that occur originally in nature may be isolated from the organisms that contain them using methods described herein or well known in the art. The non-naturally occurring polynucleotides may be prepared using various manipulations known in the field of recombinant DNA. For example, the cloned CelA polynucleotide can be modified according to methods described by Sambrook et al., 1989. The sequence can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated *in vitro*. In the production of the modified polynucleotides, for example, care should be taken to ensure that the modified polynucleotide remains within the appropriate translational reading frame (if to be expressed) or uninterrupted by translational stop signals. As a further example, a CelA-encoding nucleic acid sequence can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy preexisting ones, to facilitate further *in vitro* modification. Preferably, such mutations enhance the func-

tional activity of the mutated CelA polynucleotide. Any technique for mutagenesis known in the art can be used, including but not limited to, *in vitro* site-directed mutagenesis (Hutchinson, C., et al., 1978, J. Biol. Chem. 253:6551; Zoller and Smith, 1984, DNA 3:479-488; Oliphant et al., 1986, Gene 44:177; Hutchinson et al., 1986, Proc. Natl. Acad. Sci. U.S.A. 83:710), use of TAB linkers (Pharmacia), etc. PCR techniques are preferred for site directed mutagenesis (see Higuchi, 1989, "Using PCR to Engineer DNA", in *PCR Technology: Principles and Applications for DNA Amplification*, H. Erlich, ed., Stockton Press, Chapter 6, pp. 61-70).

The polynucleotides of the present invention may be introduced into various vectors adapted for plant or non-plant replication. These are well known in the art, thus, choice, construction and use of such vectors is well within the skill of a person skilled in the art. Possible vectors include, but are not limited to, plasmids or modified viruses of plants, but the vector system must be compatible with the host cell used. An example of a suitable vector is Ti plasmid. The insertion into a cloning vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary cohesive termini. However, if the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. An expression cassette containing cellulose synthase or recombinant molecules thereof can be introduced into host cells via silicon carbide whiskers, transformed protoplasts, transformation, e.g., *Agrobacterium* vectors (discussed below), electroporation, infection, etc., so that many copies of the gene sequence are generated. Preferably, the cloned gene is contained on a shuttle vector plasmid, which provides for expansion in a cloning cell, e.g., *E. coli*, and facile purification for subsequent insertion into an appropriate expression cell line, if such is desired. For example, a shuttle vector, which is a vector that can replicate in more than one type of organism, can be prepared for replication in both *E. coli* and *Saccharomyces cerevisiae* by linking sequences from an *E. coli* plasmid with sequences from the yeast 2m plasmid.

Transgenic plants containing the polynucleotides described herein are also within the scope of the invention. Methods for introducing exogenous polynucleotides into plant cells and regenerating transgenic plants are well known. Some are provided below.

In one embodiment, to introduce a plasmid containing a CelA coding sequence or promoter of the invention into a plant, a 1:1 mixture of plasmid DNA containing a selectable marker expression cassette and plasmid DNA containing a cellulose synthase expression cassette is precipitated with gold to form microprojectiles. The microprojectiles are rinsed in absolute ethanol and aliquots are dried onto a suitable macrocarrier such as the macrocarrier available from BioRad in Hercules, Calif. Prior to bombardment, embryogenic tissue is preferably desiccated under a sterile laminar-flow hood. The desiccated tissue is transferred to semi-solid proliferation medium. The prepared microprojectiles are accelerated from the macrocarrier into the desiccated target cells using a suitable apparatus such as a BioRad PDS-1000/HE particle gun. In a preferred method, each plate is bombarded once, rotated 180 degrees, and bombarded a second time. Preferred bombardment parameters

are 1350 psi rupture disc pressure, 6 mm distance from the rupture disc to macrocarrier (gap distance), 1 cm macrocarrier travel distance, and 10 cm distance from macrocarrier stopping screen to culture plate (microcarrier travel distance). Tissue is then transferred to semi-solid proliferation medium containing a selection agent, such as hygromycin B, for two days after bombardment.

Cellulose Synthase Protein and Fragment Thereof

A cellulose synthase of the invention is a plant protein that contains a catalytic subunit which has UDP-glucose binding activity for the synthesis of glucan from glucose, and eight transmembrane domains for localizing the cellulose synthase to the cell membrane. The cellulose synthase of the invention has eight transmembrane binding domains; two at the amino terminal and six at the carboxyl terminal. The UDP-glucose binding domain is located between transmembrane domains two and three. Examples of this protein structure are seen in the aspen cellulose synthase as well as in those of RSW1 and GhCelA. The location of the transmembrane domain may be identified as described below and as exemplified in the Example. Preferably, the cellulose synthase of the invention has an amino acid sequence of a tree cellulose synthase.

In one embodiment, the cellulose synthase protein of the invention is isolated from aspen. Aspen cellulose synthase contains about 978 amino acids and has a molecular weight of about 110 KDa and a pI of about 6.58. In one embodiment, the aspen cellulose synthase has the amino acid sequence of SEQ ID NO:2 as represented in FIG. 1. In another aspect, the invention relates to cellulose synthase of SEQ ID NO: 5.

The invention further relates to fragments of plant cellulose synthases, such as fragments containing at least one transmembrane region and/or a UDP-glucose binding domain. The transmembrane regions may be identified as described in the Example by using the method of Hoffman and Stoffel (1993).

The cellulose synthase fragment containing the UDP-glucose binding domain is functional without the presence of the rest of the protein. This separable activity is as shown in the Example. This result was surprising and unexpected because previously identified UDP-glucose binding domains were not known to be functional when isolated from other portions of the protein. Thus, a fragment of any cellulose synthase (such as PtCelA, RSW1, GhCelA and SEQ ID NO:5) that contains a UDP-glucose binding domain and is independently functional is within the scope of the invention. The function of the UDP-glucose binding domain may be determined using the assay described in the Example. The UDP-glucose binding domain of the invention is located between the second and third transmembrane region of the cellulose synthase and has conserved amino acid sequences for UDP-glucose binding, such as the sequence QVLRW and conserved D residues. The UDP-glucose binding domain and the conserved regions therein may be located in a cellulose synthase using the guidance of the present specification and the general knowledge in the art, for example Brown, 1996. In one embodiment, the UDP-glucose binding domain and the conserved regions therein may be identified by comparing the amino acid sequence of cellulose synthase of interest with the amino acid sequence of aspen cellulose synthase using the algorithms described in the specification or generally known in the art. For example, the UDP-glucose binding domain of SEQ ID NO:2 is in the position amino acids 220 to 749. The conserved QVLRW sequence is located at positions 715–719 of SEQ ID NO:2.

Polypeptides having at least 75%, preferably at least 85% and most preferably at least 95% similarity to the amino acid sequence of SEQ ID NO: 2, amino acids 220–749 of SEQ ID NO:2, SEQ ID NO:5 or its UDP-glucose binding domain using Power Blast or GAP algorithm described above. In a preferred embodiment, these polypeptides are of about the same length as the polypeptide of SEQ ID NO: 2 or amino acids 220–749 of SEQ ID NO:2. For example, the polypeptide may be from about 2–3 to about 5–7 and to about 10–15 amino acids longer or shorter. In another embodiment, the polypeptides described in this paragraph are not originally found (i.e., naturally occurring) in *Arabidopsis* or cotton. These polypeptides may be prepared by, for example, altering the nucleic acid sequence of a cloned polynucleotide encoding the protein of SEQ ID NO:2 or SEQ ID NO:5 using the methods well known in the art.

Function conservative variants of cellulose synthase are also within the scope of the invention and can be prepared by altering the sequence of a cloned polynucleotide encoding cellulose synthase or fragments thereof. Conventional methods used in the art can be used to make substitutions, additions or deletions in one or more amino acids, to provide functionally equivalent molecules. For example, a function conservative variant that has substitutions, deletions and/or additions in the amino and/or carboxyl terminus of the protein, outside of the UDP-glucose binding domain is within the scope of the invention. Preferably, variants are made that have enhanced or increased functional activity relative to native cellulose synthase. Methods of directed evolution can be used for this purpose.

The invention also includes function conservative variants which include altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a conservative amino acid substitution. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point. Particularly preferred substitutions are: (i) Lys for Arg and vice versa such that a positive charge may be maintained; (ii) Glu for Asp and vice versa such that a negative charge may be maintained; (iii) Ser for Thr such that a free —OH can be maintained; and (iv) Gln for Asn such that a free CONH₂ can be maintained. Amino acid substitutions may also be introduced to substitute an amino acid with a particularly preferable property. For example, a Cys may be introduced a potential site for disulfide bridges with another Cys. A His may be introduced as a particularly “catalytic” site (i.e., His can act as an acid or base and is the most common amino acid in biochemical catalysis). Pro may be introduced because of its particularly planar structure, which induces b-turns in the protein’s structure.

The cellulose synthase of the invention can be isolated by expressing a cloned polynucleotide encoding the cellulose synthase as well as using direct protein purification techniques. These methods will be apparent to those of skill in the art.

Polynucleotides Containing Cellulose Synthase Promoter

The present invention further relates to a cellulose synthase promoter. The promoter is a stress-inducible promoter and may be used to synthesize greater quantities of high crystalline cellulose in plant, and preferably in trees. This permits an increase in the proportion of cellulose in transgenic plants, greater strength of juvenile wood and fiber, and acceleration of overall growth rate.

In one embodiment, the promoter of the invention is from aspen and is represented in FIG. 4. The promoter sequence is located within the region of nucleotides 1–840 of SEQ ID NO:3. A person of skill in the art will appreciate that not the entire sequence is required for the promoter function and can easily identify the critical regions by looking for conserved boxes and doing routine deletion analysis. Thus, functional fragments of SEQ ID NO:1 are within the scope of the invention.

Polynucleotides that hybridize under conditions of low, medium, and high stringency to SEQ ID NO:3, and its non-coding portion are also within the scope of the invention. The hybridizable polynucleotide may be about the same length as the sequence to which it hybridizes, for example, not more than about 10 to about 20 nucleotides longer or shorter. In another embodiment, the hybridizable polynucleotide is at least about 200 nucleotides long, preferably at least about 400 nucleotides long and most preferably at least 500 nucleotides long. In yet another embodiment, the hybridizable polynucleotide comprises at least one MSRE element identified according to the method described below.

A cellulose synthase promoter of the invention typically provides tissue-specific gene regulation in xylem, but also permits up-regulation of gene expression in other tissues as well, e.g., phloem under tension stress. Furthermore, expression of cellulose synthase is localized to an area of the plant under stress.

This stress-inducible phenomenon is regulated by positive and negative mechanical stress response elements (MSREs). These MSREs upregulate (positive) or downregulate (negative) the expression of a cellulose synthase polynucleotide under stress conditions through binding of transcription factors. MSRE-regulated expression of cellulose synthase permits synthesis of cellulose with high crystallinity.

The MSREs of cellulose synthase can be modified or employed otherwise in methods to regulate expression of a polynucleotide, including a cellulose synthase, operatively linked to a promoter containing an MSRE in response to mechanical stress (e.g., tension or compression) to a transgenic plant.

Negative MSREs of a cellulose synthase promoter can be modified, removed or blocked to improve expression of a cellulose synthase, and thereby increase cellulose production and improve wood quality. Alternatively, positive MSREs can be removed or blocked to decrease expression of a cellulose synthase, which decreases cellulose production and increases lignin deposition. This is useful for increasing the fuel value of wood because lignin has a higher BTU value than cellulose. Moreover, a modified cellulose synthase promoter can be operatively linked to a polynucleotide of interest to control its expression upon mechanical stress to a plant harboring it.

The location of MSRE elements in the SEQ ID NO:3 may be identified, for example, using promoter deletion analysis, DNase Foot Print Analysis, and Southwestern screening of an expression library for an MSRE. In one embodiment, cellulose synthase promoter that has one or more portions deleted, and is operatively linked to a reporter sequence, is introduced into a plant or a plant cell. A positive MSRE is detected by observing no relative change or increase in the amount of reporter in a transgenic plant or tissue, e.g., phloem after inducing a stress to the plant, and a negative MSRE is detected by observing increases in the amount of reporter in the plant in the absence of any stress to the plant. A positive element is detected when by removing it, GUS expression goes down and by adding it kept at the same level or more. The negative element does not support, or suppress, expression of GUS and by removing it, normal or enhanced GUS expression is observed as compared to when negative element is present.

Manipulation of a MSRE binding sites and/or providing transcription factors that bind thereto, provides a mechanism to continuously produce high crystalline cellulose in woody plant cell walls of transgenic plants. For example, one having ordinary skill in the art can delete or block negative MSRE elements, or provide cDNA encoding protein(s) that bind the positive MSREs, to enable constitutive expression of a cellulose synthase without the requirement of a mechanical stress. The increased cellulose synthase, and therefore, increased cellulose content, can improve the strength properties of juvenile wood and fiber. It is also contemplated that the positive MSREs can be deleted or blocked, or cDNA in an antisense direction, which in the sense direction encodes a protein that binds a positive MSRE, can be provided, to reduce cellulose synthase activity and decrease cellulose production.

Method of Isolating Polynucleotides Encoding Cellulose Synthase

The invention further relates to identifying and isolating polynucleotides encoding cellulose synthase in plants, e.g., trees, (in addition to those polynucleotides provided in the Example and represented in FIG. 1 and FIG. 7). These polynucleotides may be used to manipulate expression of cellulose synthase with an objective to improve the cellulose content and properties of wood.

The method comprises identifying a nucleic acid fragment containing a sequence encoding cellulose synthase or a portion thereof by using a fragment of SEQ ID NOS:1 or 4 as a probe or a primer. Once identified, the nucleic acid fragment containing a sequence encoding cellulose synthase or a portion thereof is isolated.

Polynucleotides encoding cellulose synthases of the invention, whether genomic DNA, cDNA, or fragments thereof, can be isolated from many sources, particularly from cDNA or genomic libraries from plants, preferably trees (e.g. aspen, sweetgum, loblolly pine, eucalyptus, and other angiosperms and gymnosperms). Molecular biology methods for obtaining polynucleotides encoding a cellulose synthase are well known in the art, as described above (see, e.g., Sambrook et al., 1989, supra).

Accordingly, cells from any species of plant can potentially serve as a nucleic acid source for the molecular cloning of a polynucleotide encoding a cellulose synthase of the invention. The DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a DNA “library”), and preferably is obtained from a cDNA library prepared from tissues with high level expression of a cellulose synthase (e.g., xylem tissue, since cells in this tissue

evidence very high levels of expression of CelA), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from a desired cell (see, for example, Sambrook et al., 1989, supra; Glover, D. M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will not contain intron sequences. Whatever the source, a polynucleotide should be molecularly cloned into a suitable vector for its propagation.

In another embodiment for the molecular cloning of a polynucleotide encoding a cellulose synthase of the invention from genomic DNA, DNA fragments are generated from a genome of interest, such as from a plant, or more particularly a tree genome, part of which will correspond to a desired polynucleotide. The DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, as for example, by sonication. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing a desired CelA sequence may be accomplished in a number of ways. For example, if an amount of a portion of a CelA sequence or its specific RNA, or a fragment thereof, is available and can be purified and labeled, the generated DNA fragments may be screened by nucleic acid hybridization to a labeled probe (Benton and Davis, 1977, Science 196:180; Grunstein and Hogness, 1975, Proc. Natl. Acad. Sci. U.S.A. 72:3961). For example, a set of oligonucleotides corresponding to the partial amino acid sequence information obtained for a CelA protein from trees can be prepared and used as probes for DNA encoding cellulose synthase, or as primers for cDNA or mRNA (e.g., in combination with a poly-T primer for RT-PCR). Preferably, a fragment is selected that is highly unique to a cellulose synthase of the invention, such as the UDP-glucose binding regions. Those DNA fragments with substantial homology to the probe will hybridize. As noted above, the greater the degree of homology, the more stringent hybridization conditions can be used. In a specific embodiment, stringency hybridization conditions can be used to identify homologous CelA sequences from trees or other plants.

Thus, in one embodiment, a labeled cellulose synthase cDNA from, e.g., *Populus tremuloides* (PtCelA), can be used to probe a library of genes or DNA fragments from various species of plants, especially angiosperm and gymnosperm, to determine whether any bind to a CelA of the invention. Once genes or fragments are identified, they can be amplified using standard PCR techniques, cloned into a vector, e.g., pBluescript vector (Stratagene of LaJolla, Calif.), and transformed into a bacteria, e.g., DH5 α *E. coli* strain (Gibco BRL of Gaithersburg, Md.). Bacterial colonies are typically tested to determine whether any contains a cellulose synthase-encoding nucleic acid. Once a positive clone is identified through binding, it is sequenced from an end, preferably the 3' end.

cDNA libraries can be constructed in various hosts, such as lambda ZAPII, available from Stratagene, LaJolla, Calif., using poly(A)⁺ RNA isolated from aspen xylem, according to the methods described by Bugos et al. (Biotechniques 19:734-737, 1995). The above mentioned probes are used to assay the aspen cDNA library to locate cDNA which codes

for enzymes involved in production of cellulose synthases. Once a cellulose synthase sequence is located, it is then cloned and sequenced according to known methods in the art.

Further selection can be carried out on the basis of the properties of the gene, e.g., if the gene encodes a protein product having the isoelectric, electrophoretic, hydrophathy plot, amino acid composition, or partial amino acid sequence of a cellulose synthase protein of the invention, as described herein. Thus, the presence of the gene may be detected by assays based on the physical, chemical, or immunological properties of its expressed product. For example, cDNA clones or DNA clones which hybrid-select the proper mRNAs can be used to produce a protein that has similar properties known for cellulose synthases of the invention. Such properties may include, for example, similar or identical electrophoretic migration patterns, isoelectric focusing or non-equilibrium pH gel electrophoresis behavior, proteolytic digestion maps, hydrophathy plots, or functional properties (such as isolated, functional UDP-glucose binding domains).

A cellulose synthase polynucleotide of the invention can also be identified by mRNA selection, i.e., by nucleic acid hybridization followed by in vitro translation. In this procedure, nucleotide fragments are used to isolate complementary mRNAs by hybridization. Such DNA fragments may represent available, purified CelA DNA, or may be synthetic oligonucleotides designed from the partial amino acid sequence information. Functional assays (e.g., UDP-glucose activity) of the in vitro translation products of the products of the isolated mRNAs identifies the mRNA and, therefore, the complementary DNA fragments, that contain the desired sequences.

A radiolabeled CelA cDNA can be synthesized using a selected mRNA as a template. The radiolabeled mRNA or cDNA may then be used as a probe to identify homologous CelA DNA fragments from amongst other genomic DNA fragments.

It will be appreciated that other polynucleotides, in addition to a CelA of the invention can be operatively linked to a CelA promoter to control expression of the polynucleotide upon application of a mechanical stress.

Expression of CelA Polypeptides

The nucleotide sequence coding for CelA, or a functional fragment, derivative or analog thereof, including chimeric proteins, can be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. Preferably, an expression vector includes an origin of replication. The elements are collectively termed herein a "promoter." Thus, a nucleic acid encoding CelA of the invention can be operatively associated with a promoter in an expression vector of the invention. Both cDNA and genomic sequences can be cloned and expressed under control of such regulatory sequences. The necessary transcriptional and translational signals can be provided on a recombinant expression vector, or they may be supplied by the native gene encoding CelA and/or its flanking regions.

In addition to a CelAP, expression of cellulose synthase can be controlled by any promoter/enhancer element known in the art, but these regulatory elements must be functional in the host selected for expression. Promoters which may be used to control CelA polynucleotide expression include, constitutive, development-specific and tissue-specific. Examples of these promoters include 35S Cauliflower Mosaic Virus, terminal flower and 4CL-1. Thus, there are

various ways to alter the growth of a plant using different promoters, depending on the needs of the practitioner.

The nucleotide sequence may be inserted in a sense or antisense direction depending on the needs of the practitioner. For example, if augmentation of cellulose biosynthesis is desired then polynucleotides encoding, e.g., cellulose synthase, can be inserted into the expression vector in the sense direction to increase cellulose synthase production and thus cellulose biosynthesis. Alternatively, if it is desired that cellulose biosynthesis is reduced or lignin content is increased, then polynucleotides encoding, e.g., cellulose synthase, can be inserted in the antisense direction so that upon transcription the antisense mRNA hybridizes to other complementary transcripts in the sense orientation to prevent translation. In other embodiments, the polynucleotide encodes a UDP-glucose binding domain and is used in a similar manner as described.

A recombinant CelA protein of the invention, or functional fragment, derivative, chimeric construct, or analog thereof, may be expressed chromosomally, after integration of the coding sequence by recombination. In this regard, any of a number of amplification systems for plants may be used to achieve high levels of stable gene expression, as discussed above. Any of the methods previously described for the insertion of DNA fragments into a cloning vector may be used to construct expression vectors containing a gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and synthetic techniques and in vivo recombination (genetic recombination).

Expression vectors containing a nucleic acid encoding a CelA of the invention can be identified by four general approaches: (a) PCR amplification of the desired plasmid DNA or specific mRNA, (b) nucleic acid hybridization, (c) presence or absence of selection marker gene functions, (d) analyses with appropriate restriction endonucleases, and (e) expression of inserted sequences. In the first approach, the nucleic acids can be amplified by PCR to provide for detection of the amplified product. In the second approach, the presence of a foreign gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted marker gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "selection marker" gene functions (e.g., β -glucuronidase activity, resistance to antibiotics, transformation phenotype, etc.) caused by the insertion of foreign genes in the vector. In another example, if the nucleic acid encoding CelA is inserted within the "selection marker" gene sequence of the vector, recombinants containing the CelA insert can be identified by the absence of the CelA gene function. In the fourth approach, recombinant expression vectors are identified by digestion with appropriate restriction enzymes. In the fifth approach, recombinant expression vectors can be identified by assaying for the activity, biochemical, or immunological characteristics of the gene product expressed by the recombinant, provided that the expressed protein assumes a functionally active conformation.

After a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously

explained, the expression vectors which can be used include, but are not limited to those vectors or their derivatives described above.

Vectors are introduced into the desired host cells by methods known in the art, e.g., *Agrobacterium*-mediated transformation (described in greater detail below), transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; Hartmut et al., Canadian Patent Application No. 2,012,311, filed Mar. 15, 1990).

The cell into which the recombinant vector comprising the nucleic acid encoding CelA is cultured in an appropriate cell culture medium under conditions that provide for expression of CelA by the cell. In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (such as glycosylation, cleavage, e.g., of a signal sequence) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed.

Agrobacterium-Mediated Transformation and Inducing Somatic Embryos

The culture media used in the invention, and for transforming *Agrobacterium*, contain an effective amount of each of the medium components (e.g. basal medium, growth regulator, carbon source) described above. As used in describing the present invention, an "effective amount" of a given medium component is the amount necessary to cause a recited effect. For example, an effective amount of a growth hormone in the primary callus growth medium is the amount of the growth hormone that induces callus formation when combined with other medium components. Other compounds known to be useful for tissue culture media, such as vitamins and gelling agents, may also be used as optional components of the culture media of the invention.

Transformation of cells from plants, e.g., trees, and the subsequent production of transgenic plants using *Agrobacterium*-mediated transformation procedures known in the art, and further described herein, is one example of a method for introducing a foreign gene into trees. Transgenic plants may be produced by various methods, such as by the following steps: (i) culturing *Agrobacterium* in low-pH induction medium at low temperature and preconditioning, i.e., coculturing bacteria with wounded tobacco leaf extract in order to induce a high level of expression of the *Agrobacterium vir* genes whose products are involved in the T-DNA transfer; (ii) coculturing a desired plant tissue explants, including zygotic and/or somatic embryo tissues derived from cultured explants, with the incited *Agrobacterium*; (iii) selecting transformed callus tissue on a medium containing antibiotics; and (v) and converting the embryos into plantlets.

Any non-tumorigenic *A. tumefaciens* strain harboring a disarmed Ti plasmid may be used in the method of the invention. Any *Agrobacterium* system may be used. For example, Ti plasmid/binary vector system or a cointegrative vector system with one Ti plasmid may be used. Also, any marker gene or polynucleotide conferring the ability to select transformed cells, callus, embryos or plants and any other gene, such as, for example, a gene conferring resis-

tance to a disease, or one improving cellulose content, may also be used. Any promoter desired can be used, such as, for example, a PtCelAP of the invention, and those promoters as described above. A person of ordinary skill in the art can determine which markers and genes are used depending on particular needs.

For purposes of the present invention, “transformed” or “transgenic” means that at least one marker gene or polynucleotide conferring selectable marker properties is introduced into the DNA of a plant cell, callus, embryo or plant. Additionally, any gene may also be introduced.

To increase the infectivity of the bacteria, *Agrobacterium* is cultured in low-pH induction medium, i.e., any bacterium culture media with a pH value adjusted to from 4.5 to 6.0, most preferably about 5.2, and at low temperature such as for example about 19–30EC, preferably about 21–26EC. The conditions of low-pH and low temperature are among the well-defined critical factors for inducing virulence activity in *Agrobacterium* (e.g., Altmorbe et al., Mol. Plant-Microbe. Interac. 2: 301, 1989; Fullner et al., Science 273: 1107, 1996; Fullner and Nester, J. Bacteriol. 178: 1498, 1996).

The bacteria is preconditioned by coculturing with wounded tobacco leaf extract (prepared according to methods known generally known in the art) to induce a high level of expression of the *Agrobacterium vir* genes. Prior to inoculation of plant somatic embryos, *Agrobacterium* cells can be treated with a tobacco extract prepared from wounded leaf tissues of tobacco plants grown in vitro. To achieve optimal stimulation of the expression of *Agrobacterium vir* genes by wound-induced metabolites and other cellular factors, tobacco leaves can be wounded and precultured overnight. Culturing of bacteria in low pH medium and at low temperature can be used to further enhance the bacteria vir gene expression and infectivity. Preconditioning with tobacco extract and the vir genes involved in the T-DNA transfer process are generally known in the art.

Agrobacterium treated as described above is then cocultured with a plant tissue explant, such as for example zygotic and/or somatic embryo tissue. Non-zygotic (i.e., somatic) or zygotic tissues can be used. Any plant tissue may be used as a source of explants. For example, cotyledons from seeds, young leaf tissue, root tissues, parts of stems including nodal explants, and tissues from primary somatic embryos such as the root axis may be used. Generally, young tissues are a preferred source of explants.

The invention also relates to methods of altering the growth of a plant by expressing the polynucleotide of the invention, which as a result alters the growth of the plant. The polynucleotide used in the method may be a homologous polynucleotide or a heterologous polynucleotide and are described in detail above. For example, both full-length and UDP-glucose binding region containing fragments may be expressed. Additionally, depending on the aim of the method, the polynucleotide may be introduced into the plant in the sense or in the antisense orientation. Any suitable promoter may be used to provide expression. The promoter or a functional fragment thereof is operatively linked to the polynucleotide. The promoter may be a constitutive promoter, a tissue-specific promoter or a development-specific plant promoter. Examples of suitable promoters are Cauliflower Mosaic Virus 35S, 4CL, cellulose synthase promoter, PtCelAP and terminal flower promoter.

The invention further relates to a method of altering the cellulose content in a plant by expressing the polynucleotide of the invention as described above. The method may be used to increased the ratio of cellulose to lignin in the plant that have an exogenous polynucleotide of the invention introduced therein.

The invention further relates to a method for altering expression of a cellulose synthase in a plant cell by introducing into the cell a vector comprising a polynucleotide of the invention and expressing the polynucleotide. The polynucleotides and promoters described above may be used.

A method for causing stress-induced gene expression in a plant cell is also within the scope of the invention. The method comprises (i) introducing into the plant or a plant cell an expression cassette comprising a cellulose synthase promoter or a functional fragment thereof or providing a plant or a plant cell that comprises the expression cassette (The promoter of the cassette is operatively linked to a coding sequence of choice.); and (ii) applying mechanical stress to the plant to induce expression of the desired coding sequence.

A method for determining a positive mechanical stress responsive element (MSRE) in a cellulose synthase promoter is also within the scope of the invention and comprises (i) making serial deletions in the cellulose synthase promoter, such as for example, SEQ ID NO:3; (ii) introducing the deletion linked to a polynucleotide encoding a reporter sequence into a plant cell, and (iii) detecting a decrease in the amount of reporter in the plant after inducing a stress to the plant. Similarly, a method for determining a negative MSRE in a cellulose synthase promoter is provided. It comprises (i) making serial deletions in the cellulose synthase promoter, such as for example, SEQ ID NO:3; (ii) introducing the deletion linked to a polynucleotide encoding a reporter sequence into a plant cell, and (iii) detecting an increase in the amount of reporter in the plant after inducing a stress to the plant.

The following methods are also within the scope of the invention: a method for expressing cellulose synthase in a tissue-specific manner comprising transforming a plant with a tissue specific promoter operatively linked to a polynucleotide encoding a cellulose synthase; a method for inducing expression of a cellulose synthase in a plant comprising introducing into a plant a cDNA encoding a protein that binds to a positive MSRE of a cellulose synthase promoter, thereby resulting in increased expression of cellulose in the plant, wherein the binding to the positive MSRE results in expression of a cellulose synthase; a method for reducing expression of a cellulose synthase comprising introducing into a plant a cDNA in an antisense orientation, wherein the cDNA in a sense orientation encodes a protein that binds to a positive MSRE and results in expression of a cellulose synthase; a method for increasing cellulose biosynthesis in a plant comprising introducing into a plant a cDNA encoding a protein that binds to a positive MSRE of a cellulose synthase promoter, whereby binding of the protein to the positive MSRE results in expression of a cellulose synthase, and A method for reducing cellulose biosynthesis in a plant comprising introducing into a plant a cDNA in an antisense orientation, wherein the cDNA in a sense orientation encodes a protein that binds to a positive MSRE of a cellulose synthase promoter.

21

EXAMPLE

Molecular Cloning of Cellulose Synthase

This Example describes the first tree cellulose synthase cDNA (PtCelA, GenBank No. AF072131) cloned from developing secondary xylem of aspen trees using RSW1 cDNA.

Prior to the present invention, only partial clones of cellulose synthases from crop species and cotton GhCelA have been discovered, which have significant homology to each other. The present inventors have discovered and cloned a new full-length cellulose synthase cDNA, AraxCelA (GenBank No. AF062485) (FIG. 7, [SEQ ID NO: 4]), from an *Arabidopsis* primary library. AraxCelA is a new member of cellulose synthase and shows 63–85% identity and 72–90% similarity in amino acid sequence with other *Arabidopsis* CelA members.

Another cellulose synthase was cloned in aspen using a ³²P-labeled 1651-bp long EcoRI fragment of *Arabidopsis* CelA cDNA, which encodes a centrally located UDP-glucose binding domain, was used as a probe to screen about 500,000 pfu of a developing xylem cDNA library from aspen (*Populus tremuloides*) (Ge and Chiang, 1996). Four positive clones were obtained after three rounds of plaque purification. Sequencing the 3N ends of these four cDNAs showed that they were identical clones. The longest cDNA clone was fully sequenced and determined to be a full-length cDNA having a 3232 bp nucleotide sequence (FIG. 1) [SEQ ID NO: 1], which encodes a protein of 978 amino acids [SEQ ID NO: 2].

Characterization of a Cellulose Synthase from Aspen

The first AUG codon of PtCelA was in the optimum context for initiation of transcription on the basis of optimal context sequence described by Joshi (1987a) and Joshi et al. (1997). A putative polyadenylation signal (AATACA) was found 16 bp upstream of a polyadenylated tail of 28 bp, which is similar to the proposed plant structure (Joshi, 1987b). The 5N untranslated leader was determined to have 68 bp and the 3N untranslated trailer was 227 bp. Both of these regions have a typical length observed in many plant genes (Joshi, 1987a and Joshi, 1987b). This cDNA clone exhibited 90% amino acid sequence similarity with cellulose synthase from cotton (GhCelA) and 71% with cellulose synthase from *Arabidopsis* (RSW1), suggesting that this particular tree homolog also encodes a cellulose synthase.

The full length cDNA was designated PtCelA, and encodes a 110,278 Da polypeptide having an isoelectric point (pI) of 6.58 and 8 charged molecules. The hydropathy curve indicated that this particular cellulose synthase has eight transmembrane binding domains; two at the amino terminal and six at the carboxyl terminal, using the method of Hoffman and Stoffel (1993). This protein structure is analogous to those of RSW1 and GhCelA. All of the conserved domains for UDP-glucose binding, such as QVLRW and conserved D residues, are also present in a cellulose synthase of the invention, e.g., PtCelA (Brown et al., 1996). Thus, based on sequence and molecular analyses, it was concluded that PtCelA encodes a catalytic subunit which, like RSW1 in *Arabidopsis*, is essential for the cellulose biosynthesis machinery in aspen.

In situ localization of PtCelA mRNA transcripts along the developmental gradient defined by stem primary and secondary growth demonstrated that cellulose synthase expres-

22

sion is confined exclusively to developing xylem cells undergoing secondary wall thickening. This cell-type-specific nature of PtCelA gene expression was also consistent with xylem-specific activity of cellulose synthase promoter (PtCelAP) based on heterologous promoter- β -glucuronidase (GUS) fusion analysis. Overall, the results provide several lines of evidence that cellulose synthase is the gene primarily responsible for cellulose biosynthesis during secondary wall formation in woody xylem of trees, such as aspen. Previous results by the inventors (Hu et al., 1999) showed that cellulose and lignin are deposited in a compensatory fashion in wood. The discovery of a cellulose synthase in trees, such as aspen, permits the up-regulation of the protein to elevate cellulose production. Surprisingly, expression of CelA in trees suppressed lignin biosynthesis to further improve wood properties of trees.

Preparation of Transgenic Plants

The UDP-glucose binding sequence was subcloned into pBI121, which was used to prepare transgenic tobacco plants (Hu et al., 1998). The expression of a heterologous UDP-glucose binding sequence resulted in a remarkable growth-accelerating effect. This was surprising because current knowledge of the function of plant cellulose synthases teaches that a UDP-glucose sequence must remain intact with other functional domains in CelA, e.g., the transmembrane domains, in order for cellulose synthase to initiate cellulose biosynthesis. The remarkable growth and tremendous increase in plant biomass observed in transgenic tobacco was due likely to an augmented deposition of cellulose, indicating that the UDP-glucose domain alone is sufficient for genetic augmentation of cellulose biosynthesis in plants.

Genome Organization and Expression of a Novel Cellulose Synthase

To confirm that the cDNA clone of FIG. 1 [SEQ ID NO: 1] was a cellulose synthase, genomic Southern blot analysis was performed under both high and low stringency conditions using the cDNA. Genomic DNA from aspen was digested with PstI (lane P), HindIII (lane H) and EcoRI (lane E), and probed using a 1 kb ³²P-labeled fragment from the 5N end of a cellulose synthase of FIG. 1. The Southern blot suggested the presence of a small family of cellulose synthase genes in aspen genome (FIG. 2, panels a and b). Repeated screening of the aspen xylem cDNA library with various plant CelA gene-related probes always resulted in the isolation of the same cellulose synthase cDNA clone. This suggested that the cellulose synthase cDNA cloned (FIG. 1) [SEQ ID NO: 1], represents the primary and most abundant cellulose synthase-encoding gene in developing xylem of trees, such as aspen, where active cellulose deposition takes place. It also indicates that manipulation of cellulose synthase gene expression can have a profound influence on cellulose biosynthesis in trees.

In Situ Hybridization

Northern blot analysis of total RNA from the internodes of aspen seedling stems (FIG. 2, panel c) using the labeled probe (as described above) revealed the near absence of cellulose synthase transcripts in tissues undergoing primary growth (internodes 1 to 4), and that the presence of cellulose synthase transcripts occurs during the secondary growth of stem tissues (internodes 5 to 11). However, weak northern

signals in primary growth may only suggest that cellulose synthase gene expression is specific to xylem, of which there is little in primary growth tissue.

Xylogenesis in higher plants offers a unique model that involves sequential execution of cambium cell division, commitment to xylem cell differentiation, and culmination in xylem cell death (Fukuda, 1996). Although primary and secondary xylem cells originate from different types of cambia, namely procambium and inter/intrafascicular cambium, both exhibit conspicuous secondary wall development with massive cellulose and lignin deposition (Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons). To further investigate spatial and temporal cellulose synthase gene expression patterns at the cellular level, in situ hybridization was used to localize cellulose synthase mRNA along the developmental gradient defined by stem primary and secondary growth.

Localization of cellulose synthase gene transcripts (RNA) in stem at various growth stages was also observed. FIG. 3 shows transverse sections from 2nd, 4th and 6th internodes hybridized with digoxigenin (DIG)-labeled cellulose synthase antisense or sense (control) RNA probes, as described.

PtCelA transcripts were detected in young aspen stem sections by in situ hybridization with transcripts of highly variable 5N region of PtCelA cDNA (a 771 bp long fragment generated from PstI and SacI). This region was first subcloned in the plasmid vector, pGEM, -3Zf (+) (Promega) for the production of digoxigenin (DIG)-labeled transcripts using T7 (for antisense transcripts) and SP6 (for sense transcripts) RNA polymerase (DIG system: Boehringer Mannheim). Probes were subjected to mild alkaline hydrolysis by incubation in 100 mM NaHCO₃, pH 10.2 at 60° C., which produced approximately 200 bp fragments.

Aspen young stems were prepared for sectioning by fixation in 4% (w/v) paraformaldehyde in 100 mM phosphate buffer (pH 7.0) at 4° C. overnight, dehydrated through an ethanol series on ice, and embedded in Paraplast medium (Sigma). Ten µm sections were mounted on Superfrost/plus (Fisher) slides at 42° C. overnight, dewaxed and then rehydrated through a descending ethanol series. The sections were incubated with proteinase K (10 µg/ml in 100 mM Tris-HCl, 50 mM EDTA, pH 7.5) for 30 min and were post-fixed with FAA. The sections were acetylated with 0.33% (v/v) acetic anhydride in 0.1 M triethanolamine-HCl (pH 8.0) prior to hybridization. The sections were then incubated in a hybridization mixture (approximately 2 µg/ml DIG-labeled probes, 50% (v/v) formamide, 2×SSPE, 10% (w/v) dextran sulfate, 125 µg/ml tRNA, pH 7.5) at 45° C. for 12–16 hrs. Nonhybridized single-stranded RNA probe was removed by treatment with 20 µg/ml RNase A in TE buffer with 500 mM NaCl. The sections were washed at 50° C. Hybridized DIG-labelled probe was detected on sections using anti-digoxigenin antiserum at a 1:1500 dilution, as described in the manufacturer's instruction (DIG system: Boehringer Mannheim). Sections were examined by Eclipse 400 light microscope (Nikon) and photographed.

During the primary growth stage (FIG. 3, panels a and b), strong expression of cellulose synthase was found localized exclusively to primary xylem (PX) cells. At this stage, young internodes are elongating, resulting in thickening of primary xylem cells through formation of secondary walls (Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons). The concurrence of shoot elongation with high expression of cellulose synthase strongly suggests the association of cellulose synthase protein with secondary cell wall cellulose synthesis. Later stages of primary growth (FIG. 3, panel b) are characterized by the appearance of an

orderly alignment of primary xylem cells. Active cellulose biosynthesis accompanies cell elongation-induced wall thickening, as indicated by the strong expression of cellulose synthase in these primary xylem cells.

At the beginning of secondary growth in older internodes, it was observed that expression of cellulose synthase is also exclusively localized to xylem cells (FIG. 3, panel c). Instead of elongation in internodes distal to the meristematic activity, growth at this stage is mainly radial due to thickening in secondary cell walls of secondary xylem. At the same time, expression of PtCelA gene becomes localized to the secondary developing xylem cells (SX in FIG. 3, panel c), which is again consistent with the idea that PtCelA encodes a secondary cell wall cellulose synthase. At this stage, secondary xylem cells cover the elongated and differentiated primary xylem cells in which PtCelA gene expression is no longer detectable (FIG. 3, panel c). These results demonstrate that expression of PtCelA gene is xylem-specific and the cellulose synthase of FIG. 1 [SEQ ID NO: 1] encodes a cellulose synthase associated with cellulose biosynthesis in secondary walls of xylem cells. To further confirm xylem-specific expression of cellulose synthase, a cellulose synthase gene promoter sequence was cloned and characterized for regulatory activities.

Characterization of Expression Regulated by Cellulose Synthase Promoter

A 5N 1,200 bp cDNA fragment of a cellulose synthase of FIG. 1 [SEQ ID NO: 1] was used as a probe to screen an aspen genomic library for 5N regulatory sequences of a novel cellulose synthase gene, PtCelA. The library was constructed by cloning aspen genomic DNA fragments, generated from an Sau3AI partial-digest and sucrose gradient-selected, into the BamHI site of a Lambda DASH II vector (Stratagene, La Jolla, Calif.). Five positive clones were obtained from about 150,000 pfu and Lambda DNA was purified. One clone having about a 20 kb DNA insert size was selected for restriction mapping and partial sequencing. This resulted in the identification of a 5N flanking region of PtCelA gene of approximately 1 kb. This genomic fragment, designated PtCelAP (FIG. 4) [SEQ ID NO: 3], contained about 800 bp of promoter sequence, 68 bp of 5N end untranslated region and 160 bp of coding sequence. To investigate regulation of tissue-specific cellulose synthase expression at the cellular level, promoter activity was analyzed in transgenic tobacco plants by histochemical staining of a GUS protein. A PtCelAP-GUS fusion binary vector was constructed in pBI121 with the 35S promoter replaced with PtCelAP [SEQ ID NO: 3] and introduced into tobacco (*Nicotiana tabacum*) as per Hu et al. (1998).

Eleven independent transgenic lines harboring a CelAP-GUS fusion were generated. FIG. 5 shows a histochemical analysis of GUS expression driven by a cellulose synthase promoter of the invention in transgenic tobacco plants. Transverse sections from the 3rd (panel a), 5th (panel b), 7th (panel c), and 8th (panels d and f) internodes were stained from GUS activity, and fluorescence microscopy was used to visualize expression under UV radiation.

GUS staining was detected exclusively in xylem tissue of stems, roots and petioles. In stems, strong GUS activity was found localized to xylem cells undergoing primary (FIG. 5, panel a) and secondary growth (FIG. 5 panels b-d and f). GUS expression was confined to xylem cells in the primary growth stage and became more localized in developing secondary xylem cells during secondary growth. An entire

section from the 8th internode stained for GUS activity (FIG. 5, panel f). These results are consistent with the in vivo expression patterns of cellulose synthase in aspen stems. Lignin autofluorescence was visualized after UV radiation. Phloem fibers, which are also active in cellulose and lignin biosynthesis (FIG. 5, panels d and e), did not show GUS activity, suggesting that cellulose synthase gene expression is not associated with cellulose biosynthesis in cell types other than xylem. Examination of GUS activity in roots, stems, leaves, anthers and fruit also showed GUS expression in xylem tissue of all these organs suggesting that cellulose synthases of the invention are xylem-specific cellulose and expressed in all plant organs.

Characterization of promoter activity and cellular expression of a cellulose synthase of the invention from one particular source (aspen) indicated that expression produces a protein that encodes a secondary cell wall-specific cellulose synthase and is specifically compartmentalized in developing xylem cells. Characterization of the cellulose synthase gene promoter sequence not only confirms cell type-specific expression of cellulose synthase, but also provides a method for over-expressing cellulose synthase in a tissue-specific manner to augment cellulose production in xylem.

Expression of Cellulose Synthase Under Tension Stress

As described earlier, a cellulose synthase promoter of the invention is involved in a novel gene regulatory phenomenon of cellulose synthase. To further characterize a cellulose synthase of the invention, GUS expression driven by an aspen cellulose synthase promoter (PtCelAP) was observed in transgenic tobacco plants without or under tension stress. The stress was induced by bending and affixing the plants to maintain the bent position (e.g., tying) over a 40 hour period. Tangential and longitudinal sections were taken before bending, and 4 hrs, 20 hrs and 40 hrs after bending (panels a-d, respectively).

The cellulose synthase promoter-GUS fusion binary constructs showed exclusive xylem-specific expression of GUS without any tension stress (FIG. 6, panel a). However, under tension stress conditions endured by angiosperms in nature, the transgenic tobacco plants induced xylem and phloem-specific expression on the upper side of the stem within the first four hours of stress (FIG. 6, panel b).

This observation was surprising because during tension wood development fibers produce highly crystalline cellulose in order to provide essential mechanical strength to a bending stem. The present observation was the first showing of transcriptional up-regulation of a cellulose synthase, mediated through a cellulose synthase promoter that is directly responsible for development of highly crystalline cellulose in trees. Furthermore, after 20 hrs of tension stress, both xylem and phloem exhibited GUS expression, but only on the upper side of the stem that was under tensile stress, i.e., GUS expression on the lower side was inhibited (FIG. 6, panel c). With extended stress (up to 40 hrs), GUS expression was restricted to only one small region on the upper side of the stem where maximum tension stress was present (FIG. 6, panel d). Based on the observation of GUS signal in woody cells upon tension stress and the absence of GUS under compression or no stress, it was concluded that a cellulose synthase promoter of the invention has mechanical stress responsive elements (MSREs) that turn cellulose synthase genes on and off depending on the presence and type of stress to the stem.

The results indicate that positive MSREs exist in a cellulose synthase promoter of the invention to bind transcription factors in response to tension stress for regulating the expression of cellulose synthase and increasing biosynthesis of higher crystalline cellulose. This is evident based on the expression of GUS in xylem and phloem tissue at the upper side of the stem subjected to tension stress, but not when tissue on the lower side was subjected to compression or no stress. Furthermore, the tissue at the lower side of the stem, which was subjected to compression stress, showed no GUS expression, i.e., expression was turned off. This indicated the presence of negative MSREs, which bind transcription factors to turn off expression of cellulose synthase at the lower side of the stem. Negative MSREs likely suppress development of highly crystalline cellulose in normal wood.

These results provide a mechanism for genetically engineering synthesis of highly crystalline cellulose in juvenile wood for enhancing strength properties, and for synthesizing a higher percentage of cellulose in reaction wood. The positive MSREs and their cognate transcription factors are important in the synthesis of highly crystalline cellulose of high tensile strength, as are the negative MSREs and inhibition of cognate transcription factors thereto. The present invention thus provides a starting point for cloning cDNAs for the transcription factors that bind to positive and negative MSREs according to methods known in the art. Constitutive expression of cDNAs for positive MSRE transcription factors allows the continuous production of highly crystalline cellulose in transgenic trees, while expression of antisense cDNAs for negative MSRE transcription factors inhibits those transcription factors so that cellulose synthase cannot turn off. This combination will assure continuous production of highly crystalline cellulose in trees.

Genetic Engineering of Cellulose Synthase in Transgenic Plants

As discussed above, the nucleotide sequence of a cellulose synthase of the invention, e.g., PtCelA cDNA from aspen, shows significant homology with other polynucleotides encoding cellulose synthase proteins that have been suggested as authentic cellulose synthase clones. To further characterize the activity of a cellulose synthase, four constructs were prepared in a PBI121 plasmid.

1) A constitutive plant promoter Cauliflower mosaic Virus 35S was operatively linked to PtCelA (35SP-PtCelA-s) and overexpressed in transgenic plants. This causes excess production of cellulose, resulting in a reduction in lignin content. Tobacco and aspen have been transformed with this construct.

2) Cauliflower mosaic Virus 35S was operatively linked to antisense RNA from PtCelA (35S-PtCelA-a) and constitutively expressed to reduce production of cellulose and increase lignin content in transgenic plants. This negative control construct may not result in healthy plants since cellulose is essential for plant growth and development. Aspen plants have been transformed with this construct.

3) Aspen 4CL-1 promoter (Hu et al., 1998) was operatively linked to PtCelA (Pt4CLP-PtCelA) (the 35S promoter of PBI121 was removed in this construct) and expressed in a tissue-specific manner in developing secondary xylem of transgenic aspen. This expression augments the native cellulose production and reduces lignin content of angiosperm tissues. Tobacco and aspen have transformed with this construct.

4) The cytoplasmic domain of PtCelA which contains three conserved regions thought to be involved in UDP-glucose binding during cellulose biosynthesis, was linked to a 35S promoter to produce binary constructs (35S-PtCelA UDP-glucose). Expression by this promoter permits constitutive expression of a UDP glucose binding domain of PtCelA in transgenic plants. Tobacco and aspen have been transformed with this construct.

35S-GUS constructs (pBI121, ClonTech, CA) were used as controls for each experiment with the constructs. Transgenic tobacco plants were transformed with the constructs. The following table shows the general growth measurements of the T0 tobacco plants. Plants carrying a PtCelA construct grew much faster than control plants carrying a pBI121 (control) construct. In comparing developmental 4CL and constitutive 35S promoter control of PtCelA expression, the 35S was more effective, permitting faster growth of transgenic tobacco plants. The fastest growth was seen in transgenic plants carrying a 35S promoter driven UDP-G domain from PtCelA.

It is noted that T0 generation plants can have carry over effects from their tissue culture treatments. Therefore, seeds were collected for testing this growth phenomenon in T1 generations. The transgenic tobacco plants were analyzed for presence of the transferred genes and all tested positive for the respective gene constructs.

TABLE

Transgenic tobacco plant measurements after transfer in soil for about 1.5 months (N = 2)					
Construct	Height	Diameter	Internode length	No. of leaves	Longest leaf
35S-GUS	17	0.5	1	11	17
35S-PtCelA	77	1.0	6	13	37
35S-UDPG	83	1.0	6	13	37
4CLP-PtCelA	41	0.8	5	10	29

Note:

All values were measured in centimeters, excluding number of leaves.

It will be appreciated by persons of ordinary skill in the art that the examples and preferred embodiments herein are illustrative, and that the invention may be practiced in a variety of embodiments which share the same inventive concept.

BIBLIOGRAPHY

- Hu et al., 1999, Nature Biotechnology, In Press
- Whetten et al., 1998, Ann Rev Pl Physiol Pl Mol Biol, 49: 585-609
- Arioli et al., 1998, Science, 279: 717-720
- Wu et al., 1998, Pl Physiol, 117: 1125
- Hu et al., 1998, PNAS, 95: 5407-5412
- Joshi et al., 1997, PMB, 35: 993-1001
- Fukuda, 1996, Ann Rev Pl Physiol Pl Mol Biol, 47: 299-325
- Pear et al., 1996, PNAS, 93: 12637-12642
- Haigler and Blanton, 1996, PNAS, 93: 12082-12085
- Ge and Chiang, 1996, Pl Physiol, 112: 861
- Brown et al., 1996, Trends Pl Sci., 1: 149-156
- Delmer and Amor, 1995, Pl Cell, 7: 987-1000
- Hoffman and Stoffel, 1993, Biol Chem, Hoppe-Seyler 374: 166
- Joshi, 1987, NAR, 15: 6643-6653
- Joshi, 1987, NAR, 15: 9627-9640
- Timmell, 1986, Compression Wood in Gymnosperms, Springer Verlag
- Easu, K., 1960, Anatomy of Seed Plants, New York: John Wiley and Sons
- Higuchi, 1997, Biochemistry and Molecular Biology of Wood, Springer Verlag

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 3232

<212> TYPE: DNA

<213> ORGANISM: Populus tremuloides

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (69)..(3002)

<400> SEQUENCE: 1

```

gtcgaccac gcgtccgtct tgaagaata tgaagttgta aagagctggt aaagtggtaa      60
taagcaag atg atg gaa tct ggg gct cct ata tgc cat acc tgt ggt gaa      110
      Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu
          1             5             10
cag gtg ggg cat gat gca aat ggg gag cta ttt gtg gct tgc cat gag      158
Gln Val Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu
15             20             25             30
tgt agc tat ccc atg tgc aag tct tgt ttc gag ttt gaa atc aat gag      206
Cys Ser Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Asn Glu
          35             40             45

```

-continued

ggc cgg aaa gtt tgc ttg cgg tgt ggc tcg cca tat gat gag aac ttg Gly Arg Lys Val Cys Leu Arg Cys Gly Ser Pro Tyr Asp Glu Asn Leu 50 55 60	254
ctg gat gat gta gaa aag aag ggg tct ggc aat caa tcc aca atg gca Leu Asp Asp Val Glu Lys Lys Gly Ser Gly Asn Gln Ser Thr Met Ala 65 70 75	302
tct cac ctc aac gat tct cag gat gtc gga atc cat gct aga cat atc Ser His Leu Asn Asp Ser Gln Asp Val Gly Ile His Ala Arg His Ile 80 85 90	350
agt agt gtg tcc act gtg gat agt gaa atg aat gat gaa tat ggg aat Ser Ser Val Ser Thr Val Asp Ser Glu Met Asn Asp Glu Tyr Gly Asn 95 100 105 110	398
cca att tgg aag aat cgg gtg aag agc tgt aag gat aaa gag aac aag Pro Ile Trp Lys Asn Arg Val Lys Ser Cys Lys Asp Lys Glu Asn Lys 115 120 125	446
aag aaa aag aga agt cct aag gct gaa act gaa cca gct caa gtt cct Lys Lys Lys Arg Ser Pro Lys Ala Glu Thr Glu Pro Ala Gln Val Pro 130 135 140	494
aca gaa cag cag atg gaa gag aaa ccg tct gca gag gct tcg gag ccg Thr Glu Gln Met Glu Glu Lys Pro Ser Ala Glu Ala Ser Glu Pro 145 150 155	542
ctt tca att gtt tat cca att cca cgc aac aag ctc aca cca tac aga Leu Ser Ile Val Tyr Pro Ile Pro Arg Asn Lys Leu Thr Pro Tyr Arg 160 165 170	590
gca gtg atc att atg cga ctg gtc att ctg ggc ctc ttc ttc cac ttc Ala Val Ile Ile Met Arg Leu Val Ile Leu Gly Leu Phe Phe His Phe 175 180 185 190	638
aga ata aca aat cct gtc gat agt gcc ttt ggc ctg tgg ctt act tct Arg Ile Thr Asn Pro Val Asp Ser Ala Phe Gly Leu Trp Leu Thr Ser 195 200 205	686
gtc ata tgt gag atc tgg ttt gca ttt tct tgg gtg ttg gat cag ttc Val Ile Cys Glu Ile Trp Phe Ala Phe Ser Trp Val Leu Asp Gln Phe 210 215 220	734
ccc aag tgg aat cct gtc aat aga gaa acg tat atc gaa agg ctg tcg Pro Lys Trp Asn Pro Val Asn Arg Glu Thr Tyr Ile Glu Arg Leu Ser 225 230 235	782
gca agg tat gaa aga gag ggt gag cct tct cag ctt gct ggt gtg gat Ala Arg Tyr Glu Arg Glu Gly Glu Pro Ser Gln Leu Ala Gly Val Asp 240 245 250	830
ttt ttc gtg agt act gtt gat ccg ctg aag gaa ccg cca ttg atc act Phe Phe Val Ser Thr Val Asp Pro Leu Lys Glu Pro Pro Leu Ile Thr 255 260 265 270	878
gcc aat aca gtc ctt tcc atc ctt gct gtg gac tat ccc gtc gat aaa Ala Asn Thr Val Leu Ser Ile Leu Ala Val Asp Tyr Pro Val Asp Lys 275 280 285	926
gtc tcc tgc tac gtg tct gat gat ggt gca gct atg ctt tca ttt gaa Val Ser Cys Tyr Val Ser Asp Asp Gly Ala Ala Met Leu Ser Phe Glu 290 295 300	974
tct ctt gta gaa aca gct gag ttt gca agg aag tgg gtt ccg ttc tgc Ser Leu Val Glu Thr Ala Glu Phe Ala Arg Lys Trp Val Pro Phe Cys 305 310 315	1022
aaa aaa ttc tca att gaa cca aga gca ccg gag ttt tac ttc tca cag Lys Lys Phe Ser Ile Glu Pro Arg Ala Pro Glu Phe Tyr Phe Ser Gln 320 325 330	1070
aaa att gat tac ttg aaa gac aag gtt caa cct tct ttc gtg aaa gaa Lys Ile Asp Tyr Leu Lys Asp Lys Val Gln Pro Ser Phe Val Lys Glu 335 340 345 350	1118
cgT aga gca atg aaa agg gat tat gaa gag tac aaa gtc cga gtt aat Arg Arg Ala Met Lys Arg Asp Tyr Glu Glu Tyr Lys Val Arg Val Asn	1166

-continued

355				360				365								
gcc	ctg	gta	gca	aag	gct	cag	aaa	aca	cct	gaa	gaa	gga	tg	act	atg	1214
Ala	Leu	Val	Ala	Lys	Ala	Gln	Lys	Thr	Pro	Glu	Glu	Gly	Trp	Thr	Met	
			370					375					380			
caa	gat	gga	aca	cct	tg	cct	ggg	aat	aac	aca	cgt	gat	cac	cct	ggg	1262
Gln	Asp	Gly	Thr	Pro	Trp	Pro	Gly	Asn	Asn	Thr	Arg	Asp	His	Pro	Gly	
		385					390						395			
cat	gat	tca	ggt	ctt	cct	tg	gaa	ata	ctg	gga	gct	cgt	gac	att	gaa	1310
His	Asp	Ser	Gly	Leu	Pro	Trp	Glu	Ile	Leu	Gly	Ala	Arg	Asp	Ile	Glu	
	400					405					410					
gga	aat	gaa	cta	cct	cgt	cta	gta	tat	gtc	tcc	agg	gag	aag	aga	cct	1358
Gly	Asn	Glu	Leu	Pro	Arg	Leu	Val	Tyr	Val	Ser	Arg	Glu	Lys	Arg	Pro	
	415				420					425					430	
ggc	tac	cag	cac	cac	aaa	aag	gct	ggt	gca	gaa	aat	gct	ctg	gtg	aga	1406
Gly	Tyr	Gln	His	His	Lys	Lys	Ala	Gly	Ala	Glu	Asn	Ala	Leu	Val	Arg	
			435						440						445	
gtg	tct	gca	gta	ctc	aca	aat	gct	ccc	tac	atc	ctc	aat	ggt	gat	tgt	1454
Val	Ser	Ala	Val	Leu	Thr	Asn	Ala	Pro	Tyr	Ile	Leu	Asn	Val	Asp	Cys	
			450					455						460		
gat	cac	tat	gta	aac	aat	agc	aag	gct	ggt	cga	gag	gca	atg	tgc	atc	1502
Asp	His	Tyr	Val	Asn	Asn	Ser	Lys	Ala	Val	Arg	Glu	Ala	Met	Cys	Ile	
		465					470						475			
ctg	atg	gac	cca	caa	gta	ggt	cga	gat	gta	tgc	tat	gtg	cag	ttc	cct	1550
Leu	Met	Asp	Pro	Gln	Val	Gly	Arg	Asp	Val	Cys	Tyr	Val	Gln	Phe	Pro	
	480					485						490				
cag	agg	ttt	gat	ggc	ata	gat	aag	agt	gat	cgc	tac	gcc	aat	cgt	aac	1598
Gln	Arg	Phe	Asp	Gly	Ile	Asp	Lys	Ser	Asp	Arg	Tyr	Ala	Asn	Arg	Asn	
	495				500					505					510	
gta	ggt	ttc	ttt	gat	ggt	aac	atg	aaa	ggg	ttg	gat	ggc	att	caa	gga	1646
Val	Val	Phe	Phe	Asp	Val	Asn	Met	Lys	Gly	Leu	Asp	Gly	Ile	Gln	Gly	
			515						520					525		
cca	gta	tac	gta	gga	act	ggt	tgt	ggt	ttc	aac	agg	caa	gca	ctt	tac	1694
Pro	Val	Tyr	Val	Gly	Thr	Gly	Cys	Val	Phe	Asn	Arg	Gln	Ala	Leu	Tyr	
			530					535						540		
ggc	tac	ggg	cct	cct	tct	atg	ccc	agc	tta	cgc	aag	aga	aag	gat	tct	1742
Gly	Tyr	Gly	Pro	Pro	Ser	Met	Pro	Ser	Leu	Arg	Lys	Arg	Lys	Asp	Ser	
			545				550						555			
tca	tcc	tgc	ttc	tca	tgt	tgc	tgc	ccc	tca	aag	aag	aag	cct	gct	caa	1790
Ser	Ser	Cys	Phe	Ser	Cys	Cys	Cys	Pro	Ser	Lys	Lys	Lys	Pro	Ala	Gln	
		560				565						570				
gat	cca	gct	gag	gta	tac	aga	gat	gca	aaa	aga	gag	gat	ctc	aat	gct	1838
Asp	Pro	Ala	Glu	Val	Tyr	Arg	Asp	Ala	Lys	Arg	Glu	Asp	Leu	Asn	Ala	
					580					585					590	
gcc	ata	ttt	aat	ctt	aca	gag	att	gat	aat	tat	gac	gag	cat	gaa	agg	1886
Ala	Ile	Phe	Asn	Leu	Thr	Glu	Ile	Asp	Asn	Tyr	Asp	Glu	His	Glu	Arg	
				595					600					605		
tca	atg	ctg	atc	tcc	cag	ttg	agc	ttt	gag	aaa	act	ttt	ggc	tta	tct	1934
Ser	Met	Leu	Ile	Ser	Gln	Leu	Ser	Phe	Glu	Lys	Thr	Phe	Gly	Leu	Ser	
			610					615					620			
tct	gtc	ttc	att	gag	tct	aca	cta	atg	gag	aat	gga	gga	gta	ccc	gag	1982
Ser	Val	Phe	Ile	Glu	Ser	Thr	Leu	Met	Glu	Asn	Gly	Gly	Val	Pro	Glu	
			625				630						635			
tct	gcc	aac	tca	cca	cca	ttc	atc	aag	gaa	gcg	att	caa	gtc	atc	ggc	2030
Ser	Ala	Asn	Ser	Pro	Pro	Phe	Ile	Lys	Glu	Ala	Ile	Gln	Val	Ile	Gly	
			640			645							650			
tgt	ggc	tat	gaa	gag	aag	act	gaa	tg	gga	aaa	cag	att	ggt	tg	ata	2078
Cys	Gly	Tyr	Glu	Glu	Lys	Thr	Glu	Trp	Gly	Lys	Gln	Ile	Gly	Trp	Ile	
					660					665					670	
tat	ggg	tca	gtc	act	gag	gat	atc	tta	agt	ggc	ttc	aag	atg	cac	tg	2126

-continued

```

acaaaacctt tgggaattgg aatatgatcc tcggtgtagt ttccctcaag aaagcacata 3102
tcgctgtcag tatttaaatg aactgcaaga tgattgttct ctatgaagtt ttgaacagtt 3162
tgaaatgata ttatgttaaa atacagggtt tgattgtgtt gaaaaaaaa aagaaaaaaaa 3222
aaaaaaaaaa 3232

```

```

<210> SEQ ID NO 2
<211> LENGTH: 978
<212> TYPE: PRT
<213> ORGANISM: Populus tremuloides

```

```

<400> SEQUENCE: 2

```

```

Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val
 1           5           10          15
Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser
 20          25          30
Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Asn Glu Gly Arg
 35          40          45
Lys Val Cys Leu Arg Cys Gly Ser Pro Tyr Asp Glu Asn Leu Leu Asp
 50          55          60
Asp Val Glu Lys Lys Gly Ser Gly Asn Gln Ser Thr Met Ala Ser His
 65          70          75          80
Leu Asn Asp Ser Gln Asp Val Gly Ile His Ala Arg His Ile Ser Ser
 85          90          95
Val Ser Thr Val Asp Ser Glu Met Asn Asp Glu Tyr Gly Asn Pro Ile
100         105         110
Trp Lys Asn Arg Val Lys Ser Cys Lys Asp Lys Glu Asn Lys Lys Lys
115         120         125
Lys Arg Ser Pro Lys Ala Glu Thr Glu Pro Ala Gln Val Pro Thr Glu
130         135         140
Gln Gln Met Glu Glu Lys Pro Ser Ala Glu Ala Ser Glu Pro Leu Ser
145         150         155         160
Ile Val Tyr Pro Ile Pro Arg Asn Lys Leu Thr Pro Tyr Arg Ala Val
165         170         175
Ile Ile Met Arg Leu Val Ile Leu Gly Leu Phe Phe His Phe Arg Ile
180         185         190
Thr Asn Pro Val Asp Ser Ala Phe Gly Leu Trp Leu Thr Ser Val Ile
195         200         205
Cys Glu Ile Trp Phe Ala Phe Ser Trp Val Leu Asp Gln Phe Pro Lys
210         215         220
Trp Asn Pro Val Asn Arg Glu Thr Tyr Ile Glu Arg Leu Ser Ala Arg
225         230         235         240
Tyr Glu Arg Glu Gly Glu Pro Ser Gln Leu Ala Gly Val Asp Phe Phe
245         250         255
Val Ser Thr Val Asp Pro Leu Lys Glu Pro Pro Leu Ile Thr Ala Asn
260         265         270
Thr Val Leu Ser Ile Leu Ala Val Asp Tyr Pro Val Asp Lys Val Ser
275         280         285
Cys Tyr Val Ser Asp Asp Gly Ala Ala Met Leu Ser Phe Glu Ser Leu
290         295         300
Val Glu Thr Ala Glu Phe Ala Arg Lys Trp Val Pro Phe Cys Lys Lys
305         310         315         320
Phe Ser Ile Glu Pro Arg Ala Pro Glu Phe Tyr Phe Ser Gln Lys Ile
325         330         335

```

-continued

Asp	Tyr	Leu	Lys	Asp	Lys	Val	Gln	Pro	Ser	Phe	Val	Lys	Glu	Arg	Arg
			340					345					350		
Ala	Met	Lys	Arg	Asp	Tyr	Glu	Glu	Tyr	Lys	Val	Arg	Val	Asn	Ala	Leu
		355				360						365			
Val	Ala	Lys	Ala	Gln	Lys	Thr	Pro	Glu	Glu	Gly	Trp	Thr	Met	Gln	Asp
	370				375						380				
Gly	Thr	Pro	Trp	Pro	Gly	Asn	Asn	Thr	Arg	Asp	His	Pro	Gly	His	Asp
385					390					395					400
Ser	Gly	Leu	Pro	Trp	Glu	Ile	Leu	Gly	Ala	Arg	Asp	Ile	Glu	Gly	Asn
			405						410					415	
Glu	Leu	Pro	Arg	Leu	Val	Tyr	Val	Ser	Arg	Glu	Lys	Arg	Pro	Gly	Tyr
			420					425					430		
Gln	His	His	Lys	Lys	Ala	Gly	Ala	Glu	Asn	Ala	Leu	Val	Arg	Val	Ser
		435				440						445			
Ala	Val	Leu	Thr	Asn	Ala	Pro	Tyr	Ile	Leu	Asn	Val	Asp	Cys	Asp	His
	450					455						460			
Tyr	Val	Asn	Asn	Ser	Lys	Ala	Val	Arg	Glu	Ala	Met	Cys	Ile	Leu	Met
465					470						475				480
Asp	Pro	Gln	Val	Gly	Arg	Asp	Val	Cys	Tyr	Val	Gln	Phe	Pro	Gln	Arg
			485						490					495	
Phe	Asp	Gly	Ile	Asp	Lys	Ser	Asp	Arg	Tyr	Ala	Asn	Arg	Asn	Val	Val
			500					505					510		
Phe	Phe	Asp	Val	Asn	Met	Lys	Gly	Leu	Asp	Gly	Ile	Gln	Gly	Pro	Val
		515					520					525			
Tyr	Val	Gly	Thr	Gly	Cys	Val	Phe	Asn	Arg	Gln	Ala	Leu	Tyr	Gly	Tyr
	530					535					540				
Gly	Pro	Pro	Ser	Met	Pro	Ser	Leu	Arg	Lys	Arg	Lys	Asp	Ser	Ser	Ser
545					550					555					560
Cys	Phe	Ser	Cys	Cys	Cys	Pro	Ser	Lys	Lys	Lys	Pro	Ala	Gln	Asp	Pro
			565						570					575	
Ala	Glu	Val	Tyr	Arg	Asp	Ala	Lys	Arg	Glu	Asp	Leu	Asn	Ala	Ala	Ile
			580					585					590		
Phe	Asn	Leu	Thr	Glu	Ile	Asp	Asn	Tyr	Asp	Glu	His	Glu	Arg	Ser	Met
		595					600					605			
Leu	Ile	Ser	Gln	Leu	Ser	Phe	Glu	Lys	Thr	Phe	Gly	Leu	Ser	Ser	Val
	610					615					620				
Phe	Ile	Glu	Ser	Thr	Leu	Met	Glu	Asn	Gly	Gly	Val	Pro	Glu	Ser	Ala
625					630					635					640
Asn	Ser	Pro	Pro	Phe	Ile	Lys	Glu	Ala	Ile	Gln	Val	Ile	Gly	Cys	Gly
				645					650					655	
Tyr	Glu	Glu	Lys	Thr	Glu	Trp	Gly	Lys	Gln	Ile	Gly	Trp	Ile	Tyr	Gly
			660					665					670		
Ser	Val	Thr	Glu	Asp	Ile	Leu	Ser	Gly	Phe	Lys	Met	His	Cys	Arg	Gly
		675					680					685			
Trp	Arg	Ser	Ile	Tyr	Cys	Met	Pro	Val	Arg	Pro	Ala	Phe	Lys	Gly	Ser
	690					695					700				
Ala	Pro	Ile	Asn	Leu	Ser	Asp	Arg	Leu	His	Gln	Val	Leu	Arg	Trp	Ala
	705				710					715					720
Leu	Gly	Ser	Val	Glu	Ile	Phe	Phe	Ser	Arg	His	Cys	Pro	Leu	Trp	Tyr
			725					730						735	
Gly	Phe	Gly	Gly	Gly	Arg	Leu	Lys	Trp	Leu	Gln	Arg	Leu	Ala	Tyr	Ile
		740						745					750		
Asn	Thr	Ile	Val	Tyr	Pro	Phe	Thr	Ser	Leu	Pro	Leu	Ile	Ala	Tyr	Cys

-continued

755					760					765					
Thr	Ile	Pro	Ala	Val	Cys	Leu	Leu	Thr	Gly	Lys	Phe	Ile	Ile	Pro	Thr
	770					775					780				
Leu	Ser	Asn	Leu	Ala	Ser	Met	Leu	Phe	Leu	Gly	Leu	Phe	Ile	Ser	Ile
	785					790					795				800
Ile	Val	Thr	Ala	Val	Leu	Glu	Leu	Arg	Trp	Ser	Gly	Val	Ser	Ile	Glu
				805					810					815	
Asp	Leu	Trp	Arg	Asn	Glu	Gln	Phe	Trp	Val	Ile	Gly	Gly	Val	Ser	Ala
			820					825					830		
His	Leu	Phe	Ala	Val	Phe	Gln	Gly	Phe	Leu	Lys	Met	Leu	Ala	Gly	Ile
		835					840						845		
Asp	Thr	Asn	Phe	Thr	Val	Thr	Ala	Lys	Ala	Ala	Glu	Asp	Ala	Glu	Phe
	850					855					860				
Gly	Glu	Leu	Tyr	Met	Val	Lys	Trp	Thr	Thr	Leu	Leu	Ile	Pro	Pro	Thr
	865					870					875				880
Thr	Leu	Leu	Ile	Ile	Asn	Met	Ser	Gly	Cys	Ala	Gly	Phe	Ser	Asp	Ala
				885					890					895	
Leu	Asn	Lys	Gly	Tyr	Glu	Ala	Trp	Gly	Pro	Leu	Phe	Gly	Lys	Val	Phe
			900					905					910		
Phe	Ala	Phe	Trp	Val	Ile	Leu	His	Leu	Tyr	Pro	Phe	Leu	Lys	Gly	Leu
		915					920					925			
Met	Gly	Arg	Gln	Asn	Leu	Thr	Pro	Thr	Ile	Val	Val	Leu	Trp	Ser	Val
	930					935					940				
Leu	Leu	Ala	Ser	Val	Phe	Ser	Leu	Val	Trp	Val	Lys	Ile	Asn	Pro	Phe
	945					950					955				960
Val	Asn	Lys	Val	Asp	Asn	Thr	Leu	Val	Ala	Glu	Thr	Cys	Ile	Ser	Ile
				965					970					975	

Asp Cys

<210> SEQ ID NO 3

<211> LENGTH: 1010

<212> TYPE: DNA

<213> ORGANISM: Populus tremuloides

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (841)..(1008)

<220> FEATURE:

<221> NAME/KEY: misc_feature

<223> OTHER INFORMATION: 5' flanking region of PtCela coding sequence

<400> SEQUENCE: 3

```

gaattcgccc tttgaattc aggagaagat agtttccggt tcggtgaatg gctttgttca    60
cttctggtct agcaatttgc aaaagaagtt acaaacaaa tgcataattat gtaaatttaa    120
caagagatgg gttctatggt cacttatatta tgcccatcat ttgttctggg gttactcttt    180
atagtctgat tcgaagtgtc aaactgccgt ttctggtatt gcaattatgt agccataaac    240
tgtaaatcct gtacttatta gcggaccaac aaccagatat acgggatcag cgtcgtaaaa    300
gagatctcca ttctacgttt ctttctaatt tttccgtttc agtgagagaa ttaccctgat    360
acattgacat gatgattgat gattatggga accattccga tgtagacac gagaccatct    420
ggatcctgcc agttttcagt tcacatggca tctcagccca agatcatgtg tttatacgcc    480
taatgacttg tattgaaagt ttggtaagtt gaagatgtgc tctgccaac agaacccttc    540
cttaaatttc cagcaaatct ttcaaacttg gccttacacc ccgaaaatag acgtgcttct    600
acttgggttc ttgaaacca tgcaccaacc gccatacccc accaaccacc caccctcaac    660

```

-continued

cttctcttcg ccattacaaa aatgtcagta ccacctctg aaagacacca acacacccta	720
gctttgggta gggatttga tataaaaaa aggccaaaac aaaagattgg aaggaagcag	780
aggaagacc tcttgaaga attgaagttg taaagagctg gtaaagtggg aataagcaag	840
atg atg gaa tct ggg gct cct ata tgc cat acc tgt ggt gaa cag gtg	888
Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val	
1 5 10 15	
ggg cat gat gca aat ggg gag cta ttt gtg gct tgc cat gag tgt agc	936
Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser	
20 25 30	
tat ccc atg tgc aag tct tgt ttc gag ttt gaa atc aaa gag ggc cgg	984
Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Lys Glu Gly Arg	
35 40 45	
aaa gtt tgc ttg cgg tgt ggc tcg ag	1010
Lys Val Cys Leu Arg Cys Gly Ser	
50 55	

<210> SEQ ID NO 4

<211> LENGTH: 3444

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 4

gcggccgagg ttaatcgccg gttctcacia caggaatgag tttgtcctca ttaatgccga	60
tgagaatgcc cgaataagat cagtccaaga gctgagtgga cagacatgac aaatctgcag	120
agatgagatc gaattgactg ttgatggaga accgtttgtg gcatgtaacg aatgtgcatt	180
ccctgtgtgt agacctgtct atgagtacga aagacgagaa ggcaatcaag cttgtccaca	240
gtgcaaaacc cgtttcaaac gtcttaaagg aagtccaaga gttgaagggt atgaagagga	300
agatgacatt gatgatttag acaatgagtt tgagtatgga aataatggga ttggatttga	360
tcaggtttct gaaggatgt caatctctcg tcgcaactcc ggtttcccac aatctgattt	420
ggattcagct ccacctggct ctacagattcc attgctgact tacggcgacg aggacgttga	480
gatttcttct gatagacatg ctcttattgt tcctccttca cttgggtggtc atggcaatag	540
agttcatcct gtttctcttt ctgaccgac cgtggctgca catcgaaggc tgatggtacc	600
tcagaaagat cttgagggtt atggttatgg aagtgtcgtc tggaaagatc ggatggagga	660
atggaagaga aagcagaatg agaaacttca gtttgttagg catgaaggag atcctgattt	720
tgaagatggt gatgatgctg attttccaat gatggatgag ggaaggcagc cattgtctat	780
gaagatacca atcaaatcga gcaagataaa tccttaccgg atgttaattg tgctacgtct	840
tgtgattcct ggtctcttct ttcactaccg tattcttcac cccgtcaaag atgcatatgc	900
tttgtggctt atttctgtta tatgtgagat atggtttgct gtttcatggg ttcttgatca	960
gttccctaaa tggtagccta tcgagcgaga aacgtacttg gaccgactct cattaagata	1020
tgagaaagaa gggaaaccgt cgggactatc ccctgtggat gtatttgta gtacagtgga	1080
tcattgaaa gagcctcogc ttattaactgc aaataactgtc ttgtctattc ttgctgttga	1140
ttatcctgtc gataagggtg cttgttacgt atctgatgat ggtgctgcta tgcttacttt	1200
cgaagctcct tctgagaccg ctgaattcgc aaggaaatgg gttcctttct gcaagaaata	1260
ttgtattgag cctcgtgctc ccgaatggta tttctgccat aaaatggact acttgaagaa	1320
taaagtcat cccgattttg ttagggagcg gcgagccatg aagagagatt atgaagaatt	1380
caaagtaaag atcaatgctt tagtagcaac agcacagaaa gtgcctgagg atggttggac	1440
tatgcaagac ggtacacott ggcccggtaa tagtgtgcga gatcatcctg gcatgattca	1500

-continued

```

ggtcttcctt ggaagtgacg gtgttcgtga tgtcgaaaac aacgagttgc ctcgattagt 1560
ttacgtttct cgtgagaaga gacccggatt tgatcaccat aagaaggctg gagctatgaa 1620
ttccctgata cgagtctctg gggttctatc aaatgctcct taccttctga atgtcgattg 1680
tgatcactac atcaacaata gcaagctct tagagaagca atgtgtttca tgatggatcc 1740
tcagtcagga aagaaaatct gttatgttca gttccctcaa aggttcgatg ggattgatag 1800
gcacgatcga tactcaaadc gcaatgttgt gttctttgat atcaatatga aaggtttga 1860
tgggctacaa gggcctatat acgtcggtag aggttgtgtt ttcaggaggc aagcgcttta 1920
cggatttgat gcaccgaaga agaagaaggc cccacgtaag acatgcaatt gctggccaaa 1980
atggtgtctc ctatgttttg gttcaagaaa gaatcgtaaa gcaagacag tggctcgga 2040
taagaagaag aagaataggg aagcgtcaaa gcagatccac gcattagaaa atacgaaga 2100
gggcccggtg cataaagttc ttaacgtaga acagtcaacc gaggcaatgc aaatgaagtt 2160
gcagaagaaa tatgggcagt ctctgtatt tgttgcattc gcgcgtctgg agaatggagg 2220
gatggctaga aacgcaagcc cggcttctct gcttaaagaa gccatccaag tcattagtcg 2280
cggatatgaa gataaaactg aatggggaaa agagattggg tggatctatg gttctgttac 2340
cgaagatatt cttacgggtt ctaagatgca ttctcatggt tggagacatg tttattgtac 2400
accaaagtta gcggctttca aaggatcagc tccaatcaat ctttcggatc gtctccatca 2460
agttctctca tggggccttg ggtcggttga gattttcttg agtaggcatt gtcctatttg 2520
gtatggttat ggaggtgggt tgaatggct tgagcgggtg tcctacatta actctgtggt 2580
ttaccctggg acctctctac cgctcatcgt ttaactgtct ctccctgcca tctgtcttct 2640
cactggaaaa ttcatcgctc ccgagattag caactatgag agtatcctct tcatggcgct 2700
cttctcgtcg attgcaataa cgggtattct cgagatgcaa tggggcaaag ttgggatcga 2760
tgattgtgag agaaacgaac agttttgggt cattggagggt gtttctgcgc atctgtttgc 2820
tctcttccaa ggtctcctca aggttcttgc tgggtgtagc actaacttca cagtccatc 2880
aaaagcagct gatgatggag agttctctga cctttacctc ttcaaatgga cttcacttct 2940
catccctcca atgactctac tcatcataaa cgtcattgga gtcatagtcg gagtctctga 3000
tgccatcagc aatggatagc actcgtgggg accgcttttc ggaagactgt tctttgcact 3060
ttgggtcatc attcatcttt acccgttcct taaaggtttg cttgggaaac aagatagaat 3120
gccaaccatt attgtcgtct ggtccatcct cctggcctcg attcttacac ttctttgggt 3180
ccgggttaat ccgtttgttg cgaaggcgg tcctattctc gagatctgtg gtttagactg 3240
cttgtgattc gattgaccgg tggatgggtt ggtgaaaaag gtttaattcc cacggatcaa 3300
agagaggtaa gagagatatt gttttacctc taaaagactc cttcattgtg ttcattagat 3360
gatgaaaaat gaaaagaaaa agaagattta atttgttac gagaattgtt atttttgcaa 3420
gaatgtgttg tagatagcgg ccgc 3444

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1080
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: cellulose synthase

<400> SEQUENCE: 5

```

Arg Pro Arg Leu Ile Ala Gly Ser His Asn Arg Asn Glu Phe Val Leu

-continued

1	5	10	15
Ile Asn Ala Asp	Glu Asn Ala Arg	Ile Arg Ser Val	Gln Glu Leu Ser
	20	25	30
Gly Gln Thr Cys	Gln Ile Cys Arg	Asp Glu Ile Glu	Leu Thr Val Asp
	35	40	45
Gly Glu Pro Phe	Val Ala Cys Asn	Glu Cys Ala Phe	Pro Val Cys Arg
	50	55	60
Pro Cys Tyr Glu	Tyr Glu Arg Arg	Glu Gly Asn Gln	Ala Cys Pro Gln
	65	70	75
Cys Lys Thr Arg	Phe Lys Arg Leu	Lys Gly Ser Pro	Arg Val Glu Gly
	85	90	95
Asp Glu Glu Glu	Asp Asp Ile Asp	Asp Leu Asp Asn	Glu Phe Glu Tyr
	100	105	110
Gly Asn Asn Gly	Ile Gly Phe Asp	Gln Val Ser Glu	Gly Met Ser Ile
	115	120	125
Ser Arg Arg Asn	Ser Gly Phe Pro	Gln Ser Asp Leu	Asp Ser Ala Pro
	130	135	140
Pro Gly Ser Gln	Ile Pro Leu Leu	Thr Tyr Gly Asp	Glu Asp Val Glu
	145	150	155
Ile Ser Ser Asp	Arg His Ala Leu	Ile Val Pro Pro	Ser Leu Gly Gly
	165	170	175
His Gly Asn Arg	Val His Pro Val	Ser Leu Ser Asp	Pro Thr Val Ala
	180	185	190
Ala His Arg Arg	Leu Met Val Pro	Gln Lys Asp Leu	Ala Val Tyr Gly
	195	200	205
Tyr Gly Ser Val	Ala Trp Lys Asp	Arg Met Glu Glu	Trp Lys Arg Lys
	210	215	220
Gln Asn Glu Lys	Leu Gln Val Val	Arg His Glu Gly	Asp Pro Asp Phe
	225	230	235
Glu Asp Gly Asp	Asp Ala Asp Phe	Pro Met Met Asp	Glu Gly Arg Gln
	245	250	255
Pro Leu Ser Met	Lys Ile Pro Ile	Lys Ser Ser Lys	Ile Asn Pro Tyr
	260	265	270
Arg Met Leu Ile	Val Leu Arg Leu	Val Ile Leu Gly	Leu Phe Phe His
	275	280	285
Tyr Arg Ile Leu	His Pro Val Lys	Asp Ala Tyr Ala	Leu Trp Leu Ile
	290	295	300
Ser Val Ile Cys	Glu Ile Trp Phe	Ala Val Ser Trp	Val Leu Asp Gln
	305	310	315
Phe Pro Lys Trp	Tyr Pro Ile Glu	Arg Glu Thr Tyr	Leu Asp Arg Leu
	325	330	335
Ser Leu Arg Tyr	Glu Lys Glu Gly	Lys Pro Ser Gly	Leu Ser Pro Val
	340	345	350
Asp Val Phe Val	Ser Thr Val Asp	Pro Leu Lys Glu	Pro Pro Leu Ile
	355	360	365
Thr Ala Asn Thr	Val Leu Ser Ile	Leu Ala Val Asp	Tyr Pro Val Asp
	370	375	380
Lys Val Ala Cys	Tyr Val Ser Asp	Asp Gly Ala Ala	Met Leu Thr Phe
	385	390	395
Glu Ala Leu Ser	Glu Thr Ala Glu	Phe Ala Arg Lys	Trp Val Pro Phe
	405	410	415
Cys Lys Lys Tyr	Cys Ile Glu Pro	Arg Ala Pro Glu	Trp Tyr Phe Cys
	420	425	430

-continued

His Lys Met Asp Tyr Leu Lys Asn Lys Val His Pro Ala Phe Val Arg
 435 440 445

Glu Arg Arg Ala Met Lys Arg Asp Tyr Glu Glu Phe Lys Val Lys Ile
 450 455 460

Asn Ala Leu Val Ala Thr Ala Gln Lys Val Pro Glu Asp Gly Trp Thr
 465 470 475 480

Met Gln Asp Gly Thr Pro Trp Pro Gly Asn Ser Val Arg Asp His Pro
 485 490 495

Gly Met Ile Gln Val Phe Leu Gly Ser Asp Gly Val Arg Asp Val Glu
 500 505 510

Asn Asn Glu Leu Pro Arg Leu Val Tyr Val Ser Arg Glu Lys Arg Pro
 515 520 525

Gly Phe Asp His His Lys Lys Ala Gly Ala Met Asn Ser Leu Ile Arg
 530 535 540

Val Ser Gly Val Leu Ser Asn Ala Pro Tyr Leu Leu Asn Val Asp Cys
 545 550 555 560

Asp His Tyr Ile Asn Asn Ser Lys Ala Leu Arg Glu Ala Met Cys Phe
 565 570 575

Met Met Asp Pro Gln Ser Gly Lys Lys Ile Cys Tyr Val Gln Phe Pro
 580 585 590

Gln Arg Phe Asp Gly Ile Asp Arg His Asp Arg Tyr Ser Asn Arg Asn
 595 600 605

Val Val Phe Phe Asp Ile Asn Met Lys Gly Leu Asp Gly Leu Gln Gly
 610 615 620

Pro Ile Tyr Val Thr Gly Cys Val Phe Arg Arg Gln Ala Leu Tyr Gly
 625 630 635 640

Phe Asp Ala Pro Lys Lys Lys Gly Pro Arg Lys Thr Cys Asn Cys
 645 650 655

Trp Pro Lys Trp Cys Leu Leu Cys Phe Gly Ser Arg Lys Asn Arg Lys
 660 665 670

Ala Lys Thr Val Ala Ala Asp Lys Lys Lys Asn Arg Glu Ala Ser
 675 680 685

Lys Gln Ile His Ala Leu Glu Asn Ile Glu Glu Gly Arg Gly His Lys
 690 695 700

Val Leu Asn Val Glu Gln Ser Thr Glu Ala Met Gln Met Lys Leu Gln
 705 710 715 720

Lys Lys Tyr Gly Gln Ser Pro Val Phe Val Ala Ser Ala Arg Leu Glu
 725 730 735

Asn Gly Gly Met Ala Arg Asn Ala Ser Pro Ala Cys Leu Leu Lys Glu
 740 745 750

Ala Ile Gln Val Ile Ser Arg Gly Tyr Glu Asp Lys Thr Glu Trp Gly
 755 760 765

Lys Glu Ile Gly Trp Ile Tyr Gly Ser Val Thr Glu Asp Ile Leu Thr
 770 775 780

Gly Ser Lys Met His Ser His Gly Trp Arg His Val Tyr Cys Thr Pro
 785 790 795 800

Lys Leu Ala Ala Phe Lys Gly Ser Ala Pro Ile Asn Leu Ser Asp Arg
 805 810 815

Leu His Gln Val Leu Arg Trp Ala Leu Gly Ser Val Glu Ile Phe Leu
 820 825 830

Ser Arg His Cys Pro Ile Trp Tyr Gly Tyr Gly Gly Leu Lys Trp
 835 840 845

-continued

Leu Glu Arg Leu Ser Tyr Ile Asn Ser Val Val Tyr Pro Trp Thr Ser
 850 855 860
 Leu Pro Leu Ile Val Tyr Cys Ser Leu Pro Ala Ile Cys Leu Leu Thr
 865 870 875 880
 Gly Lys Phe Ile Val Pro Glu Ile Ser Asn Tyr Ala Ser Ile Leu Phe
 885 890 895
 Met Ala Leu Phe Ser Ser Ile Ala Ile Thr Gly Ile Leu Glu Met Gln
 900 905 910
 Trp Gly Lys Val Gly Ile Asp Asp Trp Trp Arg Asn Glu Gln Phe Trp
 915 920 925
 Val Ile Gly Gly Val Ser Ala His Leu Phe Ala Leu Phe Gln Gly Leu
 930 935 940
 Leu Lys Val Leu Ala Gly Val Asp Thr Asn Phe Thr Val Thr Ser Lys
 945 950 955 960
 Ala Ala Asp Asp Gly Glu Phe Ser Asp Leu Tyr Leu Phe Lys Trp Thr
 965 970 975
 Ser Leu Leu Ile Pro Pro Met Thr Leu Leu Ile Ile Asn Val Ile Gly
 980 985 990
 Val Ile Val Gly Val Ser Asp Ala Ile Ser Asn Gly Tyr Asp Ser Trp
 995 1000 1005
 Gly Pro Leu Phe Gly Arg Leu Phe Phe Ala Leu Trp Val Ile Ile His
 1010 1015 1020
 Leu Tyr Pro Phe Leu Lys Gly Leu Leu Gly Lys Gln Asp Arg Met Pro
 1025 1030 1035 1040
 Thr Ile Ile Val Val Trp Ser Ile Leu Leu Ala Ser Ile Leu Thr Leu
 1045 1050 1055
 Leu Trp Val Arg Val Asn Pro Phe Val Ala Lys Gly Gly Pro Ile Leu
 1060 1065 1070
 Glu Ile Cys Gly Leu Asp Cys Leu
 1075 1080

<210> SEQ ID NO 6
 <211> LENGTH: 56
 <212> TYPE: PRT
 <213> ORGANISM: Populus tremuloides

<400> SEQUENCE: 6

Met Met Glu Ser Gly Ala Pro Ile Cys His Thr Cys Gly Glu Gln Val
 1 5 10 15
 Gly His Asp Ala Asn Gly Glu Leu Phe Val Ala Cys His Glu Cys Ser
 20 25 30
 Tyr Pro Met Cys Lys Ser Cys Phe Glu Phe Glu Ile Lys Glu Gly Arg
 35 40 45
 Lys Val Cys Leu Arg Cys Gly Ser
 50 55

<210> SEQ ID NO 7
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: concensus sequence

<400> SEQUENCE: 7

Gln Val Leu Arg Trp
 1 5

51

What is claimed is:

1. An isolated cellulose synthase promoter comprising nucleotides from position 1 to position 840 of SEQ ID NO:3.
2. A polynucleotide comprising the promoter of claim 1, operatively linked to a polynucleotide encoding a cellulose synthase or a polynucleotide encoding a UDP-glucose binding domain.
3. A vector comprising the promoter of claim 1.
4. The vector of claim 3, wherein the cellulose synthase promoter is operatively linked to a polynucleotide encoding a cellulose synthase or a polynucleotide encoding a UDP-glucose binding domain.
5. A transgenic plant cell comprising the promoter of claim 1.
6. The plant cell of claim 5, wherein the plant cell is a tree cell.
7. The promoter of claim 1, wherein the promoter is a stress-inducible promoter.
8. A method of inducing expression of a polynucleotide in a plant comprising the polynucleotide operatively linked to the cellulose synthase promoter of claim 1, the method comprising exposing the plant to mechanical stress thereby inducing expression of the polynucleotide.
9. The method of claim 8, wherein the polynucleotide encodes a cellulose synthase, and wherein expression of the polynucleotide alters a characteristic of a plant selected from the group consisting of growth, cellulose content, lignin content, juvenile wood strength, fiber strength, and a combination thereof, as compared to a control plant.
10. The method of claim 9, wherein the altered characteristic is selected from the group consisting of accelerated growth, increased cellulose content, decreased lignin content, improved juvenile wood strength improved juvenile fiber strength, and a combination thereof.
11. A transformed plant produced by the method of claim 8.

52

12. The plant of claim 11, wherein the plant is a tree.
13. A method of altering a characteristic of a plant cell comprising the promoter of claim 1 operatively linked to a polynucleotide encoding a polypeptide, wherein the promoter comprises a negative mechanical stress response element or a positive mechanical stress response element, the method comprising exposing the plant cell to a stress to alter expression of the polypeptide, thereby altering the characteristic of the plant cell, as compared to a control plant cell.
14. A transformed plant cell produced by the method of claim 13.
15. The method of claim 13, wherein the promoter comprises a negative mechanical stress response element and the polypeptide comprises a cellulose synthase, and wherein the altered characteristic is selected from the group consisting of accelerated growth, increased cellulose content, decreased lignin content, improved strength of juvenile wood or fiber, and combinations thereof.
16. A transformed plant cell produced by the method of claim 15.
17. The plant cell of claim 16, which is from a tree.
18. The method of claim 13, wherein the promoter comprises a positive mechanical stress response element and the polypeptide comprises a cellulose synthase, and wherein the altered characteristic is selected from the group consisting of decreased cellulose content, increased lignin content, and combinations thereof.
19. A transformed plant cell produced by the method of claim 18.
20. The plant cell of claim 19, which is from a tree.
21. A transformed plant comprising the plant cell of claim 19.
22. The plant of claim 21, wherein the plant is a tree.

* * * * *