



**Michigan  
Technological  
University**

Michigan Technological University  
**Digital Commons @ Michigan Tech**

---

Department of Geological and Mining  
Engineering and Sciences Publications

Department of Geological and Mining  
Engineering and Sciences

---

2003

## Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of april 7 2001 sandstorm in China

Yingxin Gu  
*Michigan Technological University*

William I. Rose  
*Michigan Technological University*

Gregg J. Bluth  
*Michigan Technological University*

Follow this and additional works at: <https://digitalcommons.mtu.edu/geo-fp>



Part of the [Geology Commons](#), [Mining Engineering Commons](#), and the [Other Engineering Commons](#)

---

### Recommended Citation

Gu, Y., Rose, W. I., & Bluth, G. J. (2003). Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of april 7 2001 sandstorm in China. *Geophysical Research Letters*, 30(15). <http://dx.doi.org/10.1029/2003GL017405>  
Retrieved from: <https://digitalcommons.mtu.edu/geo-fp/75>

Follow this and additional works at: <https://digitalcommons.mtu.edu/geo-fp>



Part of the [Geology Commons](#), [Mining Engineering Commons](#), and the [Other Engineering Commons](#)

## Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of April 7, 2001 sandstorm in China

Yingxin Gu, William I. Rose, and Gregg J. S. Bluth

Dept. of Geological Engineering and Sciences, Michigan Technological Univ., Houghton, Michigan, USA

Received 25 March 2003; revised 24 June 2003; accepted 3 July 2003; published 8 August 2003.

[1] A thermal infrared remote sensing retrieval method developed by *Wen and Rose* [1994], which retrieves particle sizes, optical depth, and total masses of silicate particles in the volcanic cloud, was applied to an April 07, 2001 sandstorm over northern China, using MODIS. Results indicate that the area of the dust cloud observed was 1.34 million km<sup>2</sup>, the mean particle radius of the dust was 1.44 μm, and the mean optical depth at 11 μm was 0.79. The mean burden of dust was approximately 4.8 tons/km<sup>2</sup> and the main portion of the dust storm on April 07, 2001 contained 6.5 million tons of dust. The results are supported by both independent remote sensing data (TOMS) and in situ data for a similar event in 1998. This paper demonstrates that Wen and Rose's retrieval method could be successfully applied to past and future sandstorm events using IR channels of AVHRR, GOES or MODIS. **INDEX TERMS:** 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; 6015 Planetology: Comets and Small Bodies: Dust; 6061 Planetology: Comets and Small Bodies: Remote sensing; 9320 Information Related to Geographic Region: Asia. **Citation:** Gu, Y., W. I. Rose, and G. J. S. Bluth, Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of April 7, 2001 sandstorm in China, *Geophys. Res. Lett.*, 30(15), 1805, doi:10.1029/2003GL017405, 2003.

### 1. Introduction

[2] Upwelling thermal infrared radiation between 10 and 12 μm from the earth's surface is selectively scattered and absorbed by airborne particles. Volcanologists have found that clouds with silicate ash can be discriminated by using the dual thermal infrared bands found on meteorological satellites [*Prata*, 1989a], because ice or liquid water particles preferentially absorb longer wavelengths while silicates preferentially absorb shorter wavelengths (Figure 1). Silicate particles in the size range of 1–15 μm in diameter cause a negative brightness temperature difference (BTD < 0) between 10 μm and 12 μm for the volcanic cloud region with the longer wavelength channel recording a higher brightness temperature. Using BTD to discriminate the volcanic cloud from meteorological clouds is called the “split window” method [*Prata*, 1989a]. *Wen and Rose* [1994] adapted this method to retrieve particle sizes, cloud optical depth and ash mass in volcanic clouds from thermal infrared satellite data, such as the data collected by GOES (Geostationary Operational Environmental Sat-

ellites), AVHRR (Advanced Very High Resolution Radiometer), and MODIS (Moderate Resolution Imaging Spectroradiometer) detectors.


[3] Sandstorms contain large amounts of dust and sand particles which are mainly silicates and present health and transportation hazards. Many dust aerosols are small enough to be inhaled and deposit in the alveolar region of the lung which can cause serious health problems. Thick dust reduces visibility and can make it hazardous to travel by air or road. For Asian dust transported to the West Coast of North America, about 30–50% of the dust mass has a diameter of 2.5 μm [*McKendry et al.*, 2001], which represents a significant public health hazard (especially to people with chronic respiratory problems) [*Simpson et al.*, 2003].

[4] Remote sensing has been successfully used to detect and monitor the location and the trajectories of sandstorms [*Zheng et al.*, 1998, *Simpson et al.*, 2003]. *Sokolik* [2002] recently studied how the loading and composition of dust affects IR radiances observed by satellite sensors and found that mineral dust has a unique radiative signature which separates the effect of dust from that of clouds and gases. Because we know from laboratory work [*Walter and Salisbury*, 1989] that both volcanic glass and most silicate minerals including quartz and feldspar have “reststrahlen” absorption effects, related to the Si-O bonds, that will produce BTD < 0 effects, we have decided to apply the quantitative retrievals developed for volcanic clouds to dust clouds. We note that work by *Sokolik* [2002] has demonstrated that the detection of sand storms using MODIS data is optimized by using 8.6 μm data. In this study we do not use 8.6 μm data because our method is designed for volcanic clouds which also contain SO<sub>2</sub>. It is not based on the 8.6 μm band because SO<sub>2</sub> absorption is strong at that wavelength and causes an interference with ash detection. Since sandstorms do not contain anomalous SO<sub>2</sub>, it is likely that algorithms using that channel should be investigated.

[5] The MODIS was launched in December 1999 aboard the Terra satellite. It has a spatial resolution of 1 km (channels 8–36) at nadir and 36 spectral channels. Channels 31 and 32 (centered at 11.03 and 12.02 μm) are used to analyze the dust cloud in this paper.

### 2. April 07, 2001 Sandstorm in China

[6] Sandstorms are well-known phenomena during the springtime in China. The sources of the dust are from the deserts of Mongolia and China [*Zheng et al.*, 1998]. On April 6 and 7, 2001, a series of huge sandstorm events developed over Mongolia, spread to the Inner Mongolia



**Figure 1.** Schematic diagram showing how two band thermal IR transmission through meteorological and ash laden volcanic clouds is different. Bands 4 and 5 refer to AVHRR or GOES detectors; the equivalent bands for MODIS are 31 and 32 respectively.


region of China, and swept across northern China. This was the strongest sandstorm in China during the 2001 spring-time [http://www.enviroinfo.org.cn/Disasters/Sand\_storm/index.html], and affected a large area of China. The visibility in Erlianhaote City (Inner Mongolia province) was less than 50 meters, and the maximum instantaneous wind speed in that region reached 30 m/s. Large dust clouds and powerful gales made the visibility very low, many stores were closed and some airline flights had to be cancelled during that time. This sandstorm reached the U.S. two weeks later, blanketing areas from Canada to Arizona with

a layer of fine particles [http://www.noaanews.noaa.gov/stories/s624.htm].

### 3. Methodology

[7] Our retrieval of the particle sizes, optical depth, and total masses of particles from infrared two-band data follows the methods of *Wen and Rose* [1994]. Radiative transfer calculations were used with a semitransparent cloud model based on three assumptions: The shapes of the particles are spherical, the particle size distribution is uniform and monodisperse within each pixel, and the cloud forms a well-defined homogeneous single layer in each pixel [Wen and Rose, 1994]. The BTDRetrieval method assumes that the particle is within the Mie region (about 1–15 μm diameter) [Prata, 1989b]. Accurate retrievals of silicates require knowledge of refraction index data for the silicate minerals. For volcanic cloud work we use data from *Pollack et al.* [1973], but for the dust clouds we use refraction index data for Sahara dust [Volz, 1973].

[8] In situ data for this sandstorm event have not yet appeared in the literature. However, major element analysis of Chinese dust indicate the dust contains 60% wt of SiO<sub>2</sub> [Goudie, 1978]. We use the physical characteristics and the chemical composition of Asian dust observed from in situ observations for an event in April 1998 [Simpson et al., 2003]. The observed dust size distribution had a sharp peak between 1–5 μm with a mean diameter of 2 μm [Chun et al., 2001]. The chemical composition of the dust was described by SEM X-ray analysis as containing Si, Fe, Al, and Ca [McKendry et al., 2001]. The particle size distribution and the chemical composition of the Asian dust are broadly similar to andesitic volcanic ash, which could produce a strong negative BTDRetrieval signal and could discriminate the dust cloud from meteorological clouds. Using Sahara dust refraction index data, we assume that the dust particle size, total dust mass in the dust cloud and the optical



**Figure 2a.** MODIS BTDRetrieval (band 31–band 32) image at 02:40 UTC, April 07, 2001, showing the dust cloud distribution over northern China.



**Table 3.** Error Analysis of the Mass Loading Retrieval Based on Using Different Refraction Indices, Particle Size Distribution, Particle Shape, Surface Temperature and Cloud Temperature

Parameters	Resulting error range in total mass retrieval	Source
Refraction index	9–13%	<i>Wen and Rose</i> [1994] and this study
Particle size distribution	17–32%	<i>Wen and Rose</i> [1994]
Particle shape	5–20% <sup>a</sup>	<i>Krotkov et al.</i> [1999]
Surface temperature (5K change)	10–34%	<i>Schneider et al.</i> [1999]
Cloud temperature (10K change)	7–10%	this study

<sup>a</sup>The effects of particle shape are not well known. We know from *Okada et al.* [2001] that Chinese sandstorm particles have aspect ratios of 1.3 to 1.4. We also know that volcanic ash particles with similar aspect ratios (1.5) produce slight inaccuracy in UV retrievals [*Krotkov et al.*, 1999]. But we are unsure of the effects on IR retrievals.

comparison, we recalculated the mean optical depth, particle radius and mass using refraction index data from quartz [*Spitzer and Kleinman*, 1961; *Holland et al.*, 1975] and derived values of 1.16, 1.25 μm, and 8.4 million tons (Table 2). Thus the mass retrievals of the dust cloud are not significantly different if different refraction index values are used. These results may be compared with the total annual dust emission of 25 million tons and the spring season estimates of 15 million tons in the Northern China area using a “U.S. EPA formula” [*Xuan*, 1999; *Xuan et al.*, 2000]. Table 3 lists error ranges in the mass retrievals based on different refraction indices, particle size distribution, shape of the particles, surface temperature, and cloud temperature. This shows that estimates of mass using this technique have precision errors (1σ) of the order of ±10–40%, because of uncertainties and assumptions.

### 5. Conclusion

[12] The mass and sizes of the particles in large Chinese sandstorms can be retrieved using two IR bands from MODIS. Spatial mapping of the cloud with MODIS agrees with ultraviolet remote sensing data (TOMS) and with *in situ* observations of a similar event in 1998. This paper demonstrates that two band “split window” retrievals of dust events could be used to routinely quantify the intensity of the past and future sandstorm events. Because we can measure the mass and synoptic area of dust clouds, the degree of the hazard caused by the sandstorm could potentially be rapidly evaluated by remote sensing data using this method and serve as a warning system of detection. The major uncertainty for this technique comes from lack of data for IR refraction index of Chinese dust. *In situ* sample collection and the laboratory study of measuring the refraction index of Chinese dust are suggested for future work.

[13] **Acknowledgments.** The aerosol optical depth data of XiangHe site we used here was obtained from <http://aeronet.gsfc.nasa.gov>, and we thank Brent Holben for his effort in establishing and maintaining XiangHe site. This paper benefited from reviews by A. J. Prata and an anonymous reviewer. David J. Scheider designed Figure 1.

### References

Chun, Y. K., O. Boo, J. Kim, S-U. Park, and M. Lee, Synopsis, transport and physical characteristics of Asian dust in Korea, *J. Geophys. Res.*, **106**, 18,461–18,470, 2001.  
 Goudie, A. S., Dust storms and their geomorphological implications, *J. Arid Environments*, **1**, 291–310, 1978.  
 Holland, W. E., M. R. Querry, and R. M. Coveney, Measurements of spectral reflectance and optical constants of selected rock samples for

application to remote sensing of soil moisture, *NOAA Rep.*, 76022603, 78 pp., 1975.  
 Krotkov, N. A., A. J. Krueger, S. J. Schaefer, V. E. Fioletov, and J. B. Kerr, Radiative transfer modeling of the ground-based and satellite SO<sub>2</sub> and ash retrievals in volcanic clouds, *9th AMS Conference on Atmospheric Radiation*, Long Beach, CA, 1997.  
 Krotkov, N. A., D. E. Flittner, A. J. Krueger, A. Kostinski, C. Riley, W. Rose, and O. Torres, Effect of particle non-sphericity on satellite monitoring of drifting volcanic ash clouds, *J. Quant. Spectroscop. and Radiat. Transfer*, **63**, 613–630, 1999.  
 McKendry, I. G., J. P. Hacker, R. Stull, S. Sakiyama, D. Mignacca, and K. Reid, Long-range transport of Asian dust to the Lower Fraser Valley, British Columbia, Canada, *J. Geophys. Res.*, **106**, 18,361–18,370, 2001.  
 Okada, K., J. Heintzenberg, K. Kai, and Y. Qin, Shape of atmospheric mineral particles collected in three Chinese arid-regions, *Geophys. Res. Lett.*, **28**, 3123–3126, 2001.  
 Pollack, J. B., O. B. Toon, and B. N. Khare, Optical properties of some terrestrial rocks and glasses, *Icarus*, **19**, 372–389, 1973.  
 Prata, A. J., Observations of volcanic ash clouds in the 10–12 μm window using AVHRR/2 data, *Int. J. Remote Sensing*, **10**, 751–761, 1989a.  
 Prata, A. J., Infrared radiative transfer calculations for volcanic ash clouds, *Geophys. Res. Lett.*, **16**, 1293–1296, 1989b.  
 Schneider, D. J., W. I. Rose, L. R. Coke, and G. J. S. Bluth, Early evolution of a stratospheric volcanic eruption cloud as observed with TOMS and AVHRR, *J. Geophys. Res.*, **104**, 4037–4050, 1999.  
 Seftor, C. J., N. C. Hsu, J. R. Herman, P. K. Bhartia, O. Torres, W. I. Rose, D. J. Schneider, and N. Krotkov, Detection of volcanic ash clouds from Nimbus 7/total ozone mapping spectrometer, *J. Geophys. Res.*, **102**, 16,749–16,759, 1997.  
 Simpson, J. J., G. L. Hufford, R. Servranckx, J. Berg, and D. Pieri, Airborne Asian Dust: Case study of long-range transport and implications for the detection of volcanic ash, *Weather and Forecasting*, **18**, 121–141, 2003.  
 Sokolik, I. N., The spectral radiative signature of wind-blown mineral dust: Implications for remote sensing in the thermal IR region, *Geophys. Res. Lett.*, **29**(24), 2154, doi:10.1029/2002GL015910, 2002.  
 Spitzer, W. G., and D. A. Kleinman, Infrared lattice bands of quartz, *Phys. Rev.*, **121**, 1324–1335, 1961.  
 Volz, F. E., Infrared optical constants of ammonium sulfate, Sahara dust, volcanic pumice, and flyash, *Appl. Opt.*, **12**, 564–568, 1973.  
 Walter, L. S., and J. W. Salisbury, Spectral characterization of Igneous rocks in the 8- to 12-μm Region, *J. Geophys. Res.*, **94**, 9203–9213, 1989.  
 Wen, S., and W. I. Rose, Retrieval of sizes and total masses of particles in volcanic clouds using AVHRR bands 4 and 5, *J. Geophys. Res.*, **99**, 5421–5431, 1994.  
 Xuan, J., Dust emission factors for environment of Northern China, *Atmos. Environment*, **33**, 1767–1776, 1999.  
 Xuan, J., G. Liu, and K. Du, Dust emission inventory in Northern China, *Atmos. Environment*, **34**, 4565–4570, 2000.  
 Zheng, X., F. Lu, X. Fang, Y. Fang, and L. Guo, A study of dust storms in China using satellite data, in *Optical remote Sensing of the Atmosphere and Clouds*, edited by J. Wang, N. Wu, T. Ogama, and Z. Guan, *Proc. SPIE*, **3501**, 163–168, 1998.

Y. Gu, Geological Engineering and Sciences, Michigan Technological Univ., Houghton, MI, 49931 USA. (yigu@mtu.edu)  
 W. I. Rose, Geological Engineering and Sciences, Michigan Technological Univ., Houghton, MI, 49931 USA. (raman@mtu.edu)  
 G. J. S. Bluth, Geological Engineering and Sciences, Michigan Technological Univ., Houghton, MI, 49931 USA. (gbluth@mtu.edu)