

**University of New Mexico
UNM Digital Repository**

Biomedical Sciences ETDs

Electronic Theses and Dissertations

7-1-2009

Transcriptional networks of lung airway epithelial ciliogenesis

Xiaobo Sun

Follow this and additional works at: https://digitalrepository.unm.edu/biom_etds

Recommended Citation

Sun, Xiaobo. "Transcriptional networks of lung airway epithelial ciliogenesis." (2009). https://digitalrepository.unm.edu/biom_etds/107

This Thesis is brought to you for free and open access by the Electronic Theses and Dissertations at UNM Digital Repository. It has been accepted for inclusion in Biomedical Sciences ETDs by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Xiaobo Sun
Candidate

Biomedical Sciences
Department

This thesis is approved, and it is acceptable in quality
and form for publication:

Approved by the Thesis Committee:




BRIAN HJELLE

Chairperson



Kevin Harrod



Robert Rubin

**TRANSCRIPTIONAL NETWORKS OF LUNG AIRWAY
EPITHELIAL CILIOGENESIS**

BY

XIAOBO SUN

B.S., BIOTECHNOLOGY, WUHAN UNIVERSITY, 2005

THESIS

Submitted in Partial Fulfillment of the
Requirements for the Degree of

**Master of Science
Biomedical Sciences**

The University of New Mexico
Albuquerque, New Mexico

August 2009

DEDICATION

This thesis is dedicated to my parents and wife Qian who have supported me all the way since the beginning of my studies.

Also, this thesis is dedicated to those who believe in the richness of learning.

ACKNOWLEDGMENT

First of all, I would like to express my deepest sense of gratitude to my mentor and supervisor Dr. Kevin Harrod for his excellent and patient guidance in both of my experiments and thesis writings, encouragement and brilliant instructions throughout this study.

I would express my gratitude to my Scholarship sponsor from Lovelace Respiratory Research Institute and Biomedical Sciences Graduate Program of UNM.

I am also thankful to my committee members, Dr. Brian Hjelle, Dr. Robert Rubin, for their valuable opinions and advice in this study, and in revisions of this thesis.

I also would give special thanks to Dr. Wanli Lei, my experimental instructor and colleague in this study, for his constant and patient guidance in my experiments, generous assistance on editing thesis writing.

I also thank my colleague Dr. Junita Martinez, Richard Jaramillo, Kimberly Ortega and Bilan Li for sharing their experience and knowledge in this study.

**TRANSCRIPTIONAL NETWORKS OF LUNG AIRWAY
EPITHELIAL CILIOGENESIS**

BY

XIAOBO SUN

ABSTRACT OF THESIS

Submitted in Partial Fulfillment of the
Requirements for the Degree of

**Master of Science
Biomedical Sciences**

The University of New Mexico
Albuquerque, New Mexico

August 2009

**TRANSCRIPTIONAL NETWORKS OF LUNG AIRWAY
EPITHELIAL CILIOGENESIS**

BY

XIAOBO SUN

B.S., Biotechnology, Wuhan University, 2005

ABSTRACT

Motile cilia of the mammalian airway play an essential role in innate defense. The coordinated transcriptional regulation of ciliary axoneme genes remains to be elucidated. Transcription factor FOXJ1 has been shown to be important in ciliogenesis; however, direct transactivation of cilia genes by FOXJ1 has not been reported. Using a combined bioinformatics and experimental approach, here, we show a transcriptional network for cilia gene expression. FOXJ1 can directly transactivate endogenous cilia genes such as *ENKURIN*, *EFHC2*, *IFT57*, *RIBC2* and *ROPN1L* in human bronchial epithelial cells (HBEC). FOXJ1 transactivation is localized to the proximal end of the 5' flanking region of *ENKURIN* and *EFHC2* promoters. However, FOXJ1 failed to transactivate *HSPA1A* and *MNS1*, two other cilia genes, indicating there are other transcription

factor(s) involved in the ciliogenesis. Motif discovery analysis indicates ETS and RFX binding sites located in promoters of several cilia genes (*IFT57*, *HSPA1A*, *MNS1*, *RIBC2* and *ROPNIL*). QRT-PCR indicates 5 of 27 human ETS transcription factors members (ETV1, ETV5, SPDEF, SPIC and ESE1) and 4 of 7 RFX transcription factors (RFX1, RFX2, RFX5 and RFXANK) are regulated coincidentally with the differentiation of human bronchial epithelial cells and the appearance of cilia, suggesting these transcription factors are involved in gene regulations during lung airway epithelial ciliogenesis. ETS family members ESE1 and ETV1 both can transactivate endogenous EFHC2 and ENKURIN while SPIC represses EFHC2. An RFX family member RFX1 transactivates endogenous ENKURIN, HSPA1A, ROPN1L, and RFX5 transactivates EFHC2, HSPA1A and RIBC2. Furthermore, another transcription factor, GATA2, transactivates endogenous ENKURIN and EFHC2, and its transactivation also locates in the proximal end of the 5' flanking region of *ENKURIN* and *EFHC2* promoters. Our work defines a transcriptional network that regulates expression of cilia genes during airway epithelial differentiation and highlights the importance of multiple transcription factors in ciliogenesis.

TABLE OF CONTENTS

LIST OF FIGURES	x
LIST OF TABLES.....	xii
CHAPTER 1: INTRODUCTION.....	1
Human Cilia and Cilia-Related Diseases	1
Functions of Lung Epithelial Cilia and Related Diseases.....	4
Mucociliary Differentiation of Human Airway Epithelial Cells	7
Cilia Genes in Mucociliary Differentiation of HBEC Cells	8
FOXJ1 Transcription Factor in Ciliogenesis	12
ETS Transcription Factor Family	13
RFX Transcription Factor Family.....	15
General Features of Transcription Factor GATA2 and other GATA factors	16
Hypothesis and Rationale	17
CHAPTER 2: MATERIALS AND METHODS	19
Cell Cultures	19
Bioinformatic Analysis of Cilia Gene Promoters	19
Construction of Protein Expression Vectors.....	20
Construction of Luciferase-Reporter Plasmids	23
Maxiprep of Plasmids	26
Transfection and Reporter Gene Assay	26
Quantitative Real-time PCR	27
Chromatin Immunoprecipitation Assay.....	28

CHAPTER 3: RESULTS	32
FOXJ1 in human airway epithelial differentiation and ciliogenesis.....	32
ETS transcription factor family in human airway epithelial differentiation and ciliogenesis.....	39
RFX transcription factor family in human airway epithelial differentiation and ciliogenesis.....	44
GATA2 in Human Lung Airway Epithelial Differentiation and Ciliogenesis	47
CHAPTER 4: DISCUSSION	55
REFERENCE.....	65

LIST OF FIGURES

Figure 1. Axonemal Structure of Motile 9+2 Cilia.....	3
Figure 2. Scanning electron microscope image of lung trachea epithelium.....	5
Figure 3. Mucociliary differentiation of HBEC cells during ALI culture	9
Figure 4. List of the 37 cilia genes.....	10
Figure 5. The members and their homologs of ETS family	14
Figure 6. Matrices of the binding sequences of ETS family and RFX family....	21
Figure 7. Promoter structures of <i>ENKURIN</i> and <i>EFHC2</i> in luciferase reporter vectors.....	25
Figure 8. Expression of <i>FOXJ1</i> , <i>EFHC2</i> , <i>ENKURIN</i> and <i>IFT57</i> were up-regulated during lung epithelial differentiation and ciliogenesis.....	33
Figure 9. FOXJ1 transactivates <i>IFT57</i> , <i>EFHC2</i> and <i>ENKURIN</i> promoters.	35
Figure 10. Effects of FOXJ1 on endogenous cilia gene expression	36
Figure 11. <i>Cis</i> -element(s) responsive for FOXJ1 transactivation were located in the proximal promoter regions of <i>ENKURIN</i>	37
Figure 12. <i>Cis</i> -element(s) responsive for FOXJ1 transactivation were located in the proximal promoter regions of <i>EFHC2</i>	38
Figure 13. Direct binding of FOXJ1 to the proximal promoter of <i>ENKURIN</i>	40
Figure 14. Expression level changes of ETS transcription factors during lung epithelial ciliogenesis.....	42
Figure 15. Expression level changes of ETS factors during the initiation of NHBE cell differentiation and ciliogenesis	43

Figure 16. Numbers of ETS binding motifs in proximal cilia gene promoters ..	45
Figure 17. ETV1 and ESE1 transactivate endogenous <i>EFHC2</i> and <i>ENKURIN</i> , while SPIC represses <i>EFHC2</i>	46
Figure 18. Expression level changes of RFX factors during the initiation of NHBE cell differentiation and ciliogenesis.	48
Figure 19. Numbers of RFX binding motifs in proximal cilia gene promoters..	49
Figure 20. RFX1 transactivates endogenous <i>ENKURIN</i> , <i>HSPA1A</i> and <i>ROPNIL</i> ; and RFX5 transactivates endogenous <i>EFHC2</i> , <i>HSPA1A</i> and <i>RIBC2</i>	50
Figure 21. GATA2 transactivates <i>EFHC2</i> and <i>ENKURIN</i> promoters.....	52
Figure 22. GATA2 transactivates endogenous <i>EFHC2</i> and <i>ENKURIN</i>	53
Figure 23. GATA2 transactivates proximal regions of <i>ENKURIN</i> and <i>EFHC2</i> promoters.	54
Figure 24. Transcriptional network in ciliogenesis.....	64

LIST OF TABLES

Table 1. PCR primers for amplifying cDNAs of transcription factors	22
Table 2. PCR primers for amplifying proximal promoter and 5'UTR exon of cilia genes.	
.....	24
Table 3. qRT-PCR primer sites for ETS, RFX family and cilia genes.....	29

Chapter 1: Introduction

Human Cilia and Cilia-Related Diseases

Cilium is an organelle found in eukaryotic cells, which projects from the cell surface and is membrane bounded. It contains a microtubule cytoskeleton, the ciliary axoneme, surrounded by the ciliary membrane. The ciliary axoneme derives from and keeps the nine-fold symmetry of centriole, which is nearly identical to, and often becomes a ciliary basal body (1).

There are two major patterns of mammalian ciliary axoneme. Those in multiciliated epithelial cells are motile 9+2 type, possessing dynein arms. The other pattern is single, nonmotile 9+0 primary cilia which are found on virtually every cell in the body (2). These nonmotile 9+0 cilia do not assemble the central microtubule which appears in the motile 9+2 cilia and lack of the molecular motors, termed axonemal dyneins (both ODA and IDA). Primary cilia function as various specialized sensory structures with chemosensory, mechanosensory, or positional sensory activity. They incorporate specific receptors and channels into their ciliary membrane and thus can respond to either mechanical stimuli or defined ligands. The presence of primary cilium is important for cell cycle, cell differentiation and controlling tissue homeostasis (1). Loss of primary cilia leads to abnormal cell function, cell division and can cause diseases such as polycystic kidney disease (3, 4, 5, 6, 7). In addition, unique motile 9+0 cilia possessing dynein arms with left-right dynein (8, 9) are found at the mammalian embryonic node during development, determining left-right asymmetry of the body (10, 11, 12, 13).

The ciliary axoneme of motile cilia grows from and continues the nine-fold symmetry of the centriole, which is polymerized from $\alpha\beta$ tubulin heterodimers and the precursor of a ciliary basal body. The protein complex of axonemes is attached to the microtubules, including outer and inner dynein arms (ODAs and IDAs), the radial spokes, and the central pair projections. The ODAs and IDAs are molecular motors with ATPase activity, and can change their conformation alternatively by utilizing the energy of ATP hydrolysis to move along their neighboring outer microtubule doublets. This activity generates the force to make the microtubule doublets slide with respect to one another, producing a sliding motion (14, 15). The radial spokes resist this sliding motion which is converted into a local bending. The highly extensible interdoublets protein, nexin, keeps adjacent doublets together during this process (Figure 1).

The wave-like or beating motion of motile cilia is used either for locomotion such as in spermatozoa, or in the case of cilia along the respiratory tract and nodal cilia, to generate mucosal or fluid flow (16-18). Non-motile (or primary) cilia are usually associated with various sensory functions and present singly on cells (16, 19, 20). Their functions are cell type-dependent, for example, specifically responsible for vision, olfaction and mechanosensation in various tissue settings.

Ciliary defects are classified into primary and secondary ciliary defects. Primary ciliary dyskinesia is a genetic disorder causing dysfunctional motility of cilia and impaired mucociliary clearance. Secondary ciliary defects are caused by a variety of pathogens, toxins and trauma. These ciliary defects can lead to a number of human




Figure 1. Axonemal Structure of Motile 9+2 Cilia: Cilia grow from basal body composed of nine microtubule triplets. The microtubule fibers in an axoneme are in a characteristic 9+2 array, with a peripheral group of nine pairs of microtubule doublets (subfiber A and B) surrounding two singlet microtubules. The microtubule doublets are held together by nexin links and joined to a central sheath by radial spokes. Two dynein arms emerge from each subfiber A and slide along neighboring subfiber B to yield a local bending.

(Source: http://en.wikipedia.org/wiki/File:Eukaryotic_cilium_diagram_en.svg)

diseases including polycystic kidney disease (PKD), retinal degeneration, anosmia, laterality defects, chronic respiratory problems, *situs inversus*, hydrocephalus, or infertility (17, 21-23) and other ailments such as Bardet–Biedl, Alstrom and Meckel syndromes. These ailments extend the ciliary disorder to obesity, diabetes, liver fibrosis, hypertension, heart malformations, skeletal anomalies (e.g. polydactyly), cognitive impairment and developmental defects such as exencephaly (17, 21, 24-29).

Functions of Lung Epithelial Cilia and Related Diseases

The lung and respiratory passage are constantly exposed to respiratory pathogens, toxins and foreign particles, and have evolved a very effective defense system. Lung epithelium exerts protective effects through several ways: by providing a physical barrier; through the secretion of factors that mediate immunity, inflammation, and antioxidant defense; and through mucociliary clearance. Mucociliary clearance plays an important role among these defense mechanisms for clearance of foreign particles and prevention of bacterial colonization or viral spreading. The pseudostratified columnar respiratory epithelium consists of ciliated, mucus-producing and basal cells (30). Cilia line the nasal cavity, paranasal sinuses, Eustachian tubes and middle ear of the upper respiratory tract. The lower respiratory tract is lined by ciliated cells from the trachea to the terminal bronchioles. These epithelial cells lining the airway are multi-ciliated which posses approximately 100-200 9+2 motile cilia on each cell (Figure 2).

Because the basal foot processes of individual cilia in each cell point in a similar




Figure 2. Scanning electron microscope image of lung trachea epithelium: These lung epithelial cells are both ciliated and non-ciliated. The rod-like cilia on multiciliated cells is different in size from the sand-like microvilli on non-ciliated cell surface.

(Source: <http://remf.dartmouth.edu/images/mammalianLungSEM/source/9.html>)

direction (31), the effective ciliary stroke for each cell beats in a common direction.

Mucus secreted by respiratory epithelium acts as a physical barrier for capture of inhaled pathogens and particles, while the tips of cilia interact with the mucus layer to propel it along the airway surface from the lung and nasal cavities (32, 33).

Respiratory ciliary defects contribute to a wide range of human diseases such as asthma, cystic fibrosis and chronic obstructive pulmonary disease. Primary respiratory ciliary defects include primary ciliary dyskinesia (PCD) which results in chronic sinusitis, bronchiectasis, rhinitis, otitis and “glue ear” with associated hearing loss (36, 37).

A variety of pathogens, toxins and trauma could cause the secondary ciliary defects and abnormal functions. During acute viral infections such as coronavirus, mucociliary clearance has been shown to be impaired (38), which might be due to the cytopathic loss of epithelial cilia (38-42). When infected by various common cold viruses or, RSV, a paramyxovirus, ciliated lung epithelial cells are lost. Cilia may also become withdrawn into the cell body as was seen in the nasal epithelium during an episode of coronavirus infection (43-46, 55, 56). Interestingly, infection of several respiratory viruses is unique in ciliated and nonciliated airway epithelial cells (43). In addition to viral infections, bacterial infections also compromise ciliated respiratory epithelial cells. For example, when infecting the respiratory epithelium, *Pseudomonas aeruginosa* will secrete a toxic substance named pyocyanin that will immobilize the tracheal cilia (54). Infection with *Mycoplasma pneumoniae*, *Bordetella pertussis* or *B. bronchiseptica* will cause derangement of ciliary attachment (50-52).

Mucociliary Differentiation of Human Airway Epithelial Cells

The commitment of an undifferentiated airway epithelial cell to a ciliated cell and subsequent maturation to the ciliated phenotype occurs through undefined molecular pathways. In the developing human lung, ciliated epithelial cells arise from undifferentiated cells in the trachea during the canalicular stage, and, as lung development proceeds, cilia progressively appear in large and then small airways (61). Ciliogenesis in airway epithelial cells is a multistep process, in which precursor structures called “amorphous bodies” give rise to centrioles that subsequently transform to basal bodies. The basal bodies then migrate and dock at the apical membrane where they give rise to axonemes on the ciliated cells (62, 63). Although some regulators have been identified for the primary ciliogenesis (64), few molecular regulators of ciliogenesis in motile ciliated cells have been identified.

Primary human bronchial epithelial cells (HBEC) can be harvested from lung tissues and cultured in an air-liquid interface (ALI), which will promote their mucociliary differentiation. In the first week of ALI culture, the cells form a stratified squamous epithelial layer. By the end of the second week, those squamous cells are no longer flattened in appearance but without other obvious morphological changes. Although cilia could not be discerned on cells of this stage, α -tubulin are observed to concentrate on the surface of individual cells, indicating that ciliary protein constituents are being synthesized and transported, and that ciliogenesis has initiated. In other cells in this culture mode, there is high concentration of MUC5B, which is the main secreted mucin by secretory cells (57), suggesting the initiation of secretory

cell differentiation of non-ciliated cells. By the end of the third week, many of the cells have arranged in the pseudostratified layer with a columnar morphology. At the end of both third and fourth week, cilia are observed on the apical membrane surface of the cells, and there is a large increase of acidic mucin on the surface of epithelium, giving rise to the characteristic features of the predominant epithelial subpopulations of ciliated and secretory cells in the airways (58, Figure 3).

Cilia Genes in Mucociliary Differentiation of HBEC Cells

Axonemes purified from primary ALI cultures of HBEC cells have been studied by proteomic analysis, through which 200 proteins have been found (59). In another study, changes in gene expression during differentiation of airway epithelial cells have been examined by Affymetrix cDNA microarray analysis, and 1488 genes were found to display two or more-fold increase at early stage of ciliogenesis. By comparing the proteomics data with the microarray data, there are 37 overlapping genes between these two data sets (Figure 4, 58). Thus these genes are very likely to have potential roles in cilia formation or functions, and it is notable that most of them have very similar expression profiles, with a dramatic increase between days 2 and 8 of ALI culture.

Among these 37 genes, a number of genes encode proteins that are components of the ciliary axoneme, such as those encoding light, intermediate or heavy chains of axonemal dyneins (*DNAH9, DNAI1, DNAI2, DNALII*), and tubulins (*TUBA3, TUBA4, TUBB2*).





Figure 3. Mucociliary differentiation of HBEC cells during ALI culture on day 21.

Acetylated α -tubulin is immunostained to mark ciliary axonemes (red) and nuclei are stained with DAPI (blue) respectively. Ciliated cells are denoted with red apical staining to show axonemal structures.

A.



B.

Affymetrix ID	Gene	Description	Maximum Fold Increase	Protein	Protein Accession
237314_at	C10orf63	Chromosome 10 open reading frame 63	5.36	Mus musculus male testis cDNA	AK017056
220581_at	C6orf97	Chromosome 6 open reading frame 97	22.04	cDNA FLJ23305 FIS	Q9H5M3
231728_at	CAPS	Calypsoin	8.17	CAYP Human calypsoin	Q13938
209194_at	CENP2	Centrin, EF-hand protein, 2	2.52	Human calretinin	P41208
218876_at	CGI-38	Brain-specific protein	73.18	Brain-specific protein	Q9YH00
204576_s_at	CLUAP1	Clusterin-associated protein 1	3.37	KIAA0643 protein	Q75138
240857_at	DNAH9	Dynein, axonemal, heavy polypeptide 9	6.83	Dynein heavy chain	Multiple
220125_at	DNAI1	Dynein, axonemal, intermediate polypeptide 1	4.64	Dynein intermediate chain	Multiple
220636_at	DNAI2	Dynein, axonemal, intermediate polypeptide 2	9.75	Dynein intermediate chain	Multiple
205186_at	DNAI1	Dynein, axonemal, light intermediate polypeptide 1	12.24	Dynein light chain	Multiple
239733_at	DYDC2	DYDy30 domain containing 2	5.49	Similar to Riken 4933428D01	BC007374
219833_s_at	EFHC1	EF-hand domain (C-terminal) containing 1	7.52	FLJ10466 FIS	Q9NNW6
220591_s_at	EFHC2	EF-hand domain (C-terminal) containing 2	18.79	FLJ22601 FIS	Q9H653
213679_at	FU13946	Hypothetical protein FU13946	2.71	FLJ13946	Q9H849
132516_at	FZD3377	KRAB domain containing protein	3.10	KRAB10	BC007384
205498_at	GHR	Growth hormone receptor	3.39	Growth hormone receptor	AA457170
200890_s_at	HSP90A	Heat shock 70 kD protein 1A	2.36	70-kD heat shock proteins	Multiple
221800_s_at	IFT57	Intraflagellar transport 57 homolog (Chlamydomonas)	4.04	MHS482	AAK13588
222519_s_at	IFT81	Intraflagellar transport 81 homolog (Chlamydomonas)	2.15	CDV-1 protein	Q9UNV8
219372_at	IFT88	Intraflagellar transport 88 homolog (Chlamydomonas)	3.49	TG737	Q13099
204703_at	IQC6	Homo sapiens IQ motif containing E (IQCE), mRNA	3.45	KIAA1023	Q9UPX7
204202_at	LRRK23	Leucine-rich repeat containing 23	10.61	B7 protein	Q92977
204783_at	MLF1	Mlyoid leukemia factor 1	6.09	Fusion gene	AA499997
219703_at	MNS1	Meiosis-specific nuclear structural protein 1	22.83	Unnamed protein product	BAA92077
206197_at	NMES1	Non-metastatic cells 5, protein expressed in non-metastatic cells kinase	7.98	Human nucleoside diphosphate kinase	PS6597
205037_at	RAB4	RAB, member of RAS oncogene family-like 4	3.12	Putative GTP-binding protein	AAH00566
222742_s_at	RAB5	RAB, member of RAS oncogene family-like 5	2.59	Unknown protein	AAH04522
206526_at	RIBC2	RIB43A domain with coiled-coils 2	6.22	Unknown protein	AAH03024
223609_at	ROPN1L	Ropponen 1-like	21.82	AKAP-associated sperm protein	AAC59587
220105_at	RTRD1	Rhabdoid tumor deletion region gene 1	3.28	Rhabdoid tumor deletion region protein	Q9UHP6
205406_s_at	SP17	Sperm autoantigenic protein 17	4.45	Human sperm surface protein Sp17	Q15306
210033_s_at	SPAG6	Sperm-associated antigen 6	33.05	Sperm filamentous protein SPAG6	Q75602
210223_at	TEKT2	Tektin 2 (testicular)	4.30	H-Tektin-T	Multiple
209118_s_at	TUBA3	Tubulin, alpha 3	14.37	alpha-Tubulin	Multiple
207490_at	TUB44	Tubulin, alpha 4	2.84	alpha-Tubulin	Multiple
208977_x_at	TUBB2	Tubulin, beta, 2	2.81	beta-Tubulin	Multiple
223636_at	ZMYND12	Zinc finger, MYND domain containing 12	4.74	cDNA DKFZp434N2435	AL136858

Figure 4. A). This diagram shows the overlapping genes found to be regulated in ciliogenesis by microarray analysis with proteins known to be found in the cilia axoneme by proteomic analyses. The microarray analyses (Left) identified genes displaying two or more fold increases during mucociliary differentiation of HBEC cells. The proteomic analyses (Right) identified genes encoding proteins found in purified ciliary axoneme. **B).** The list of the 37 overlapping genes and their encoding proteins (Andrea J. Ross. etc. 2007. *Am J Respir Cell Mol Biol.*)

Another number of genes (*TEK2*, *CENT2*, *SPAG6*, *NME5*, *SPA17*, *IFT57*, *IFT81*, and *IFT88*) encode proteins associated with cilia, flagella or microtubules. For example, IFT57, IFT81, IFT88 are component proteins of the intraflagellar transport machinery.

Calcium homeostasis is important for many aspects of ciliary and flagellar functions, and proteins encoded by some genes (*ENKURIN*, *CAPS*, *EFHC1*, *EFHC2*) bind calcium ions or are involved in calcium transport. Enkurin is a calmodulin protein co-localizing with calcium channels, which is suggested to be an adaptor to localize calcium signal transduction machinery to calcium channels (60). *EFHC2* encodes a protein very similar to EFHC1 protein, and is proposed to be a calcium protein associated with the axoneme and plays a role in ciliary axoneme assembly and stability.

In addition, several genes encode proteins with different functions. For example, HSPA1A is a member of heat shock protein 70 family, and in conjunction with other heat shock proteins, stabilizes existing proteins against aggregation to mediate the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin –proteasome pathway (91). *ROPN1L* encodes a sperm protein, which interacts with A-kinase anchoring protein AKAP3 and allows ROPN1L to be targeted to specific subcellular compartments (92). MNS1 is a meiosis-specific structural protein expressed at the pachytene stage during spermatogenesis, and thought to be involved in the control of meiotic division and germ cell differentiation through regulation of pairing and recombination during

meiosis (89, 90). Other genes (*RABL4*, *RABL5*, *RIBC2*) encode proteins with unknown functions in cilia.

FOXJ1 Transcription Factor in Ciliogenesis

FOXJ1 (also called HFH-4) is a forkhead box (f-box) transcription factor expressed in ciliated cells located in the upper and lower airway, choroid plexus, ependyma, oviducts, testis and embryonic node (65-68). Colocalization of FOXJ1 and β -tubulin IV but not Clara cell secretory protein confirmed that FOXJ1 is expressed in ciliated epithelial cells rather than nonciliated columnar epithelial cells in the respiratory tract (73). Much evidence indicates a direct relationship between *Foxj1* expression and ciliogenesis: During the murine lung development, the expression of *Foxj1* immediately precedes the appearance of cilia in airway epithelial cells, which is in the late pseudoglandular stage at embryonic day 15.5 (65). In the infection of some respiratory viruses such as the paramyxovirus respiratory syncytial virus (RSV), loss of cilia is associated with loss of *Foxj1* expression, and during airway repair, the appearance of ultrastructural component of ciliogenesis is associated with re-expression of *Foxj1* (69, 74). It is noteworthy that interruption of *Foxj1* gene in genetically engineered mice leads to the loss of left-right dynein (*Dnahc11*) expression and absence of cilia (66, 70). In airway cells that have been already committed to ciliated cell phenotype, *Foxj1* promotes the expression of capastatin, an inhibitor of the protease calpain, leading to the stabilization of cytoskeletal anchoring proteins ezrin and EBP-50. Ezrin and EBP50 both are associated with basal bodies of

cilia in the pulmonary epithelium. Thus, Foxj1 helps the docking of basal bodies to the apical membrane and the induction of mechanisms of axoneme assembly (71, 72). Although mutations in the FOXJ1 gene have not yet been identified in human diseases with abnormalities of cilia formation (75), FOXJ1 is still considered to be a potential master regulator of ciliogenesis in airway epithelial cells.

ETS Transcription Factor Family

ETS is one of the largest families of transcription factors, and all members are identified to have a highly conserved DNA binding domain, the ETS domain, a winged helix-turn-helix structure that binds to DNA sites with a central GGA motif sequence. As well as DNA binding functions, the ETS domain is also involved in protein-protein interactions suggesting transcriptional co-regulation with other transcription factors. ETS factors act as a transcriptional activator, repressor, or both (78). There are 28 ETS genes in human and they are divided into 11 subfamilies (Figure 5, [80]). The extensive conservation in DNA-binding ETS domain among these genes leads to a substantial of redundancy in their DNA binding capacity. It is thought that interactions with other proteins are one way in which specific binding to DNA is achieved (79). There is limited similarity outside the ETS DNA binding domain. This sequence divergence contributes to the individual properties of ETS-domain transcription factors, altering their DNA binding specificity and interactions with other transcription factors (76, 77). These unique properties, coupled with distinct expression pattern, culminate in their distinct biological roles. The ETS family is present through out the body and is involved in wide variety of functions

Subfamily	Members and their homologs
ELF	ELF1, NERF/ELF2, MEF/ELF4
ELG	GABP α , ELG
ERG	ERG, FLI1, FEV
ERF	ERF/PE2, ETV3/PE1
ESE	ESE1/ESX/ELF3, ESE2/ELF5, ESE3/EHF
ETS	ETS1, ETS2, POINTED
PDEF	PDEF/SPDEF/PSE
PEA3	PEA3/E1AF/ETV4, ERM/ETV5, ER81/ETV1, ER71/ETV2
SPI	PU.1/SPI, SPIB, SPIC
TCF	ELK1, SAP1/ELK4, NET/SAP2/ELK3, LIN
TEL	TEL/ETV6, TEL2/ETV7, YAN

Figure 5. The members and their homologs of ETS family and they are divided into 11 sub-families. (Gutierrez-Hartman A.et.al. 2007. *Trends Endocrinol Metab*).

including the regulation of cellular differentiation, cell cycle control, cell migration, cell proliferation, apoptosis and angiogenesis. Multiple Ets factors have been reported to be associated with cancer, such as through gene fusion (81, 82). It is worthy to mention that ETS family members have been proven to be expressed in stratified epithelial cells, including the lung epithelium (85, 86)

Motif discovery analysis of the promoters of mouse lung gene subsets as well as their human homologues, which are found to transactivate several genetic programs during lung embryogenesis and lung cell differentiation (88, 89), identified ETS family members as candidate regulators in both sets of promoters. This analysis suggests that ETS family potentially regulates crucial lung functions and development (88).

RFX Transcription Factor Family

RFX proteins belong to the winged-helix family of transcription factors. They share a characteristic 76-amino acid DNA binding domain that binds to an X-box motif, and is conserved in many eukaryotes (99, 100). Five RFX genes (RFX1,-2,-3,-4 and -5) have been identified in mice and human genome (101). RFX factors are involved in the regulation of cell cycle (93, 94, 95), brain development and neuronal functions (96, 97), and ciliogenesis (98).

In mice and humans, RFX-1 is the first gene found as a candidate major histocompatibility complex (MHC) class II promoter binding protein (103). It also has been proved to be a transactivator of hepatitis B virus enhancer (104). It is primarily found in the human brain with high expression in cerebral cortex and Purkinje cells

(105). RFX-2 and -4 both have highest expression in human testis (106, 107). In mice, RFX-4 has been implicated in dorsal patterning of brain development and may regulate circadian rhythm in human (112, 113). RFX-3 is expressed in ciliated cells including pancreatic endocrine cells, ependymal cells and neuronal cells (108-110). RFX-5 is a transcriptional activator of MHCII genes. It also mediates the formation of an enhanceosome complex containing RFXANK, RFXAP, CREB, and CIITA (111).

The roles of RFX family in ciliogenesis of human airway epithelial cells remain largely unknown. Similar to the discovery of ETS family members by motif discovery, our lab has recently identified binding sites of RFX family members over-represented in lung gene networks in mouse and in their human homologue promoters. These findings predicate that RFX family members are candidate transcriptional regulators in mammal lung development and functions (88).

General Features of Transcription Factor GATA2 and other GATA factors

The GATA family of transcription factors controls multiple developmental processes by regulating tissue-specific gene expression. The zinc finger DNA binding domains bind to W(A/T)GATAR(A/G) motifs of gene promoter regulatory elements (118, 119).

GATA1, GATA2 and GATA3 regulate the development of different hematopoietic lineages---erythroid, hematopoietic progenitor and T-lymphoid, respectively (120-122). Similarly, GATA4, GATA5 and GATA6 have been shown to be involved in cardiac, genitourinary, and multiple endodermal developmental events

(123-125).

GATA2 was first cloned from chicken reticulocyte and shown to be present all along the development of erythroid cells (126). It is also expressed in other hematopoietic cells, neurons and cells of developing heart, liver, pituitary and in trophoblasts (127-132).

The roles of GATA transcription factors in human lung epithelial cells differentiation and ciliogenesis remain elucidated. Our lab has shown that binding motifs of GATA3 and GATA4 are over-represented in the gene promoters of mouse and human lung genes, respectively (88). These findings predicate that there might be multiple *cis*-regulatory elements of entire GATA family functioning in both organisms, and GATA family may have roles in lung epithelial cell differentiation and ciliogenesis.

Hypothesis and Rationale

The lung epithelial cells are highly specialized, differentiated cells originating from common progenitors in the lung during development. They have unique functions involved in airway host defense, gas exchange and ion transport (88, 136). The lung epithelium is a pseudostratified layer consisting of basal cells, secretory cells and columnar ciliated cells. Mucociliary clearance is an important host defense mechanism, and defects in mucociliary clearance are associated with various respiratory disorders and diseases.

Transcription of cilia genes is significantly increased during lung epithelial differentiation and ciliogenesis. Transcription factor FOXJ1 has been identified as the

only one transcription factor during *in vitro* differentiation of air way epithelial cells. However, direct transactivation of cilia genes by FOXJ1 was not elucidated. In addition, through motif discovery and bioinformatics analyses, multiple binding sites for other families of transcription factors, including RFX, ETS and GATA, were located in many of the cilia gene proximal promoter regions.

In this thesis, we hypothesize that FOXJ1, ETS, RFX and GATA transcription factors regulate the expressions of cilia genes in human bronchial epithelial cells, alone or together.

To verify this, we address the following aims:

Aim 1: Analyze the expression patterns of ETS and RFX family transcription factors during human bronchial epithelial cells differentiation and ciliogenesis, and locate binding motifs of these transcription factors in the 5' proximal promoters of cilia genes.

Aim 2: Experimentally assess transactivation of distinct cilia genes by the transcription factors FOXJ1, ETS, RFX and GATA2 through reporter assay, followed by qRT-PCR detection of the changes of endogenous cilia genes.

Aim 3: Show the direct binding of FOXJ1 onto the promoters of selected cilia genes.

Chapter 2: Materials and Methods

Cell Cultures

Cells were kept at 37°C with 5% CO₂. Normal human bronchial epithelial (NHBE) cells were purchased from Lonza and plated onto collage-coated 100mm dishes (Becton-Dickinson, Biocoat) and grown in BEGM media (Lonza) to approximately 80% confluence (~4-5days). Then the cells were plated onto collagen coated transwell inserts (Corning) with BEGM plus DMEM (Sigma-Aldrich) media in the basolateral and apical chambers. NHBE cells are allowed to grow to 100% confluency (~3-7 days) at which time the media in the chamber was removed and the cells were grown under air-liquid interface (ALI) for differentiation (14-28 days). HBEC3KT, kindly provided by Dr. John. D. Minna (137), were maintained in L-Glutamine containing Keratinocyte-SFM medium (GIBCO, Invitrogen) in flask coated with FNC mix (AthenaESTM's).

Bioinformatic Analysis of Cilia Gene Promoters

Proximal promoters of 37 cilia genes were acquired from human genome build 36 (v.1, March 2006) through UCSC genome browser. For each gene, the proximal promoter was defined as 1000 base pairs upstream of transcriptional start site and the downstream 5'UTR exon sequences were included to accommodate for the possible existence of additional *cis*-regulatory elements.

The ExPlain tool of TRANSFAC Suite (BIOBASE Knowledge Library) was used to predict transcription factor binding motifs on the cilia gene promoters. The profile

containing a matrix for the binding motifs of a set of transcription factors (ETS family for example) was created by choosing them from existing matrices (Figure 6), and the cut-offs value was set to be custom CSS (core similarity score) 0.75 and MSS (matrix similarity score) 0.85. Potential binding motifs of specific transcription factors on cilia gene promoters were analyzed and searched by uploading promoter sequences from FASTA files, and match them with a specific matrix using the Match Workflow Wizard.

Construction of Protein Expression Vectors

Total RNAs were isolated with TRIzol (Invitrogen) according to the protocol provided by manufacturer. RNAs were extracted from NHBE cells under ALI culture for 0, 2, 4 days and 3 weeks. Five hundred nanograms of total RNA were used to synthesize the first-strand cDNA with Quanti-Tech Reverse Transcription Kit (Qiagen).

Full length coding sequences of transcription factors FOXJ1, ETV1, ESE1, SPIC, SPDEF, ETV5, RFX1, RFX5 and GATA2 were amplified and flag-tagged through PCR with FastStart *Taq* DNA polymerase (Roche). The primers used are outlined in Table 1 (restriction enzyme sites are shown in italics, and the antisense of FLAG-tag coding sequences are underlined).

PCR products were cloned into pTarget T Vector (Promega). Competent NEB-5 α cells (NEB) were transformed with ligation products and selected through white/blue colonies in Agar plates containing 100ug/ml Ampicillin. White colonies were cultured

a)



b)



Figure 6. Matrices of the binding sequences of a) ETS family (identifier: V\$ETS_Q6) and b) RFX family (identifier: V\$RFX_Q6), for predicting their potential binding sites on cilia gene promoters.

Table 1. PCR primers for amplifying cDNAs of transcription factors

Name	Sequences
FOXJ1 upstream	5'-aatggcgagagactggctgcg-3'
Downstream	5'-aaagctttacttat <u>cgtcg</u> tat <u>cctgt</u> aat <u>ccaagaaggccccacgcgtggc</u> -3' (<i>HindIII</i>)
GATA3 upstream	5'-aagatctatggagggtgacggcggacc-3' (<i>BglIII</i>)
Downstream	5'-at <u>cttagattactt</u> at <u>cgtcg</u> tat <u>cctgt</u> aat <u>ccatggcggtgaccatgc</u> -3' (<i>XbaI</i>)
ETV1 upstream	5'-at <u>accggtccatggatggatttatgaccagcaag</u> -3' (<i>AgeI</i>)
Downstream	5'-at <u>ctcgagttacttatcgtcg</u> tat <u>cctgt</u> aat <u>catacacgtacgtccgtttagg</u> -3' (<i>XhoI</i>)
ETV5 upstream	5'-at <u>accggtccatggacgggtttatgatcagcaag</u> -3' (<i>AgeI</i>)
Downstream	5'-at <u>ctcgagttacttatcgtcg</u> tat <u>cctgt</u> aat <u>cgtaagcaaaggcctcggcatagg</u> -3' (<i>XhoI</i>)
ESE1 upstream	5'-at <u>accggtccatggctgcaacctgtgagattag</u> -3' (<i>AgeI</i>)
Downstream	5'-at <u>atcgattacttatcgtcg</u> tat <u>cctgt</u> aat <u>cgttccgactctggagaacctttc</u> -3' (<i>Clal</i>)
SPIC upstream	5'-at <u>accggtccatgacgtgtgt</u> aa <u>acaagacaag</u> -3' (<i>AgeI</i>)
Downstream	5'-at <u>ctcgagttacttatcgtcg</u> tat <u>cctgt</u> aat <u>cgcaatcatggatgttc</u> -3' (<i>XhoI</i>)
SPDEF upstream	5'-at <u>accggtccatggcagcgc</u> ag <u>cccggtctg</u> -3' (<i>AgeI</i>)
Downstream	5'-at <u>atcgattacttatcgtcg</u> tat <u>cctgt</u> aat <u>cgatgggtgcacgaactggtag</u> -3' (<i>Clal</i>)
RFX1 upstream	5'-aa <u>agcttatggcaacacaggcg</u> tata <u>c-3</u> ' (<i>HindIII</i>)
Downstream	5'-at <u>cttagattacttatcgtcg</u> tat <u>cctgt</u> aat <u>cgctggaggcagcgcctgcac</u> -3' (<i>XbaI</i>)
RFX5 upstream	5'-aa <u>agcttatggcagaagatgaggctg</u> -3' (<i>HindIII</i>)
Downstream	5'-at <u>cttagattacttatcgtcg</u> tat <u>cctgt</u> aat <u>ctgggggtgttgctttgggtc</u> -3' (<i>XbaI</i>)

in 2ml 2xYT medium overnight at 37°C with shaking at 250rpm. Plasmids were extracted with GenElute Plasmid Miniprep Kit (Sigma-Aldrich) according to the manufacture's protocol, and verified through restriction enzymatic cleavage analysis and sequencing.

Construction of Luciferase-Reporter Plasmids


Genomic DNAs were extracted from A549 cells with TRIzol (Invitrogen) according to the protocol provided by the manufacture and used as templates in the PCR with primers to amplify proximal promoters (1000 bases upstream) and 5' UTR exon of cilia genes *ENKURIN*, *EFHC2* and *IFT57*. The primers used are in Table 2 (restriction enzyme sites were shown in italics).

The PCR products were purified with QIAquick PCR Purification Kit (Qiagen) and digested, then ligated into pGL3-Basic vectors (Promega) treated with corresponding restriction enzymes. All the reporter plasmids were verified through sequencing. Reporter plasmids containing truncated proximal promoters and 5' UTR of *ENKURIN* and *EFHC2* were also constructed and were digested with NheI/HindIII and BstEII/HindIII respectively. Two truncated fragments (distal 400bp and proximal 700bp) of *ENKURIN* promoters as well as proximal 600bp of *EFHC2* promoters were acquired (Figure 7). The cohesive end (BstEII) of proximal 600bp fragment of *EFHC2* promoter was converted to blunt end through T4 polymerase treatment before HindIII digestion. Distal 400bp *ENKURIN* promoter, proximal 700bp *ENKURIN* promoter, proximal 600bp *EFHC2* promoters were inserted into pGL3-Basic vector treated with NheI, NheI/HindIII, SmaI/HindIII respectively. Proximal 450bp and

Table 2. PCR primers for amplifying proximal promoter and 5'UTR exon of cilia genes.

Name	Sequence
<i>ENKURIN</i> upstream	5'-atctcgaggagtcggaccgggaag-3' (XhoI)
Downstream	5'-atagatctggccaccaaattgactcct-3' (BgII)
<i>EFHC2</i> upstream	5'-actcgagaattataaaaaacgacggctc-3' (XhoI)
Downstream	5'-aaagcttacgttgccctggagacgggaag-3' (HindIII)
<i>IFT57</i> upstream	5'-acccgggcattccaccacttacctcatt-3' (XmaI)
Downstream	5'-aagatctcgccgccagtagccacgac-3' (HindIII)

a)



b)




Figure 7. a) Promoter structures of *ENKURIN* in luciferase reporter vectors. b) Promoter structures of *EFHC2* in luciferase reporter vectors.

250bp *ENKURIN* promoter were further acquired by treating the reporter plasmid containing proximal 700bp *ENKURIN* promoter with PvuII/HindIII, PstI/HindIII, then ligated to pGL3-Basic vector treated with SmaI/HindIII. All reporter plasmids were propagated in NEB-5 α cells (NEB) and verified through restriction enzymatic analysis and sequencing.

Maxiprep of Plasmids

Large amounts (~1-2 μ g/ μ l) of plasmids were acquired using Spin Doctor BAC Prep Kit& Spin Doctor Super Clean BAC Prep Kit (Gerard Biotech). The procedure was according to the protocol provided by the manufacturer with modifications as follows: after the first centrifugation, 0.8ml endotoxin removal resin (Promega) was added to supernatants and shaken for 10mins at room temperature. The resin was removed by centrifuging at 4000rpm shortly. The DNA pellets in the last step was re-suspended in 500 μ l Tris-HCl (pH8.0) and kept in 57° water bath. After re-suspension, each tube was centrifuged at maximum speed for 5mins and transferred to a new tube for subsequent transfections.

Transfection and Reporter Gene Assay

Immortalized hBEC3KT cells were seeded onto 6-well plate (2.5×10^5 cells per well, for endogenous gene transcription assay) or 48-well plate (2×10^4 cells per well, for reporter assay) 24 h before the transfection. FuGENE HD (Roche) was used for transfection according to the protocol provided by manufacturer. Typically, for each well of 6-well plate, 250ng of expression vectors, and empty expression vectors

(CMV3) were added to a total 500ng in 100 μ l serum-free GIBCO Opti-MEM medium (Invitrogen) with 2 μ l FuGENE HD. For each well of 48-well plate, 100 ng of reporter plasmid, 25 ng of transcription factor expression vector, and empty expression vector were added to a total 200ng in 17ul Opti-MEM medium with 1 μ l FuGENE HD. Each transfection of a 48-well plate was carried out in duplicates. Twenty-four hours after transfection, for 6-well plate, cells were washed cold PBS, and RNA was isolated for RT and qRT-PCR analysis with TRIzol (Invitrogen) according to the manufacturer's protocol. For 48-well plate, 24 h after transfection, cells were washed once with cold PBS followed by adding 100 μ l of 1xlysis buffer (Promega) per well, and kept at -70°C for 1 h. After 30 min of rotation at room temperature, 5 μ l of lysates was mixed with 25 μ l of substrate (Promega) to detect luciferase activity with Turner.

Quantitative Real-time PCR

Total RNAs isolated from transfected hBEC3KT and differentiated NHBE cells on day 0, 2, 4 and three weeks were extracted and reverse transcribed as aforementioned. In parallel, GAPDH CDS was amplified as an internal control with primers GS (atcactgccacccagaagac) and primer GR (ttactcctggaggccatgtg). To detect mRNA level of ETS family and RFX family, and their changes during NHBE cell differentiation, qRT-PCR forward and reverse primers were designed. Besides, to identify the effects of transcription factors on endogenous cilia gene expression after transfection, qRT-PCR primers for cilia genes (*ENKURIN*, *EFHC2*, *IFT57*, *HSPA1A*, *MNS1*, *RIBC2* and *ROPN1L*) were also designed (Table 3). QuantiTect SYBR Green

PCR Kit (Qiagen) was used with forward and reverse primers (final concentration 300nM each) according to the manufacturer's protocol in an ABI 7300 real-time PCR system (Applied Biosystem). The quantitative PCR results were analyzed using ABI's 7300 System SDS software, RQ Study.

Chromatin Immunoprecipitation Assay

To verify whether FOXJ1 directly binds to the proximal 250bp *ENKURIN* promoter and regulates its expression, chromatin immunoprecipitation assay was performed with ChIP-IT Express Enzymatic Kit (Active Motif) according to the manufacturer's protocol. In brief, twelve wells of NHBE cells cultured for 3 weeks were digested with 0.25% trypsin-EDTA solution and washed once with cold PBS, then fixed with 1% formaldehyde for 10 min at room temperature. After centrifugation and washing with cold PBS once, the residual formaldehyde was neutralized with glycine (0.125 M). Released nuclei from collected cells were enzymatically treated at 37 °C for 10 min in 400µl volume. 50µl of sheared chromatins were used for immunoprecipitation in 500µl volume with normal goat serum as a control, or polyclonal antibody against FOXJ1 (Santa Cruz). *ENKURIN* proximal 250bp promoter was amplified with EPF (gcacagtcaacgacgcttagcaat) and EPR (tctccgtcccttcttcactgctt). Surfactant protein D was used as a negative control to verify the specific enrichment of *ENKRUIN* proximal promoter by FOXJ1 antibody, and amplified with primers SPDF (tggaatgacaggcttgtggagaa) and primer SPDR (aggatattggcagcatgagggtct). Specific enrichment of *ENKURIN* proximal promoter was

quantified with software described in qRT-PCR section, and normalized to nonspecific precipitation of last exon of surfactant protein D.

Table 3. qRT-PCR primer sites for ETS, RFX family and cilia genes.

Name	Sequence
ELF1 upstream primer	agctagaaccagtaccatgcagga
downstream primer	agacacaaccactggaacctgggt
ELF2 upstream primer	cctgaaggtaaaacagattgagcc
downstream primer	acatcgagaccatctggctaaca
ELF4 upstream primer	actacccaccatgctgtct
downstream primer	tggatgttgcggcactgaaga
GABP- α upstream primer	gcctacgtgaactatgagaaactcagtcg
downstream primer	tgccttgaactttacaaatcatgtccc
ELG upstream primer	cgtatccgtccagctaacaacttgc
downstream primer	cagctcctggagaaatggctgat
FLI1 upstream primer	aggagtggatcaatcagccagtga
downstream primer	caccagcttgctgcatttgcataac
FEV upstream primer	tctttcaaggacgggaaaccc
downstream primer	agaaaactgccacagctggatct
ERF upstream primer	gatgaattacgacaagctgagccg
downstream primer	gaaattgaacttgttaggtgaaccgt
ETV3 upstream primer	ccgcaggaaatgcaaaccaca
downstream primer	tgaaggatccttgttaatgtatctg
ESE1 upstream primer	acagcaacatgacacctacgagaagc
downstream primer	ttgccaaacttgttagacgagtgc
ESE2 upstream primer	agagtactaccctgcctttagca
downstream primer	cacacatggcgcttagtccagtat
ESE3 upstream primer	tgc当地gcccacacaaa
downstream primer	tcctgggttctgtctgggtcaa
ETS1 upstream primer	ttatcagctggacaggagatggct
downstream primer	aggccacggctcagttctcataa
ETS2 upstream primer	tgccagtcattcatcagctggact
downstream primer	gcttcgttagttcatctggct

Table 3 (Cont.)

SPDEF upstream primer	tcctcaaggagggttgctactcaagc
downstream primer	acctgggctgagtcctcaattt
ETV4 upstream primer	atgaattacgacaagctgagccgc
downstream primer	aacttgtacacgtaacgctaccagc
ETV5 upstream primer	atgaactatgacaagctgagccgc
downstream primer	aggcatctgggtcacagacaat
ETV1 upstream primer	gataaacttagccgttactccgc
downstream primer	ggcctctggatcacacacaact
ETV2 upstream primer	tagcagctgcattccgttggact
downstream primer	aattcatgccggcttctcttc
SPI upstream primer	actggagggtgtcgacggcga
downstream primer	actgtacaggcggatcttcttct
SPIB upstream primer	tctggacagcgagtcggatgag
downstream primer	aactggtacaggcgcagcttctt
SPIC upstream primer	gtacaacccactttctccagcaa
downstream primer	atgcacatctccggattatacaggg
ELK1 upstream primer	ttggaggcctgtctggaggctgaa
downstream primer	agcttccgatttcaggtttggg
ELK4 upstream primer	ttatgacaaaactcagccgagccct
downstream primer	tgtacacaaaacttctgaccattcac
ELK3 upstream primer	tggaggcgccttagtccagttgct
downstream primer	tggcatgtggccattaagcagtgt
ETV6 upstream primer	ccctgcgccactactacaactaa
downstream primer	tcggccactcatgatttcatctgg
ETV7 upstream primer	gccaagatttcgagttgtggat
downstream primer	ttataatagtggcgcagggcacga
ERG upstream primer	ctctccacggtaatgcgtatgtaga
downstream primer	ggcctagtttaattttcgtagc
RFX1 upstream primer	tgcgcaggcttactctcaca
downstream primer	cctcagccgtctcatagtttcca
RFX2 upstream primer	gatcgctgtatggagagttcaa
downstream primer	gctcatgcccattgtcatttgt
RFX3 upstream primer	agtggtaacacagacaggcacatct
downstream primer	ttctccaaactctggaaatgtctcg
RFX4 upstream primer	actgtcagagaatactggacactg
downstream primer	gccgacaatgttccaccacgt
RFX5 upstream primer	tcctgctacagcagcatctcatct
downstream primer	ttggtttagatgaccgtcccgag
RFXANK upstream primer	tcaccaccgaagccgactct
downstream primer	tggttctcgatcacctgttgcact

Table 3 (Cont.)

RFXAP upstream primer	ggatcgctgcgaagacactacttt
downstream primer	ctgcttgtcttgctcca
<i>ENKURIN</i> upstream primer	tccgatgaagaaaggaggcagtt
downstream primer	agcctctgtcggtatcttctt
<i>EFHC2</i> upstream primer	agctacccaagaattgcccaccta
downstream primer	tatcgaacaggttagccatccgctt
<i>IFT57</i> upstream primer	acagatgtcgagaatggaggcata
downstream primer	cacttctgtgtgtgcatttgggt
<i>HSPA1A</i> upstream primer	aatctcagagccgagccgacaga
downstream primer	atctccacctgcccgtgtggaa
<i>RIBC2</i> upstream primer	agctctggacagcagcaacctca
downstream primer	tgtgtgaaatagtctccgtgggt
<i>ROPN1L</i> upstream primer	agactcagatgtctcccttggaa
downstream primer	agacccatcatgcccgttccctg
<i>MNS1</i> upstream primer	gagcatgtacaaaacttaacttaggc
downstream primer	cctgaactttcaccaaggcaga

Chapter 3: Results

FOXJ1 in human airway epithelial differentiation and ciliogenesis

FOXJ1 is a master regulator during the ciliogenesis; however, few target genes of FOXJ1 have been identified thus far. Mutation of the FOXJ1 gene is not correlated with abnormality of ciliogenesis (75). The expression of thirty-seven cilia genes, which encode axonemal components of cilia, increases significantly during human lung epithelial ciliogenesis (58), corresponding to the increased expression of FOXJ1. These genes are likely transcriptional targets of FOXJ1.

By quantitative RT-PCR, we assessed the relative mRNA levels of FOXJ1 as well as several cilia genes (*ENKURIN*, *EFHC2*, *IFT57*) during differentiation of NHBE cells under ALI cultures (0 day and the 21 day). We found that the expression of FOXJ1 and these cilia genes increased significantly in NHBE cells at day 21 of differentiation, when compared to undifferentiated NHBE cells lacking a ciliated cell population on day 0 (Figure 8). It is noteworthy that FOXJ1 expression was highly up-regulated.

To test whether ENKURIN, EFHC2 and IFT57 are transcriptional regulation targets of FOXJ1, we constructed an expression vector of FOXJ1 through cloning of coding sequence into pTarget expression plasmid; and reporter plasmids containing the proximal promoters (1000 bases upstream of transcriptional start site and 5' UTR sequences) of *ENKURIN*, *EFHC2* and *IFT57* were constructed with pGL3 to direct expression of the reporter gene luciferase. FOXJ1 expression vectors and the reporter




Figure 8. Expression of *FOXJ1*, *EFHC2*, *ENKURIN* and *IFT57* were up-regulated during lung epithelial differentiation and ciliogenesis. Fully differentiation of NHBE cells cultured under ALI situation will achieved after 21 days. Total RNAs were extracted from D0 and D21 ALI cells for quantitative RT-PCR detection of the relative mRNA levels of *FOXJ1* and three cilia genes, mRNA levels of D0 were set as 1.

plasmids were cotransfected into HBEC3KT cells in 48-well plate and 24 hours later, cellular lysates were mixed with luciferase substrates for reporter assay. Results showed FOXJ1 can markedly transactivate the *ENKURIN*, *EFHC2* and *IFT57* promoters (Figure 9).

Overexpression of FOXJ1 in HBEC3KT cells is able to induce the expression of ciliary axoneme genes *EFHC2*, *ENKURIN*, *RIBC2* and *ROPN1L*, while *IFT57*, *HSPA1A* and *MNS1* were not induced dramatically (Figure 10), indicating FOXJ1 can only selectively transactivate the ciliary axoneme genes *EFHC2*, *ENKURIN*, *RIBC2* and *ROPN1L*. The inability of FOXJ1 to induce expression of endogenous *IFT57* while capable of transactivating the *IFT57* promoter will be discussed later.

To identify the FOXJ1 responsive *cis*-element(s) located in *EFHC2* and *ENKURIN* proximal promoters, we truncated their proximal promoters and cloned them to pGL3 plasmid as described, and then cotransfected with FOXJ1 expression plasmid in HBEC3KT cells. *Cis*-elements responsive to FOXJ1 transactivation in the *ENKURIN* promoter were located in the proximal 250bp sequences, while the distal 400bp of the promoter didn't contain *cis*-elements responsive to FOXJ1 (Figure 11). Similarly, the proximal 600bp of *EFHC2* promoter could be transactivated by FOXJ1 (Figure 12). These results indicate that the *cis*-elements responsive to FOXJ1 transactivation are located in the proximal region of *ENKURIN* and *EFHC2* promoters.

Next, chromatin immunoprecipitation (ChIP) was employed to analyze if FOXJ1 could directly bind to the 250bp proximal region of *ENKURIN* promoter. Twelve




Figure 9. FOXJ1 transactivates *IFT57*, *EFHC2* and *ENKURIN* promoters.

Reporter plasmids containing 1000bp proximal promoter and 5' UTR of *IFT57*, *EFHC2* and *ENKURIN* were co-transfected with FOXJ1 expression plasmid respectively. Luciferase activities were measured 24h later. Luciferase activity of reporter plasmids co-transfected with control expression plasmids (empty CMV3) was set as 1.





Figure 10. Effects of FOXJ1 on endogenous cilia gene expression. Total RNAs were extracted from HBEC3KT cells transfected with FOXJ1 expression vectors. The mRNA levels of cilia genes (*EFHC2*, *ENKURIN*, *HSPA1A*, *IFT57*, *MNS1*, *RIBC2* and *ROPN1L*) were assayed by quantitative RT-PCR. Control groups were transfected with empty CMV3 expression plasmid, whose mRNA levels of cilia genes were set as 1.

A)



B)





Figure 11. Cis-element(s) responsive for FOXJ1 transactivation were located in the proximal promoter regions of ENKURIN. A) ENKURIN promoter (proximal 1000bp+5'UTR) were truncated at 700 bp, 450 bp, 250 bp proximal to the transcriptional start site, and cloned into the luciferase reporter gene construct. In addition the distal 400 bp of the proximal promoter was also cloned into the luciferase expression construct. B) Subsequent cotransfection with FOXJ1 expression plasmid was used to determine promoter activity as measured by luciferase activity assay.

A)

Promoter structure of *EFHC2*



B)




Figure 12. Cis-element(s) responsive for FOXJ1 transactivation were located in the proximal promoter regions of *EFHC2*. A) *EFHC2* truncated proximal 600bp fragment was cloned to pGL3 luciferase- reporter plasmid. B) Subsequent co-transfection with FOXJ1 expression plasmid.

wells of NHBE cells under ALI culture for three weeks were treated as described for ChIP analysis. Chromatin precipitates were enzymatically sheared into 200bp~400bp pieces. Goat polyclonal antibody against FOXJ1 (Santa Cruz) was used for specific enrichment of *ENKURIN* proximal promoter and non-immunized goat serum was used as negative control. Quantitative PCR showed that the FOXJ1 polyclonal antibody, but not the normal goat sera, specifically enriched the proximal *ENKURIN* promoter (Figure 13), indicating *ENKURIN* is an authentic transcriptional target of FOXJ1, and the *cis*-elements for FOXJ1 transactivation are located in the proximal *ENKURIN* promoter.

ETS transcription factor family in human airway epithelial differentiation and ciliogenesis

Although FOXJ1 was the only transcription factor identified by microarray to be increased during the differentiation of NHBE cells (58), it failed to transactivate several endogenous cilia genes. We analyzed the transcript levels of other transcription family members such as ETS proteins, which are expressed in stratified epithelium such as the pulmonary airways (85, 86). In lung epithelial cells, ETS family members are involved in transcriptional regulation and differentiation of several differentiated components of the airways (83, 84, 87). Previously, this laboratory has discovered ETS binding motifs overrepresented in the promoters of cilia genes. ETS binding motifs were found on promoters of mouse lung gene subsets as well as their human homologues, regulating the lung cell differentiation (88, 89).




Figure 13. Direct binding of FOXJ1 to the proximal promoter of *ENKURIN*.

Sheared chromatins from NHBE cells of three-week ALI culture were co-immunoprecipitated with goat polyclonal antibody against FOXJ1 or normal goat serum. The specific enrichment of *ENKURIN* proximal promoter sequence was quantified by quantitative PCR as detailed in *Materials and Methods*.

Thus ETS family members are potential transcriptional regulators involved in lung epithelial ciliogenesis in addition to FOXJ1. Quantitative RT-PCR was employed to detect the changes of expression patterns of all twenty-eight ETS family members in differentiated, ciliated NHBE cells under ALI culture. Results showed that the expression of ETV1, ETV2, ETV6, ETV7, ESE1, ESE2, ESE3, SPDEF and SPIC were significantly increased during the differentiation, while the expression of ETV4 and ETV5 were decreased. The level of other ETS factors did not show significant changes during ciliogenesis, and some, such as SPI, SPIB and ELG, were expressed in very low level in both unciliated and ciliated NHBE cells, indicating they are not likely involved in the lung epithelial ciliogenesis (Figure. 14).

The initial stage of ciliogenesis occurs between 2d and 4d apical upon exposure to the air. To further define more acutely the temporal induction of ETS family members, we assessed the expression level changes of ETS factors, during this stage to further identify those that are most likely involved in regulation of ciliogenesis. As Figure 15 shows, by quantitative RT-PCR of identification of the eleven dramatically changed ETS factors in Day 21 of ALI cells, the mRNA level of eight (ESE1, ETV1, ESE2, ESE3, SPIC, ETV2, ETV6, SPDEF) were significantly up-regulated during the initial stage of epithelial differentiation, while ETV5 significantly down-regulated. These results suggest that these nine ETS factors, are most likely involved in regulation the initiation of NHBE differentiation.

To identify transcription factors in regulation of ciliary axoneme genes, motif discovery analysis as detailed in *Methods and Materials* was utilized to examine the





Figure 14. Expression level changes of ETS transcription factors during lung epithelial ciliogenesis. Total RNAs were extracted from unciliated (ALI Day 0) and ciliated (ALI Day 21) NHBE cells. The mRNA levels of ETS transcription factors were detected by quantitative RT-PCR. For comparison, the mRNA levels in unciliated cells (D0) were set as 1.




Figure 15. Expression level changes of ETS factors during the initiation of NHBE cell differentiation and ciliogenesis. Total RNAs were extracted from NHBE cells under ALI culture at Day 0, Day 2 and Day 4, followed by quantitative RT-PCR to detect the mRNA levels of chosen ETS genes (heatmap, deep red: mRNA levels increased over 2 fold; deep green, mRNA levels decreased over 2 fold). Day 0 was set as calibrator (black).

proximal promoter (proximal 1000bp plus 5' UTR) of cilia genes to predict ETS binding sites. As shown in Figure 16, eight and seven ETS binding motifs were identified in *ROPN1L* and *HSPA1A* promoters among all cilia genes, respectively. Four ETS sites were identified in the proximal promoter region of *ENKURIN* as well. So we picked up *ROPN1L*, *HSPA1A* as well as four other cilia genes of interests (*ENKURIN*, *EFHC2*, *IFT57* and *MNS1*) for further analyses to test if their expressions are regulated by the five ETS factors. HBEC3KT cells were transfected with expression plasmids of ETV1, ESE1, SPDEF, SPIC and ETV5 in 6-well plates. Twenty four hours after transfection, mRNA level of these cilia genes were quantified by quantitative RT-PCR. We found that ETV1 and ESE1 were able to transactivate endogenous *EFHC2* and *ENKURIN*, and SPIC repressed the expression of *EFHC2* (Figure 17). These five ETS factors had no effects on expression of endogenous *ROPN1L*, *HSPA1A*, *MNS1* and *IFT57* (data not shown).

RFX transcription factor family in human airway epithelial differentiation and ciliogenesis

The RFX family of transcription factors is associated with cilia formation in many eukaryotic species including human being (102, 114-117). However, their roles in ciliogenesis of lung epithelial cells remain largely unknown. Recently, our lab has identified RFX binding motifs in promoters of human and mouse gene subsets that are associated with fully differentiated lung epithelium, suggesting RFX factors are possibly involved in lung epithelial differentiation (88). No information exists regarding the expression or function of RFX family transcription factors in human




Figure 16. Numbers of ETS binding motifs in proximal promoters of cilia genes.

The numbers of ETS binding motifs on proximal 1000bp plus 5' UTR region of cilia gene promoters were predicted by motif discovery analysis as detailed in *Methods and Materials*.




Figure 17. ETV1 and ESE1 transactivate endogenous *EFHC2* and *ENKURIN*, while SPIC represses *EFHC2*. HBEC3KT cells were transfected with expression plasmids of ETS factors (ETV1, ESE1, SPIC, SPDEF, ETV5). The effects of these transcription factors on mRNA levels of *EFHC2* and *ENKURIN* were analyzed by quantitative RT-PCR. The mRNA levels in cells transfected with empty CMV3 plasmid was set as calibrator (whose signal was set as 1).

airway epithelium. Similar to ETS family of transcription factors, we first identified RFX family members during the initial stage of differentiation and ciliogenesis of NHBE cells by quantitative RT-PCR. As Figure 18 shows, RFX1, RFX3, RFXAP and RFX5 were up-regulated significantly during differentiation and ciliogenesis, while other RFX members did not show obvious changes, indicating these RFX factors may be associated with the initiation of NHBE differentiation and ciliogenesis.

To identify the possible transcriptional regulation targets of RFX factors, we analyzed the proximal promoters of cilia gene promoters for RFX binding motifs. The numbers of binding motifs in seven cilia gene promoters (*MNS1*, *RIBC2*, *ROPNIL*, *HSPA1A*, *ENKURIN*, *EFHC2* and *IFT57*) are shown in Figure 19, in which *MNS1* and *RIBC2* have most among all cilia genes.

We tested effects of RFX1 and RFX5 on the endogenous expression of the seven cilia genes. As showed in Figure. 20, RFX1 transactivated endogenous *ENKURIN*, *HSPA1A* and *ROPNIL*, and RFX5 trans-activated endogenous *EFHC2*, *HSPA1A* and *RIBC2* (Figure 20). No obvious trans-activating effects were observed on other cilia genes.

GATA2 in Human Lung Airway Epithelial Differentiation and Ciliogenesis

GATA2 regulates the development of hematopoietic progenitor, and is expressed in many cell types such as cells of developing heart, liver, pituitary and trophoblasts (121, 127-132) and has been shown to be an important regulator of differentiated




Figure 18. Expression level changes of RFX factors during the initiation of NHBE cell differentiation and ciliogenesis. Total RNAs were extracted from NHBE cells under ALI culture at Day 0, Day 2 and Day 4. The mRNA levels of RFX factors were quantified by quantitative RT-PCR (deep red: >2 fold upregulation; deep green: >2 fold down-regulation). The mRNA level on Day 0 was set as calibrator (black).




Figure 19. Numbers of RFX binding motifs in proximal promoters of cilia genes.

RFX binding motifs in proximal 1000bp plus 5' UTR region of the seven cilia gene promoters were identified through motif discovery as detailed in *Methods and Materials*.




Figure 20. RFX1 transactivates endogenous *ENKURIN*, *HSPA1A* and *ROPN1L*; and RFX5 transactivates endogenous *EFHC2*, *HSPA1A* and *RIBC2* (over 1.5 fold change). HBEC3KT cells were transfected with RFX1 and RFX5 expression plasmids. Their effects on mRNA levels of the seven cilia genes (*MNS1*, *RIBC2*, *ROPN1L*, *HSPA1A*, *ENKURIN*, *EFHC2* and *IFT57*) were analyzed by quantitative RT-PCR. The mRNA levels of the cells transfected with empty expression plasmid (CMV3) was set as calibrator (whose signal was set as 1).

epithelium in embryogenesis. Its role in human airway epithelial cells differentiation

is still unclear. Previously, we have identified multiple GATA binding motifs in the promoters of gene subsets associated with lung epithelial differentiation. To verify this presumption, HBEC3KT cells were cotransfected with expression plasmids of GATA2 and reporter plasmids containing *ENKURIN* or *EFHC2* promoters directing the repression of reporter gene luciferase. The reporter assay showed GATA2 transactivated both *ENKURIN* and *EFHC2* promoters (Figure 21).

The GATA2 transactivation on the endogenous *ENKURIN* and *EFHC2* gene promoters was assessed in HBEC3KT cells following transfection of a GATA2 expression plasmid. The mRNA levels of *ENKURIN* and *EFHC2* were detected 24 hours later. As showed in Figure. 22, both endogenous promoters were transactivated by GATA2. To further locate the *cis*-element(s) responsive to transactivation, GATA2 expression vector and reporter plasmids containing truncated promoter fragments of *ENKURIN* and *EFHC2* were cotransfected as aforementioned. Results showed that GATA2 significantly trans-activated proximal *ENKURIN* (700bp and 400bp) and *EFHC2* (600bp) promoters, indicating the *cis*-element(s) responsive for transactivation of GATA2 were located in the proximal promoter regions of the two cilia genes (Figure 23).




Figure 21. GATA2 transactivates *EFHC2* and *ENKURIN* promoters. Reporter plasmids containing 1000bp proximal promoter and 5' UTR of *EFHC2* or *ENKURIN* was co-transfected with GATA2 expression plasmid. Luciferase activity was measured 24h after co-transfection. Reporter plasmid co-transfected with empty CMV3 expression plasmids was set as control, whose activity was set as 1.




Figure 22. GATA2 transactivates endogenous *EFHC2* and *ENKURIN*.

HBEC3KT cells were transfected with empty vector (CMV3) or GATA2 expression vector respectively. The mRNA levels of *EFHC2* or *ENKURIN* were quantified by qRT-PCR. The mRNA levels in cells transfected with empty expression vector (CMV3) was set as 1.




Figure 23. GATA2 transactivates proximal regions of *ENKURIN* and *EFHC2* promoters. HBEC3KT cells was transfected with GATA2 expression vector and reporter plasmids containing truncated distal 400bp, proximal 700bp, proximal 450bp of *ENKURIN* promoter, as well as proximal 600bp of *EFHC2* promoter. Luciferase activities were measured 24 hours after transfection. Cells transfected with empty CMV3 expression plasmid were set as value of one (1).

Chapter 4: Discussion

Ciliated cells in human airways are important components of host defense as part of the mucociliary clearance mechanism. Ciliary defects are caused by an ever growing recognition of genetic disorders and are often characterized by chronic or recurrent infections of the respiratory tract. In addition, viral infections lead to dysfunction of cilia which likely contributes in part to the overall presentation of lung disease. The transcriptional regulation of human airway epithelial ciliogenesis is still largely undefined. In this study, transcription factors that confer the coordinated expression of genes that encode ciliary axoneme structural proteins in ciliogenesis are identified and transcriptional targets of each transcription factor are defined.

FOXJ1 is a master regulator of motile cilia and is up-regulated dramatically during differentiation of NHBE cells *in vitro*. Quantitative RT-PCR data showed that the expression of several cilia genes (*ENKURIN*, *EFHC2*, *IFT57*, *MNS1*, *RIBC2*, *ROPN1L* and *HSPA1A*) are markedly increased during differentiation of human airway epithelia. We found that FOXJ1 transactivated the promoters of several cilia genes in reporter gene assays, as well as transactivated the endogenous cilia genes of human differentiated airway epithelial cells. Using studies of truncated promoters, the *cis*-elements responsive for transactivation of FOXJ1 were located into the proximal promoter regions of *ENKURIN* and *EFHC2*. Chromatin immunoprecipitation assays provided further support that FOXJ1 binds an unknown *cis*-active element in proximal promoter region of *ENKURIN*.

Although FOXJ1 endogenously transactivated the promoters of several cilia genes, it failed to transactivate other cilia genes, such as MNS1 and IFT57. Previous studies have indicated that overexpression of FOXJ1 alone does not confer ciliogenesis or differentiation of unciliated lung epithelia. These findings suggest that other transcription factors are also important in regulation of ciliogenesis of lung epithelial cells. Through motif discovery, multiple binding sites for ETS, RFX and GATA transcription factors were mapped in the promoters of a number of cilia genes, indicating cilia genes are potential targets of these transcription factors.

The twenty-eight factors of ETS transcription family are involved in a variety of mammalian developmental process at the cellular, tissue and organ level, including hematopoiesis, immune function, lymph/angiogenesis, neurogenesis/neuromuscular function, spermatogenesis, early embryonic patterning, and development of extraembryonic tissues (138-149). All members of this transcription factor family have the ETS DNA-binding domain which binds to sequences that contain a central GGA motif. However, individual ETS domain protein can select nucleotides over an 11-base-pair sequences centered on this GGA motif. ETS transcription factors are regulated, in part, by phosphorylation. Extracellular signals are transduced through a series of mitogen-activated protein kinases (MAPKs) which result in activation of downstream proteins such as ERK, JNK and p38, which in turn phosphorylate and regulate the activity of ETS transcription factors (78). Although the overall functions and roles of ETS family in lung epithelial differentiation is still unknown by now, ETS family members have been shown to be involved in the transcriptional regulation

and differentiation of mammalian lung and its epithelium differentiation. For example, aberrant expression of ELF5 disrupts lung branching and inhibits epithelial differentiation (83); In adult mouse lung, ETV5 is expressed by alveolar epithelial cells and regulates caveolin-1 transcription in mouse lung epithelial cell lines (84); In human lung epithelial cells, ESE-1, -2, -3 have regulatory effects on expression of the host defense genes lysozyme, chitinase and secretoglobins (87).

Many RFX factors are involved in cilia formations and functions in different eukaryotes. For example, in *C. elegans*, the only known RFX factor to date regulates *daf-19* which is a key regulator of ciliogenesis (102). In zebra fish, *rfx-2* is specifically expressed in multiciliated cells of the pronephros and loss of its expression results in cyst formation and loss of multicilia (114). *Rfx2, foxj1* and *ift88* are recently shown to be regulated by FGF signaling which regulates cilia length in diverse epithelial during embryo development (151). Most importantly, in mice, RFX-3 functions in ciliogenesis are conserved. It controls the growth of mouse embryonic node cilia, and loss of RFX-3 leads to significantly truncated but functional node cilia (115). RFX-3 loss-of-function also results in hydrocephalus with differentiation defects of ciliated ependymal cells of the choroid plexus and subcommisural organ (116). Moreover, RFX-3 mutant mice show primary ciliary growth defects on islet cells and subsequent insulin secretion failure and impaired glucose tolerance (117).

Although ETS and RFX families are associated with cilia formation in many cell types and in many other simpler organisms, their roles in ciliogenesis of lung

epithelial cell subsets are still unknown and remain to be elucidated. We have performed quantitative RT-PCR to determine the expression levels of all twenty eight ETS factors and six RFX factors during the differentiation and ciliogenesis of NHBE cells under ALI culture. Among the twenty-eight ETS family members, the expression of nine (ETV1, ETV2, ETV6, ETV7, ESE1, ESE2, ESE3, SPDEF and SPIC) were significantly increased during the differentiation, while ETV4 and ETV5 were decreased. ESE1, ETV1, SPIC and SPDEF were most significantly up-regulated while ETV5 down-regulated most. RFX1 and RFX5 were the only RFX factors dramatically up-regulated, RFX3 and RFXAP were also up-regulated to some extent, while RFX2 and RFXANK slightly down-regulated. Based on these findings, we focused our studies on the five ETS factors (ESE1, ETV1, SPIC, SPDEF, ETV5) and the two RFX factors (RFX1 and RFX5) as putative transcriptional regulators in the initiation of ciliogenesis. Motif discovery analysis of cilia gene promoters identified ETS or RFX binding motifs in the proximal 1.2 kb region upstream of the first exon, suggesting putative binding sites for these transcription factors. Among the 37 cilia genes, promoters of *MNS1*, *RIBC2* and *ROPNIL* have the most RFX binding motifs, and promoters of *ROPNIL*, *HSPA1A* and *ENKURIN* had the most ETS binding motifs, suggesting these gene promoters are most likely to have potential RFX and ETS binding sites for transcription regulation. These five cilia genes as well as *EFHC2* and *IFT57* were selected to assess the transactivation ability of ectopic ETS or RFX in endogenous gene expression studies. Among the five ETS transcription factors (ETV1, ESE1, SPIC, SPDEF, ETV5), ETV1 and ESE1 were shown to transactivate both

EFHC2 and *ENKURIN* expression, and SPIC repressed *EFHC2* expression. None of the ETS factors has obvious effects on the expression of *ROPNIL*, *HSPA1A*, *MNS1* and *IFT57*. The transcriptional regulation of these ETS factors on the cilia genes may be directly or indirectly, cooperative or independent, which requires further study. Although *HSPA1A* and *ROPNIL* have most ETS binding motifs, these binding motifs are not functional or they require other factors that are not yet apparent.

RFX1 and RFX5 overexpression was also assessed for the ability to regulate endogenous cilia structural gene expression. *ENKURIN*, *HSPA1A* and *ROPNIL* were transactivated by RFX1, and *EFHC2*, *HSPA1A* and *RIBC2* by RFX5. *IFT57* and *MNS1* were not regulated by either RFX1 or RFX5. Our findings provide the first evidence for ETS- or RFX-mediated cilia gene induction in any metazoan system.

Our motif discovery also identified GATA elements in the promoters of ciliary axoneme genes. Although GATA family members GATA4 and GATA6 are involved in regulation of mammalian organogenesis by transactivating BMP-4 (150), so far, GATA6 is the only known GATA factor important in lung-specific gene expression and development (133-135). In this study, we investigate the effects of GATA2 on cilia gene expressions. Over-expression of GATA2 in HBEC cells led to the transactivation of *ENKURIN* and *EFHC2*, and this transactivation was located to proximal promoter region of *ENKURIN* and *EFHC2*. Transactivation of endogenous cilia genes in lung epithelial cells further supports GATA2 as a direct transactivator of those cilia genes.

The genomic program conferring differentiation, development, or other complex

cell changes operates in part through the coordinated regulation of gene expression. This program is executed by the recognitions of *cis*-regulatory DNAs by regulated transcription factors and thus the control of specific gene expression. These regulatory inputs and functional outputs of genes constitute a network-like architecture commonly referred as transcriptional regulatory networks. Transcriptional regulatory networks provide specific causal links between genomic expression patterns and biological processes such as differentiation or development (152, 153). Many transcription factors are expressed across multiple tissues and organs. For example, in adult mammals, ETS family of transcription factors are widely expressed in a variety of tissues including hematopoietic tissue, vasculature, brain and central nervous system, mammary gland, endometrium, ovaries, testes, kidneys and lungs (140). Responding to stimuli, these transcription factors turn on/off the transcription of the appropriate genes which in turn allows changes of cell morphology or activity needed for cell fate determination and cellular differentiation. Although most ETS-domain proteins have been shown to be transcriptional activators, a considerable number have been shown to be repressors, including ERF, YAN and TEL. For example, ETS factor TEL1 has been shown to be a transcriptional activator and repressor, requiring homo-dimerization or oligomerization (156). It also interacts with another ETS factor Fli1, inhibiting Fli1 transactivation activity and thereby preventing the activation of target genes (157,158). Therefore, the cellular specificity required for tissue specific phenotypic of functional processes likely involves multiple, unique transcription factor combinations to confer a coordinated gene expression network. Here we show

that multiple transcription factors from ETS (ETV1, ESE1, SPIC, SPDEF, ETV5), RFX (RFX1, RFX5) and GATA (GATA2) family can regulate cilia gene expressions together with FOXJ1 in ciliogenesis. A diagram illustrating the overlapping and unique transcription factor-gene promoter pairs is depicted in Figure 24.

Besides transcriptional regulation, increasing evidence indicates that post-transcriptional regulation plays a very important role in regulating steady-state mRNA abundance or mRNA translation and, in turn, determining cell or tissue specificity. After a gene is transcribed, the stability and distribution of the different transcripts is regulated (post-transcriptional regulation) by means of RNA binding proteins (RBP) that control the various steps and rates of the transcripts. Small RNA, either encoded as polycistronic RNA elements expressed and processed into 21-24 bp microRNAs (miRNAs), or small RNAs processed from the introns of expressed protein-encoding genes is another type of post-transcriptional regulation of genomic expression.

Much information regarding the expression of miRNA has been elucidated in recent years. First, a primary miRNA transcript (pri-miRNA), which can be many kilobases long and forms a stem-loop secondary structure, is transcribed from the genome. This pri-miRNA is recognized by a protein complex including the enzyme Drosha in nucleus, and excised as a ~60nt hairpin called pre-miRNA. This pre-miRNA is transported into cytoplasm and processed by an enzyme called Dicer, leaving a 22nt double-stranded RNA duplex which is then taken up by a protein complex RISC. Within RISC, the double strand RNA duplex will be separated, and a

guide miRNA strand will remain bound to RISC and be the element recognized by target RNAs. Depending on the degree with which the guide miRNA strand matches the target mRNA, the target RNA is inhibited through two different mechanisms. If the match is imperfect, protein translation of that RNA will be inhibited by RISC complex. On the other hand, if the match is perfect, the target RNA will be cleaved by Aragonaute (Ago) protein which is a component of RISC complex. MiRNAs play key roles in mammalian gene expression and cellular processes, especially differentiation and development. In Dicer knock-out mice in which all miRNA productions are inhibited, developing embryos die at an early embryonic stage and are depleted of pluripotent stem cells (159), supporting miRNA is crucial in proper embryogenesis and stem cell development. Conditional knockout of Dicer in a tissue or organ of interests has demonstrated that miRNAs play roles in morphogenesis of several organs including the lungs, limbs, and muscles, in T cell differentiation, and in skin development (160-165). Considering the crucial roles of miRNAs in regulation of gene expression and cellular differentiation, it is not surprising that misregulation of miRNAs are linked with a variety of human diseases such as cancers. In cilia dysfunction polycystic kidney disease (PKD), the functional patterns of predicted miRNA targets and differentially expressed mRNAs are similar, suggesting an important role of miRNAs in specific pathways underlying PKD (155). It is noteworthy to mention that in mice within which Dicer is conditionally knocked out in lung epithelium, normal branching is arrested and a few large epithelial pouches appear in these mutant lungs. Significantly, these initial phenotypes are apparent

before an increase in epithelial cell death is observed, proposing that miRNAs play specific roles in lung epithelial morphogenesis independent of its requirement in cell survival (160). In the ciliogenesis of lung epithelial cells, some cilia genes may be also regulated by post-transcriptional regulation, especially miRNA mechanism. In the future study, techniques such as RNA immunoprecipitation on a chip can be applied to investigate the expression patterns of predicted miRNAs targeting cilia genes, and their role in regulating cilia gene expressions, which may shed lights on elucidating the whole process of lung epithelial differentiation during development.




Figure 24. Transcriptional network in ciliogenesis: The transcriptional factors and their cilia gene targets are connected. (arrow: transactivation; bar: inhibition)

Reference

1. Peter Satir, Soren Tvorup Chrsitensen. 2007. Overview of Structure and Function of Mammalian Cilia. *Annu. Rev. Physiol.* 69:377-400
2. Wheatley DN. 1982. *The Centriole: A Central Enigma of Cell Biology.* Amsterdam: Elsevier. 232 pp.
3. Fonte VG, Searls RL, Hilfer SR. 1971. The relationship of cilia with cell division and differentiation. *J. Cell Biol.* 49:226–29
4. Schwartz EA, Leonard ML, Bizios R, Bowser SS. 1997. Analysis and modeling of the primary cilium bending response to fluid shear. *Am. J. Physiol.* 272:F132–3
5. Praetorius HR, Spring KA. 2001. Bending the MDCK cell primary cilium increases intracellular calcium. *J. Membr. Biol.* 184:71–79
6. Praetorius HR, Spring KA. 2003. Removal of the MDCK cell primary cilium abolishes flow sensing. *J. Membr. Biol.* 191:69–76
7. Liu W, Murcia NS, Duan Y, Weinbaum S, Yoder BK, et al. 2005. Mechanoregulation of intracellular Ca²⁺ concentration is attenuated in collecting duct of monocilium-impaired *orpk* mice. *Am. J. Physiol. Renal Physiol.* 289:F978–88
8. Supp DM, Witte DP, Potter SS, Brueckner M. 1997. Mutation of an axonemal dynein affects left-right asymmetry in *inversus viscerum* mice. *Nature* 389:963–96

9. Supp DM, Brueckner M, Kuehn MR, Witte DP, Lowe LA, et al. 1999. Targeted deletion of the ATPbinding domain of left-right dynein confirms its role in specifying development of left-right asymmetries. *Development* 126:5495–504
10. Nonaka S, Shiratori H, Saijoh Y, Hamada H. 2002. Determination of left-right patterning of the mouse embryo by artificial nodal flow. *Nature* 418:96–99
11. McGrath J, Somlo S, Makova S, Tian X, Brueckner M. 2003. Two populations of node monocilia initiate left-right asymmetry in the mouse. *Cell* 114:61–73
12. Nakaya MA, Biris K, Tsukiyama T, Jaime S, Rawls JA, Yamaguchi TP. 2005. *Wnt3a* links left-right determination with segmentation and anteroposterior axis elongation. *Development* 132:5425–36
13. Tanaka Y, Okada Y, Hirokawa N. 2005. FGF-induced vesicular release of Sonic hedgehog and retinoic acid in leftward nodal flow is critical for left-right determination. *Nature* 435:172–77
14. Satir P. 1997. Cilia and related microtubular arrays in the eukaryotic cell. In *Handbook of Physiology*, ed. JF Hoffman, JD Jamieson, pp. 787–817. New York: Oxford Univ. Press
15. Okada Y, Takeda S, Tanaka Y, Belmonte JC, Hirokawa N. 2005. Mechanism of nodal flow: a conserved symmetry breaking event in left-right axis determination. *Cell* 121:633–44
16. Scholey, J.M. (2003) Intraflagellar transport. Annu. Rev. *Cell Dev. Biol.* 19, 423–443
17. Ibanez-Tallon, I. et al. (2003) To beat or not to beat: roles of cilia in

- development and disease. *Hum. Mol. Genet.* 12, 27–35
18. Cole, D.G. (2003) The intraflagellar transport machinery of *Chlamydomonas reinhardtii*. *Traffic* 4, 435–442
19. Davenport, J.R. and Yoder, B.K. (2005) An incredible decade for the primary cilium: a look at a once-forgotten organelle. *Am. J. Physiol. Renal Physiol.* 289, 1159–1169
20. Pazour, G.J. and Witman, G.B. (2003) The vertebrate primary cilium is a sensory organelle. *Curr. Opin. Cell Biol.* 15, 105–110
21. Pan, J. et al. (2005) Cilium-generated signaling and cilia-related disorders. *Lab. Invest.* 85, 452–463
22. Afzelius, B.A. (2004) Cilia-related diseases. *J. Pathol.* 204, 470–477
23. Pazour, G.J. and Rosenbaum, J.L. (2002) Intraflagellar transport and cilia-dependent diseases. *Trends Cell Biol.* 12, 551–555
24. Beales, P.L. (2005) Lifting the lid on Pandora's box: the Bardet-Biedl syndrome. *Curr. Opin. Genet. Dev.* 15, 315–323
25. Zhang, Q. et al. (2003) Loss of the Tg737 protein results in skeletal patterning defects. *Dev. Dyn.* 227, 78–90
26. Ansley, S.J. et al. (2003) Basal body dysfunction is a likely cause of pleiotropic Bardet–Biedl syndrome. *Nature* 425, 628–633
27. Hearn, T. et al. (2005) Subcellular localization of ALMS1 supports involvement of centrosome and basal body dysfunction in the pathogenesis of obesity, insulin resistance, and type 2 diabetes.

28. Kulaga, H.M. et al. (2004) Loss of BBS proteins causes anosmia in humans and defects in olfactory cilia structure and function in the mouse. *Nat. Genet.* 36, 994–998
29. Nishimura, D.Y. et al. (2004) Bbs2-null mice have neurosensory deficits, a defect in social dominance, and retinopathy associated with mislocalization of rhodopsin. *Proc. Natl. Acad. Sci. U. S. A.* 101, 16588–16593
30. Nettesheim, P., J.S. Koo, and T. Gray. 2000. Regulation of differentiation of the tracheobronchial epithelium. *J. Aerosol. Med.* 13:207–218.
31. Sleigh, M.A., J.R. Blake, and N. Liron. 1988. The propulsion of mucus by cilia. *Am. Rev. Respir. Dis.* 137:726–741.
32. Knowles, M.R., and R.C. Boucher. 2002. Mucus clearance as a primary innate defense mechanism for mammalian airways. *J. Clin. Invest.* 109:571–577.
33. Widdicombe, J.H. 2002. Regulation of the depth and composition of airway surface liquid. *J. Anat.* 201: 313–318.
34. Schidlow, D.V. 1994. Primary ciliary dyskinesia (the immotile cilia syndrome). *Ann. Allergy* 73:457–470.
35. Camner, P., B. Mossberg, and B.A. Afzelius. 1983. Measurements of tracheobronchial clearance in patients with immotile-cilia syndrome and its value in differential diagnosis. *Eur. J. Respir. Dis. Suppl.* 127:57–63.

36. W. Stannard, M.B. Ch.B. 2006. Ciliary Functions and the Role of Cilia in Clearance. *Journal of Aerosol Medicine*. 19: 110-115
37. Afzelius, B.A., B. Mossberg, and S. Bergstrom. 2001. Immotile cilia syndrome (primary ciliary dyskinesia), including Kartagener syndrome. In The metabolic and molecular bases of inherited disease. C.R. Scriver, A.L. Beaudet, W.S. Sly, and D. Valle, eds. New York: McGraw Hill, 4817–4827.
38. Pedersen, M., Y. Sakakura, B. Winther, et al. 1983. Nasal mucociliary transport, number of ciliated cells, and beating pattern in naturally acquired common colds. *Eur. J. Respir. Dis. Suppl.* 128:355–365.
39. Wilson, R., E. Alton, A. Rutman, et al. 1987. Upper respiratory tract viral infection and mucociliary clearance. *Eur. J. Respir. Dis.* 70:272–279.
40. Chilvers, M.A., M. McKean, A. Rutman, et al. 2001. The effects of coronavirus on human nasal ciliated respiratory epithelium. *Eur. Respir. J.* 18:965–970.
41. Carson, J.L., A.M. Collier, and S.S. Hu. 1985. Acquired ciliary defects in nasal epithelium of children with acute viral upper respiratory infections. *N. Engl. J. Med.* 312:463–468.
42. Rautiainen, M., J. Nuutinen, H. Kiukaanniemi, et al. 1992. Ultrastructural changes in human nasal cilia caused by the common cold and recovery of ciliated epithelium. *Ann. Otol. Rhinol. Laryngol.* 101:982–987.
43. Thompson C, Barclay W, Zambon M, et al. 2006. Infection of Human Airway Epithelium by Human and Avian Strains of Influenza A Virus. *Journal of Virology*. 80: 8060-8068

44. D. Look, M. Walter, M. Williamson, et al. 2001. Effects of Paramyxoviral Infection on Airway Epithelial Cell Foxj1 Expression, Ciliogenesis, and Mucociliary Function. *American Journal of Pathology*. 159: 2055-2069.
45. J. Nicholls, L. Poon, K. Lee, et al. 2003. Lung pathology of fatal severe acute respiratory syndrome. *Lancet*. 361: 1773–78.
46. S. Philippou · P. Otto · P. Reinhold, et al. 1999. Respiratory syncytial virus-induced chronic bronchiolitis in experimentally infected calves. *Virchows Arch*. 436:617–621
50. Carson JL, Collier AM, Hu DC. 1980. Ultrastructural observations on cellular and subcellular aspects of experimental *Mycoplasma pneumoniae* disease. *Infect Immun*. 29: 11 117–11 124.
51. Muse K. 1980. Host cell membrane perturbations during experimental *Bordetella pertussis* infection. *Proc 7th Eur Conf El Micr Haag*. 2: 432–433.
52. Bemis DA, Wilson SA. 1985. Influence of potential virulence determinants on *Bordetella bronchiseptica* induces ciliostasis. *Infect Immun*. 50: 3–42.
54. Reimer A, Klementsson K, Ursing J, Wretlind B. 1980. The mucociliary activity of the respiratory tract. *Acta Otolaryngol*. 90:462–469.
55. Rautiainen M, Nuutinen J, Kiukaanniemi H, Collan Y. 1992. Ultrastructural changes in human nasal cilia caused by the common cold and recovery of ciliated epithelium. *Ann Otol Rhinol Laryngol* . 101: 982–987.
56. Afzelius BA. 1994. Ultrastructure of human nasal epithelium during an episode

- of coronavirus infection. *Virchows Arch.* 424: 295–300.
57. Bernacki SH, Nelson AL, Abdullah L, Sheehan JK, Harris A, Davis CW, Randell SH. 1999. Mucin gene expression during differentiation of human airway epithelia *in vitro*:Muc4 and muc5b are strongly induced. *Am J Respir Cell Mol Biol.* 20:595–604.
58. Andrea J. Ross, Lisa A Dailey, Luisa E Brighton, etc. 2007. Transcriptional Profiling of Mucociliary Differentiation in Human Airway Epithelial Cells. *Am J Respir Cell Mol Biol.* 37: 169-185.
59. Ostrowski LE, Blackburn K, Radde KM, Moyer MB, Schlatzer DM, Moseley A, Boucher RC. (2002). A proteomic analysis of human cilia: Identification of novel components. *Mol Cell Proteomics.* 1:451–465.
60. Sutton KA, Jungnickel MK, Wang Y, Cullen K, Lambert S, Florman HM. 2004. Enkurin is a novel calmodulin and trpc channel binding protein in sperm. *Dev Biol.* 274:426–435.
61. Gaillard DA, Lallement AV, Petit AF, and Puchelle ES. 1989. In vivo ciliogenesis in human fetal tracheal epithelium. *Am J Anat* 185: 415–428.
62. Dirksen ER. 1991. Centriole and basal body formation during ciliogenesis revisited. *Biol Cell* 72: 31–38.
63. Sorokin SP. 1968. Reconstructions of centriole formation and ciliogenesis in mammalian lungs. *J Cell Sci* 3: 207–230.
64. Pazour GJ, Dickert BL, Vucica Y, Seeley ES, Rosenbaum JL, Witman GB, and Cole DG. 2000. Chlamydomonas IFT88 and its mouse homologue, polycystic

- kidney disease gene tg737, are required for assembly of cilia and flagella. *J Cell Biol* 151: 709–718.
65. Blatt EN, Yan XH, Wuerffel MK, Hamilos DL, and Brody SL. 1999. Forkhead transcription factor HFH-4 expression is temporally related to ciliogenesis. *Am J Respir Cell Mol Biol* 21: 168–176.
66. Brody SL, Yan XH, Wuerffel MK, Song SK, and Shapiro SD. 2000. Ciliogenesis and left-right axis defects in forkhead factor HFH-4-null mice. *Am J Respir Cell Mol Biol* 23: 45–51.
67. Hackett BP, Brody SL, Liang M, Zeitz ID, Bruns LA, and Gitlin JD. 1995. Primary structure of hepatocyte nuclear factor/forkhead homologue 4 and characterization of gene expression in the developing respiratory and reproductive epithelium. *Proc Natl Acad Sci USA* 92: 4249–4253.
68. Lim L, Zhou H, and Costa RH. 1997. The winged helix transcription factor HFH-4 is expressed during choroid plexus epithelial development in the mouse embryo. *Proc Natl Acad Sci USA* 94: 3094–3099.
69. Look DC, Walter MJ, Williamson MR, Pang L, You Y, Sreshta JN, Johnson JE, Zander DS, and Brody SL. 2001. Effects of paramyxoviral infection on airway epithelial cell Foxj1 expression, ciliogenesis, and mucociliary function. *Am J Pathol* 159: 2055–2069.
70. Chen J, Knowles HJ, Hebert JL, and Hackett BP. 1998. Mutation of the mouse hepatocyte nuclear factor/forkhead homologue 4 gene results in an absence of cilia and random left-right asymmetry. *J Clin Invest* 102:1077–1082.

71. Yingjian Y, Tao H, Edward J R, Steven L B. 2004. Role of f-box factor foxj1 in differentiation of ciliated airway epithelial cells. *Am J Respir Cell Mol Biol* 286: 650-657.
72. Brigitte N G, Xiulan G, Brian H. 2004. Foxj1 regulates basal body anchoring to the cytoskeleton of ciliated pulmonary epithelial cells. *J Cell Sci* 117: 1329-1337
73. Tichelaar, J. W., S. E. Wert, R. H. Costa, S. Kimura, and J. A. Whitsett. 1999. HNF-3/forkhead homologue-4 (HFH-4) is expressed in ciliated epithelial cells in the developing mouse lung. *J. Histochem. Cytochem.* 47:823–831.
74. Tristam DA, Hicks WJ, Hard R. 1998. Respiratory syncytial virus and human bronchial epithelium. *Arch Otolaryngol Head Neck Surg.* 124:777–783.
75. Maiti AK, Bartoloni L, Mitchison HM, Meeks M, Chung E, Spiden S, Gehrig C, Rossier C, DeLozier-Blanchet CD, Blouin J, et al. 2000. No deleterious mutations in the Foxj1 (alias HFH-4) gene in patients with primary ciliary dyskinesia (PCD). *Cytogenet Cell Genet.* 90:119–122.
76. Shore, P. *et al.* 1996. Determinants of DNA-binding specificity of ETS-domain transcription factors. *Mol. Cell. Biol.* 16, 3338–3349 .
77. Fitzsimmons, D. *et al.* 1996. Pax-5 (BSAP) recruits Ets protooncogene family proteins to form functional ternary complexes on a B-cell-specific promoter. *Genes Dev.* 10, 2198–2211.
78. Sharrocks AD. 2001. The ETS-domain transcription factor family. *Nat Rev Mol Cell Biol* 2 (11): 827–37.

79. Verger A, Duterque-Coquillaud M. 2002. When Ets transcription factors meet their partners. *Bioessays* **24** (4): 362–70.
80. Gutierrez-Hartman A, Duval DL, Bradford AP. 2007. ETS transcription factors in endocrine systems. *Trends Endocrinol Metab* **18** (4): 150–8.
81. Ida K, Kobayashi S, Taki T, Hanada R, Bessho F, Yamamori S, Sugimoto T, Ohki M, Hayashi Y. 1995. EWS-FLI-1 and EWS-ERG chimeric mRNAs in Ewing's sarcoma and primitive neuroectodermal tumor. *Int J Cancer* **63** (4): 500–4.
82. Peeters P, Raynaud SD, Cools J, Wlodarska I, Grosgeorge J, Philip P, Monpoux F, Van Rompaey L, Baens M, Van den Berghe H, Marynen P. 1997. Fusion of TEL, the ETS-variant gene 6 (ETV6), to the receptor-associated kinase JAK2 as a result of t(9;12) in a lymphoid and t(9;15;12) in a myeloid leukemia. *Blood* **90** (7): 2535–40.
83. Metzger DE, Stahlman MT, Shannon JM. 2008. Misexpression of ELF5 disrupts lung branching and inhibits epithelial differentiation. *Dev Biol*. 1;320(1):149-60.
84. Kathuria H, Cao Y, Hinds A, Ramirez MI, Williams MC. 2007. ERM is expressed by alveolar epithelial cells in adult mouse lung and regulates caveolin-1 transcription in mouse lung epithelial cell lines. *J Cell Biochem*. 102(1):13-27.
85. Kola,I. et al. 1993. The Ets1 transcription factor is widely expressed during

- murine embryo development and is associated with mesodermal cells involved in morphogenetic processes such as organ formation. *Proc. Natl Acad. Sci. USA*, 90, 7588–7592.
86. Kathuria,H. et al. 2004. Transcription of the caveolin-1 gene is differentially regulated in lung type I epithelial and endothelial cell lines: a role for ETS proteins in epithelial cell expression. *J. Biol. Chem.* 279, 30028–30036.
87. W. Lei, K. Harrod. et al. 2007. Transactivation of lung lysozyme expression by Ets family member ESE-1. *Am J Physiol Lung Cell Mol Physiol* 293: 1359–1368.
88. Martinez. MJ, Smith .AD, Harrod. KS. 2007. Computational prediction of novel components of lung transcriptional networks. *Bioinformatics*. 23(1): 21-9.
89. Furukawa K, Hotta Y. et al. 1994. cDNA cloning and functional characterization of a meiosis-specific protein (MNS-1) with apparent nuclear association. *Chromosome Res.* 2(2): 99-113.
90. Furukawa K, Hotta Y. et al. 1995. Meiosis specific transcription and functional proteins. *Adv Biophys.* 31: 101-15
91. “Entrez Gene: HSPA1A heat shock 70kDa protein 1A.”
92. “Entrez Gene: ROPN1L ropporin 1-like.”
93. Wu, S. Y., and McLeod, M. 1995. The sak1+ gene of *Schizosaccharomyces pombe* encodes an RFX family DNA-binding protein that positively regulates cyclic AMP-dependent protein kinase-mediated exit from the mitotic cell cycle.

- Mol. Cell. Biol* 15, 1479–1488.
94. Huang, M., Zhou, Z., and Elledge, S. J. 1998. The DNA replication and damage checkpoint pathways induce transcription by inhibition of the Crt1 repressor. *Cell* 94, 595–605.
95. Otsuki, K., Hayashi, Y., Kato, M., Yoshida, H., and Yamaguchi, M. 2004. Characterization of dRFX2, a novel RFX family protein in *Drosophila*. *Nucleic Acids Res* 32, 5636–5648.
96. Ma, K., Zheng, S., and Zuo, Z. 2006. The transcription factor regulatory factor X1 increases the expression of neuronal glutamate transporter type 3. *J. Biol. Chem* 281, 21250–21255.
97. Zhang, D., Stumpo, D. J., Graves, J. P., DeGraff, L. M., Grissom, S. F., Collins, J. B., Li, L., Zeldin, D. C., and Blackshear, P. J. 2006. Identification of potential target genes for RFX4_v3, a transcription factor critical for brain development. *J. Neurochem* 98, 860–875.
98. Senti G, Swoboda P. 2008. Distinct isoforms of the RFX Transcription factor DAF-19 regulate ciliogenesis and maintenance of synaptic activity. *Mol Biol of the Cell*. 19 (12): 5517-28.
99. Zhang, D., Stumpo, D. J., Graves, J. P., DeGraff, L. M., Grissom, S. F., Collins, J. B., Li, L., Zeldin, D. C., and Blackshear, P. J. 2006. Identification of potential target genes for RFX4_v3, a transcription factor critical for brain development. *J. Neurochem* 98, 860–875.
100. Gajiwala KS, Chen H, Cornille F, Roques BP, Reith W, Mach B, Burley SK.

2000. Structure of the winged-helix protein hRFX1 reveals a new mode of DNA binding. *Nature*. 403:916-921.
101. Emery P, Durand B, Mach B, Reith W. 1996. RFX proteins, a novel family of DNA binding proteins conserved in the eukaryotic kingdom. *Nucleic Acids Res*. 24:803-807.
102. Swoboda P, Adler HT, Thomas JH. 2000. The RFX-type transcription factor DAF-19 regulates sensory neuron cilium formation in *C. elegans*. *Mol Cell*. 5:411-421.
103. Reith W, Barras E, Satola S, Kobr M, Reinhart D, Sanchez CH, Mach B. 1989. Cloning of the major histocompatibility complex class II promoter binding protein affected in a hereditary defect in class II gene regulation. *Proc Natl Acad Sci U S A*. 86(11):4200-4204.
104. Siegrist CA, Durand B, Emery P, David E, Hearing P, Mach B, Reith W. 1993. RFX1 is identical to enhancer factor C and functions as a transactivator of the hepatitis B virus enhancer. *Mol Cell Biol*. 13:6375-6384.
105. Ma K, Zheng S, Zuo Z. 2006. The transcription factor regulatory factor X1 increases the expression of neuronal glutamate transporter type 3. *J Biol Chem*. 281(30):21250-21255.
106. Wolfe SA, van Wert J, Grimes SR. 2006. Transcription factor RFX2 is abundant in rat testis and enriched in nuclei of primary spermatocytes where it appears to be required for transcription of the testis-specific histone H1t gene. *J Cell Biochem*. 99(3):735-746.

107. Morotomi-Yano K, Yano K, Saito H, Sun Z, Iwama A, Miki Y. 2002. Human regulatory factor X 4 (RFX4) is a testis-specific dimeric DNA-binding protein that cooperates with other human RFX members. *J Biol Chem.* 277(1):836-842.
108. Morotomi-Yano K, Yano K, Saito H, Sun Z, Iwama A, Miki Y. 2002. Human regulatory factor X 4 (RFX4) is a testis-specific dimeric DNA-binding protein that cooperates with other human RFX members. *J Biol Chem.* 277(1):836-842.
109. Blackshear PJ, Graves JP, Stumpo DJ, Cobos I, Rubenstein JL, Zeldin DC. 2003. Graded phenotypic response to partial and complete deficiency of a brain-specific transcript variant of the winged helix transcription factor RFX4. *Development.* 130(19):4539-4552.
110. Ait-Lounis A, Baas D, Barras E, Benadiba C, Charollais A, Nlend Nlend R, Liegeois D, Meda P, Durand B, Reith W. 2007. Novel function of the ciliogenic transcription factor RFX3 in development of the endocrine pancreas. *Diabetes.* 56(4):950-959.
111. Reith W, Mach B. 2001. The bare lymphocyte syndrome and the regulation of MHC expression. *Annu Rev Immunol.* 19:331-373.
112. Blackshear PJ, Graves JP, Stumpo DJ, Cobos I, Rubenstein JL, Zeldin DC. 2003. Graded phenotypic response to partial and complete deficiency of a brain-specific transcript variant of the winged helix transcription factor RFX4. *Development.* 130:4539-4552.
113. Araki R, Takahashi H, Fukumura R, Sun F, Umeda N, Sujino M, Inouye ST,

- Saito T, Abe M. 2004. Restricted expression and photic induction of a novel mouse regulatory factor X4 transcript in the suprachiasmatic nucleus. *J Biol Chem.* 279:10237-10242.
114. Liu Y, Pathak N, Kramer-Zucker A, Drummond IA. 2007. Notch signaling controls the differentiation of transporting epithelia and multiciliated cells in the zebrafish pronephros. *Development.* 134:1111-1122.
115. Bonnafé E, Touka M, AitLounis A, Baas D, Barras E, Ucla C, Moreau A, Flamant F, Dubruille R, Couble P, *et al.* 2004. The transcription factor RFX3 directs nodal cilium development and left-right asymmetry specification. *Mol Cell Biol.* 24:4417-4427.
116. Baas D, Meiniel A, Benadiba C, Bonnafé E, Meiniel O, Reith W, Durand B. 2006. A deficiency in RFX3 causes hydrocephalus associated with abnormal differentiation of ependymal cells. *Eur J Neurosci.* 24:1020-1030.
117. Ait-Lounis A, Baas D, Barras E, Benadiba C, Charollais A, Nlend Nlend R, Liegeois D, Meda P, Durand B, Reith W. 2007. Novel function of the ciliogenic transcription factor RFX3 in development of the endocrine pancreas. *Diabetes.* 56:950-959.
118. Bresnick, E. H., Martowicz, M. L., Pal, S., and Johnson, K. D. 2005. Developmental control via GATA factor interplay at chromatin domains. *J. Cell Physiol.* 205, 1-9
119. Molkentin, J. D. 2000. The zinc finger-containing transcription factors GATA-4, -5, and -6. Ubiquitously expressed regulators of tissue-specific gene

- expression. *J. Biol. Chem.* 275, 38949-38952
120. Pevny, L., Simon, M. C., Robertson, E., Klein, W. H., Tsai, S. F., D'Agati, V., Orkin, S. H., and Costantini, F. 1991. Erythroid differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription factor GATA-1. *Nature* 349, 257-260
121. Tsai, F. Y., Keller, G., Kuo, F. C., Weiss, M., Chen, J., Rosenblatt, M., Alt, F. W., and Orkin, S.H. 1994. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature* 371, 221-226
122. Pandolfi, P. P., Roth, M. E., Karis, A., Leonard, M. W., Dzierzak, E., Grosveld, F. G., Engel, J.D., and Lindenbaum, M. H. 1995. Targeted disruption of the GATA3 gene causes severe abnormalities in the nervous system and in fetal liver haematopoiesis. *Nat. Genet.* 11, 40-44
123. Molkentin, J. D., Lin, Q., Duncan, S. A., and Olson, E. N. 1997. Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev.* 11, 1061-1072
124. Molkentin, J. D., Tymitz, K. M., Richardson, J. A., and Olson, E. N. 2000. Abnormalities of the genitourinary tract in female mice lacking GATA5. *Mol. Cell. Biol.* 20, 5256-5260
125. Morrisey, E. E., Tang, Z., Sigrist, K., Lu, M. M., Jiang, F., Ip, H. S., and Parmacek, M. S. 1998. GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.* 12,

126. Ng, Y. K., George, K. M., Engel, J. D., and Linzer, D. I. 1994. GATA factor activity is required for the trophoblast-specific transcriptional regulation of the mouse placental lactogen I gene. *Development* 120, 3257-3266
127. Yamamoto, M., Ko, L. J., Leonard, M. W., Beug, H., Orkin, S. H., and Engel, J. D. 1990. Activity and tissue-specific expression of the transcription factor NF-E1 multigene family. *Genes Dev.* 4, 1650-1662
128. Craven, S. E., Lim, K. C., Ye, W., Engel, J. D., de Sauvage, F., and Rosenthal, A. 2004. Gata2 specifies serotonergic neurons downstream of sonic hedgehog. *Development* 131, 1165-1173.
129. Minegishi, N., Ohta, J., Yamagiwa, H., Suzuki, N., Kawauchi, S., Zhou, Y., Takahashi, S., Hayashi, N., Engel, J. D., and Yamamoto, M. 1999. The mouse GATA-2 gene is expressed in the para-aortic splanchnopleura and aorta-gonads and mesonephros region. *Blood* 93, 4196-4207
130. Nardelli, J., Thiesson, D., Fujiwara, Y., Tsai, F. Y., and Orkin, S. H. 1999. Expression and genetic interaction of transcription factors GATA-2 and GATA-3 during development of the mouse central nervous system. *Dev. Biol.* 210, 305-321
131. Orlic, D., Anderson, S., Biesecker, L. G., Sorrentino, B. P., and Bodine, D. M. 1995. Pluripotent hematopoietic stem cells contain high levels of mRNA for c-kit, GATA-2, p45 NF-E2, and c-myb and low levels or no mRNA for c-fms and the receptors for granulocyte colony-stimulating factor and interleukins 5

- and 7. *Proc. Natl. Acad. Sci. U S A* 92, 4601-4605
132. Steger, D. J., Hecht, J. H., and Mellon, P. L. 1994. GATA-binding proteins regulate the human gonadotropin alpha-subunit gene in the placenta and pituitary gland. *Mol. Cell. Biol.* 14, 5592-5602
133. Bruno,M.D. et al. 2000. GATA-6 activates transcription of surfactant protein A. *J. Biol. Chem.* 275, 1043–1009.
134. Yang,H. et al. 2002. GATA6 regulates differentiation of distal lung epithelium. *Development.* 129, 2233–2246.
135. Liu,C. et al. 2003. Inhibition of alveolarization and altered pulmonary mechanics in mice expressing GATA-6. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 285. L1246–L1254.
136. Perl,A.K. et al. 2002. Conditional gene expression in the respiratory epithelium of the mouse. *Transgenic Res.* 11, 21–29.
137. Ramirez RD, Sheridan S, Girard L, et al. 2004. Immortalization of human bronchial epithelial cells in the absence of viral oncoproteins. *Cancer Res.* 64:9027–34.
138. Jedlicka P, Sui X, Sussel L, Gutierrez-Hartmann A. 2009. Ets Transcription Factors Control Epithelial Maturation and Transit and Crypt-Villus Morphogenesis in the Mammalian Intestine. *Am J Pathol.* 174.4
139. Oikawa T, Yamada T. 2003. Molecular biology of the Ets family of transcription factors. *Gene.* 303:11–34
140. Maroulakou IG, Bowe DB. 2000. Expression and function of Ets transcription

- factors in mammalian development: a regulatory network.
Oncogene.19:6432–6442
- 141.Chen C, Ouyang W, Grigura V, Zhou Q, Carnes K, Lim H, Zhao GQ, Arber S, Kurpios N, Murphy TL, Cheng AM, Hassell JA, Chandrashekhar V, Hofmann MC, Hess RA, Murphy KM. 2005. ERM is required for transcriptional control of the spermatogonial stem cell niche. *Nature*. 436:1030–1034
142. Donnison M, Beaton A, Davey HW, Broadhurst R, L’Huillier P, Pfeffer PL. 2005.Loss of the extraembryonic ectoderm in Elf5 mutants leads to defects in embryonic patterning. *Development*, 132:2299–2308
143. Georgiades P, Rossant J. 2006. Ets2 is necessary in trophoblast for normal embryonic anteroposterior axis development. *Development*, 133:1059–1068
144. Hippenmeyer S, Huber RM, Ladle DR, Murphy K, Arber S. 2007. ETS transcription factor Erm controls subsynaptic gene expression in skeletal muscles. *Neuron*, 55:726–740
145. Livet J, Sigrist M, Stroebel S, De Paola V, Price SR, Henderson CE, Jessell TM, Arber S. 2002. ETS gene Pea3 controls the central position and terminal arborization of specific motor neuron pools. *Neuron*, 35:877–892
146. O’Leary DA, Noakes PG, Lavidis NA, Kola I, Herzog PJ, Ristevski S. 2007.Targeting of the ETS factor GABPalpha disrupts neuromuscular junction synaptic function. *Mol Cell Biol*, 27:3470–3480
147. Ristevski S, O’Leary DA, Thornell AP, Owen MJ, Kola I, Herzog PJ. 2004. The

- ETS transcription factor GABPalpha is essential for early embryogenesis. *Mol Cell Biol*, 24:5844–5849
148. Rosmarin AG, Resendes KK, Yang Z, McMillan JN, Fleming SL. 2004. GA-binding protein transcription factor: a review of GABP as an integrator of intracellular signaling and protein-protein interactions. *Blood Cells Mol Dis*, 32:143–154
149. Zhou J, Chehab R, Tkalcevic J, Naylor MJ, Harris J, Wilson TJ, Tsao S, Tellis I, Zavarsk S, Xu D, Lapinskas EJ, Visvader J, Lindeman GJ, Thomas R, Ormandy CJ, Hertzog PJ, Kola I, Pritchard MA. 2005. Elf5 is essential for early embryogenesis and mammary gland development during pregnancy and lactation. *EMBO J*, 24:635–644
150. Nemer G, Nemer M. 2003. Transcriptional activation of BMP-4 and regulation of mammalian organogenesis by GATA-4 and -6. *Dev Biol*. 254(1):131-48
151. Neugebauer JM, Amack JD, Peterson AG, Bisgrove BW, Yost HJ. 2009. FGF signalling during embryo development regulates cilia length in diverse epithelia. *Nature*. 10.1038.
152. Levine M, Davidson EH. 2005. Gene regulatory networks for development. *Proc Natl Acad Sci USA* 102:4936–4942.
153. Oliveri P, Tu Q, Davidson EH. 2008. Global regulatory logic for specification of an embryonic lineage. *Proc Natl Acad Sci USA* 105:5955–5962.
154. Whitsett J.A. 2002. Intrinsic and innate defenses in the lung: intersection of

- pathways regulating lung morphogenesis, host defense, and repair. *J. Clin. Invest.* 109, 565–569.
155. Pandev P, Brors B, Srivastava PK, etc. 2008. Microarray-based approach identifies microRNAs and their target functional patterns in polycystic kidney disease. *BMC Genomics*. 9: 624.
156. Lopez RG, Carron C, Oury C, Gardellin P, Bernard O and Ghysdael J. 1999. TEL is a sequence-specific transcriptional repressor. *J Biol Chem* 274 (42): 30132-8
157. Bassuk AG and Leiden JM. 1995. A direct physical association between ETS and AP-1 transcription factors in normal human T cells. *Immunity* 3(2):223-37
158. Kwiatkowski BA, Bastian LS, Bauer Jr TR, Tsai S, Zielinska-Kwiatkowska AG and Hickstein DD. 1998. The ets family member Tel binds to the Fli-1 oncoprotein and inhibits its transcriptional activity. *J Biol Chem* 273 (28): 17525-30
159. Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, et al. 2003. Dicer is essential for mouse development. *Nat Genet* 35:215-7.
160. Harris KS, Zhang Z, McManus MT, Harfe BD, Sun X. 2006. Dicer function is essential for lung epithelium morphogenesis. *Proc Natl Acad Sci USA* 103:2208-13.
161. Harfe BD, McManus MT, Mansfield JH, Hornstein E, Tabin CJ. 2005. The RNaseIII enzyme Dicer is required for morphogenesis but not patterning of the vertebrate limb. *Proc Natl Acad Sci USA*;102:10898-903.

162. O'Rourke JR, Georges SA, Seay HR, Tapscott SJ, McManus MT, Goldhamer DJ, et al. 2007. Essential role for Dicer during skeletal muscle development. *Dev Biol*;311:359-68.
163. Muljo SA, Ansel KM, Kanellopoulou C, Livingston DM, Rao A, Rajewsky K. 2005. Aberrant T cell differentiation in the absence of Dicer. *J Exp Med*;202:261-9.
164. Andl T, Murchison EP, Liu F, Zhang Y, Yunta-Gonzalez M, Tobias JW, et al. 2006. The miRNA-processing enzyme dicer is essential for the morphogenesis and maintenance of hair follicles. *Curr Biol*;16:1041-9.
165. Yi R, O'Carroll D, Pasolli HA, Zhang Z, Dietrich FS, Tarakhovsky A, et al. 2006. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat Genet*;38: 356-62.