

University of New Mexico UNM Digital Repository

Research Supported by El Centro

El Centro de la Raza

2015

Disparities in cervical cancer mortality rates as determined by the longitudinal hyperbolastic mixed-effects type II model.

Juliette Tayou Guemmegne

Mohammad A. Tabatabai

Jean-Jacques Kengwoung-Keumo

Wayne M. Eby

Sejong Bae

See next page for additional authors

Follow this and additional works at: https://digitalrepository.unm.edu/el_centro_research

Recommended Citation

Tayou Guemmegne, Juliette; Mohammad A. Tabatabai; Jean-Jacques Kengwoung-Keumo; Wayne M. Eby; Sejong Bae; Uppender Manne; Mona Fouad; Edward E. Partridge; and Karan P. Singh. "Disparities in cervical cancer mortality rates as determined by the longitudinal hyperbolastic mixed-effects type II model.." (2015). https://digitalrepository.unm.edu/el_centro_research/11

This Article is brought to you for free and open access by the El Centro de la Raza at UNM Digital Repository. It has been accepted for inclusion in Research Supported by El Centro by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Authors

Juliette Tayou Guemmegne, Mohammad A. Tabatabai, Jean-Jacques Kengwoung-Keumo, Wayne M. Eby, Sejong Bae, Uppender Manne, Mona Fouad, Edward E. Partridge, and Karan P. Singh

Conferences:

- **First Annual de la Raza Conference., March 20th 2015**
- **The New Mexico Public Health Association, March 31-April 1st**

Title: Disparities in cervical cancer mortality rates as determined by the longitudinal hyperbolastic mixed-effects type II model.

We analyze the dynamics of cervical cancer mortality rates for African American and White women residing in 13 states located in the eastern half of the United States of America from 1975 through 2010. Despite decreasing trends in cervical cancer mortality rates for both races, racial disparities in mortality rates still exist. In all 13 states, Black women had higher mortality rates at all times. The degree of disparities and pace of decline in mortality rates over time differed among these states. In all 13 states, cervical cancer mortality rates for both racial groups have fallen. Disparities in the pace of decline in mortality rates in these states may be due to differences in the rates of screening for cervical cancers. Of note, the gap in cervical cancer mortality rates between Black women and White women is narrowing.

Paper's authors:

Tabatabai, Mohammad A.; Kengwoung-Keumo, Jean-Jacques; Eby, Wayne M.; Bae, Sejong; Guemmegne, Juliette T.; Manne, Upender; Fouad, Mona; Partridge, Edward E.; Singh, Karan P